

23	HO ₂ CCHCH ₂ S	SO ₂ Cl ₂	OH	F	32	184.5-186 dec	M	C ₈ H ₁₆ N ₂ O ₈ S ₃ ^d	-17.8 (2)
24	NH ₂ COCHCH ₂ S NH ₂ SO ₂ CH ₃	CO(CH ₂) ₂ CO ₂ H	NH ₂	(5)	57	186.5-187.5 dec	D	C ₁₄ H ₂₂ N ₄ O ₈ S ₂	-122.3 (1)
25	NH ₂ COCHCH ₂ S NHCO(CH ₂) ₂ CO ₂ H	CO(CH ₂) ₂ Cl	NH ₂	(6)	72	185.5-186.5 dec	N	C ₁₂ H ₂₀ Cl ₂ N ₄ O ₈ S ₂	-116.7 (3)
26	NH ₂ COCHCH ₂ S NHCO(CH ₂) ₂ Cl NHCONH ₂	CONH ₂	NH ₂	II	70	211.5 dec	O	C ₈ H ₁₆ N ₄ O ₈ S ₂	-54.6 (3)

^a (1) Piperazinium. ^b See Experimental Section for the letters; (1) the optical isomerism and synthetic method are described in the text; (2) except for the use of 1 equiv of NEt₃, the procedure was similar to that given in ref 3b, method P; (3) see ref 3b, method Q; (4) prepared from L-cystine dimethyl ester dihydrochloride and carbobenzoxyglycine in the presence of NEt₃ and N,N'-dicyclohexylcarbodiimide; (5) prepared from L-3,3'-dithiobis(2-amino propionamide) dihydrochloride and succinic anhydride in the presence of NaHCO₃ and aqueous THF (ref 1, 2); (6) similar to variation 5, except for the use of 3-chloropropionyl chloride and H₂O as the reaction medium. ^c A = 30% aqueous EtOH, B = EtOAc, C = 50% aqueous MeOH, D = H₂O, E = EtOH, F = DMF-H₂O, G = MeOH, H = EtOAc-MeOH, I = EtOAc-Skellysolve B, J = 2-PrOH-EtOAc-Skellysolve B, K = DMF-MeOH, L = 50% aqueous EtOH, M = 2-PrOH-H₂O, N = DMSO-MeOH, O = 5% aqueous MeOH. ^d N and S analyses. ^e c 1, in all cases; (1) 1 N NaOH, (2) H₂O, (3) DMSO, (4) DMF, (5) MeOH, (6) EtOH. ^f All compounds except 4, 14, and 23 analyzed correctly for C, H, N.

TABLE II
COMPARISON OF THE RATE AND EXTENT OF REDUCTION
OF VISCOSITY OF MUCOPROTEIN SOLUTION^a

Compd	% decrease in viscosity		
	3 min	30 min	60 min
L-N-Sulfamylcysteine (5)	20	27	30
L-3-Mercapto-2-ureidopropionamide (6)	22	26	27
L-3-Mercapto-2-methanesulfonamido- propionamide (7)	24	30	30
2-Acetamido-N-(L-1-carboxy-2-mercapto- ethyl)-3-mercapto-DL-propionamide (8)	18	28	30
2-Acetamido-N-(L-1-carbamoyl-2-mercapto- ethyl)-3-mercapto-DL-propionamide (9) ^b	23	29	30
N-Acetyl-L-cysteine ^c	11	20	25

^a See ref 3b, Table II. ^b Saturated solution of 0.036 M, instead of the usual 0.05 M, was used. ^c Included as reference material.

L. L-3-(Diphenylmethylthio)-2-(methanesulfonamido)propionic Acid (17).—A mixture of 6 g (0.0073 mole) of 16 was slurried with 150 ml of 33% aqueous MeOH while acidifying with 1 N HCl. The compound was isolated by extracting with EtOAc, concentrating, and recrystallizing from EtOAc-Skellysolve B; yield 3.7 g (69%).

M. Compound 17 may be isolated directly from the EtOAc extract of procedure K in an over-all improved yield of 68% by adding Skellysolve B and seeding.

N. L-3-(Diphenylmethylthio)-2-(methanesulfonamido)propionamide (18).—A solution of 15.2 g (0.04 mole) of 15 and 175 ml of MeOH saturated at 15° with NH₃ was allowed to stand for 2 days. The solid was collected; yield 9.3 g (64%) in three crops.

Acknowledgment.—The author is grateful to Dr. A. L. Sheffner and Mr. L. W. Jacobs for the mucolytic data, to Mr. J. P. Catlett and Mr. W. F. Kavanaugh for technical assistance, to Dr. J. R. Corrigan and Dr. W. T. Comer for their interest and advice in this work, and to Mr. C. M. Combs, Mr. C. I. Kennedy, and Mr. J. G. Schmidt for analytical and spectral data.

Amides of N-Acylcysteines as Potential Amino Acid Antagonists in Bacteria

WALTER A. ZYGMUNT AND TELLIS A. MARTIN

Meal Johnson Research Center, Evansville, Indiana 47721

Received May 9, 1969

Recently, we reported¹ on the activity of 21 cysteine or cystine analogs as potential amino acid antagonists in bacteria. Of these, N-acetyl-L-cysteine, N-propionyl-L-cysteine, L-cysteine hydantoin, and L-cystine hydantoin were the most effective inhibitors of L-cysteine utilization.

As an extension of these studies, 38 additional analogs were tested as inhibitors of cysteine or cystine utilization by *Leuconostoc mesenteroides*, a cysteine-cystine-dependent bacterium, and by *Escherichia coli*, an organism able to synthesize all its amino acid requirements. Most of these analogs² were amides of cysteines or cystines.

(1) W. A. Zygmunt and T. A. Martin, *J. Med. Chem.*, **11**, 623 (1968).

(2) (a) T. A. Martin, D. H. Causey, A. L. Sheffner, A. G. Wheeler, and J. R. Corrigan, *ibid.*, **10**, 1172 (1967); (b) T. A. Martin and A. L. Sheffner, U. S. Patent 3,340,147 (1967).

TABLE I
 COMPARATIVE GROWTH INHIBITION IN *L. mesenteroides* BY VARIOUS CYSTEINE AND CYSTINE ANALOGS

Test compd ^a	Concn, $\mu\text{g/ml}$					
	25	50	100	200	400	800
N-Acetyl-L-cysteine ³	5	9	30	68	80	84
L-2-Acetamido-3-mercapto- propionamide ²	7	66	92	94	100	100
L-3-Mercapto-2-ureidopropionamide	55	86	90	100	100	100
L-3-Mercapto-2-methanesulfonamido- propionamide	14	27	72	73	79	83
L-2-Acetamido-3-mercapto-N- phenylpropionamide ²	6	29	45	88	88	88
2-Acetamido-N-(L-1-carboxy-2- mercaptoethyl)-3-mercapto-DL- propionamide	84	100	100	100	100	100
2-Acetamido-N-(L-1-carbamoyl-2- mercaptoethyl)-3-mercapto-DL- propionamide	74	100	100	100	100	100
L-2-Amino-3-benzylthiopropionamide hydrochloride ²	7	10	20	54	65	53
L-2-Amino-3-(diphenylmethylthio)- propionamide ²	5	12	16	18	48	69

^a Several of the compounds listed are described elsewhere: T. A. Martin, *J. Med. Chem.*, **12**, 950 (1969). ² Other analogs tested which required concentrations $>800 \mu\text{g/ml}$ for 50% growth inhibition were: L-2-amino-3-mercaptopropionamide hydrochloride, L-2-propionamido-3-mercaptopropionamide, L-2-acetamido-3-mercapto-N-methylpropionamide, L-3-(benzylthio)-2-formamidopropionamide, L-2-acetamido-3-benzylthiopropionamide, L-2-acetamido-3-benzylthio-N-methylpropionamide, DL-2-acetamido-3-(benzylthio)-N-(2-hydroxyethyl)propionamide, L-2-benzamido-3-(benzylthio)propionamide, 2-acetamido-3-(benzylthio)-N-[L-2-(benzylthio)-1-methoxycarbonyl-ethyl]-DL-propionamide, 2-acetamido-3-(benzylthio)-N-[L-2-(benzylthio)-1-carboxyethyl]-DL-propionamide, L-3-(diphenylmethylthio)-2-formamido-N-phenylpropionamide, L-3-(diphenylmethylthio)-2-(methanesulfonamido)propionic acid, L-S-diphenylmethyl-N-methanesulfonylcysteine methyl ester, L-3-(diphenylmethylthio)-2-methanesulfonamidopropionamide, L-2-acetamido-3-acetylthiopropionamide, N,N'-diacetyl-L-cystine piperazinium salt (R. Marshall, M. Winitz, S. M. Birnbaum, and J. P. Greenstein, *J. Am. Chem. Soc.*, **79**, 4538 (1957)), L-N,N'-diacetylcystine dimethyl ester (H. Heymann, T. Ginsberg, Z. R. Gulick, E. A. Konopka, and R. L. Mayer, *ibid.*, **81**, 5125 (1959)), N,N'-di(isovaleryl)-L-cystine, N,N'-di(isovaleryl)-L-cystine dimethyl ester, N,N'-dibenzoyl-L-cystine, L-N,N'-di(N-carbobenzoxymethyl)cystine dimethyl ester, L-3,3'-dithiobis(2-aminopropionamide) dihydrochloride, L-3,3'-dithiobis(2-acetamidopropionamide), L-3,3'-dithiobis(2-propionamidopropionamide), L-3,3'-dithiobis(2-acetamido-N-methylpropionamide), L-3,3'-dithiobis[2-(3-chloropropionamido)propionamide], L-3,3'-dithiobis(2-benzamidopropionamide), L-3,3'-dithiobis[2-(3-(carboxypropionamido))propionamide], L-3,3'-dithiobis(2-ureidopropionamide).

The data in Table I list N-acetyl-L-cysteine³ and eight of the 38 compounds tested which at a final concentration of $800 \mu\text{g/ml}$ inhibited growth of *L. mesenteroides* by at least 50%. Interestingly, the amide of N-acetyl-L-cysteine (L-2-acetamido-3-mercaptopropionamide) is about a fivefold more effective inhibitor of *L. mesenteroides* growth than is the carboxyl analog. The three additional most effective growth inhibitors were 2-acetamido-N-(L-1-carboxy-2-mercaptoethyl)-3-mercapto-DL-propionamide, 2-acetamido-N-(L-1-carbamoyl-2-mercaptoethyl)-3-mercapto-DL-propionamide, and L-3-mercapto-2-ureidopropionamide.

The nine most active compounds (Table II) were tested at equimolar concentrations in the presence of several levels of L-cysteine hydrochloride. With the exception of L-2-amino-3-(diphenylmethylthio)propionamide and L-3-mercapto-2-methanesulfonamidopropionamide, the growth inhibition observed with the remaining analogs decreased markedly with increasing levels of cysteine.

The growth inhibition found with L-3-mercapto-2-methanesulfonamidopropionamide was reversible with added L-cysteine but only at lower levels of inhibitor. Inhibition of *L. mesenteroides* growth by L-2-amino-3-(diphenylmethylthio)propionamide appears to be non-specific and not easily reversed by cysteine even at lower levels of inhibitor.

L-2-Amino-3-mercaptopropionamide hydrochloride

 TABLE II
 EFFECT OF VARYING CYSTEINE CONCENTRATIONS
 ON GROWTH INHIBITION OF *L. Mesenteroides*
 BY AMINO ACID ANALOGS

L-Cysteine HCl, $\mu\text{g/ml}$	Test compd, 0.003 M ^a								
	A	B	C	D	E	F	G	H	I
1	81	93	98	75	88	97	96	58	69
10	34	55	82	90	41	53	77	60	67
100	1	2	25	88	32	22	20	48	60
1000	3	2	3	82	37	1	1	9	61

^a A, N-acetyl-L-cysteine; B, L-2-acetamido-3-mercaptopropionamide; C, 3-mercapto-2-ureidopropionamide; D, L-3-mercapto-2-methanesulfonamidopropionamide; E, L-2-acetamido-3-mercapto-N-phenylpropionamide; F, 2-acetamido-N-(L-1-carboxy-2-mercaptoethyl)-3-mercapto-DL-propionamide; G, 2-acetamido-N-(L-1-carbamoyl-2-mercaptoethyl)-3-mercapto-DL-propionamide; H, L-2-amino-3-benzylthiopropionamide; I, L-2-amino-3-(diphenylmethylthio)propionamide.

and L-3,3'-dithiobis(2-aminopropionamide) dihydrochloride were inactive as amino acid antagonists or as substitutes for L-cysteine or L-cystine in the growth of *L. mesenteroides*. Both compounds *per se* had only 5–10% microbiological growth replacement activity (based on an equivalent weight basis) in the absence of added cysteine or cystine. With the former compound this activity increased to 40% in the presence of $1.5 \mu\text{g/ml}$ of L-cysteine hydrochloride (a level required for about half-maximal growth). No such stimulation of the latter compound utilization, however, was noted with the further addition of L-cysteine.

L-2-Amino-3-mercaptopropionamide hydrochloride effectively inhibited growth of *E. coli* in a chemically defined medium and caused complete inhibition at 100

(3) (a) T. A. Martin, J. R. Corrigan, and C. W. Waller, *J. Org. Chem.*, **30**, 2839 (1965); (b) T. A. Martin and C. W. Waller, U. S. Patent, 3,184,505 (1965).

$\mu\text{g/ml}$. Of the remaining compounds tested in *E. coli* moderate growth inhibition (80–90%) was found with L-3,3'-dithiobis(2-aminopropionamide) dihydrochloride (300 $\mu\text{g/ml}$) and L-2-amino-3-(diphenylmethylthio)propionamide (600 $\mu\text{g/ml}$).

None of the 38 compounds tested showed any significant cysteine–cystine replacement activity for growth of *L. mesenteroides*. In summary, none of the 38 compounds tested showed significant growth inhibition of *L. mesenteroides* and *E. coli*. In most instances, this inhibition was readily reversed by the addition of cysteine.

The microbiological assay and testing procedures were similar to those previously described.¹

Antiamoebic, Antimalarial, and Anthelmintic Effects of Distal Hydrazine Analogs of Azacrine, Quinacrine, and 7-[3-(Octylamino)propyl]amino}benz[c]acridine^{1,2}





EDWARD F. ELSLAGER AND DONALD F. WORTH

Department of Chemistry, Medical and Scientific Affairs Division, Parke, Davis & Company, Ann Arbor, Michigan 48106

Received April 24, 1969


An array of basically substituted 9-aminoacridines,^{3–11} 7-aminobenz[c]acridines,^{3,4,9,10,12–14} and aminobenzonaphththyridines^{4,9,15–18} exhibit noteworthy antiprotozoal, anthelmintic, antibacterial, and antitumor properties. Among them, quinacrine (I),^{3–6} 3-chloro-9-[(4-(diethylamino)-1-methylbutyl)amino]acridine 10-oxide dihydrochloride (II),⁷ 7-[3-(octylamino)propyl]amino}benz[c]acridine dihydromintic activ-

chloride (III),¹³ and azacrine (IV)¹⁵ have been demonstrated to have appreciable antiprotozoal and anthel-



ity in man. It was therefore of interest to synthesize representative {[3-(2,2-dialkylhydrazino)alkyl]amino}acridines, benz[c]acridines, and benzo[b][1,5]naphththyridines to enable a determination of the effects of a distal hydrazine moiety on antiprotozoal and anthelmintic activity.

The condensation of 6,9-dichloro-2-methoxyacridine with 2-(3-aminopropyl)-1,1-dimethylhydrazine,¹⁹ 1-[(3-aminopropyl)amino]piperidine,¹⁹ and 1-[(3-aminopropyl)amino]-4-methylpiperazine¹⁹ in phenol afforded 6-chloro-9-[[3-(2,2-dimethylhydrazino)propyl]amino]-2-methoxyacridine dihydrochloride (Va) (55%), 6-chloro-2-methoxy-9-[[3-(piperidinoamino)propyl]amino]acridine dihydrochloride (Vb) (38%), and 6-chloro-



2-methoxy-9-[(3-[(4-methyl-1-piperazinyl)amino]propyl)amino]acridine trihydrochloride (Vc) (40%), respectively (Table I, procedures I and II). 7-[3-(Piperidinoamino)propyl]amino}benz[c]acridine dihydrochloride (VIa) and 7-[3-[(4-methyl-1-piperazinyl)amino]propyl]amino}benz[c]acridine hydrochloride (VIb) (68%) were obtained in a similar

(19) E. F. Elslager, E. A. Weinstein, and D. F. Worth, *ibid.*, **7**, 493 (1964).

(1) This is paper X of a series on synthetic amoebicides and paper XVII of a series relating to antimalarial substances. For the previous paper, see L. M. Werbel, E. F. Elslager, A. A. Phillips, D. F. Worth, P. J. Islip, and M. C. Neville, *J. Med. Chem.*, **12**, 521 (1969).

(2) This is communication III of a series on anthelmintic drugs. For paper II, see D. B. Capps, O. D. Bird, E. F. Elslager, Z. B. Gavrilis, J. A. Roush, P. E. Thompson, and J. W. Vaitkus, *J. Heterocyclic Chem.*, **5**, 355 (1968).

(3) A. Albert, "The Acridines," 2nd ed, Edward Arnold, London, 1966.

(4) For a review, see E. F. Elslager in "Medicinal Chemistry," A. Burger, Ed., 3rd ed, Interscience Division of John Wiley and Sons, Inc., New York, N. Y., 1969.

(5) For a review, see P. B. Russell in "Medicinal Chemistry," A. Burger, Ed., 2nd ed, Interscience Publishers, Inc., New York, N. Y., 1960, pp 814–850.

(6) For a review, see O. D. Standen in "Experimental Chemotherapy," Vol. I, R. J. Schnitzer and F. Hawking, Eds., Academic Press, New York, N. Y., 1963, pp 701–892.

(7) E. F. Elslager, R. E. Bowman, F. H. Tendick, D. J. Tivey, and D. F. Worth, *J. Med. Pharm. Chem.*, **5**, 1159 (1962).

(8) E. F. Elslager and F. H. Tendick, *ibid.*, **5**, 1153 (1962).

(9) N. B. Ackerman, D. K. Haldorsen, F. H. Tendick, and E. F. Elslager, *ibid.*, **11**, 315 (1968).

(10) E. F. Elslager, F. W. Short, and F. H. Tendick, *J. Heterocyclic Chem.*, **5**, 599 (1968).

(11) E. F. Elslager, F. H. Tendick, and L. M. Werbel, *J. Med. Chem.*, **12**, 600 (1969).

(12) E. F. Elslager, A. M. Moore, F. W. Short, M. J. Sullivan, and F. H. Tendick, *J. Am. Chem. Soc.*, **79**, 4699 (1957).

(13) F. W. Short, E. F. Elslager, A. M. Moore, M. J. Sullivan, and F. H. Tendick, *ibid.*, **80**, 223 (1958).

(14) E. F. Elslager, F. W. Short, M. J. Sullivan, and F. H. Tendick, *ibid.*, **80**, 451 (1958).

(15) D. M. Besly and A. A. Goldberg, *J. Chem. Soc.*, 2448 (1958).

(16) E. F. Elslager and F. H. Tendick, *J. Med. Pharm. Chem.*, **5**, 546 (1962).

(17) P.-L. Chien and C. C. Cheng, *ibid.*, **11**, 164 (1968).

(18) E. F. Elslager, F. H. Tendick, and S. C. Perricone, 1969, unpublished data.