

1-[2-[(5-Cyanopyridin-2-yl)amino]-ethylamino]acetyl-2-(*S*-pyrrolidine-carbonitrile): A Potent, Selective, and Orally Bioavailable Dipeptidyl Peptidase IV Inhibitor with Antihyperglycemic Properties

Edwin B. Villhauer,^{*,†} John A. Brinkman,[†]
Goli B. Naderi,[†] Beth E. Dunning,[‡]
Bonnie L. Mangold,[§] Manisha D. Mone,[‡]
Mary E. Russell,[‡] Stephen C. Weldon,[‡] and
Thomas E. Hughes[‡]


Novartis Institute for Biomedical Research,
556 Morris Avenue, Summit, New Jersey 07901

Received March 26, 2002

Abstract: Dipeptidyl peptidase IV (DPP-IV) inhibition has the potential to become a valuable therapy for type 2 diabetes. We report the first use of solid-phase synthesis in the discovery of a new DPP-IV inhibitor class and a solution-phase synthesis that is practical up to the multikilogram scale. One compound, NVP-DPP728 (**2**), is profiled as a potent, selective, and short-acting DPP-IV inhibitor that has excellent oral bioavailability and potent antihyperglycemic activity.

Introduction. Dipeptidyl peptidase IV (DPP-IV, EC 3.4.14.5) is a ubiquitous yet highly specific serine protease that cleaves N-terminal dipeptides from polypeptides with L-proline or L-alanine at the penultimate position.¹ The biological activities of many circulating regulatory peptides are altered or abolished by the action of DPP-IV in vitro.² However, in part because of the multiplicity of enzymes exhibiting DPP-IV-like activity,³ the in vivo role of DPP-IV in mediating the cleavage and determining the action of most substrates has yet to be established. One exception is with the incretin known as glucagon-like peptide-1 (GLP-1), the most potent insulinotropic hormone known.⁴ Numerous studies with DPP-IV⁵ and DPP-IV inhibitors^{6–9} support a principal role of DPP-IV in the inactivation of GLP-1 in vivo. More importantly, the contribution of DPP-IV catalytic activity to blood glucose control through GLP-1 inactivation has recently been confirmed.¹⁰ Because of multiple benefits of GLP-1 augmentation, DPP-IV inhibition has been recognized as a mechanistic approach of potential value in the treatment of type 2 diabetes.¹¹ By extending the duration of action of GLP-1, one would stimulate insulin secretion, inhibit glucagon release,¹² and slow gastric emptying;¹³ each a benefit in the control of glucose homeostasis. DPP-IV inhibition, through the preservation of active GLP-1 levels, has the potential to slow or even prevent the progression of type 2 diabetes by stimulating insulin gene expression and biosynthesis, increasing the expression of the β -cell's

Chart 1



glucose-sensing mechanism and promoting genes involved in the differentiation (neogenesis) of β -cells.¹⁴ GLP-1 may play a role in acutely suppressing appetite in humans¹⁵ and may play a role in mediating peripheral glucose uptake.¹⁶ Since the blood glucose lowering effects of GLP-1 are dependent on elevated blood glucose and abate as glucose levels return to normal, the incidence of hypoglycemia during treatment with a DPP-IV inhibitor is expected to be very low.¹⁷

With few exceptions,^{18–20} DPP-IV inhibitors resemble the P2–P1 dipeptidyl substrate cleavage product, where the P-1 site contains a proline mimic.²¹ A straightforward replacement of the normally cleaved P-1 substrate amide (R in Chart 1) with an electrophile provides both irreversible (R = P(O)(OPh)₂, CO–NH–O–COR') and reversible (R = B(OH)₂, H, CN) inhibitors.²² Low nanomolar inhibition and chemical stability adequate for oral administration are obtained only with nitrile replacement of a substrate P-1 site amide (X_{aa}-(2*S*)-cyanopyrrolidines^{23–25} and X_{aa}-(4*R*)-cyanothiazolidines²⁶). Cyclohexylglycine-(2*S*)-cyanopyrrolidine **1** is one of the more potent, selective, and stable representatives of this nitrile class (*K_i* of 1.4 nM, >1000-fold selectivity over closely related peptidases, and *t*_{1/2} stability of >48 h at pH 7.4).²⁴

Until recently, a constant in DPP-IV inhibitor design had been an L-amino acid with a protonatable N-terminal primary amine in the P-2 site. Noticing that *N*-methylglycine was recognized in the substrate P-2 site,^{21a} we were curious to investigate whether structurally more complicated N-substituted glycines would be tolerated at the P-2 site. We were gratified to find that a number of diverse P-2 site N-substituted glycines provided potent inhibition when combined with a (2*S*)-cyanopyrrolidide in the P-1 site (**8** in Scheme 1).²⁷ Intensive evaluation of this class²⁸ has led to the selection of the slow binding inhibitor **2** (NVP-DPP728)²⁹ as a clinical development candidate for type 2 diabetes. Herein, a tandem resin–solution parallel synthesis that led to the discovery of the P-2 site of **2** is described along with a solution-based method for multigram synthesis. Additionally, we report on the pharmacologic profile of this selective DPP-IV inhibitor, which exhibits excellent potency and oral bioavailability.


Chemistry. The preparation of a library of N-substituted 2-(*S*)-pyrrolidinecarbonitriles **8** has been carried out in a tandem five-step solid-phase and three step solution-phase sequence starting from commercially available Fmoc-protected Rink amide AM resin **4** as

* To whom correspondence should be addressed. Phone: 908-277-7208. Fax: 908-277-2405. E-mail: edwin.villhauer@pharma.novartis.com.


[†] Chemistry Department, Metabolic and Cardiovascular Diseases Research.

[‡] Metabolic Diseases Pharmacology Department, Metabolic and Cardiovascular Diseases Research.

[§] Drug Metabolism & Pharmacokinetics Department.

Scheme 1^a

^a Reagents: (i) 20% piperidine/DMF; (ii) Fmoc-proline, DIC, DMF; (iii) 20% piperidine/DMF; (iv) BrCH₂COOH, DIC, DMF; (v) RNH₂, DMSO; (vi) 95% TFA/H₂O; (vii) TFAA, THF; (viii) NH₃/MeOH.

Scheme 2^a

^a Reagents: (i) BrCH₂COBr, Et₃N, CH₂Cl₂, and DMAP for **10a** while ClCH₂COCl, K₂CO₃, and THF for **10b**; (ii) **10a** for **11a** and **10b** for **11b**, TFAA, CH₂Cl₂; (iii) **11a**, THF, **12a** for **3** and **11b**, THF, **12b** for **2**; (iv) excess HCl/THF for di-HCl of **2** and di-HCl if **3** and 1 equiv of ethanolic HCl for mono-HCl of **2**.

described in Scheme 1.³⁰ Successive deprotection of **4** with piperidine, 1,3-diisopropylcarbodiimide (DIC) coupling with Fmoc-protected proline, deprotection with piperidine, and finally DIC coupling with bromoacetic acid provided the resin-bound α -bromoacetyl prolinamide **5**. Analogous to Zuckermann's synthesis for solid-phase peptoid libraries,³¹ **5** was treated with a diverse array of over 200 primary aliphatic amines to provide a library of discrete, resin-bound N-substituted glycine-2-(S)-pyrrolidinecarboxamides (**6**). Trifluoroacetic acid (TFA) resin cleavage of **6** followed by amide dehydration with trifluoroacetic anhydride (TFAA) afforded N-substituted N-trifluoroacetylated-2-(S)-pyrrolidinecarboxamides **7**. Deacetylation of **7** with ammonia in methanol provided the product library of N-substituted 2-(S)-pyrrolidinecarboxamides (**8**) in a 1 to 1 mixture with trifluoroacetamide.³² The commercially available 2-(2-aminoethylamino)-5-nitropyridine (**12a**) provided resin-derived **3** (IC₅₀ of 20 \pm 3 nM in the DPP-IV Caco-2 assay) as one of the few low nanomolar DPP-IV inhibitors from this library effort. Compound **3** served as our starting point for the structure-activity relationship (SAR) effort that led to the title compound **2**.²⁸ A solution-based preparation of **2** and **3** had been carried out in three steps beginning with L-prolinamide (**9**) as shown in Scheme 2. Coupling of **9** with either bromo-

Table 1. DPP-IV Inhibition and Selectivity Assays^a

	Caco-2 ^b	rat plasma ^b	human plasma ^b	PPCE ^c	DPP-II ^d
1	2.0 \pm 0.3	2.8 \pm 0.2	3.2 \pm 0.19	41 000 \pm 14 000	102 000 \pm 20 000
2	22.0 \pm 2.0	6.0 \pm 1.0	7.0 \pm 1.7	190 000 \pm 46 000	110 000 \pm 5800
3	8.0 \pm 3.0	17 \pm 0.3	8.7 \pm 0.8	16 000 \pm 1200	12 000 \pm 580

^a Values are IC₅₀ (nM) expressed as the mean \pm SD of three independent determinations. Procedures are described in Supporting Information. ^b Primary DPP-IV assays. ^c Extract from human erythrocytes. ^d Extract from bovine kidney homogenate.

acetyl bromide or chloroacetyl chloride provided **10a** or **10b**, respectively.

Amide dehydration of **10a** and **10b** with trifluoroacetic anhydride produced **11a** and **11b** as solids that were stable for months at room temperature. Coupling of bromide **11a** with an excess of **12a** provided **3**, which was isolated as the dihydrochloride salt. The coupling of commercially available 5-cyano-2-chloropyridine with excess ethylenediamine provided **12b**. Reaction of **11b** with excess **12b** provided **2**, which was isolated as either the mono- or the dihydrochloride salt. The monohydrochloride **2** possessed a solubility of >100 mg/mL in distilled water and crystallized as a hemihydrate trans-amide rotomer with (S) chirality, as evidenced by the X-ray crystallographic analysis.³³ In solution, **2** was a mixture of cis- and trans-amide rotomers according to NMR. With minor modifications, the present solution synthesis has provided **2** on the 100 kg scale.

Results and Discussion. Compounds **1–3** were evaluated in vitro for their inhibition of DPP-IV extracted from Caco2 cells as well as from rat and human plasma (Table 1). Since under neutral and basic aqueous conditions the P-2 site amine can nucleophilically attack the carbon of the pyrrolidine-nitrile to form an inactive cyclic amidine,²⁸ the stability of **2** was examined under assay conditions. Under the assay conditions employed, this intramolecular cyclization was slow ($t_{1/2}$ > 2 days), resulting in less than 1% of **2** converting during the time frame of the experiment. As shown in Table 1, compound **2** potentially inhibited both human and rat plasma DPP-IV and human epithelial cell-surface DPP-IV (IC₅₀ = 7, 6, and 22 nM, respectively). Also, **2** was highly selective for DPP-IV over closely related peptidases, post-proline-cleaving enzyme (PPCE) and DPP-II³⁴ (Table 1). In addition, the in vitro specificity of **2** was profiled in over 100 receptor and enzyme assays and no significant binding was observed (10 μ M).

In vivo evaluation of **2** in rat⁹ and human³⁵ has supported the connection between DPP-IV inhibition and an improvement in oral glucose tolerance through an increase in active GLP-1 levels. We also found that **2** rapidly and effectively improved the metabolic profile in nonhuman primates. Oral administration of **2** (1 μ mol/kg) significantly reduced plasma glucose levels (38% reduction in the 0–90 min glucose AUC, p < 0.05) in cynomolgus monkey compared to the control during an oral glucose tolerance test (OGTT) (Figure 1). Additionally, peak glucose levels are significantly reduced in treated animals compared to control (98 \pm 4 vs 88 \pm 3 mg/dL, p < 0.05). When administered 30 min before an OGTT study, **2** maximally inhibited plasma DPP-IV activity (89%) 25 min postdose and provided a \geq 70% DPP-IV inhibition throughout the study.

Pharmacokinetic evaluation of **2** was performed in male Sprague-Dawley rats and male cynomolgus mon-




Figure 1. Incremental area under the glucose curve (from 0 to 90 min) during oral glucose tolerance tests (OGTTs) performed in seven anesthetized cynomolgus monkeys following oral administration of vehicle or NVP-DPP728 (**2**) (1 $\mu\text{mol/kg}$, po), mean \pm SEM, (*) $p < 0.05$. Vehicle is 0.5% carboxymethylcellulose in 0.2% Tween 80. Experimental procedure is detailed in Supporting Information.

keys. After an oral dose of 10 $\mu\text{mol/kg}$, C_{max} was 3.65 μM in rat and 7.56 μM in monkey. Absolute bioavailability was high in both rat and monkey at $\geq 74\%$. The steady-state volumes of distribution are similar in rat (670 mL/kg) and monkey (841 mL/kg), suggesting that **2** is distributed principally in the body fluids. Clearance of **2** from plasma is moderate at about $28 \pm 1.5 \text{ mL min}^{-1} \text{ kg}^{-1}$ in the rat and $21.9 \pm 3.1 \text{ mL min}^{-1} \text{ kg}^{-1}$ in the monkey. After an oral dose in monkey of 1 $\mu\text{mol/kg}$, **2** provided a half-life of 0.85 h and inhibited plasma DPP-IV activity by $> 50\%$ for 4 h. A 100 mg oral dose of **2** in humans provided a similar half-life of 0.85 h, a $> 80\%$ inhibition of plasma DPP-IV activity for ~ 4 h, a significant increase in active GLP-1 levels, and an improvement in metabolic control.³⁵ As a reversible DPP-IV inhibitor possessing a relatively short half-life, **2** might most effectively be taken with a meal when GLP-1 secretion is at its maximal rate.

In summary, we report here the use of combined solid-phase and solution-phase chemistry to discover a potent, selective, and short-duration DPP-IV inhibitor that also has excellent oral bioavailability. The favorable pharmacokinetic profile for NVP-DPP728 (**2**) led to the selection of this compound for further study in a clinical setting for type 2 diabetes.³⁵ Profiling of this new class of DPP-IV inhibitors is under study and will be reported in due course.

Acknowledgment. The authors thank Dr. Phillip E. Fanwick from Purdue University for conducting the X-ray crystallographic analysis of the monohydrochloride of **2**. The authors thank Prasad Kapa and Eric M. Loeser for crystalline monohydrochloride of **2**, B rk Balkan, Elina Dunn, Elizabeth D. Graham, Brenda H. Hamilton, Lori A. Kwasnik, Xue Li, Robert H. Tullman, Michele Valentin, Kathy Ramos, and R. Eric Walter for their assistance in the in vivo analysis of **2**, and Robert C. Anderson and Philip A. Bell for helpful discussions.

Supporting Information Available: Experimental procedures including characterization data for all compounds, biological methods, ORTEP drawing, and atomic coordinate information for **2**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Yaron, A.; Nadier, F. Proline-Dependent Structural and Biological Properties of Peptides and Proteins. *Crit. Rev. Biochem. Mol. Biol.* **1993**, *28*, 31–81. (b) Demuth, H.; Heins, J. Catalytic mechanism of dipeptidyl peptidase-IV. In *Dipeptidyl Peptidase-IV (CD26) in Metabolism and the Immune Response*; Fleischer, B., Ed.; Landes, R. G.: Austin, TX, 1995; Vol. 3, pp 1–35.
- (2) Mentlein, R. Dipeptidyl-peptidase IV (CD26)-role in the inactivation of regulatory peptides. *Regul. Pept.* **1999**, *85*, 9–24.
- (3) (a) Jacotot, E.; Callebaut, C.; Blanco, J.; Krust, B.; Neubert, K.; Barth, A.; Hovanessian, A. G. Dipeptidyl-peptidase IV-B, a novel form of cell-surface-expressed protein with dipeptidyl-peptidase IV activity. *Eur. J. Biochem.* **1996**, *239*, 248–258. (b) Schneider, B. L.; Thevananther, S.; Moyer, M. S.; Walters, H. C.; Rinaldo, P.; Devarajan, P.; Sun, A. Q.; Dawson, P. A.; Ananthanarayanan, M. Cloning and Characterization of a Novel Peptidase from Rat and Human Ileum. *J. Biol. Chem.* **1997**, *272*, 31006–31015. (c) Duke-Cohan, J. S.; Gu, J.; McLaughlin, D. F.; Xu, Y.; Freeman, G. J.; Schlossman, S. F. Attractin (DPPT-L), a member of the CUB family of cell adhesion and guidance proteins, is secreted by activate human T lymphocytes and modulate immune cell interactions. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, *95*, 11336–11341. (d) Niedermeyer, J.; Enenkel, B.; Park, J. E.; Lenter, M.; Rettig, W. J.; Damm, K.; Schnapp, A. Mouse fibroblast-activation protein-conserved *Fap* gene organization and biochemical function as a serine protease. *Eur. J. Biochem.* **1998**, *254*, 650–654. (e) Chiravuri, M.; Schmitz, T.; Yardley, K.; Underwood, R.; Dayal, Y.; Huber, B. T. A Novel Apoptotic Pathway in Quiescent Lymphocytes Identified by Inhibition of a Post-Proline Cleaving Aminodipeptidase: A Candidate Target Protease, Quiescent Cell Proline Dipeptidase. *J. Immunol.* **1999**, *163*, 3092–3099.
- (4) (a) Holst, J. J. Glucagon-like peptide-1 (GLP-1) a newly discovered GI hormone. *Gastroenterology* **1994**, *107*, 1048–1055. (b) Orsakov, C. Glucagon-like peptide-1, a new hormone of the enteroinular axis. *Diabetologia* **1992**, *35*, 701–711. (c) Drucker, D. J. Glucagon-Like Peptides. *Diabetes* **1998**, *47*, 159–169.
- (5) (a) Mentlein, R.; Gallwitz, B.; Schmidt, W. E. Dipeptidyl-peptidase IV hydrolyzes gastric inhibitory polypeptide, glucagon-like peptide-1 (7–36), peptide histidine methionine and is responsible for their degradation in human serum. *Eur. J. Biochem.* **1993**, *214*, 829–835. (b) Deacon, C. F.; Johnsen, A. H.; Holst, J. J. Degradation of glucagon-like peptide-1 by human plasma in vitro yields an N-terminally truncated peptide that is a major endogenous metabolite in vivo. *J. Clin. Endocrinol. Metab.* **1995**, *80*, 952–957. (c) Kieffer, T. J.; McIntosh, C. H. S.; Pederson, R. A. Degradation of glucose-dependent insulinotropic polypeptide and truncated glucagon-like peptide-1 in vitro and in vivo by dipeptidyl peptidase IV. *Endocrinology* **1995**, *136*, 3585–3596. (d) Lene, H.; Deacon, C. F.; Orskov, C.; Holst, J. J. Glucagon-like peptide-1-(7–36) amide is transformed to glucagon-like peptide-1-(9–36) amide by dipeptidyl peptidase IV in the capillaries supplying the L cells of the porcine intestine. *Endocrinology* **1999**, *140*, 5356–5363.
- (6) Ahren, B.; Holst, J. J.; Martensson, H.; Balkan, B. Improved glucose tolerance and insulin secretion by inhibition of dipeptidyl peptidase IV. *Eur. J. Pharmacol.* **2000**, *404* (1/2), 239–245.
- (7) Deacon, C. F.; Hughes, T. E.; Holst, J. J. Dipeptidyl peptidase IV inhibition potentiates the insulinotropic effect of glucagon-like peptide 1 in the anesthetized pig. *Diabetes* **1998**, *47*(5), 764–769.
- (8) (a) Pederson, R. P.; White, H. A.; Schlenzig, D.; Pauly, R. P.; McIntosh, C. R. P.; Demuth, H. U. Improved glucose tolerance in Zucker fatty rats by oral administration of the dipeptidyl peptidase IV inhibitor isoleucine thiazolidide. *Diabetes* **1998**, *47*(8), 1253–1258. (b) Pauly, R. P.; Demuth, H.-U.; Rosche, R.; Schmidt, J.; White, H. A.; Lynn, F.; McIntosh, C. H. S.; Pederson, R. A. Improved glucose tolerance in rats treated with the dipeptidyl peptidase IV (CD26) inhibitor Ile-thiazolidide. *Metabolism* **1999**, *3*, 385–389.
- (9) Balkan, B.; Kwasnik, L.; Miserendino, R.; Holst, J. J.; Li, X. Inhibition of dipeptidyl peptidase IV with NVP-DPP728 increases plasma GLP-1 (7–36 amide) concentrations and improves oral glucose tolerance in obese Zucker rats. *Diabetologia* **1999**, *42* (11), 1324–1331.
- (10) Marguet, D.; Baggio, L.; Kobayashi, T.; Bernard, A.-M.; Pierres, M.; Nielsen, P. F.; Ribet, U.; Wantanabe, T.; Drucker, D. J.; Wagtman, N. Enhanced insulin secretion and improved glucose tolerance in mice lacking CD26. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 6864–6879.
- (11) Holst, J. J.; Deacon, C. F. Inhibition of the activity of Dipeptidyl-Peptidase IV as a treatment for Type 2 Diabetes. *Diabetes* **1998**, *47*, 1663–1670.
- (12) (a) Ahren, B.; Larsson, J.; Holst, J. J. Effects of glucagon-like peptide-1 on islet function and insulin sensitivity in non-insulin dependent diabetes mellitus. *J. Clin. Endocrinol. Metab.* **1997**, *82*, 473–478. (b) Rachman, J.; Borrow, B. A.; Levy, J. C.; Turner, R. C. Near normalization of diurnal glucose concentrations by

- continuous administration of glucagon-like peptide 1 (GLP-1) in subjects with NIDDM. *Diabetologia* **1997**, *40*, 205–211. (c) Nauck, M. A.; Wollschlaeger, D.; Werner, J.; Holst, J. J.; Orskov, C.; Creutzfeldt, W.; Willms, B. Effects of subcutaneous glucagon-like peptide 1 (GLP-1 [7–36 amide]) in patients with NIDDM. *Diabetologia* **1996**, *39*, 1546–1553. (d) Drucker, D. J. Biological actions and therapeutic potential of the glucagon-like peptides. *Gastroenterology* **2002**, *122* (2), 531–544.
- (13) (a) Wettergren, A.; Schjoldager, B.; Martensen, P. E.; Myhre, J.; Christiansen, J.; Holst, J. J. Truncated GLP-1 (proglucagon 72–107 amide) inhibits gastric and pancreatic functions in man. *Dig. Dis. Sci.* **1993**, *38*, 665–673. (b) Nauck, M. A.; Niederreithholz, U.; Ettler, R.; Holst, J. J.; Orskov, C.; Ritzel, R.; Schmiguel, W. H. Glucagon-like peptide-1 inhibition of gastric-emptying outweighs its insulinotropic effects in healthy humans. *Am. J. Physiol.* **1997**, *273*, E981–E988. (c) Nauck, M. A. Is GLP-1 an incretin hormone? *Diabetologia* **1999**, *42*, 373–379. (d) Egan, J.; Clocquet, A. R.; Elahi, D. The Insulinotropic Effect of Acute Exendin-4-Administered to Humans: Comparison of Non-diabetic State to Type 2 Diabetes. *J. Clin. Endocrinol. Metab.* **2002**, *87* (3), 1282–1290.
- (14) (a) Habener, J. F. Glucagonlike peptide-1 agonist stimulation of β -cell growth and differentiation. *Curr. Opin. Endocrinol. Diabetes* **2001**, *8* (2), 74–81. (b) Ling, Z.; Wu, D.; Zambare, Y.; Flamez, D.; Drucker, D. J.; Pipeleers, D. G.; Schuit, F. C. Glucagon-like peptide 1 receptor signaling influences topography of islet cells in mice. *Virchows Arch.* **2001**, *438* (4), 382–387. (c) Aspelun, F.; Egan, J. M.; Slezak, L. A.; Sritharan, K. C.; Elahi, D.; Andersen, D. K. Glucagon-like peptide-1 and exendin-4 improve glucose tolerance and induce islet cell growth in pancreatitis in rats. *Surg. Forum* **2000**, *51*, 40–42. (d) Kemp, D. M.; Habener, J. F.; Insulinotropic hormone glucagon-like peptide 1 (GLP-1) activation of insulin gene promoter inhibited by p38 mitogen-activated protein kinase. *Endocrinology* **2001**, *142* (3), 1179–1187. (e) Parkes, D. G.; Pittner, R.; Jodka, C.; Smith, P.; Young, A. Insulinotropic actions of exendin-4 and glucagon-like peptide-1 in vivo and in vitro. *Metab., Clin. Exp.* **2001**, *50* (5), 583–589.
- (15) Flint, A.; Raben, A.; Ersboll, A. K.; Holst, J. J.; Astrup, A. The effect of physiological levels of glucagon-like peptide-1 on appetite, gastric emptying, energy and substrate metabolism in obesity. *Int. J. Obes.* **2001**, *25*, 781–792.
- (16) (a) Doyle, M. E.; Egan, J. M. Glucagon-like peptide-1. *Recent Prog. Horm. Res.* **2001**, *56*, 377–399. (b) Villanueva-Peñacarrillo, M. L.; Puente, J.; Redondo, A.; Clemente, F.; Valverde, I. Effect of GLP-1 treatment on GLUT2 and GLUT4 expression in type 1 and type 2 rat diabetic models. *Endocrine* **2001**, *15* (2), 241–248.
- (17) Juana, F.; Valdeolmillos, M. Glucose-dependent stimulatory effect of glucagon-like peptide 1(7–36) amide on the electrical activity of pancreatic β -cells recorded in vivo. *Diabetes* **1999**, *48* (4), 754–757.
- (18) These nonsubstrate DPP-IV inhibitors provide low micromolar IC_{50} 's and, except in refs 3a and 3b, contain a protonatable amine. (a) Shimazawa, R.; Takayama, H.; Kato, F.; Kato, M.; Hashimoto, Y. Nonpeptide small-molecular inhibitors of dipeptidyl peptidase IV; *N*-phenylphthalimide analogs. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 559–562. (b) Miyachi, H.; Kato, M.; Kato, F.; Hashimoto, Y. Novel potent nonpeptide aminopeptidase N inhibitors with a cyclic imide skeleton. *J. Med. Chem.* **1998**, *41* (3), 263–265. (c) Yamada, M.; Okagaki, C.; Higashijima, T.; Tanaka, S.; Ohnuki, T.; Sugita, T. A potent dipeptide inhibitor of dipeptidyl peptidase IV. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 1537–1540. (d) Coppola, G. M.; Zhang, Y. L.; Schuster, H. F.; Russell, M. E.; Hughes, T. E. 1-Aminomethylisquinoline-4-carboxylates as novel dipeptidylpeptidase IV inhibitors. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 1555–1558.
- (19) Irreversible oligopeptides with the N-terminal X–Pro–X sequence (low micromolar IC_{50}). Rahfeld, J.; Schierhorn, M.; Hartrodt, B.; Neubert, K.; Heins, J. Are Diprotin A and (Ile-Pro-Ile) and Diprotin B (Val-Pro-Leu) Inhibitors or Substrates of Dipeptidyl Peptidase IV? *Biochim. Biophys. Acta* **1991**, *1076*, 314–316.
- (20) Suicide substrate (latent quinoniminium methide electrophile) large cyclopeptide. Nguyen, C.; Blanco, J.; Mazaleyrat, J. P.; Krust, B.; Callebaut, C.; Jacotot, E.; Hovanessian, A. G.; Wakselman, M. Specific and Irreversible Cyclopeptide Inhibitors of Dipeptidyl Peptidase IV Activity of the T-Cell Activation Antigen CD26. *J. Med. Chem.* **1998**, *41*, 2100–2110.
- (21) (a) Heins, J.; Welker, P.; Schönlein, C.; Born, I.; Hartrodt, B.; Neubert, K.; Tsuru, D.; Barth, A. Mechanism of Proline-Specific Proteinases; (I) Substrate Specificity of Dipeptidyl Peptidase IV from Pig Kidney and Proline-Specific Endopeptidase from *Flavobacterium meningosepticum*. *Biochim. Biophys. Acta* **1988**, *954*, 161–169. (b) Barth, A.; Heins, J.; Fischer, G.; Neubert, K.; Schneeweiss, B. Dipeptidyl-peptidase IV: mechanism and specificity of the substrate cleavage. *Mol. Cell. Regul. Enzyme Act.* *Part 1–3* **1984**, *46*, 297–339. (c) Schön, E.; Born, I.; Demuth, H.-U.; Faust, J.; Neubert, K.; Steinmetzer, T.; Barth, A.; Ansorge, S. Dipeptidyl Peptidase IV in the Immune System. *Biol. Chem. Hoppe-Seyler* **1991**, *372*, 305–311.
- (22) For comprehensive reviews on DPP-IV inhibitors, please see the following. (a) Villhauer, E. B.; Coppola, G. M.; Hughes, T. E. DPP-IV inhibition and Therapeutic Potential. *Annu. Rep. Med. Chem.* **2001**, *36*, 191–200. (b) Augustyns, K.; Bal, G.; Thonus, G.; Belyaev, A.; Zhang, X. M.; Bollaert, W.; Lambeir, A. M.; Durinx, C.; Goossens, F.; Haemers, A. The Unique Properties of Dipeptidyl-Peptidase IV (DPP-IV/CD26) and the Therapeutic Potential of DPP-IV Inhibitors. *Curr. Med. Chem.* **1999**, *6*, 311–327. (c) Snow, R. J.; Bachovchin, W. W. Boronic acid inhibitors of dipeptidyl peptidase IV: a new class of immunosuppressive agents. *Adv. Med. Chem.* **1995**, *3*, 149–177.
- (23) Jenkins, P. D.; Jones, D. M. DP-IV-serine protease inhibitors. Patent WO 95/15309, June 5, 1995.
- (24) Ashworth, D. M.; Atrash, B.; Baker, G. R.; Baxter, A. J.; Jenkins, P. D. 2-Cyanopyrrolidides as potent, stable inhibitors of dipeptidyl peptidase IV. *Bioorg. Med. Chem. Lett.* **1996**, *6* (10), 1163–1166.
- (25) Li, J.; Wilk, E.; Wilk, S. Aminoacylpyrrolidine-2-nitriles: Potent and stable inhibitors of dipeptidyl-peptidase IV (CD26). *Arch. Biochem. Biophys.* **1995**, *323* (1), 148–154.
- (26) Ashworth, D. M.; Atrash, B.; Baker, G. R.; Baxter, A. J.; Jenkins, P. D. 4-Cyanothiazolidides as very potent, stable inhibitors of dipeptidyl peptidase IV. *Bioorg. Med. Chem. Lett.* **1996**, *6* (22), 2745–2748.
- (27) Villhauer, E. B. N-(Substituted glycoly)-2-cyanopyrrolidides, Pharmaceutical compositions containing them and their use in inhibiting dipeptidyl peptidase-IV. U.S. Patent 6,011,155, January 4, 2000.
- (28) The SAR of N-substituted glycine-2-(S)-cyanopyrrolidines as DPP-IV inhibitors will be presented in a full report in due course.
- (29) Hughes, T. E.; Mone, M. D.; Russell, M. E.; Weldon, S. C.; Villhauer, E. B. NVP-DPP728 (1-[[[2-[(5-Cyanopyridin-2-yl)-amino]acetyl]-2-cyano-(S)-pyrrolidine], a Slow-Binding Inhibitor of Dipeptidyl Peptidase IV. *Biochemistry* **1999**, *36*, 111597–11603.
- (30) (a) Villhauer, E. B.; Anderson, R. C.; Balkan, B.; Barilla, D.; Brinkman, J. A.; Dunn, E.; Dunning, B.; Graham, E. D.; Gu, H.; Gutierrez, C. M.; Hamilton, B. H.; Kwasnik, L. A.; Li, X.; Mangold, B. L.; Maniara, W. M.; Miserendino-Molteni, R.; Mone, M.; Naderi, G. B.; Ramos, K. L.; Russell, M. E.; Rothenberg, T. L.; Tullman, R. H.; Valentin, M.; Walter, R. E.; Weldon, S. C.; Hughes, T. E. N-Substituted glycoly 2-cyanopyrrolidines as a new family of DPP-IV inhibitor and their potential use in type 2 diabetes, Presented at the 221st National Meeting of the American Chemical Society, San Diego, 2001; Abstract MEDI-343. (b) Brinkman, J. A.; Villhauer, E. B.; Naderi, G. B.; Hughes, T. E.; Mone, M.; Russell, M. E.; Weldon, S. C. Design and synthesis of N-substituted glycoly 2-cyanopyrrolidines as a new class of DPP-IV inhibitors. Presented at the 222nd National Meeting of the American Chemical Society, Chicago, IL, 2001; Abstract MEDI-39.
- (31) Zuckermann, R. N.; Martin, E. J.; Spellmeyer, D. C.; Stauber, G. B.; Shoemaker, K. R.; Kerr, J. M.; Gianine, M.; Figliozzi, D. A.; Goff, M. A.; Siani, M. A.; Simon, R. J.; Banville, S. C.; Brown, E. G.; Wang, L.; Richter, L. S.; Moos, W. H. Discovery of nanomolar ligands for 7-transmembrane G-protein-coupled receptors from a diverse N(substituted)glycine peptoid library. *J. Med. Chem.* **1994**, *37*, 2678–2685.
- (32) Experimental data for individual compounds from tandem solid-phase/solution-phase library can be obtained from Supporting Information.
- (33) ORTEP drawing of NVP-DPP728 (2) and atomic coordinate information can be obtained from Supporting Information.
- (34) PPCE and DPP-II are the standard enzymes in DPP-IV selectivity studies. (a) Belyaev, A.; Zhang, X.; Augustyns, K.; Lambeir, A. M.; De Meester, I.; Vedernikova, I.; Scharpé, S.; Haemers, A. Structure–Activity Relationship of Diaryl Phosphonate Esters as Potent Irreversible Dipeptidyl Peptidase IV Inhibitors. *J. Med. Chem.* **1999**, *42*, 1041–1052. Other related enzymes are quite new, and their assays are not yet well validated. (b) Sedo, A.; Malik, R. Dipeptidyl peptidase IV-like molecules: homologous proteins or homologous activities. *Biochim. Biophys. Acta* **2001**, *1550*, 107–116.
- (35) (a) Rothenberg, P.; Kalbag, J.; Smith, H. T.; Gingerich, R.; Nedelman, J.; Villhauer, E.; McLeod, J.; Hughes, T. Treatment with a DPP-IV Inhibitor, NVP-DPP728, Increases Prandial Intact GLP-1 Levels and Reduces Glucose Exposure in Humans. *Diabetes* **2000**, *49* (1), A39. (b) Åhrén, B.; Simonsson, E.; Larsson, H.; Landin-Olsson, M.; Torgeirsson, H.; Jansson, P. A.; Sandqvist, M.; Båvenholm, P.; Efendic, S.; Eriksson, J. W.; Dickinson, S.; Holmes, D. Inhibition of dipeptidyl peptidase IV improves metabolic control over a 4 week study period in type 2 diabetes. *Diabetes Care* **2002**, *25* (5), 869–875.