

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/47356198>

Stereospecificity of the Dehydratase Domain of the Erythromycin Polyketide Synthase

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · OCTOBER 2010

Impact Factor: 12.11 · DOI: 10.1021/ja107344h · Source: PubMed

CITATIONS

31

READS

30

6 AUTHORS, INCLUDING:



Chiara Valenzano

University of Cambridge

6 PUBLICATIONS 223 CITATIONS

SEE PROFILE



Young-Ok You

George Mason University

6 PUBLICATIONS 152 CITATIONS

SEE PROFILE



Ashish Garg

Brown University

7 PUBLICATIONS 166 CITATIONS

SEE PROFILE

Stereospecificity of the Dehydratase Domain of the Erythromycin Polyketide Synthase

Chiara R. Valenzano,[†] Young-Ok You,[†] Ashish Garg,[†] Adrian Keatinge-Clay,[§] Chaitan Khosla,[‡] and David E. Cane^{*,†}

Department of Chemistry, Box H, Brown University, Providence, Rhode Island 02912-9108, Departments of Chemical Engineering, Chemistry, and Biochemistry, Stanford University, Stanford, California 94305, and Department of Chemistry and Biochemistry, The University of Texas at Austin, 1 University Station A5300, Austin, Texas 78712-0165

Received August 15, 2010; E-mail: David_Cane@brown.edu

Abstract: The dehydratase (DH) domain of module 4 of the 6-deoxyerythronolide B synthase (DEBS) has been shown to catalyze an exclusive *syn* elimination/*syn* addition of water. Incubation of recombinant DH4 with chemoenzymatically prepared *anti*-(2*R*,3*R*)-2-methyl-3-hydroxypentanoyl-ACP (**2a**-ACP) gave the dehydration product **3**-ACP. Similarly, incubation of DH4 with synthetic **3**-ACP resulted in the reverse enzyme-catalyzed hydration reaction, giving an ~3:1 equilibrium mixture of **2a**-ACP and **3**-ACP. Incubation of a mixture of propionyl-SNAC (**4**), methylmalonyl-CoA, and NADPH with the DEBS β -ketoacyl synthase-acyl transferase [KS6][AT6] didomain, DEBS ACP6, and the ketoreductase domain from tylactone synthase module 1 (TYLS KR1) generated *in situ* *anti*-**2a**-ACP that underwent DH4-catalyzed *syn* dehydration to give **3**-ACP. DH4 did not dehydrate *syn*-(2*S*,3*R*)-**2b**-ACP, *syn*-(2*R*,3*S*)-**2c**-ACP, or *anti*-(2*S*,3*S*)-**2d**-ACP generated *in situ* by DEBS KR1, DEBS KR6, or the rifamycin synthase KR7 (RIFS KR7), respectively. Similarly, incubation of a mixture of (2*S*,3*R*)-2-methyl-3-hydroxypentanoyl-*N*-acetylcysteaminethioester (**2b**-SNAC), methylmalonyl-CoA, and NADPH with DEBS [KS6][AT6], DEBS ACP6, and TYLS KR1 gave *anti*-(2*R*,3*R*)-**6**-ACP that underwent *syn* dehydration catalyzed by DEBS DH4 to give (4*R*,5*R*)-(E)-2,4-dimethyl-5-hydroxy-hept-2-enoyl-ACP (**7**-ACP). The structure and stereochemistry of **7** were established by GC-MS and LC-MS comparison of the derived methyl ester **7**-Me to a synthetic sample of **7**-Me.

Of the more than 2000 nonaromatic polyketides, the vast majority contain one or more disubstituted or trisubstituted double bonds, most of which have *E* (*trans*) geometry.¹ Moreover, essentially all polyketides that do not themselves display a double bond are biosynthesized by way of one or more unsaturated polyketide chain elongation intermediates. Thus although 6-deoxyerythronolide B (**1**, 6-dEB), the parent aglycone of the erythromycin family of antibiotics, does not have any double bonds in the final macrolactone, the responsible modular polyketide synthase (PKS), 6-dEB synthase (DEBS), does in fact harbor a dehydratase domain in module 4, termed DEBS DH4 (Figure 1).^{2,3}

Direct evidence for the intermediacy of an unsaturated polyketide in erythromycin biosynthesis first came from disruption of the NADPH-binding motif of the ER4 domain, resulting in accumula-




Figure 1. Proposed tetraketide substrate and pentaketide intermediates of DEBS module 4. The module has a KR, a DH, and an ER domain in addition to the obligate KS, AT, and ACP domains.


tion of a derivative of the corresponding (*E*)- $\Delta^{6,7}$ -anhydro-6-dEB by mutants of the erythromycin producer *Saccharopolyspora erythraea*.⁴ Although the stereochemistry of the substrate for the DEBS DH4 dehydratase is not known, the responsible ketoreductase, DEBS KR4, is predicted to generate the (3*R*)-diastereomer of the 2-methyl-3-hydroxyacyl-ACP pentaketide, as deduced from the presence of a Leu-Ala-Asp triad closely correlated with the formation of (3*R*)-3-hydroxyacyl-ACP polyketide intermediates.⁵ Indeed, the vast majority of KR domains that are paired with a DH domain appear to harbor a conserved “Leu-Asp-Asp” motif.^{5a,b} DEBS KR4 is also predicted to belong to the class of nonpimerizing ketoreductases, which would give rise to a (2*R*)-methyl group in the reduced product.^{5c}

To establish the substrate specificity and stereochemical course of the DEBS DH4-catalyzed dehydration we used a chemoenzymatic strategy to prepare the requisite ACP-bound substrate and product analogues, **2a**-ACP and **3**-ACP. To this end the free acids **2a** and **3** were each converted to the corresponding -SCoA thioesters, **2a**-SCoA and **3**-SCoA, and thence to *anti*-(2*R*,3*R*)-2-methyl-3-hydroxypentanoyl-ACP6 (**2a**-ACP) and the expected dehydration product, (*E*)-2-methylpent-2-enoyl-ACP (**3**-ACP), from DEBS *apo*-ACP6 using the phosphopantetheinyl transferase Sfp (Scheme 1A).⁶ The two ACP derivatives, which were readily distinguished by reversed-phase LC-ESI(+)-MS, both exhibited the expected molecular weights.⁷ The structures were each confirmed by the MS² phosphopantetheinate (PPant) ejection method which gave **2a**-pant, *m/z* 375.33, and **3**-pant, *m/z* 357.3, each with the predicted MW, as well as MS³ analysis of each of the characteristic PPant ejection fragments.⁸

Incubation of recombinant DEBS DH4⁹ with **2a**-ACP resulted in formation of the predicted dehydration product **3**-ACP, as

[†] Brown University.[‡] Stanford University.[§] The University of Texas at Austin.

Scheme 1. (A) Synthesis and Analysis of ACP-Bound Substrates and (B) DEBS DH4-Catalyzed Interconversion of **2a-ACP** and **3-ACP**




established by direct monitoring by LC-ESI(+)-MS³, including detection of the corresponding intact acyl-ACP and PPant ejection fragments for both **2a-ACP** and **3-ACP** (Scheme 1B). Similarly, incubation of DEBS DH4 with **3-ACP** resulted in the reverse enzyme-catalyzed hydration reaction, giving an ~3:1 equilibrium mixture of **2a-ACP** and **3-ACP**.¹⁰

We also carried out combinatorial incubations using mixtures of recombinant PKS domains in order to generate *in situ* each of the four diastereomers of **2a–2d-ACP** (Scheme 2).¹¹ In this manner, a mixture of the DEBS [KS6][AT6] didomain, DEBS ACP6, and TYLS KR1, the ketoreductase domain from module 1 of the tylactone synthase, was incubated with propionyl-SNAC (**4**), methylmalonyl-CoA, and NADPH to produce *anti*-(2R,3R)-**2a-ACP**.^{11b} Addition of recombinant DEBS DH4, either simultaneously with or subsequent to the formation of **2a-ACP**, resulted in dehydration to yield exclusively the predicted (*E*)-2-methylpent-2-enoyl-ACP (**3-ACP**), as confirmed by GC-MS analysis of the corresponding acid **3** and comparison with synthetic **3**.¹² By contrast, DEBS DH4 did not dehydrate either *syn*-(2S,3R)-**2b-ACP** or *syn*-(2R,3S)-**2c-ACP** generated by DEBS KR1 or KR6, respectively,^{11a,c} to either *E*-**3-ACP** or the corresponding *Z*-isomer **5-ACP**, nor did DEBS DH4 dehydrate *anti*-(2S,3S)-**2d-ACP** produced by recombinant RIFS KR7,¹³ the KR domain from module 7 of the rifamycin synthase.

In further confirmation of the stereochemistry of the dehydration reaction, incubation of DEBS DH4 with *anti*-(2R,3R,4S,5R)-2,4-dimethyl-3,5-dihydroxyheptanoyl-ACP (**6-ACP**) generated *in situ* from **2b-SNAC**, methylmalonyl-CoA, and NADPH by DEBS [KS6][AT6] + ACP6 + TYLS KR1, as previously described,^{11b}

Scheme 2. Stereochemistry of DEBS DH4-Catalyzed Dehydration



gave exclusively *E*-**7-ACP** (Scheme 2). The structure and stereochemistry of **7-ACP** were determined by chiral GC-MS and LC-MS analysis of the derived methyl ester **7-Me**, obtained by basic hydrolysis and treatment of the liberated acid with TMS-diazomethane, and comparison with an authentic synthetic standard of **7-Me**.¹⁴

Sequence alignments of the DEBS DH4 domain with numerous PKS and FAS DH domains reveal conserved ²⁴⁰⁹HXXXGXXXXP and ²⁵⁷¹D(A/V)(V/A)(A/L)(Q/H) motifs.² Site-directed mutagenesis of the conserved active site His2409 of the DEBS DH4 domain abolished DEBS activity in *Sac. erythraea*^{15a} while the analogous His mutation also inactivates the homologous DH2 domain of the picromycin synthase.^{15b} Together the conserved His and Asp residues comprise the catalytic dyad of the dehydratase, in which the active site His acts as a general base while the Asp2571, located 4.1 Å from H2409 at the base of the substrate tunnel, is thought to serve as a general acid.^{9,16,17}

Our results establish definitively that the DEBS DH4 domain catalyzes a *syn* elimination of water during erythromycin biosynthesis. The prototype dehydration catalyzed by the DH domain of the yeast FAS to give the characteristic disubstituted (*E*)-enoyl-ACP intermediates of fatty acid biosynthesis also takes place with net *syn* stereochemistry,¹⁸ as do the dehydrations catalyzed by the DH domains of module 2 of nanchangmycin synthase¹⁹ and module 2 of tylactone synthase.^{11b} Indeed, the significant levels of overall sequence identity (>40%) and similarity (>55%) and the presence of the conserved motifs containing the catalytic dyad in more than 50 DH domains from a wide range of modular PKS systems strongly suggest that the formation of all (*E*)-unsaturated polyketide intermediates involves a common *syn* dehydration mechanism.

Acknowledgment. This work was supported by NIH Grants GM22172 (D.E.C.) and CA66736 (C.K.) and a Welch Foundation Grant (A.K.C.).

Supporting Information Available: Experimental procedures, LC-ESI(+)-MS³, and GC-MS data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Shiomi, K.; Omura, S. In *Macrolide Antibiotics. Chemistry, Biology, and Practice*, 2nd ed.; Omura, S., Ed.; Academic Press: San Diego, CA, 2002; pp 1–56.
- (2) (a) Donadio, S.; Staver, M. J.; McAlpine, J. B.; Swanson, S. J.; Katz, L. *Science* **1991**, 252, 675–679. Donadio, S.; Staver, M. J.; McAlpine, J. B.; Swanson, S. J.; Katz, L. *Gene* **1992**, 115, 97–103. (b) Cortes, J.; Haydock, S. F.; Roberts, G. A.; Beviitt, D. J.; Leadlay, P. F. *Nature* **1990**, 348, 176–178.
- (3) Each PKS module typically has three core catalytic domains: the β -ketoacyl-ACP synthase (KS), the acyltransferase (AT), and the acyl carrier protein (ACP). Most modules also carry specific combinations of ketoreductase (KR), dehydratase (DH), and enoylreductase (ER) tailoring domains.
- (4) Donadio, S.; McAlpine, J. B.; Sheldon, P. J.; Jackson, M.; Katz, L. *Proc. Natl. Acad. Sci. U.S.A.* **1993**, 90, 7119–7123.
- (5) (a) Reid, R.; Piagentini, M.; Rodriguez, E.; Ashley, G.; Viswanathan, N.; Carney, J.; Santi, D. V.; Hutchinson, C. R.; McDaniel, R. *Biochemistry* **2003**, 42, 72–79. (b) Caffrey, P. *ChemBioChem* **2003**, 4, 654–657. (c) Keatinge-Clay, A. T. *Chem. Biol.* **2007**, 14, 898–908. Keatinge-Clay, A. T.; Stroud, R. M. *Structure* **2006**, 14, 737–748.
- (6) Quadri, L. E.; Weinreb, P. H.; Lei, M.; Nakano, M. M.; Zuber, P.; Walsh, C. T. *Biochemistry* **1998**, 37, 1585–1595.
- (7) Although it is not essential to the application of the PPant ejection methodology, **2a-ACP** and **3-ACP** were cleanly resolved under the LC conditions used.
- (8) Dorrestein, P. C.; Bumpus, S. B.; Calderone, C. T.; Garneau-Tsodikova, S.; Aron, Z. D.; Straight, P. D.; Kolter, R.; Walsh, C. T.; Kelleher, N. L. *Biochemistry* **2006**, 45, 12756–12766. (b) Meluzzi, D.; Zheng, W. H.; Hensler, M.; Nizet, V.; Dorrestein, P. C. *Biolorg. Med. Chem. Lett.* **2008**, 18, 3107–3111.
- (9) Keatinge-Clay, A. *J. Mol. Biol.* **2008**, 384, 941–953.
- (10) The equilibrium for the FAS catalyzed dehydration to a disubstituted enoyl-ACP strongly also favors the hydrated form by ~3:1. Cf. Brown, A.; Affleck, V.; Kroon, J.; Slabas, A. *FEBS Lett.* **2009**, 583, 363–368.

- (11) (a) Castonguay, R.; He, W.; Chen, A. Y.; Khosla, C.; Cane, D. E. *J. Am. Chem. Soc.* **2007**, *129*, 13758–13769. (b) Castonguay, R.; Valenzano, C. R.; Chen, A. Y.; Keatinge-Clay, A.; Khosla, C.; Cane, D. E. *J. Am. Chem. Soc.* **2008**, *130*, 11598–11599. (c) Valenzano, C. R.; Lawson, R. J.; Chen, A. Y.; Khosla, C.; Cane, D. E. *J. Am. Chem. Soc.* **2009**, *131*, 18501–18511.
- (12) The *E*-isomer **3** was cleanly resolved from the *Z*-isomer **5**, and the two compounds were not interconverted under the conditions of the reaction.
- (13) You, Y.-O. Cane, D. E., unpublished observations.
- (14) Although DH4 processed both **2a-ACP** and **6-ACP**, DEBS DH4 does not dehydrate the corresponding SNAC thioester analog, *anti*-(2*R*,3*R*)-**2a-SNAC** (ref 9). This observation is consistent with the presence of a presumptive ACP-binding region on the DH surface adjacent to the entrance to the substrate binding tunnel, a feature also found in bacterial FabZ proteins (ref 17).
- (15) (a) Bevitt, D. J.; Staunton, J.; Leadlay, P. F. *Biochem. Soc. Trans.* **1993**, *21*, 30S. (b) Wu, J.; Zaleski, T. J.; Valenzano, C.; Khosla, C.; Cane, D. E. *J. Am. Chem. Soc.* **2005**, *127*, 17393–17404.
- (16) In the corresponding Type II FAS FabZ domains, a conserved Glu serves the same role as this Asp residue (ref 17).
- (17) Kimber, M. S.; Martin, F.; Lu, Y.; Houston, S.; Vedadi, M.; Dharamsi, A.; Fiebig, K. M.; Schmid, M.; Rock, C. O. *J. Biol. Chem.* **2004**, *279*, 52593–52602. Swarnamukhi, P. L.; Sharma, S. K.; Bajaj, P.; Surolia, N.; Surolia, A.; Suguna, K. *FEBS Lett.* **2006**, *580*, 2653–2660. Zhang, L.; Liu, W.; Hu, T.; Du, L.; Luo, C.; Chen, K.; Shen, X.; Jiang, H. *J. Biol. Chem.* **2008**, *283*, 5370–5379.
- (18) Sedgwick, B.; Morris, C.; French, S. J. *J. Chem. Soc., Chem. Commun.* **1978**, 193–194.
- (19) Guo, X.; Liu, T.; Valenzano, C.; Deng, Z.; Cane, D. E. *J. Am. Chem. Soc.* **2010**, in press (DOI: 10.1021/ja1073432).

JA107344H