

Catalytic Enantioselective Synthesis of Prostaglandin E₁ Methyl Ester Using a Tandem 1,4-Addition-Aldol Reaction to a Cyclopenten-3,5-dione Monoacetal

Leggy A. Arnold, Robert Naasz, Adriaan J. Minnaard, and Ben L. Feringa*

Department of Organic and Molecular Inorganic Chemistry
Stratingh Institute, University of Groningen
Nijenborgh 4, 9747 AG Groningen, The Netherlands

Received March 28, 2001

Revised Manuscript Received May 4, 2001

Conjugate addition reactions are among the most important carbon-carbon bond formation reactions in organic synthesis,¹ and considerable progress has been made in the development of asymmetric Michael additions and 1,4-additions of organometallic reagents.² Recently, highly enantioselective copper-catalyzed conjugate addition reactions of diorganozinc reagents to enones have been reported.³ Among the various chiral ligands introduced for this purpose phosphoramidite **4**, developed in our laboratories, shows nearly complete stereocontrol in the reaction of (function-alized) dialkylzinc (R_2Zn) reagents with six-, seven- and eight-membered cycloalkenones.⁴ On the basis of this methodology, catalytic routes are now available to enantiomerically pure products, embedding cyclohexane and larger rings in their structure.⁵ In contrast, the catalytic enantioselective 1,4-addition to 2-cyclopentenone is a major challenge, particularly because chiral cyclopentane structures are ubiquitous in natural products. Employing TADDOL-based phosphoramidite ligands we obtained up to 62% ee when the Et_2Zn addition to 2-cyclopentenone was run in the presence of molecular sieves.⁶ Furthermore, with using chiral bidentate phosphoramidite ligands, the enantioselectivity improved to 83%.⁷ Chan⁸ reached 89% ee using a diphosphite ligand, whereas Pfaltz⁹ enhanced the enantioselectivity in this addition to 94%. Recently Hoveyda¹⁰ reported ee values up to 97% using a chiral peptide-based phosphine ligand in the 1,4-addition of diethylzinc to 2-cyclopentenone. Although these catalysts give excellent enantioselectivities, the isolated yields for the 3-substituted cyclopentanones are often moderate. Possible reasons are the lower reactivity of 2-cyclopentenone in comparison with other cyclic enones, the side-reactions of the resulting zinc enolate with the starting material and the high volatility of the 1,4-addition product. Performing the reaction in the presence of an aldehyde increases the yield considerably.^{4,6,11}

(1) (a) Perlmutter, P. In *Conjugate Addition Reactions in Organic Synthesis*; Tetrahedron Organic Chemistry Series, No. 9; Pergamon: Oxford, 1992. (b) Tomioka, K.; Nagaoka, Y. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag: Berlin/Heidelberg, 1999; Vol. 3; Chapter 31.1.

(2) For a review, see: Sibi, M. P.; Manyem, S. *Tetrahedron* **2000**, 56, 8033.

(3) For reviews, see: (a) Krause, N. *Angew. Chem., Int. Ed.* **1998**, 37, 283.

(b) Krause, N. Hoffmann-Röder, A. *Synthesis* **2001**, 171.

(4) (a) Feringa, B. L.; Pineschi, M.; Arnold, L. A.; Imbos, R.; de Vries, A. H. *M. Angew. Chem., Int. Ed. Engl.* **1997**, 36, 2620. (b) Feringa, B. L. *Acc. Chem. Res.* **2000**, 33, 346.

(5) (a) Naasz, R.; Arnold, L. A.; Pineschi, M.; Keller, E.; Feringa, B. L. *J. Am. Chem. Soc.* **1999**, 121, 1104. (b) Naasz, R.; Arnold, L. A.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2001**, in press.

(6) Keller, E.; Maurer, J.; Naasz, R.; Schrader, T.; Meetsma, A.; Feringa, B. L. *Tetrahedron: Asymmetry* **1998**, 9, 2409.

(7) Mandoli, A.; Arnold, L. A.; Salvadori, P.; Feringa, B. L.; *Tetrahedron: Asymmetry* **2001**, in press.

(8) Yan M.; Chan, A. S. C. *Tetrahedron Lett.* **1999**, 40, 6645.

(9) Escher, I. H.; Pfaltz, A. *Tetrahedron* **2000**, 56, 2879.

(10) Degradó, S. J.; Mizutani, H.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2001**, 123, 755.

(11) Kitamura, M.; Miki, T.; Nakano, K.; Noyori, R. *Tetrahedron Lett.* **1996**, 37, 5141.

Scheme 1




Table 1. Results of Tandem 1,4-Addition-Aldol Reactions According to Scheme 1

entry	enone	R_2Zn	$R''CHO$	prod.	yield [%] ^a	ee (3a-f) [%] ^b
1	1a	Et	Ph	2a	67	87
2	1a	<i>n</i> -Bu	Ph	2b	64	87
3	1b	Et	Ph	2c	76	94
4	1b	<i>n</i> -Bu	Ph	2d	69	94
5	1b	Et	<i>p</i> -Br-Ph	2e	69	96
6	1b	<i>n</i> -Bu	<i>p</i> -Br-Ph	2f	64	97

^a Isolated Yields. ^b Determined with HPLC (Daicel CHIRAL PAK-AD).

We report here the highly enantioselective catalytic tandem 1,4-addition-aldol reaction of dialkylzinc reagents to cyclopenten-3,5-dione monoacetals in the presence of aldehydes. These compounds show a higher reactivity, and the heavily functionalized products are less volatile. The usefulness of this new method is illustrated by the total synthesis of (-)-PGE₁ methyl ester in seven steps using achiral starting materials and only a catalytic amount of a chiral copper complex.



Monoacetals **1a** and **1b** were employed in the tandem 1,4-addition-aldol reaction with various aldehydes and dialkylzinc reagents (Scheme 1).¹² The catalyst was prepared in situ from 2 mol % $Cu(OTf)_2$ and 4 mol % (*S,R,R*)-phosphoramidite **4**.

Full conversion was reached after 16 h to provide exclusively *trans* substituted cyclopentanones **2a-f** in isolated yields up to 76% (Table 1). Excellent stereocontrol is also observed in the subsequent aldol step, as for the hydroxy ketones **2a-2f** diastereomeric ratios higher than 95:5 were measured. The configuration of the main product was determined by NOESY-NMR. The adducts **2a-f** were converted into the corresponding diketones **3a-f** in good yields to give single diastereomers suitable for ee determination by chiral HPLC. The enantioselectivity strongly depends on the acetal moiety present in the starting material as 87% ee for enone **3a** (entry 1) and 94% ee for enone **3c** (entry 3) was obtained. The use of different dialkylzinc reagents, however, has no influence on the selectivity of this reaction (entries 3 and 4). The structure of the aldehyde has a minor influence: the use of benzaldehyde and *p*-bromo benzaldehyde shows ee values of 94% and 97%, respectively (entries 4 and 6).

We have demonstrated therefore, that in the presence of 2 mol % of [(*S,R,R*)-**4**] $Cu(OTf)_2$ nearly complete stereocontrol over the formation of three consecutive stereocenters in this tandem 1,4-addition-aldol reaction is achieved, providing multifunctional cyclopentanones. These results inspired us to demonstrate the

(12) (a) Yoshida, Z.; Kimura, M.; Yoneda, S. *Tetrahedron Lett.* **1975**, 16, 1001. (b) All compounds exhibited spectroscopic data (¹H NMR, ¹³C NMR, HRMS) in accordance with the structures. Details of the synthesis of **1a**, **1b**, and **5** will be published in due course.

Scheme 2

Scheme 3^a

^a Key: (a) (1) 3 equiv Bu₄NF (1 M in THF), methylpropionate, DMSO, 80 °C, 20 min; (2) Ac₂O, DMAP, pyridine, 20 min; (b) 5 mol % Pd(CH₃CN)₂Cl₂, THF, 3 h; (c) K₂CO₃, MeOH, 18 h; (d) (NH₄)₂Ce(NO₃)₆, MeCN, borate-HCl buffer (pH = 8), 60 °C, 2 h.

usefulness of this catalytic method by applying it to the synthesis of (-)-PGE₁ methyl ester.¹³

The initial approach we followed for this catalytic asymmetric total synthesis is reminiscent of the three component coupling reaction introduced by Noyori et al.,¹⁴ a methodology which gives access to a variety of prostaglandins.¹⁵ However, the use of the required dialkenylzinc reagents instead of the previously used dialkylzincs did not lead to product formation. For this reason we developed a new strategy involving the introduction of the saturated α -chain with a functionalized zinc reagent and the ω -chain via an unsaturated aldehyde. The synthesis starts with enone **1b**, aldehyde **5**,^{12b,16} and the functionalized zinc reagent **6**¹⁷ (Scheme 2). In the presence of 3 mol % of the catalyst we obtained compound **7** in 60% yield as the only product as a mixture of diastereomers (ratio 83:17) which differ in the configuration at the exocyclic stereocenter bearing the hydroxy functionality. This one-pot procedure is carried out with an enone and an enal. To differentiate between these, the unsaturated aldehyde is equipped with a removable silyl substituent, exploiting the fact that β -disubstituted enones are not reactive in the 1,4-addition under these conditions. Reduction of the ketone moiety

of **7** proceeds with 95% stereoselectivity using Zn(BH₄)₂ in ether at -30 °C. Compound **8** was isolated after chromatography as a single isomer in 63% yield with an ee of 94%. In the next step the silyl substituent was removed using Bu₄NF in THF/DMSO to give compound **9** (Scheme 3). This concept comprises a novel protection and deprotection sequence for enones suitable for the catalytic 1,4-addition with dialkylzincs. The cleavage of vinyl carbon-silicon bonds with Bu₄NF was developed by Nozaki.¹⁸ However, under the normal reaction conditions hydrolysis of compound **9** was observed to be caused by water in the commercial THF solution of Bu₄NF. Adding first sacrificial methylpropionate to remove the water by hydrolysis and only afterwards **8**, the desilylated compound **9** was obtained as the only product and used without further purification. Acetylation of **9** afforded **10** in 71 % yield over two steps.

The 1,3-allylic transposition of **10** with a catalytic amount of Pd(CH₃CN)₂Cl₂ in THF proceeded with reasonable yield and full retention of configuration¹⁹ to give allylic acetate **11** with the required stereochemistry. After deacetylation in the presence of K₂CO₃ in MeOH, compound **12** was obtained in excellent yield. The last step is the deprotection of the ketone functionality to provide the labile β -hydroxy ketone moiety of the prostaglandin. This conversion was realized using a catalytic amount of (NH₄)₂-Ce(NO₃)₆ under nearly neutral conditions.²⁰ In this way PGE₁ methyl ester²¹ is obtained in 7% overall yield with 94% optical purity in seven steps from **1b**.

In conclusion we have demonstrated that cyclopenten-3,5-dione monoacetals give highly enantioselective tandem 1,4-addition-aldol reactions in the presence of dialkylzinc reagents and aldehydes using a catalytic amount of Cu(OTf)₂ and phosphoramidite ligand **4**. Furthermore this reaction is the key step in a short total synthesis of PGE₁ methyl ester, comprising a new route to this natural product.

Acknowledgment. Financial support by the ministry of economic affairs (EET grant) is gratefully acknowledged.

Supporting Information Available: Experimental details (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA015900+

(18) Oda, H.; Sato, M.; Morizawa, Y.; Oshima, K.; Nozaki, H. *Tetrahedron* **1985**, *41*, 3257.

(19) For the Pd(II)-mediated allylic acetate transposition in a modified prostaglandin intermediate, see: Grieco, P. A.; Takigawa, T.; Bongers, S. L.; Tanaka, H. *J. Am. Chem. Soc.* **1980**, *102*, 7588.

(20) Markó, I. E.; Ates, A.; Gautier, A.; Leroy, B.; Plancher, J. -M.; Quesnel, Y.; Vanherck, J.-C. *Angew. Chem., Int. Ed.* **1999**, *38*, 3207.

(21) The analytical and spectral data (TLC, HPLC, ¹H NMR, ¹³C NMR, CD, MS) [α]_D²³ -51° (c 1.0, CH₃OH) of **13** were identical with those of authentic material (Sigma) [α]_D²³ -54° (c 1.0, CH₃OH).

(13) Corey, E. J.; Cheng, X.-M. *The Logic of Chemical Synthesis*; Wiley: New York, 1989, Chapter 11.

(14) Suzuki, M.; Kawagishi, T.; Suzuki, T.; Noyori, R. *Tetrahedron Lett.* **1982**, *23*, 4057.

(15) Noyori, R. *Asymmetric Catalysis in Organic Synthesis*; Wiley & Sons: New York, 1994, Chapter 4.

(16) Magriotis, P. A.; Kim, K. D. *Tetrahedron Lett.* **1990**, *31*, 6137.

(17) Langer, F.; Waas, J.; Knochel, P. *Tetrahedron Lett.* **1993**, *34*, 5261.