

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/238126921>

Ternary Equilibrium for the System Water/Methyl Isobutyl Ketone/2Ethyl2-(hydroxymethyl)-1,3-propanediol

ARTICLE in JOURNAL OF CHEMICAL & ENGINEERING DATA · APRIL 1981

Impact Factor: 2.04 · DOI: 10.1021/je00024a600

CITATION

1

READS

21

1 AUTHOR:



Francisco J. Lozano

Tecnológico de Monterrey

34 PUBLICATIONS 367 CITATIONS

SEE PROFILE

Ternary Equilibrium for the System Water/Methyl Isobutyl Ketone/2-Ethyl-2-(hydroxymethyl)-1,3-propanediol

Francisco Lozano

J. Chem. Eng. Data, **1981**, 26 (2), 131-133 • DOI: 10.1021/je00024a600 • Publication Date (Web): 01 November 2004

Downloaded from <http://pubs.acs.org> on May 6, 2009

More About This Article

The permalink <http://dx.doi.org/10.1021/je00024a600> provides access to:

- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article



ACS Publications
High quality. High impact.

the pressure approaches the solvent critical point. Therefore, in this exercise, the fugacity coefficients calculated by using a modified Redlich-Kwong equation of state (7) consistent with the vapor-liquid equilibria data of this study were used to yield values of the Henry's law constant more consistent with the criteria set forth by Renon et al. (2).

Acknowledgment

The Phillips Petroleum Co. supplied the isopentane bath fluid for the studies. We also thank Mr. Ray Martin for extensive assistance with the experimental apparatuses. Special thanks are due Dr. Pervaiz Nasir for many helpful discussions. We are indebted to Dr. S. C. Hwang of Exxon Research and Engineering Co. for assistance in calculating fugacity coefficients of hydrogen in methane.

Literature Cited

- (1) Benham, Alvin L.; Katz, Donald L. *AIChE J.* **1957**, *3*, 33.
- (2) Beutler, D.; Renon, H. *AIChE J.* **1978**, *24*, 1122.
- (3) Dymond, J. H.; Smith, E. B. "The Virial Coefficients of Gases"; Clarendon Press: Oxford, 1969.
- (4) Fastowsky, M. G.; Gonikberg, V. G. *J. Phys. Chem. (Moscow)* **1970**, *14*, 427.

- (5) Freeth, F. A.; Verschoyle, T. T. H. *Proc. R. Soc. London, Ser. A.* **1931**, *130*, 453.
- (6) Goodwin, R. D. *NBS Tech. Note (U.S.)* **1974**, No. 653.
- (7) Gray, R. D., Jr. "Correlation of H_2 /Hydrocarbon VLE Using Redlich-Kwong Variants"; paper presented at the 70th Annual AIChE Meeting, New York, 1977.
- (8) Hayduk, W.; Laudie, H. *AIChE J.* **1973**, *19*, 1233.
- (9) Hwang, S. C., Exxon Research and Engineering Co., Florham Park, NJ, private communication, Aug 1980.
- (10) Kirk, B. S.; Ziegler, W. T. *Adv. Cryog. Eng.* **1965**, *10*, 160-70.
- (11) Krichevsky, I. R.; Kasarnovsky, J. S. *J. Am. Chem. Soc.* **1935**, *57*, 2168.
- (12) Levitskaya, E. P. *J. Tech. Phys. (Leningrad)* **1941**, *11*, 197.
- (13) Likhter, A. I.; Tikhonovich, N. P. *Zh. Tekh. Fiz.* **1940**, *10*, 1201-6.
- (14) Madison, J. J. *Anal. Chem.* **1958**, *30*, 1859.
- (15) Mraw, S. C.; Kobayashi, R. *J. Chromatogr. Sci.* **1977**, *15*, 191.
- (16) Mraw, S. C.; Hwang, S. C.; Kobayashi, R. *J. Chem. Eng. Data* **1978**, *23*, 135.
- (17) Nakahara, T.; Hirata, M. *J. Chem. Eng. Jpn.* **1969**, *2*, 137.
- (18) Orentlicher, M.; Prausnitz, J. M. *Chem. Eng. Sci.* **1964**, *19*, 775.
- (19) Pletsch, H.; Kohle, E. U. **1958**, *11*, 702.
- (20) Purcell, J. E.; Ettre, L. S. *J. Gas Chromatogr.* **1965**, *2*, 69.
- (21) Reid, R. C.; Prausnitz, J. M.; Sherwood, T. K. "The Properties of Gases and Liquids", 3rd ed.; McGraw-Hill, New York, 1977.
- (22) Sagara, H.; Arai, Y.; Saito, S. *J. Chem. Eng. Jpn.* **1972**, *5*, 339.
- (23) Steckel, F.; Tsai, F. *J. Chem. Ind. (Moscow)* **1939**, *16*, 24.
- (24) Yorizane, M.; Yoshimura, S.; Masuoka, H.; Toyama, A. *Proc. Int. Cryog. Eng. Conf.* **1968**, *1*, 57-68.

Received for review April 21, 1980. Accepted October 22, 1980. The Gas Research Institute and the Gas Processors Association are acknowledged for their support of this project.

Ternary Equilibrium for the System Water/Methyl Isobutyl Ketone/2-Ethyl-2-(hydroxymethyl)-1,3-propanediol

Francisco J. Lozano

Departamento de Ingeniería Química, División de Estudios de Posgrado, Facultad de Química, U.N.A.M., México 20, D. F., México

Liquid-liquid equilibrium has been determined for the ternary system water/methyl isobutyl ketone/2-ethyl-2-(hydroxymethyl)-1,3-propanediol at 25 and 70 °C. The latter compound is known commercially as trimethylolpropane.

Cyclohexanol (1) has been tested as a solvent for extracting 2-ethyl-2-(hydroxymethyl)-1,3-propanediol (TMP), but the two liquid phase region for this system is not very large. It is the purpose of the present paper to show the behavior of another solvent in the extraction of TMP.

Experimental Procedure

The water was purified by distilling it in the presence of potassium permanganate. The methyl isobutyl ketone was supplied by Merck as a spectroscopic reagent. The TMP was purified by extracting impurities with methyl isobutyl ketone followed by drying at 100 °C for 24 h. The melting point for the final TMP was 57-57.5 °C.

The experiments were carried out in a small glass stirring vessel of 100-mL volume, provided with a circulating jacket. The circulating-water temperature was controlled by a recirculating bath. A PTFE-covered magnetic bar was used for stirring.

The points on the binodal curve were first obtained by using the titration method. Tie lines were obtained secondly by preparing overall mixtures of the three components, which would yield an overall composition within the two liquid phase region, stirring them for 4 h, and allowing the mixture to settle for 6 h.

Table I. Liquid-Liquid Equilibrium Composition Data ($T = 25^\circ\text{C}$)

binodal curve (wt fraction)					
water		TMP		MIBK	
0.981				0.019	
0.877		0.097		0.026	
0.779		0.195		0.026	
0.675		0.290		0.035	
0.568		0.380		0.052	
0.467		0.464		0.069	
0.358		0.536		0.106	
0.253		0.584		0.163	
0.166		0.585		0.249	
0.081		0.495		0.424	
0.046		0.382		0.572	
0.033		0.291		0.676	
0.028		0.195		0.777	
0.029		0.097		0.874	
0.021				0.979	
tie lines (wt fraction)					
water-rich phase			ketone-rich phase		
water	TMP	MIBK	water	TMP	MIBK
0.869	0.110	0.021	0.021	0.008	0.971
0.573	0.378	0.049	0.019	0.037	0.944
0.477	0.458	0.065	0.022	0.048	0.930
0.282	0.574	0.144	0.022	0.078	0.900
0.168	0.582	0.250	0.025	0.122	0.853
0.119	0.556	0.325	0.024	0.186	0.790
0.073	0.487	0.440	0.027	0.227	0.746
0.034	0.316	0.650	0.028	0.247	0.725
					distribution coeff, ketone/water
					0.071
					0.097
					0.105
					0.136
					0.210
					0.335
					0.466
					0.782

At the end of each experiment, samples were taken from both phases and weighed, and then both water and solvent were




Figure 1. Two liquid phase region at 25 °C for the system water/methyl isobutyl ketone/trimethylolpropane.




Figure 2. Two liquid phase region at 70 °C for the system water/methyl isobutyl ketone/trimethylolpropane.

Table II. Liquid-Liquid Equilibrium Composition Data ($T = 70^\circ\text{C}$)

binodal curve (wt fraction)		
water	TMP	MIBK
0.983		0.017
0.788	0.188	0.024
0.663	0.299	0.038
0.538	0.378	0.084
0.385	0.458	0.157
0.217	0.435	0.348
0.150	0.340	0.510
0.119	0.264	0.617
0.081	0.183	0.736
0.040		0.960

tie lines (wt fraction)						distribution coeff, ketone/ water
water-rich phase			ketone-rich phase			
water	TMP	MIBK	water	TMP	MIBK	
0.716	0.242	0.042	0.048	0.042	0.910	0.174
0.529	0.385	0.086	0.060	0.084	0.856	0.218
0.371	0.455	0.174	0.064	0.106	0.830	0.233
0.312	0.469	0.219	0.081	0.154	0.765	0.328
0.231	0.439	0.330	0.086	0.174	0.740	0.396

evaporated to dryness at 100°C . Using these data and the binodal curve, we obtained the tie lines. The error in the determination of the equilibrium diagram was within 5%.

Results

Experimental data for the equilibrium of the ternary system

water/methyl isobutyl ketone/TMP in the two liquid phase region were obtained. The binodal curve as well as the tie lines were determined, and the distribution coefficients were calculated. The results are presented in Tables I and II. In Figures 1 and 2 the ternary diagrams at 25 and 70°C are shown, respectively.

Conclusions

The two liquid phase region for this system is larger than the one for the cyclohexanol as a solvent (1), even at 70°C . Nevertheless, the distribution coefficients for the present system are less than those for the cyclohexanol system. Clearly the ketone does not offer a great advantage over cyclohexanol, but it has to be taken into account that TMP is normally present with a formate salt in commercial process solutions, and this will tend to increase the distribution coefficient and enlarge the two liquid phase region.

Literature Cited

- (1) Castañeda, J. M.; Lozano, F. J.; Trejo S. *J. Chem. Eng. Data*, following paper in this issue.

Received for review April 24, 1980. Accepted November 4, 1980.

Ternary Equilibrium for the System Water/Cyclohexanol/2-Ethyl-2-(hydroxymethyl)-1,3-propanediol

Jaun M. Castañeda, Francisco J. Lozano,* and Sergio Trejo

Departamento de Ingeniería Química, División de Estudios de Posgrado, Facultad de Química, U.N.A.M., México 20, D. F., México

Liquid-liquid equilibrium at 25 and 35°C has been determined for the ternary system water/cyclohexanol/2-ethyl-2-(hydroxymethyl)-1,3-propanediol. The latter is known commercially as trimethylolpropane.

The ability of cyclohexanol to extract trimethylolpropane from water was determined at two temperatures for the purpose of exploring its capability to serve as a commercial process solvent.

Experimental Procedure

The water was purified by distilling it in the presence of potassium permanganate. Cyclohexanol was fractionally distilled in the presence of calcium oxide. The melting point obtained was 25.1°C , and a chromatographic analysis showed a purity of 99.9%. The 2-ethyl-2-(hydroxymethyl)-1,3-propanediol (TMP) was purified by extracting impurities with methyl isobutyl ketone followed by drying at 100°C for 36 h. The melting point for the final TMP was $57-57.5^\circ\text{C}$.

The experiments were carried out in a small glass stirring vessel of 100-mL volume, provided with a cooling jacket. The cooling-water temperature was controlled by means of a recirculating constant-temperature bath. A PTFE-covered mag-

netic bar was used for stirring.

To determine the time needed for the mixture to reach equilibrium, we measured the refractive index of one of the phases as a function of time; the minimum contact time to achieve equilibrium was 2 h. Nevertheless, stirring was kept for 4 h, and the mixture was allowed to settle for 6 h more.

Equilibrium data were obtained by preparing mixtures of known overall composition, stirring for 4 h, and settling for 6 h at constant temperature (25 and 35°C , $\pm 0.1^\circ\text{C}$).

At the end of each experiment, samples were taken from both phases and were analyzed by means of nuclear magnetic resonance (NMR) (1). This analytical method allowed us to measure the proton contribution of each functional group in the molecules present, and from there the mass fraction of each component in the phase was determined. The accuracy of the analytical method was tested by comparing the binary equilibrium for the system water/cyclohexanol, whose data were taken from the literature; the error was within 2%. Determination of the composition in a ternary mixture at equilibrium using the NMR method has been successfully used elsewhere (2, 3).

Results

In the present work, experimental data for the composition at equilibrium of the system water/cyclohexanol/TMP at the two