

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/244418975>

A New Method For the Construction of α-Diazo ketones

ARTICLE *in* THE JOURNAL OF ORGANIC CHEMISTRY · APRIL 1995

Impact Factor: 4.72 · DOI: 10.1021/jo00112a064

CITATIONS

34

READS

16

5 AUTHORS, INCLUDING:



Mark Gleave

Epoch BioSciences

18 PUBLICATIONS 270 CITATIONS

SEE PROFILE



R Jason Herr

AMRI

38 PUBLICATIONS 1,050 CITATIONS

SEE PROFILE


A New Method for the Construction of α -Diazoketones

Douglass F. Taber,* D. Mark Gleave, R. Jason Herr, Kimberly Moody, and Michael J. Hennessy

Department of Chemistry and Biochemistry, University of Delaware, Newark, Delaware 19716


Received December 6, 1994

α -Diazoketones **3** undergo a variety of useful reactions, making them valuable synthetic intermediates.^{1,2} Unsymmetrical α -diazoketones have been prepared^{3ab} by condensing the requisite acid chloride with a diazoalkane. As diazoalkanes are difficult to prepare and to purify, we have now developed an alternative method for the regioselective construction of α -diazoketones, based on the alkylation of benzoylacetone **1** ($R = H$). We now report the key observation that diazo transfer^{3,4} to a diketone **2** proceeds with selective *debenzoylation*⁵ to provide the desired unsymmetrical α -diazoketone **3** (eq 1).



α -Alkylation of benzoylacetone (**1a**) using potassium carbonate in the presence of tetra-*n*-butylammonium bromide in toluene⁶ produced a variety of α -substituted diketones **2**, as shown in Table 1. Alternatively, ben-

zoylacetone may be γ -alkylated as its dianion⁷ to provide diketone **1b**, which is then α -alkylated to provide a new unsymmetrical diketone **2e** (eq 2).



Various methods of debenzoylation/diazo transfer were then attempted, including the often-successful combination of NaH and methanesulfonyl azide^{3b} in a variety of solvents, but with mediocre results. The use of *p*-nitrobenzenesulfonyl azide (*p*-NBSA)⁸ and DBU,⁹ however, produced α -diazoketones **3** in much higher yields (cf. Table 1).¹⁰ In summary, we have developed a new method for the regioselective construction of unsymmetrical α -diazoketones using the inexpensive benzoylacetone as a precursor. This opens the way for the exploration of synthetic applications of these versatile intermediates.

Experimental Section¹¹

Preparation of 1-Benzoyl-2-undecanone (1b). To a solution of 4.4 mmol of LDA [prepared from 0.62 mL (4.4 mmol) of diisopropylamine and 1.93 mL (4.4 mmol, 2.28 M solution in hexanes) of *n*-BuLi] in 4 mL of anhydrous THF at -78°C under nitrogen was added dropwise a solution of 0.386 g (2.4 mmol) of benzoylacetone in 2 mL of anhydrous THF. The mixture was then warmed to -20°C and stirred for 3 h, after which 0.38 mL (2.2 mmol) of *n*-bromooctane was added dropwise over 10 min, and the mixture was then slowly warmed to rt and stirred for 12 h. The mixture was then diluted with 20 mL of saturated aqueous NH_4Cl and extracted twice with 8 mL portions of ethyl acetate. The combined organic extracts were then washed with 15 mL of brine solution, dried over Na_2SO_4 , and concentrated *in vacuo*. The crude residue was then chromatographed to produce **1b** as a colorless, partially enolic oil (0.462 g, 84%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.50; ^1H NMR (250 MHz, CDCl_3) δ 7.88 (d, 2H, $J = 7.0$ Hz), 7.60–7.40 (m, 3H), 6.17 (s, 0.92H), 4.08 (s, 0.16H), 2.58 (t, 0.08H, $J = 7.3$ Hz), 2.42 (t, 0.92H, $J = 7.3$ Hz), 1.75–1.60 (m, 2H), 1.45–1.05 (m, 2H), 0.88 (t, 3H, $J = 6.3$ Hz); ^{13}C NMR (62.9 MHz, CDCl_3) δ up: 196.9, 183.3, 135.0, 39.1, 31.8, 29.4, 29.3, 29.2 (2), 25.7, 22.6; down: 132.1, 128.4, 126.9, 95.9, 14.0; IR (film) 3058, 2836, 1614, 1574, 1264, 1080, 762, 689 cm^{-1} ; EI MS m/z (rel intensity) 274 (M^+ , 1), 162 (41), 161 (17), 147 (38), 120 (12), 105 (100), 77 (26), 69 (61), 55 (14); HRMS (calcd for $\text{C}_{18}\text{H}_{26}\text{O}_2$) 274.1933, found

(1) For general reviews on the use of α -diazoesters and α -diazoketones in synthesis, see: (a) Doyle, M. P. In *Homogeneous Transition Metal Catalysts in Organic Synthesis*; Moser, W. R., Slocum, D. W., Eds., ACS Advanced Chemistry Series 230, American Chemistry Society, Washington, D.C., 1992; Ch. 30. (b) Taber, D. F.; Askani, R. In *Comprehensive Organic Synthesis*; Pergamon Press: Oxford, 1991; Vol. 6, p 103.

(2) For some synthetic applications of α -diazoketones see: (a) Wolff Rearrangement/Arndt–Eistert synthesis: Tsuji, T.; Nishida, S. *J. Am. Chem. Soc.* **1988**, *110*, 2157. (b) Insertion reactions involving cleavage of heteroatom–H bonds: (i) Boduraw, C. C.; Boyer, B. D.; Brennan, J.; Runnell, C. A.; Burks, J. E.; Carr, M. A.; Doeke, C. W.; Eckrich, T. M.; Fisher, J. W.; Gardener, J. P.; Graves, B. J.; Hines, P.; Hoying, R. C.; Jackson, B. G.; Kinnick, M. D.; Kochert, C. D.; Lewis, J. S.; Luke, W. D.; Moore, L. L.; Morin, J. M., Jr.; Nist, R. L.; Prather, D. E.; Sparks, D. L.; Vladuchick, W. C. *Tetrahedron Lett.* **1989**, *30*, 2321. (ii) Moody, C. J.; Taylor, R. J. *J. Chem. Soc., Perkin Trans. 1* **1989**, 721. (iii) Bagheri, V.; Doyle, M. P.; Taunton, J.; Claxton, E. E. *J. Org. Chem.* **1988**, *53*, 6158. (c) Ylide formation: Padwa, A.; Carter, S. P.; Nimmesgern, H.; Stull, P. D. *J. Am. Chem. Soc.* **1988**, *110*, 2894. (d) Elimination/enone formation: Duggleby, P. McC.; Holt, G.; Hope, M. A.; Lewis, A. *J. Chem. Soc., Perkin Trans. 1* **1972**, 3020. (e) Enolate formation: Masamune, S.; Mori, S.; Van Horn, D.; Brookes, D. W. *Tetrahedron Lett.* **1979**, *19*, 665. (f) Reaction of acetylenes with α,β -unsaturated α -diazoketones: Danheiser, R. L.; Brisbois, R. G.; Kowalczyk, J. K.; Miller, R. F. *J. Am. Chem. Soc.* **1990**, *112*, 3093. (g) Formation of dialkyl phosphonates: Polozov, A. M.; Polezhoeva, N. A.; Mustaphin, A. H.; Khotinen, A. V.; Arbuzov, B. A. *Synthesis* **1990**, 515. (h) Formation of tertiary α -aminoketones: West, F. G.; Glaeske, K. W.; Naidu, B. N. *Synthesis* **1993**, *10*, 977.

(3) For reviews of diazoketone preparation, see: (a) Regitz, M.; Mass, G. In *Diazo Compounds: Properties and Synthesis*, Academic Press: New York, 1986. (b) Taber, D. F. *Comprehensive Organic Synthesis*; Pattenden, G., Ed., Pergamon Press: Oxford, 1991; Vol. 3, p 1045. For more recent reagents for diazo transfer, see: (c) Ghosh, S.; Datta, I. *Synth. Commun.* **1991**, *21*, 191. (d) McGuiness, M.; Shechter, H. *Tetrahedron Lett.* **1990**, *31*, 4987. (e) Kumar, S. M. *Synth. Commun.* **1991**, *21*, 2121.

(4) Formylation is the most common method for α -methylene activation: (a) Hendrickson, J. B.; Wolf, W. A. *J. Org. Chem.* **1968**, *33*, 3610. (b) Taber, D. F.; Ruckle, R. E.; Hennessy, M. J. *J. Org. Chem.* **1986**, *51*, 4077. (c) For the use of trifluoroacetylation, see Danheiser, R. L.; Miller, R. F.; Brisbois, R. G.; Park, S. Z. *J. Org. Chem.* **1990**, *55*, 1959.

(5) The use of a benzoyl activating group in the synthesis of α -diazoketones is preceded: Metcalf, B. W.; Jund, K.; Burkhart, J. P. *Tetrahedron Lett.* **1980**, *21*, 15.

(6) Choudhary, A.; Baumstark, A. L. *Synthesis* **1989**, 688.

(7) Huckin, S. N.; Weiler, L. *J. Am. Chem. Soc.* **1974**, *96*, 1082.
















(8) Reagan, M. T.; Nickon, A. *J. Am. Chem. Soc.* **1968**, *90*, 4096.

(9) For the use of DBU to promote diazo transfer, see: Koteswar Rao, Y.; Nagarajan, M. *J. Org. Chem.* **1989**, *54*, 5678.

(10) In some cases the use of MsN_3 instead of *p*-NBSA is advantageous due to the ease of removal of MsNH_3 and MsNH_2 versus *p*-NBSA and *p*-nitrobenzenesulfonamide from the crude reaction mixture (see ref 3b).

(11) For a summary of general experimental procedures see: Taber, D. F.; Houze, J. B. *J. Org. Chem.* **1994**, *59*, 4004.

Table 1. α -Alkylation and Diazo Transfer of Diketones 1

Entry	Diketone	Alkylation Product	Yield (%)	Diazoketone	Yield (%)
1			77		72
2			59 ^a		70
3			90		83
4			72		63
5			61		70

^a This reaction was performed using 10 mol % *n*Bu₄NI.

274.1925. Anal. Calcd for C₁₈H₂₆O₂: C, 78.79%; H, 9.55%. Found: C, 78.80%; H, 9.84%.

3-Benzoyl-2-butanone (2a). A mixture of 8.88 g (64.3 mmol) of anhydrous, powdered K₂CO₃, 0.052 g (0.16 mmol) of *n*-Bu₄NBr and 2.34 g (14.4 mmol) of benzoylacetone in 30 mL of anhydrous toluene under nitrogen was heated at reflux for 3.5 h. The mixture was then cooled to 40 °C for the dropwise addition of 1.00 mL (16.1 mmol) of iodomethane. The mixture was then stirred at 40 °C for an additional 18 h. The mixture was cooled to 0 °C and filtered, and the residue was washed twice with 100 mL portions of petroleum ether. The combined filtrate was concentrated, and the residue was chromatographed to produce **2a** as a yellow oil (1.96 g, 77%). TLC *R_f* (5% ethyl acetate/petroleum ether) 0.32; bp 80 °C/1 mmHg; ¹H NMR (250 MHz, CDCl₃) δ 7.97 (d, 2H, *J* = 7.0 Hz), 7.65–7.35 (m, 3H), 4.49 (q, 1H, *J* = 6.9 Hz), 2.15 (s, 3H), 1.45 (d, 3H, *J* = 6.8 Hz); ¹³C NMR (62.9 MHz, CDCl₃) δ up: 133.1, 128.4, 128.1, 55.6, 27.7, 13.0; down: 204.3, 197.0, 135.5; IR (film) 3063, 2987, 1722, 1674, 1596, 1449 cm⁻¹; EI MS *m/z* (rel intensity) 176 (M⁺, 1), 134 (9), 133 (15), 105 (100), 77 (53), 51 (21); HRMS (calcd for C₁₁H₁₂O₂) 176.0837, found 176.0833.

3-Benzoyl-2-undecanone (2b): TLC *R_f* (20% ethyl acetate/petroleum ether) = 0.64; bp 120 °C/1 mmHg; ¹H NMR (250 MHz, CDCl₃) δ 7.92 (d, 2H, *J* = 7.1 Hz), 7.51–7.37 (m, 3H), 4.39 (t, 0.9 Hz, *J* = 7.0 Hz), 3.88 (t, 0.1 Hz, *J* = 7.0 Hz), 2.36 (s, 0.3H), 2.07 (s, 2.7H), 2.00–1.82 (m, 2H), 1.30–1.05 (m, 12H), 0.79 (t, 3H, *J* = 6.9 Hz); ¹³C NMR (62.9 MHz, CDCl₃) δ up: 133.4, 128.6, 128.4, 63.2, 27.6, 13.8; down: 204.1, 196.3, 136.4, 31.6, 29.3, 29.1, 29.0, 28.9, 27.5, 22.4; IR (film) 3063, 2925, 1723, 1678, 1597, 1448, 1357 cm⁻¹; EI MS *m/z* (rel intensity) 162 (M⁺, 21), 133 (13), 105 (100), 77 (30), 55 (10); HRMS (calcd for C₁₈H₂₆O₂) 274.1933, found 274.1845.

Methyl (Z)-8-Benzoyl-9-oxo-5-decenoate (2c): TLC *R_f* (20% ethyl acetate/petroleum ether) = 0.35; bp 150 °C/1 mmHg; ¹H NMR (250 MHz, CDCl₃) δ 7.98 (d, 2H, *J* = 5.1 Hz), 7.65–7.35 (m, 3H), 5.45–5.25 (m, 2H), 4.47 (t, 1H, *J* = 7.2 Hz), 3.66 (s, 3H), 2.73 (t, 2H, *J* = 6.9 Hz), 2.30 (t, 2H, *J* = 7.4 Hz), 2.15 (s, 3H), 2.11 (m, 2H), 1.67 (m, 2H); ¹³C NMR (62.9 MHz, CDCl₃) δ up: 203.1, 195.7, 173.4, 136.0, 32.8, 26.4, 26.0, 24.1; down: 133.3, 131.0, 128.4, 128.2, 125.7, 62.0, 50.9, 28.0; IR (film) 3008, 2951, 1732, 1676, 1596, 1448 cm⁻¹; EI MS *m/z* (rel intensity) 302 (M⁺,

1), 259 (5), 227 (5), 161 (13), 140 (34), 105 (100), 77 (52); HRMS (calcd for C₁₈H₂₂O₄) 302.1516, found 302.1518. Anal. Calcd for C₁₈H₂₂O₄: C, 71.50%; H, 7.33%. Found: C, 71.24%; H, 7.29%.

(E)-3-Benzoyl-6,10-dimethylundeca-5,9-dien-2-one (2d): TLC *R_f* (5% ethyl acetate/petroleum ether) = 0.18; bp 150 °C/1 mmHg; ¹H NMR (250 MHz, CDCl₃) δ 7.97 (d, 2H, *J* = 7.0 Hz), 7.65–7.40 (m, 3H), 5.10–4.95 (m, 2H), 4.46 (t, 1H, *J* = 7.2 Hz), 2.70 (t, 2H, *J* = 7.2 Hz), 2.15 (s, 3H), 2.05–1.85 (m, 4H), 1.62 (s, 3H), 1.61 (s, 3H), 1.55 (s, 3H); ¹³C NMR (62.9 MHz, CDCl₃) δ up: 203.9, 196.2, 138.0, 136.4, 131.2, 39.4, 27.6, 26.2; down: 133.4, 128.6, 128.5, 123.8, 119.8, 62.9, 28.0, 25.4, 17.5, 15.9; IR (film) 3061, 2922, 1722, 1681, 1597, 1448 cm⁻¹; EI MS *m/z* (rel intensity) 298 (M⁺, 2), 255 (6), 161 (9), 105 (100), 77 (36), 69 (31); HRMS (calcd for C₂₀H₂₆O₂) 298.1933, found 298.1943.

2-Benzoyl-3-dodecanone (2e): TLC *R_f* (10% ethyl acetate/petroleum ether) = 0.38; ¹H NMR (250 MHz, CDCl₃) δ 7.97 (d, 2H, *J* = 7.3 Hz), 7.65–7.40 (m, 3H), 4.49 (q, 1H, *J* = 7.1 Hz), 2.55–2.30 (m, 2H), 1.52 (m, 2H), 1.44 (d, 3H, *J* = 6.9 Hz), 1.30–1.05 (m, 12H), 0.86 (t, 3H, *J* = 6.3 Hz); ¹³C NMR (62.9 MHz, CDCl₃) δ up: 133.5, 128.8, 128.6, 56.2, 14.0, 13.5; down: 207.1, 197.3, 136.1, 40.6, 31.8, 29.3, 29.2 (2), 28.9, 23.5, 22.6; IR (film) 2926, 1712, 1678, 1596, 1450 cm⁻¹; EI MS *m/z* (rel intensity) 288 (M⁺, 1), 134 (81), 133 (16), 105 (100), 77 (40), 71 (16), 57 (22); HRMS (calcd for C₁₉H₂₈O₂) 288.2089, found 288.2094. Anal. Calcd for C₁₉H₂₈O₂: C, 79.12%; H, 9.78%. Found: C, 78.85%; H, 9.81%.

3-Diazobutan-2-one (3a). To a solution of 1.82 g (10.3 mmol) of diketone **2a** in 40 mL of anhydrous CH₂Cl₂ at 0 °C under nitrogen was added 3.09 mL (20.7 mmol) of DBU, followed by the dropwise addition of a solution of 4.71 g (20.7 mmol) of *p*-NBSA in 10 mL of anhydrous CH₂Cl₂. The mixture was warmed to rt and stirred for 1 h, after which the solvent was removed *in vacuo*. The product was isolated by fractional distillation (87–89 °C/90 mmHg) to produce **3a** as a yellow oil (0.73 g, 72%). The spectral data of **3a** were identical to the known compound.¹² ¹H NMR (250 MHz, CDCl₃) δ 1.96 (s, 3H), 2.25 (s, 3H); ¹³C NMR (62.9 MHz, CDCl₃) δ up: 24.5, 7.4; down: 190.8; IR (film) 2929, 2072, 1638, 1328, 1296 cm⁻¹.

(12) (a) Regitz, M.; Menz, F. *Chem. Ber.* **1961**, *101*, 2622. (b) Diels, O.; Pfäumer, K. *Ber. Dtsch. Chem. Ges.* **1915**, *58*, 223.

3-Diazoundecan-2-one (3b). TLC R_f (20% ethyl acetate/petroleum ether) = 0.55; ^1H NMR (250 MHz, CDCl_3) δ 2.32 (t, 2H, J = 7.6 Hz), 2.23 (s, 3H), 1.46 (m, 2H), 1.40–1.15 (m, 10H), 0.88 (t, 3H, J = 6.9 Hz); ^{13}C NMR (62.9 MHz, CDCl_3) δ up: 191.1, 67.3, 31.7, 29.1, 29.0, 28.7, 27.0, 22.5, 22.2; down: 25.3, 14.0; IR (film) 2926, 2066, 1644, 1367, 1330 cm^{-1} ; EI MS m/z (rel intensity) 168 ($\text{M}^+ - \text{N}_2$, 3), 125 (6), 112 (21), 97 (65), 84 (21), 71 (28), 69 (100), 55 (93); HRMS (calcd for $\text{C}_{11}\text{H}_{20}\text{O}$) 168.1512, found 168.1514.

Methyl (Z)-8-Diazo-9-oxo-5-decenoate (3c): TLC R_f (20% ethyl acetate/petroleum ether) = 0.21; ^1H NMR (250 MHz, CDCl_3) δ 5.60 (m, 1H), 5.43 (m, 1H), 4.13 (q, 2H, J = 7.1 Hz), 3.08 (d, 2H, J = 7.4 Hz), 2.28 (t, 2H, J = 2.8 Hz), 2.25 (s, 3H), 2.12 (m, 2H), 1.70 (m, 2H), 1.26 (t, 3H, J = 7.1 Hz); ^{13}C NMR (62.9 MHz, CDCl_3) δ up: 190.0, 173.4, 66.7, 32.9, 26.1, 24.3, 19.7; down: 132.5, 123.4, 51.1, 25.1; IR (film) 2951, 2071, 1736, 1639, 1330 cm^{-1} ; EI MS m/z (rel intensity) 196 ($\text{M}^+ - \text{N}_2$, 3), 164 (6), 122 (33), 107 (25), 95 (100), 85 (21), 79 (44), 55 (23); HRMS (calcd for $\text{C}_{11}\text{H}_{16}\text{O}_3$) 196.1094, found 196.1099.

(E)-2-Diazo-6,10-dimethylundeca-5,9-dien-2-one (3d): TLC R_f (5% ethyl acetate/petroleum ether) = 0.12; ^1H NMR (250 MHz, CDCl_3) δ 5.20–5.00 (m, 2H), 3.05 (d, 2H, J = 7.5 Hz), 2.24 (s, 3H), 2.05 (m, 4H), 1.68 (s, 3H), 1.64 (s, 3H), 1.60 (s, 3H); ^{13}C NMR (62.9 MHz, CDCl_3) δ up: 190.5, 140.1, 131.6, 67.4, 39.4, 26.3, 20.4; down: 123.6, 117.2, 25.5 (2), 17.5, 15.9; IR (film) 2923,

2069, 1642, 1366, 1326 cm^{-1} ; EI MS m/z (rel intensity) 192 ($\text{M}^+ - \text{N}_2$, 3), 149 (4), 124 (27), 109 (42), 81 (57), 69 (100); HRMS (calcd for $\text{C}_{13}\text{H}_{20}\text{O}$) 192.1509, found 192.1514.

2-Diazo-3-dodecanone (3e): TLC R_f (10% ethyl acetate/petroleum ether) = 0.57; ^1H NMR (250 MHz, CDCl_3) δ 2.46 (t, 2H, J = 7.3 Hz), 1.95 (s, 3H), 1.63 (m, 2H), 1.40–1.10 (m, 12H), 0.88 (t, 3H, J = 6.9 Hz); ^{13}C NMR (62.9 MHz, CDCl_3) δ up: 194.8, 62.0, 37.7, 31.8, 29.6, 29.3 (2), 29.2 (2), 24.8, 22.6; down: 14.0, 8.0; IR (film) 2925, 2067, 1644, 1463, 1283 cm^{-1} ; EI MS m/z (rel intensity) 182 ($\text{M}^+ - \text{N}_2$, 2), 112 (35), 97 (38), 84 (18), 69 (100), 55 (63); HRMS (calcd for $\text{C}_{12}\text{H}_{22}\text{O}$) 182.1672, found 182.1671.

Acknowledgment. We thank the National Institutes of Health (GM 42056) for support for this work.

Supplementary Material Available: ^1H and ^{13}C spectra for compounds **1b**, **2a–e**, and **3b–e** (20 pages). This material is contained in libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS. Ordering information is given on any current masthead page.

JO942059S