

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51352902>

Quantitative Analysis of Pyroglutamic Acid in Peptides

ARTICLE *in* JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY · DECEMBER 1999

Impact Factor: 2.91 · DOI: 10.1021/jf991238d · Source: PubMed

CITATIONS

13

READS

18

3 AUTHORS, INCLUDING:



Yoshio Suzuki

Juntendo University

30 PUBLICATIONS 203 CITATIONS

SEE PROFILE

A SINGLE-STEP PURIFICATION FOR GLYCINE BETAINE DETERMINATION IN PLANT EXTRACTS BY ISOCRATIC HPLC, by Marie-Anne Bessieres,* Yves Gibon, Jean Claude Lefeuvre, and François Larher. *J. Agric. Food Chem.* **1999**, *47*, 3718.

JF991161T

10.1021/jf991161t

Published on Web 11/13/1999

Under HPLC Apparatus and Procedure, anion-exchange should be cation-exchange. In Figure 3, LH-RH is represented by circles and bombesin by triangles. Under Analysis of Bioactive Peptides, Tsunashima should be Tsunasawa, as should be the reference given under Literature Cited. Under Analysis of Industrially Prepared Wheat Gluten Hydrolysate, the correct gluten hydrolysate content is 0.486 mmol/g. The correct page range for the Tsunasawa et al. (1998) reference is 778–783.

JF991238D

10.1021/jf991238d

Published on Web 12/20/1999