

Nitrogen-Concentration Dependence on Photocatalytic Activity of $\text{TiO}_{2-x}\text{N}_x$ Powders

Hiroshi Irie, Yuka Watanabe, and Kazuhito Hashimoto*

Research Center for Advanced Science and Technology, The University of Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo 153-8904, Japan

Received: January 31, 2003

The oxidation power of the $\text{TiO}_{2-x}\text{N}_x$ powders with low nitrogen concentrations (<0.02) was evaluated by the decomposition of gaseous 2-propanol (IPA) under the same absorbed photon number, 1.4×10^{14} quanta $\cdot\text{cm}^{-2}\cdot\text{s}^{-1}$, of visible (Vis) or ultraviolet (UV) light. Regardless of the x value, the quantum yield values from irradiating with Vis light was lower than with UV light, which suggests that the isolated narrow band formed above the valence band is responsible for the Vis light response in the present nitrogen doped TiO_2 . In addition, increasing the nitrogen concentration when irradiating with UV light lowered the quantum yields, indicating that the doping sites could also serve as recombination sites.

1. Introduction

Titanium dioxide (TiO_2) is an efficient photocatalyst^{1–3} but is activated only by irradiating with ultraviolet (UV) light. A current area of research in this field is to modify TiO_2 so that it is sensitive to visible (Vis) light. One approach is to substitute Cr, Fe, or Ni, etc., for a Ti site.^{4–6} Anpo et al.^{5,6} substituted Cr^{3+} or V^{3+} (V^{4+}) at lattice positions of Ti^{4+} in TiO_2 by a metal ion-implantation method. They showed that the absorption band of the Cr^{3+} -doped TiO_2 shifted to the visible light region and that irradiating with Vis light (wavelength > 450 nm) photocatalytically decomposed NO into N_2 , O_2 , and N_2O . Another approach is to form Ti^{3+} sites by introducing an oxygen vacancy in TiO_2 .^{7,8} Ando et al.⁸ reported the “oxygen vacancy-typed visible light sensitive TiO_2 ” in which anatase TiO_2 was fabricated and treated under a reductive plasma (H_2 plasma).

In 1986, Sato⁹ reported that the calcination of NH_4Cl or NH_4OH involving titanium hydroxide caused the photocatalytic sensitization of TiO_2 into the Vis light region. He concluded that the prepared powders were NO_x -doped TiO_2 and that the sensitization of these TiO_2 powders was due to the NO_x impurity. Similarly, Noda et al.¹⁰ prepared yellow-colored anatase TiO_2 powders from a $(\text{NH}_2)_2\cdot\text{H}_2\text{O}$ and TiCl_4 solution and concluded that the Vis light absorption was due to the oxygen vacancies. Recently, several groups reported Vis light sensitive TiO_2 -based powders and thin films.^{11–14} Sakatani et al.¹¹ prepared nitrogen-doped TiO_2 powders by a hydrolytic process, using a TiCl_3 solution and an ammonia solution, and reported that irradiating with Vis light in the presence of the powdered photocatalyst caused the photocatalytic decomposition of acetaldehyde to CO_2 . They concluded that the nitrogen atoms (or ions) were doped at the interstitial sites and that these interstitial nitrogens were responsible for the Vis light response. Ihara et al.¹² also prepared a Vis light sensitive TiO_2 photocatalyst by a hydrolytic process, using a $\text{Ti}(\text{SO}_4)_2$ solution and an ammonia solution. They found that irradiating with Vis light in the presence of the prepared powder caused the photocatalytic decomposition of 2-propanol to CO_2 and asserted that the oxygen vacancies were responsible for the Vis light response.

Sakatani et al. and Ihara et al. insisted on a different conclusion for Vis light response, although they reported similar fabrication methods for Vis light sensitive TiO_2 -based photocatalysts. Asahi et al.^{13,14} theoretically calculated the band structure of the nitrogen-doped TiO_2 and its Vis light induced photocatalysis for powders and thin films. Asahi insisted that the oxygen sites were substituted by nitrogen atoms and that these nitrogens were responsible for the Vis light sensitivity. In addition, Asahi concluded that the Vis light sensitivity of the nitrogen-doped TiO_2 was due to the narrowing of the band gap by mixing the N 2p and O 2p states. We have previously reported¹⁵ that nitrogen-doped TiO_2 powders irradiated with Vis light photocatalytically decomposed CH_3CHO and 2-propanol into CO_2 with different product distributions than irradiating with UV light. These experimental results suggested that the isolated N 2p narrow band that forms above the O 2p valence band was responsible for the visible light response and in the nitrogen-doped TiO_2 the oxygen sites were substituted by nitrogen.

In the present paper, the band structure of the $\text{TiO}_{2-x}\text{N}_x$ with small values of x (<0.02) was experimentally investigated by comparing the quantum efficiency of decomposing gaseous 2-propanol (IPA) under the same absorbed photon number of Vis or UV light.

2. Experimental Section

$\text{TiO}_{2-x}\text{N}_x$ powders were prepared by annealing anatase TiO_2 powder (ST01, Ishihara Sangyo Kaisha Ltd.) under NH_3 flow at 550, 575, and 600 °C for 3 h. TiO_2 powder (ST01) was also annealed under an airflow at 550 °C for 3 h as a reference sample.


An X-ray diffractometer (XRD), X-ray photoemission spectroscopy (XPS), and UV–Vis light absorbance spectroscopy using the diffuse reflection method characterized the samples. The N 1s and O 1s core levels on the surface were measured by XPS in order to investigate the initial states of the NH_3 and air annealed powders. According to Saha et al.,¹⁶ the peak at 396 eV is derived from Ti–N bonds. Therefore, the x values (nitrogen concentrations) were estimated by comparing the product of the 396 eV peak area multiplied by the nitrogen sensitive factor to the product of the 531 eV peak area (O 1s, Ti–O bonds) multiplied by the oxygen sensitive factor.

* To whom correspondence should be addressed. Telephone: +81-3-5452-5080. Fax: +81-3-5452-5083. E-mail: kazuhito@fchem.chem.t.u-tokyo.ac.jp.

TABLE 1: Vis and UV Lights Intensity^a

x in $\text{TiO}_{2-x}\text{N}_x$	Vis light/ ($\text{mW}\cdot\text{cm}^{-2}$)	UV light/ ($\text{mW}\cdot\text{cm}^{-2}$)	x in $\text{TiO}_{2-x}\text{N}_x$	Vis light/ ($\text{mW}\cdot\text{cm}^{-2}$)	UV light/ ($\text{mW}\cdot\text{cm}^{-2}$)
0	0.645*	0.102	0.011	0.304	0.102
0.0050	0.645	0.110	0.019	0.102	0.095

^a Absorbed photon number = 1.0×10^{14} quanta/($\text{cm}^2\cdot\text{s}$) (for the value marked with an asterisk, 0 quanta/($\text{cm}^2\cdot\text{s}$)).

Figure 1. N 1s XPS spectra of $\text{TiO}_{2-x}\text{N}_x$ and TiO_2 powders.

The photocatalytic oxidation activities were evaluated by the decomposition of gaseous IPA when irradiating with Vis (400–530 nm) or UV (300–400 nm) light. The Vis and UV lights were obtained by a Xe lamp (Hayashi Tokei, Luminar Ace 210) with glass filters (Toshiba Glass, B-47, L-42 and C-40C) and by a black fluorescent light bulb (Toshiba Lighting & Technology) with a glass filter (Toshiba Glass, UV-D36B), respectively. Except when TiO_2 was irradiated with Vis light, the Vis or UV light intensity was calibrated for all the other sample powders to absorb 1.4×10^{14} quanta· $\text{cm}^{-2}\cdot\text{s}^{-1}$, as shown in Table 1.

The irradiation area was approximately 4.9 cm^2 . A 300 mg sample of either $\text{TiO}_{2-x}\text{N}_x$ or TiO_2 powders was uniformly spread over the irradiation area in a quartz-made vessel, which had a volume of 500 mL. About 300 ppm of the reactant gas was injected. The samples were kept in the dark once the gas concentration remained constant, which implied that the IPA gas finished adsorbing onto the powder surface. The component gas concentrations were measured using a gas chromatograph (Shimadzu model GC-8A).

3. Results

3.1. Sample Characterization. The powders annealed under NH_3 flow at 550, 575, and 600 °C were pale yellow, yellow, and dark green, respectively. The XRD patterns indicated that all the powders were a homogeneous anatase (TiO_2) phase because only the peaks from the anatase phase and not TiN were observed. The powder annealed under airflow was white and identified as the homogeneous anatase phase.

Figure 1 shows the N 1s XPS spectra of $\text{TiO}_{2-x}\text{N}_x$ and TiO_2 powders. Since peaks at 396 eV, which have previously been found to result from Ti–N bonds,¹⁶ are observed for the powders annealed under NH_3 , it was determined that the oxygen sites were substituted by nitrogen atoms. Since the XRD did not indicate the formation of TiN bonds, it was determined that O–Ti–N bonds formed.¹⁵ Therefore, these powders were described as $\text{TiO}_{2-x}\text{N}_x$. In contrast, the air annealed samples




Figure 2. (a, upper) Optical absorbance spectra of $\text{TiO}_{2-x}\text{N}_x$ and TiO_2 powders. (b, lower) Plots of the square root of the Kubelka–Munk function against the photon energy of $\text{TiO}_{2-x}\text{N}_x$ and TiO_2 powders.




Figure 3. Changes of IPA, acetone, and CO_2 concentrations as a function of time in the presence of $\text{TiO}_{1.995}\text{N}_{0.005}$ powders under Vis light irradiation ($0.645 \text{ mW}\cdot\text{cm}^{-2}$).

TABLE 2: x Values in $\text{TiO}_{2-x}\text{N}_x$

annealing condition	x in $\text{TiO}_{2-x}\text{N}_x$	annealing condition	x in $\text{TiO}_{2-x}\text{N}_x$
air 550 °C	0	NH_3 575 °C	0.011
NH_3 550 °C	0.0050	NH_3 600 °C	0.019

did not display a peak at 396 eV and is TiO_2 . The peak around 400 eV is the chemisorbed N_2 molecules, which adsorb onto the surface.^{16–19} The estimated x values are listed in Table 2.

Figure 2a shows the optical absorbance spectra obtained by the diffuse reflection of the $\text{TiO}_{2-x}\text{N}_x$ and TiO_2 . Noticeable shifts of the absorption shoulders into the Vis light region were observed for $\text{TiO}_{2-x}\text{N}_x$. For $x = 0.019$, the increase in absorbance at wavelengths greater than 550 nm corresponds to Ti^{3+} since NH_3 decomposes into N_2 and H_2 at around 550 °C²⁰ and H_2 is a reducing gas. TiO_2 is an indirect gap semiconductor,²¹ and the band gaps of the $\text{TiO}_{2-x}\text{N}_x$ and TiO_2 can be estimated from the tangent lines in the plots of the square root of the Kubelka–Munk functions against the photon energy,²² as shown in Figure 2b. The tangent lines, which are extrapolated to $Q^{1/2} = 0$, indicate the band gaps of the $\text{TiO}_{2-x}\text{N}_x$ and TiO_2 are $E_g = 3.2 \text{ eV}$, as shown in Figure 2b.

3.2. Photocatalytic Activity. Figure 3 shows the changes of IPA, acetone, and CO_2 concentrations as a function of time in the presence of $\text{TiO}_{1.995}\text{N}_{0.005}$ powders while irradiating with Vis light ($0.645 \text{ mW}\cdot\text{cm}^{-2}$). Both acetone and CO_2 were produced when irradiating with Vis light, but the acetone further decomposed into CO_2 . It was concluded that irradiating with




Figure 4. CO_2 concentration changes for the $\text{TiO}_{2-x}\text{N}_x$ ($x = 0, 0.0050, 0.011, 0.019$) powders under Vis or UV light irradiation.




Figure 5. x value dependence of the quantum efficiency.

Vis light in the presence of the photocatalyst completely decomposed the injected IPA because the evolved CO_2 concentration was 900 ppm, which was three times the initial IPA concentration.

Similar experiments were conducted with the other photocatalysts. Figure 4 shows the CO_2 concentration changes for the $\text{TiO}_{2-x}\text{N}_x$ ($x = 0, 0.0050, 0.011, 0.019$) powders under Vis or UV irradiation, as listed in Table 1. As shown in Figure 4, irradiating TiO_2 ($\text{TiO}_{2-x}\text{N}_x$, $x = 0$) with Vis light did not generate CO_2 as TiO_2 is not Vis light sensitive. CO_2 , however, was evolved from all the other sample powders by irradiating with Vis light. The CO_2 concentration, 900 ppm, indicated that irradiating with Vis light in the presence of the photocatalysts completely decomposed IPA.²³ Irradiating the $\text{TiO}_{2-x}\text{N}_x$ ($x = 0.0050, 0.011, 0.019$) powders with Vis light initially caused a lower CO_2 generation rate. As the x values increased, a longer exposure to Vis light was required to decompose IPA. Irradiating with either UV or Vis light in the presence of $\text{TiO}_{2-x}\text{N}_x$ ($x = 0.0050, 0.011, 0.019$) powders completely decomposed IPA.

Figure 5 shows the x value dependence on the quantum efficiency. Assuming that $\text{C}_3\text{H}_8\text{O} + 5\text{H}_2\text{O} + 18\text{h}^+ \rightarrow 3\text{CO}_2 + 18\text{H}^+$, the quantum efficiency (QY) was calculated²⁴ using the following equation, $\text{QY} = (\text{initial } \text{CO}_2 \text{ generation rate}) / (\text{calculated } \text{CO}_2 \text{ generation rate})$. The absorbed photon numbers were equivalent, $1.4 \times 10^{14} \text{ quanta} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$, for all the samples. The calculated CO_2 generation rate was determined to be $1.9 \times 10^{-10} \text{ mol} \cdot \text{s}^{-1}$. The initial CO_2 generation rate was calculated using the conventional least-squares method and the data in Figure 4. When irradiating with either Vis or UV light, the QY values decreased as x increased. The reason will be discussed later.

4. Discussion

Irradiating $\text{TiO}_{2-x}\text{N}_x$ with UV light resulted in a higher QY value than irradiating with Vis light. This trend is plausible because the TiO_2 oxygen lattice sites substituted by nitrogen atoms form an isolated narrow band above the valence band. Irradiating with UV light excites electrons in both the valence band and the narrow band, but irradiating with Vis light only excites electrons in the narrow band. If substituting nitrogen

for the oxygen sites in TiO_2 narrows the band gap, then the QY values would be the same whether irradiating with Vis or UV light. The observations and Figure 2b do not support the narrowing of the band gap but are consistent with an isolated narrow band above the valence band, which produces the Vis light sensitive photocatalyst. Asahi et al.¹³ calculated the band structure of $\text{TiO}_{2-x}\text{N}_x$, where $x = 0.25$ and 0.12 , i.e., when 12.5 and 6 at. % of the oxygen sites were substituted by nitrogen, and reported that mixing the N 2p and O 2p states narrowed the band gap. In our studies the x value in the $\text{TiO}_{2-x}\text{N}_x$ powders was much lower and at most was 0.02 (1 at. %). It is plausible that the band structure of the $\text{TiO}_{2-x}\text{N}_x$ with lower values of x (<0.02) should differ from higher x values (>0.12).

The QY values of the $\text{TiO}_{2-x}\text{N}_x$ ($x = 0.0050, 0.011, 0.019$) powders when irradiating with Vis light decreased as the x values increased. When annealing TiO_2 powder under an NH_3 atmosphere, the oxygen sites were partially replaced by nitrogen atoms while TiO_2 was simultaneously reduced.²⁰ These changes caused an increase in oxygen vacancy and the amount of Ti^{3+} . The oxygen vacancy state in anatase TiO_2 is below the lower end of the conduction band at 0.75–1.18 eV⁷ and acts as a recombination center for holes and electrons.²⁵ As the NH_3 annealing temperature was increased (with increasing x values), the more TiO_2 reduction was reduced and the greater the number of oxygen vacancies, which explains why the QY values were lower as x increased. In fact, $\text{Ti}_{0.981}\text{N}_{0.019}$ powders had an absorbance greater than 550 nm, which corresponds to Ti^{3+} ¹⁰ (Figure 2a). The same trend was also observed when $\text{TiO}_{2-x}\text{N}_x$ ($x = 0.0050, 0.011, 0.019$) powders were irradiated with UV light because the oxygen vacancies acted as recombination centers for holes and electrons.

5. Conclusion

$\text{TiO}_{2-x}\text{N}_x$ ($x = 0, 0.0050, 0.011, 0.019$) powders were prepared by annealing under an NH_3 flow at 550, 575, and 600 °C. XRD and XPS confirmed that the prepared powders had nitrogen substituted at some of the oxygen sites in TiO_2 , which formed a narrow N 2p band above the valence band. It was determined that the narrow N 2p band was responsible for the visible light sensitivity by considering the QY values, which were calculated by the CO_2 generation rate by decomposing IPA in Vis or UV light at a constant photon number ($1.4 \times 10^{14} \text{ quanta} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$). When irradiating with Vis light, the QY values decreased as x increased. This is because the oxygen vacancies, which promote the recombination of holes and electrons, increased with x . When irradiating with UV light, the QY values decreased as the nitrogen concentration increased, indicating that the doping sites could serve as recombination centers, too.

Acknowledgment. This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas (Grant 417) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of the Japanese Government.

References and Notes

- (1) Heller, A. *Acc. Chem. Res.* **1995**, 28, 141.
- (2) Linsebigler, A. L.; Lu, G. Q.; Yates J. T. *Chem. Rev.* **1995**, 95, 735.
- (3) Fujishima, A.; Hashimoto, K.; Watanabe, T. *TiO₂ Photocatalysis, Fundamentals and Applications*; BKC, Inc.: Tokyo, 1999.
- (4) Borgarello, E.; Kiwi, J.; Gratzel, M.; Pelizzetti, E.; Visca, M. J. *Am. Chem. Soc.* **1982**, 104, 2996.
- (5) Yamashita, H.; Ichihashi, Y.; Takeuchi, M.; Kishiguchi, S.; Anpo, M. J. *Synchrotron Radiat.* **1999**, 6, 451.
- (6) Anpo, M.; Takeuchi, M. *Int. J. Photoenergy* **2001**, 3, 1.

- (7) Nakamura, I.; Negishi, N.; Kutsuna, S.; Ihara, T.; Sugihara, S.; Takeuchi, K. *J. Mol. Catal. A: Chem.* **2000**, *161*, 205.
- (8) Ando, K.; Ando, M.; Fujii, M. *Photocatalysis* **2001**, *6*, 20.
- (9) Sato, S. *Chem. Phys. Lett.* **1986**, *123*, 126.
- (10) Noda, H.; Oikawa, K.; Ogata, T.; Matsuki, K.; Kamata, H. *Chem. Soc. Jpn.* **1986**, *8*, 1084.
- (11) Sakatani, Y.; Koike, H. Japan Patent, P2001-72419A, 2001.
- (12) Ihara, T.; Ando, M.; Sugihara, S. *Photocatalysis* **2001**, *5*, 19.
- (13) Asahi, R.; Morikawa, T.; Ohwaki, T.; Aoki, K.; Taga, Y. *Science* **2001**, *293*, 269.
- (14) Morikawa, T.; Asahi, R.; Ohwaki, T.; Taga, Y. PCT WO 01/10553 A1, 2001.
- (15) Irie, H.; Kako, T.; Nakajima, A.; Watanabe, T.; Hashimoto, K. *J. Photochem. Photobiol. A*, submitted for publication.
- (16) Saha, N. C.; Tompkins, H. G. *J. Appl. Phys.* **1992**, *72*, 3072.
- (17) Fuggle, J. C.; Umbach, E.; Menzel, D.; Wandelt, K.; Brundle, C. R. *Solid State Commun.* **1978**, *27*, 65.
- (18) Wu, H. Z.; Chou, T. C.; Mishra, A.; Anderson, D. R.; Lampert, J. K.; Gujrathi, S. C. *Thin Solid Films* **1990**, *191*, 55.
- (19) Shinn N. D.; Tsang, K. L. *J. Vac. Sci. Technol., A* **1991**, *9* (3), 1558.
- (20) Shin, C. H.; Bugli, G.; Djega-Mariadassou, G.; *J. Solid State Chem.* **1991**, *95*, 145.
- (21) Koffyberg, F. P.; Dwight, K.; Wold, A. *Solid State Commun.* **1979**, *30*, 433.
- (22) Kim, Y. I.; Atherton, S. J.; Brigham, E. S.; Mallouk, T. E. *J. Phys. Chem.* **1993**, *97*, 11802.
- (23) The CO₂ concentration in the presence of TiO_{1.981}N_{0.019} powder is around 300 ppm in Figure 4, but the complete IPA decomposition and the formation of 900 ppm of CO₂ was confirmed.
- (24) Hashimoto, K.; Fujishima, A. *TiO₂ Photocatalysis: Its Application to Self-Cleaning, Anti-bacterial and Air Purifying Materials*; 1998.
- (25) The oxygen vacancy state (Ti 3d impurity level) is not visible light sensitive. In fact, hydrogen-reduced TiO₂ (TiO_{2-x}) powders, which have gray color, are not visible light sensitive. The N 2p narrow band exists much deeper than the Ti 3d donor level, and the N 2p level has a lot more oxidation power than the oxygen vacancy level. Therefore, the origin of visible light sensitivity in TiO_{2-x}N_x is the N 2p narrow band level and not the oxygen vacancy level.