

Excitation Spectrum of the N Intermediate in the Photocycle of Bacteriorhodopsin

Hiroyuki Ohtani* and Osayasu Kikuchi

Department of Biomolecular Engineering, Tokyo Institute of Technology, 4259, Nagatsuta, Midori-ku, Yokohama 226-8501, Japan

Received: May 19, 1999; In Final Form: July 22, 1999

We obtained the absorption spectrum of the N intermediate that plays an important role in the proton uptake during the photocycle of bacteriorhodopsin. The reported spectra of N that were reconstituted from the transient difference absorption spectrum depended on the assumed kinetic models such as sequential $M \rightarrow N \rightarrow O$ reaction and $M \rightleftharpoons N \rightleftharpoons O$ quasi-equilibrium. In this work we measured the excitation spectrum of N by monitoring the characteristic absorption of Q (a photoproduct of N) at 664 nm. The λ_{max} and ϵ_{max} of N were 559 ± 3 nm and $(4.30 \pm 0.75) \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$, respectively. The present spectrum is consistent with the results of the recent SVD analysis. This work provides independent evidence that Q is photochemically formed from N.



1. Introduction

Halobacterium salinarum (*H. salinarum*) is a halophilic archaeobacterium.¹ Under anaerobic conditions it synthesizes a photoactive protein called bacteriorhodopsin (bR) that has a retinal chromophore covalently bound to the ϵ -amino group of Lys-216 via a protonated Schiff base ($-\text{C}=\text{NH}^+-\text{Lys}$). The all-trans isomer of bR (bR₅₆₈, $\lambda_{\text{max}} = 568$ nm) functions as a light-driven proton pump from the cytoplasmic side to the external side.² The proton pump is coupled with a cyclic photoreaction (photocycle) of bR₅₆₈ via five states K, L, M, N, and O.³ The photocycle is initiated by the retinal isomerization from all-trans to 13-cis conformation in the excited state of bR₅₆₈.⁴ The photon energy is stored in the cisoid chromophore and is used for the changes in the conformation and the charge distribution in the protein.

The N state was reported in the earliest photocycle scheme as N₅₂₀ ($\lambda_{\text{max}} = 520$ nm),³ and Hwang et al.⁵ and Shkrob and Rodionov⁶ supported its formation. However, it has been a ghost species that appears only in high pH suspension because it has no visible absorption spectrum remarkably distinguished from bR₅₆₈ though K, L, M, and O have. Kouyama et al.⁷ clarified long-lived intermediates P₅₆₀ and R₃₅₀ that form between M and bR₅₆₈.^{8,9} They attributed the absorption at 560 and 350 nm to the α - and β -bands of N, respectively, and pointed out that the N state plays an important role on the uptake of a proton from the cytoplasmic side.⁷

The chromophore structure of N has been clarified to be a 13-cis, 14-s-trans, C₁₅=N₁₆ anti (trans) protonated Schiff base.¹⁰ The electronic structure of N, however, has not necessarily been established. Some authors reconstituted its spectrum from the measured difference absorption spectra.^{3,7,11,12} The estimation of the concentration of N at delay time t ($[\text{N}]_t$), which is essential to the reconstitution, is difficult even though both the excitation density and the reaction quantum yield were strictly known because of the complicated photocycle of bR₅₆₈. For example, Váró et al.¹³ proposed the following two kinetic models.

Their model (1) requires us to know many rate constants for the estimation of $[\text{N}]_t$, and the model (2), furthermore, requires us to know the equilibrium constant between bR_I and bR_{II}. Thus, the reconstitution of the spectrum of N is not an easy task and



the obtained spectrum is strongly model-dependent. Gergely and Váró¹² recently obtained the model-independent spectrum of N by using the singular value decomposition (SVD) method.

We recently found that the detection of Q (a photoproduct of N^{13,14}) is efficient for the quantitative analysis of N rather than the direct detection of N itself, because Q exhibits an intense red-shifted absorption band and a fluorescence quantum yield much higher than that of N.^{15–17} In this work we measured the excitation spectrum by monitoring the absorbance change (ΔA^N) due to $\text{N} \xrightarrow{h\nu} \text{Q}$. Thus the model-independent absorption spectrum of N was obtained.

2. Experimental Section

The purple membrane (PM) was prepared from *H. salinarum* (ET1-001) by standard procedures.¹⁸ Absorption spectra of the samples were measured with a spectrophotometer (JASCO V-570 equipped with an integration sphere unit V-469) before and after photolysis experiments. An excitation light source (430–700 nm, 250–350 μJ , 5 ns) was an optical parametric oscillator pumped by a Nd:YAG laser (MOPO 700, Spectra-Physics). The excitation intensity (3.5–5.0 mJ/cm²) was set so as not to induce the saturation of the transient absorption due to the formation of the $\text{bR}_{568} \leftrightarrow \text{K}$ photostationary state in the pulse width. A continuous wave (CW) xenon lamp (150 W, Hamamatsu Photonics L2274) was used for a probe light source with appropriate glass filters. The CW light was also used as a background light source for the formation of the photostationary state purple membrane (PSPM) including N. The transmitted probe light was detected with a photomultiplier (Hamamatsu Photonics R3788-02) coupled with a triple monochromator ($f = 257$ mm, Acton Research Corp., SpectraPro-275). Signals




Figure 1. Excitation spectra of bR₅₆₈ and N in an alkaline PSPM monitored by the transient absorption of M (open circles, $\lambda_{\text{probe}} = 408$ nm) and N (solid triangles, $\lambda_{\text{probe}} = 664$ nm), respectively. Conditions: pH 10, 17 μM protein, 22 °C. Key: solid line, absorption spectrum of bR₅₆₈; dashed line, the bR₅₆₈ minus M difference spectrum reported by Fukuda and Kouyama.²⁰

(256–512 shots) were stored and averaged with a digital oscilloscope (Textronics TPS360).

The apparatus was calibrated with a light-adapted PM (LAPM) at pH 7 as standard. The characteristic absorption of M at 408 nm ($\Delta A_{408}^{\text{LA}}(\lambda_{\text{ex}}, t)$) was measured, extrapolated at $t = 0$, and normalized by the photon number of the excitation pulsed light. The photon number at each λ_{ex} was measured by an energy meter (Terahertz Technologies Inc. P-444 pyroelectric probe, DigiRad R-752 Universal laser radiometer) just before and after the measurement. The calibration constant $\alpha(\lambda_{\text{ex}})$ was set so as to make $\Delta A_{408}^{\text{LA}}(\lambda_{\text{ex}}, 0)$ reproduce the molar extinction coefficient of bR₅₆₈ (ϵ^{bR}) at λ_{ex} .

$$\Delta A_{408}^{\text{LA}}(\lambda_{\text{ex}}, 0) = \alpha(\lambda_{\text{ex}}) \epsilon^{\text{bR}}(\lambda_{\text{ex}}) \quad (3)$$

3. Results and Discussion

The measured sample suspension was a PSPM at pH 10. The PSPM was composed of bR₅₆₈ and the stationarily accumulated intermediates M and N with lifetimes elongated by alkalization. The sample suspension and the standard (pure bR₅₆₈, LAPM at pH 7) were alternatively measured at the same λ_{ex} . The absorbance change was normalized by the photon number of the excitation pulsed light for each measurement.

Open circles in Figure 1 show the corrected excitation spectrum of the PSPM monitored at 408 nm ($\Delta A_{408}^{\text{PS}}(\lambda_{\text{ex}}, 0)/\alpha(\lambda_{\text{ex}})$). It is in good agreement with the absorption spectrum of bR₅₆₈ (solid line) in the ≥ 455 nm region. The $\Delta A_{408}^{\text{PS}}(\lambda_{\text{ex}}, 0)/\alpha(\lambda_{\text{ex}})$ in the < 450 nm region was negative because not only bR₅₆₈ but also the accumulated M was excited (see eq 4). The absorbance change at 408 nm due to the excitation of N was neglected.

$$\begin{aligned} \Delta A_{408}^{\text{PS}}(\lambda_{\text{ex}}, 0) &= \Delta A_{408}^{\text{bR/M}}(\lambda_{\text{ex}}, 0) + \Delta A_{408}^{\text{M/Y}}(\lambda_{\text{ex}}, 0) \\ &= (\epsilon_{408}^{\text{M}} - \epsilon_{408}^{\text{bR}})[\text{M}]_{t=0} + (\epsilon_{408}^{\text{Y}} - \epsilon_{408}^{\text{M}})[\text{Y}]_{t=0} \quad (4) \end{aligned}$$

Here, Y denotes the photoproduct of M with a bacteriorhodopsin-like absorption spectrum ($\lambda_{\text{max}} = 570$ nm) which corresponds to P₅₈₅ found at low temperatures.^{14,19} Thus, the term $\Delta A_{408}^{\text{M/Y}}(\lambda_{\text{ex}}, 0)$ in eq 4 gives the negative value because $\epsilon_{408}^{\text{M}}$ is much larger than $\epsilon_{408}^{\text{Y}}$. The excitation spectrum of N in the PSPM was measured by monitoring the characteristic absorption




Figure 2. Kinetics in absorbance change at 664 nm following the pulsed excitation of an alkaline PSPM at pH 10: curve 1, $\lambda_{\text{ex}} = 555$ nm; curve 2, $\lambda_{\text{ex}} = 440$ nm; curves 3 and 4, the absorbance changes due to the recovery of bR₅₆₈ ($\tau = 26$ ms) at $\lambda_{\text{ex}} = 555$ and 440 nm, respectively; curve 5, summation of the absorbance changes due to the decay of a photoproduct of M ($\tau = 17$ ms) and the recovery of bR₅₆₈ at $\lambda_{\text{ex}} = 440$ nm.

of Q at 664 nm.¹⁶ The measured $\Delta A_{664}^{\text{PS}}(\lambda_{\text{ex}}, t)$ was composed of $\Delta A_{664}^{\text{bR/M,N}}(\lambda_{\text{ex}}, t)$, $\Delta A_{664}^{\text{N/Q}}(\lambda_{\text{ex}}, t)$, and $\Delta A_{664}^{\text{M/Y}}(\lambda_{\text{ex}}, t)$ due to the photocycles of bR₅₆₈, N, and M, respectively, given by the following equations.

$$\begin{aligned} \Delta A_{664}^{\text{PS}}(\lambda_{\text{ex}}, t) &= \Delta A_{664}^{\text{bR/M,N}}(\lambda_{\text{ex}}, t) + \Delta A_{664}^{\text{N/Q}}(\lambda_{\text{ex}}, t) + \\ &\quad \Delta A_{664}^{\text{M/Y}}(\lambda_{\text{ex}}, t) \quad (5) \\ &= (\epsilon_{664}^{\text{M}}[\text{M}]_t + \epsilon_{664}^{\text{N}}[\text{N}]_t)I - \epsilon_{664}^{\text{bR}}([\text{M}]_t + \\ &\quad [\text{N}]_t)I + (\epsilon_{664}^{\text{Q}} - \epsilon_{664}^{\text{N}})[\text{Q}]_tI + \\ &\quad (\epsilon_{664}^{\text{Y}} - \epsilon_{664}^{\text{M}})[\text{Y}]_tI \\ &\approx -\epsilon_{664}^{\text{bR}}([\text{M}]_t + [\text{N}]_t)I + \\ &\quad \epsilon_{664}^{\text{Q}}[\text{Q}]_tI + \epsilon_{664}^{\text{Y}}[\text{Y}]_tI \quad (6) \end{aligned}$$

Here, $\epsilon_{664}^{\text{M}}$ and $\epsilon_{664}^{\text{N}}$ are negligibly small.

When λ_{ex} was set in the ≥ 470 nm region (see curve 1 in Figure 2), the measured kinetics $\Delta A_{664}^{\text{PS}}(\lambda_{\text{ex}}, t)$ was composed of a slow bleaching recovery (curve 3) and a fast absorption decay. The former and the latter were attributed to the recovery of bR₅₆₈ and the decay of Q to N,^{15,16} respectively. The absorbance change $\Delta A_{664}^{\text{M/Y}}(\lambda_{\text{ex}}, t)$ was negligibly small in the ≥ 470 nm λ_{ex} region. The net absorbance change due to the formation of Q at 0.5 ms ($\Delta A_{664}^{\text{N/Q}}(\lambda_{\text{ex}}, 0.5\text{ms})$) was obtained by the subtraction of the $\Delta A_{664}^{\text{bR/M,N}}(\lambda_{\text{ex}}, t)$ from the measured absorbance change.

When λ_{ex} was set in the < 470 nm region, the third component appeared in the measured $\Delta A_{664}^{\text{PS}}(\lambda_{\text{ex}}, t)$ (curve 2). The slow absorption decay was attributed to the decay of Y, i.e., $\Delta A_{664}^{\text{M/Y}}(\lambda_{\text{ex}}, t)$. The accumulated M in the PSPM was pumped by the < 470 nm pulsed light. Both the contribution of the third component and the recovery of bR₅₆₈ were subtracted from the measured $\Delta A_{664}^{\text{PS}}(\lambda_{\text{ex}}, t)$.

Thus, we obtained the excitation spectrum of N ($\Delta A_{664}^{\text{N/Q}}(\lambda_{\text{ex}}, 0.5\text{ms})/\alpha(\lambda_{\text{ex}})$) shown by solid triangles in Figure 1. N exhibits a λ_{max} at 559 ± 4 nm. Hereafter we denote N as N₅₅₉. We could reproduce the N₅₅₉ minus bR₅₆₈ difference spectrum reported by Fukuda and Kouyama²⁰ when we set the ϵ_{max} of N₅₅₉ at $43\,050\text{ M}^{-1}\text{ cm}^{-1}$. Thus we obtained the ϵ_{max} of N₅₅₉ ($(4.30 \pm 0.75) \times 10^4\text{ M}^{-1}\text{ cm}^{-1}$). The reported spectra of N distributed around our spectrum (see Figure 3). Our ϵ_{max} is smaller than that reported by Kouyama et al.⁷ ($(4.5\text{--}5.3) \times 10^4$




Figure 3. Absorption spectrum of N: curves 1 and 2, calculated for $\epsilon_{580}^N/\epsilon_{580}^{bR} = 0.78$ and 0.60, respectively, according to the unidirection and no-back-reaction scheme (Kouyama et al.⁷); curve 3, this work; curve 4, calculated according to $M_{II} \rightleftharpoons N \rightleftharpoons O$ scheme by Váró et al.¹¹; curve 5, obtained by using the SVD method (Gergely and Váró¹²).




Figure 4. Absorption spectra of 13-cis pigments and bR₅₆₈: curve 1, N₅₅₉ (this work); curve 2, L₅₄₃ (Shichida et al.²¹); curve 3, bR₅₅₀; curve 4, bR₅₆₈.

$M^{-1} \text{ cm}^{-1}$, $\lambda_{\text{max}} = 555\text{--}560 \text{ nm}$) and is much higher than that reported by Váró et al.¹¹ ($3.3 \times 10^4 M^{-1} \text{ cm}^{-1}$ at 568 nm). Our spectrum resembles the recent one reported by Gergely and Váró¹² ($4.6 \times 10^4 M^{-1} \text{ cm}^{-1}$ at 562 nm). Their spectrum was obtained by using the SVD analysis.

Absorption spectra of 13-cis pigments are shown in Figure 4. The ϵ_{max} of N₅₅₉ is nearly equal to that of L₅₄₃ ($4.3 \times 10^4 M^{-1} \text{ cm}^{-1}$ at 543 nm²¹) and smaller than that of bR₅₅₀ (the 13-cis isomer of bR₅₆₈, $4.8 \times 10^4 M^{-1} \text{ cm}^{-1}$ at 550 nm). Here, the spectrum of bR₅₅₀ was calculated from the spectrum of DAPM according to the results of photocurrent measurement (trans:13-cis = 55:45).²²

The λ_{max} of N₅₅₉ is longer than those of L₅₄₃ and bR₅₅₀. Pigments with an all-trans chromophore show the remarkable

red shift of λ_{max} by the protonation of the carboxyl group of Asp-85. For example, the λ_{max} of O with COOH ($624 \pm 4 \text{ nm}$, H. Ohtani and S. Kanematsu, unpublished) is longer than that of bR₅₆₈ with COO⁻. The red-shifted spectrum of N₅₅₉ is consistent with the previous observations that Asp-85 is protonated in the N₅₅₉ state.^{23,24} The difference in the λ_{max} between L₅₄₃ and bR₅₅₀ may be due to the difference in the retinal conformation, i.e., the distance between the protonated Schiff base and its counteranion (COO⁻ of Asp-85), because L₅₄₃ and bR₅₅₀ have C=N anti and C=N syn conformations, respectively.^{25,26}

Acknowledgment. The authors express their sincere thanks to Professor Masamichi Fujihira, Tokyo Institute of Technology (TIT), for his encouragement throughout this work and fruitful discussions. The authors thank Professor Ichiro Ohkura and Dr. Toshiaki Kamachi, TIT, for their help in the measurements by using an optical parametric oscillator.

References and Notes

- (1) Stoeckenius, W. *Acc. Chem. Res.* **1980**, *13*, 337.
- (2) Lozier, R. H.; Niederberger, W.; Bogomolni, R. A.; Hwang, S.; Stoeckenius, W. *Biochim. Biophys. Acta* **1976**, *440*, 545.
- (3) Lozier, R. H.; Bogomolni, R. A.; Stoeckenius, W. *Biophys. J.* **1975**, *15*, 955.
- (4) Tsuda, M.; Glaccum, M.; Nelson, B.; Ebrey, T. G. *Nature* **1980**, *287*, 351.
- (5) Hwang, S.-B.; Korenbrot, J. I.; Stoeckenius, W. *Biochim. Biophys. Acta* **1978**, *509*, 300.
- (6) Shkrob, M. M.; Rodionov, A. V. *Bioorg. Khim.* **1978**, *4*, 500.
- (7) Kouyama, T.; Nasuda-Kouyama, A.; Ikegami, A.; Mathew, M. K.; Stoeckenius, W. *Biochemistry* **1988**, *27*, 5855.
- (8) Dancsházy, Z.; Govindjee, R.; Nelson, B.; Ebrey, T. G. *FEBS Lett.* **1986**, *209*, 44.
- (9) Drachev, L. A.; Kaulen, A. D.; Skulachev, V. P.; Zorina, V. V. *FEBS Lett.* **1986**, *209*, 316.
- (10) Fodor, S. P. A.; Ames, J. B.; Gebhard, R.; van den Berg, E. M. M.; Stoeckenius, W.; Lugtenburg, J.; Mathies, R. A. *Biochemistry* **1988**, *27*, 7097.
- (11) Váró, G.; Duschul, A.; Lanyi, J. K. *Biochemistry* **1990**, *29*, 3798.
- (12) Gergely, C.; Zimányi, L.; Váró, G. *J. Phys. Chem.* **1997**, *101*, 9390.
- (13) Váró, G.; Lanyi, J. K. *Biochemistry* **1990**, *29*, 2241.
- (14) Balashov, S. P. *Isr. J. Chem.* **1995**, *35*, 415.
- (15) Ohtani, H.; Itoh, H.; Shimura, T. *FEBS Lett.* **1992**, *305*, 6.
- (16) Yamamoto, N.; Naramoto, S.; Ohtani, H. *FEBS Lett.* **1992**, *314*, 345.
- (17) Ohtani, H.; Tsukamoto, Y.; Sakoda, Y.; Hamaguchi, H. *FEBS Lett.* **1995**, *359*, 65.
- (18) Oesterhelt, D.; Stoeckenius, W. In *Methods of Enzymology, Biomembranes Part A*; Freischer, S., Packer, L., Eds.; Academic Press: New York, 1974; Vol. 31, p 667.
- (19) Yamamoto, N. Thesis, Tokyo Institute of Technology, 1994.
- (20) Fukuda, K.; Kouyama, T. *Biochemistry* **1992**, *31*, 11740.
- (21) Shichida, Y.; Matuoka, S.; Hidaka, Y.; Yoshizawa, T. *Biochim. Biophys. Acta* **1983**, *723*, 240.
- (22) Koyama, K. *Photochem. Photobiol.* **1997**, *66*, 784.
- (23) Gerwert, K.; Souvignier, G.; Hess, B. *Proc. Natl. Acad. Sci. U.S.A.* **1990**, *87*, 9774.
- (24) Pfefferlé, J.-M.; Maeda, A.; Sasaki, J.; Yoshizawa, T. *Biochemistry* **1991**, *30*, 6548.
- (25) Smith, S. O.; Myers, A. B.; Pardo, J. A.; Winkel, C.; Mulder, P. J. J.; Lugtenburg, J.; Mathies, R. *Proc. Natl. Acad. Sci. U.S.A.* **1984**, *81*, 2055.
- (26) Fodor, S. P. A.; Pollard, W. T.; Gebhard, R.; van den Berg, E. M. M.; Lugtenburg, J.; Mathies, R. A. *Proc. Natl. Acad. Sci. U.S.A.* **1988**, *85*, 2156.