

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/236045133>

Comparative Antimalarial Activities and ADME Profiles of Ozonides (1,2,4-trioxolanes) OZ277, OZ439, and Their 1,2-Dioxolane, 1,2,4-Trioxane, and 1,2,4,5-Tetraoxane Isosteres

ARTICLE in JOURNAL OF MEDICINAL CHEMISTRY · MARCH 2013

Impact Factor: 5.45 · DOI: 10.1021/jm400004u · Source: PubMed

CITATIONS

20

READS

43

15 AUTHORS, INCLUDING:



Xiaofang Wang

University of Nebraska Medical Center

23 PUBLICATIONS 612 CITATIONS

SEE PROFILE



Susan A Charman

Monash University (Australia)

121 PUBLICATIONS 3,449 CITATIONS

SEE PROFILE



Eileen Ryan

University of Melbourne,

27 PUBLICATIONS 606 CITATIONS

SEE PROFILE



Jessica Steuten

Monash University (Australia)

9 PUBLICATIONS 116 CITATIONS

SEE PROFILE

Comparative Antimalarial Activities and ADME Profiles of Ozonides (1,2,4-trioxolanes) OZ277, OZ439, and Their 1,2-Dioxolane, 1,2,4-Trioxane, and 1,2,4,5-Tetraoxane Isosteres

Xiaofang Wang,[†] Yuxiang Dong,[†] Sergio Wittlin,^{‡,§} Susan A. Charman,^{||} Francis C. K. Chiu,^{||} Jacques Chollet,^{‡,§} Kasiram Katneni,^{||} Janne Mannila,^{||} Julia Morizzi,^{||} Eileen Ryan,^{||} Christian Scheurer,^{‡,§} Jessica Steuten,^{||} Josefina Santo Tomas,^{‡,§} Christopher Snyder,^{‡,§} and Jonathan L. Vennerstrom^{*,†}


[†]College of Pharmacy, University of Nebraska Medical Center, 986025 Nebraska Medical Center, Omaha, Nebraska, United States

[‡]Swiss Tropical and Public Health Institute, Socinstrasse 57, CH-4002 Basel, Switzerland

[§]University of Basel, CH-4003 Basel, Switzerland

^{||}Centre for Drug Candidate Optimisation, Monash Institute of Pharmaceutical Sciences, Monash University (Parkville Campus), 381 Royal Parade, Parkville, Victoria 3052, Australia

ABSTRACT: To ascertain the structure–activity relationship of the core 1,2,4-trioxolane substructure of dispiro ozonides OZ277 and OZ439, we compared the antimalarial activities and ADME profiles of the 1,2-dioxolane, 1,2,4-trioxane, and 1,2,4,5-tetraoxane isosteres. Consistent with previous data, both dioxolanes had very weak antimalarial properties. For the OZ277 series, the trioxane isostere had the best ADME profile, but its overall antimalarial efficacy was not superior to that of the trioxolane or tetraoxane isosteres. For the OZ439 series, there was a good correlation between the antimalarial efficacy and ADME profiles in the rank order trioxolane > trioxane > tetraoxane. As we have previously observed for OZ439 versus OZ277, the OZ439 series peroxides had superior exposure and efficacy in mice compared to the corresponding OZ277 series peroxides.



The discovery of artemisinin from *Artemisia annua* in the early 1970s gave rise to the semisynthetic artemisinins¹ and, more recently, to structurally diverse synthetic peroxide² antimalarials. Although the precise mechanism of action of these drugs is still not fully understood, it is hypothesized that the pharmacophoric peroxide bond³ undergoes reductive activation by heme released during parasite hemoglobin digestion⁴ to produce carbon-centered radicals that alkylate heme and parasite proteins.^{2a,5} The first synthetic peroxide drug development candidates were fenozan B07⁶ and arteflene,⁷ of which the latter progressed to a phase II clinical trial before further development was discontinued⁸ (Figure 1).

In 2012, ozonide OZ277 (**1a**),⁹ also known as arterolane maleate,¹⁰ was approved for the Indian market as a combination product with piperazine phosphate (Synriam) for the treatment of malaria. More recently, the next-generation ozonide OZ439 (**1b**)¹¹ has completed phase IIa clinical trials (Figure 2). In addition to these ozonides (1,2,4-trioxolanes),

1,2,4-trioxane PA1103/SAR116242¹² and 1,2,4,5-tetraoxane RKA 182¹³ were identified as synthetic peroxide drug development candidates.⁸ To ascertain the structure–activity relationship of the core 1,2,4-trioxolane substructure in dispiro 8'-alkyl (**1a**) and 8'-aryl (**1b**) 1,2,4-trioxolanes, we now describe the synthesis and comparative antimalarial activities and ADME profiles of the 1,2-dioxolane (**2**), 1,2,4-trioxane (**3**), and 1,2,4,5-tetraoxane (**4**) isosteres (Figure 2). This study is the first to provide a head-to-head comparison of these four peroxide heterocycles within a common structural framework.

CHEMISTRY

Ozonides **1a** and **1b** were obtained via Griesbaum coozonolysis¹⁴ as previously described.^{2c,9,11a} The key step in the synthesis of 1,2-dioxolanes **2a** and **2b** was fragmentation of triethylsilylperoxy ketals **7** and **8** with SnCl₄¹⁵ to form the corresponding peroxy-carbenium ions that underwent annulation¹⁶ with 2-methyleneadamantane¹⁷ to form dioxolane esters **9** and **10** (Scheme 1). Subsequent hydrolysis followed by either amide or ether bond formation afforded **2a** and **2b**, respectively. Dioxolanes **2a** and **2b** were obtained as single diastereomers and assigned a *cis* configuration based on the stereochemistry observed in similar reactions.¹⁶




Figure 1. Artemisinin and first generation synthetic peroxides.

Received: January 2, 2013

Published: March 15, 2013







Figure 2. Next generation synthetic peroxides and OZ277 and OZ439 isosteres.

Scheme 1^a




^aReagents and conditions: (a) 30% H₂O₂, I₂, MeOH, rt, 24 h; (b) Et₃SiOTf, Et₃N, DMF, 0 °C to rt, 24 h; (c) 2-methyleneadamantane, 1 N SnCl₄ in CH₂Cl₂, -78 to -30 °C, 12 h; (d) 15% aq KOH, EtOH, 60 °C, 20 h, then AcOH; (e) (1) HOSu, EDCI, DMF, rt, 24 h, (2) 1,2-diamino-2-methylpropane, CHCl₃, rt, 4 h, (3) *p*-TsOH, ether; (f) powdered NaOH, (Bu)₄NHSO₄, *N*-(2-chloroethyl)morpholine HCl, CH₃CN, rt to 60 °C, 12 h, then MSA, CH₂Cl₂/ether, rt.

1,2,4-Trioxane 3a was obtained by conversion of trioxane acid 11¹⁸ into the corresponding HOSu active ester followed by amide bond formation (Scheme 2). The synthesis of trioxane 3b (Scheme 2) began with conversion of ketophenol 12 into alkylidene acetate 13 followed by epoxidation and regioselective


perhydrolysis¹⁹ to form β -hydroperoxy alcohol 14, predominately as its trans isomer.¹⁸ Acid-catalyzed condensation of 14 with 2-adamantanone followed by acetate hydrolysis (Scheme 2) yielded trioxane phenol 15, which was alkylated with *N*-(2-chloroethyl)morpholine to afford 3b. 1,2,4,5-Tetraoxanes 4a and 4b (Scheme 3) were obtained by condensation of

Scheme 2^a



^aReagents and conditions: (a) HOSu, EDCI, DMF, rt, 24 h; (b) 1,2-diamino-2-methylpropane, CHCl₃, rt, 5 h, then *p*-TsOH, EtOAc; (c) Ph₃P=CH₂, 4-(4-hydroxyphenyl)cyclohexanone, rt for 1 h, then reflux for 12 h; (d) Ac₂O, pyridine, CH₂Cl₂, 0 °C to rt, 12 h; (e) peracetic acid, CHCl₃, NaOAc, 0 °C to rt, 12 h; (f) 50% H₂O₂, MoO₂(acac)₂, Et₂O, MgSO₄, rt, 12 h; (g) 2-adamantanone, CSA, CH₂Cl₂, rt, 12 h; (h) 15% aq KOH, EtOH/THF, 50 °C, 4 h, then AcOH; (i) see (f) for Scheme 1.

Scheme 3^a




^aReagents and conditions: (a) HOSu, EDCI, 1,2-diamino-2-methylpropane, CH₃CN/CH₂Cl₂, rt, 16 h, then MSA, EtOAc; (b) see (f) for Scheme 1.

tetraoxane acid 16¹³ with the corresponding diamine and *O*-alkylation of tetraoxane phenol 17¹⁸ with *N*-(2-chloroethyl)morpholine, respectively. Peroxides 1a–3a were isolated and tested as their tosylate salts; the remaining peroxides were isolated and tested as their mesylate salts.

■ ANTIMALARIAL ACTIVITY

The activity data shown in Table 1 reveals several SAR trends. Consistent with our previous data¹⁷ demonstrating that 1,2-dioxolanes react with iron(II) primarily by two-electron reduction to form inactive diol reaction products rather than one-electron reduction to form carbon-centered radicals, 2a and 2b were 2 orders of magnitude less potent than the corresponding ozonides 1a and 1b. Erhardt et al.²⁰ demon-

Table 1. Activity of Dispiro Peroxides against *P. falciparum* in Vitro and *P. berghei* in Vivo (1×30 mg/kg po)


compd	X	IC ₅₀ (nM) ^a	K1/NF54	in vivo act. (%) ^b	survival (days)	cures ^c
control	—	—	—	0	6–7	—
1a ^d	O	1.8/1.6	—	99.9	11	0/5
2a	CH ₂	170/440	—	ND ^d	ND	ND
3a	OCH ₂	19/29	—	99.1	15	0/3
4a	OO	0.90/1.7	—	99.5	9	0/3
1b ^d	O	2.8/3.4	—	>99.9	>30	5/5
2b	CH ₂	370/1,200	—	ND	ND	ND
3b	OCH ₂	2.2/2.4	—	99.2	27	3/5
4b	OO	2.9/4.0	—	99.3	15	0/3
AS ^d	—	3.4/4.2	—	92	9	0/5
CQ ^d	—	120/9.9	—	99.9	10	0/5
MF ^d	—	7.2/14	—	99.6	22	0/5

^aMean from $n = 2-3$. Individual measurements differed by less than 50%. ^bIndividual measurements differed by less than 10%. ^cNo detectable parasites at 30 days postinfection. ^dArtesunate (AS), chloroquine (CQ), and mefloquine (MF) data from Vennerstrom et al.⁹ and Charman et al.^{11a} ^dND = not determined.

strated that β -scission reactions that form carbon-centered radicals are accelerated by the adjacent oxygen atom present in ozonides (1,2,4-trioxolanes), but absent in 1,2-dioxolanes. Although the trioxane (3a) and tetraoxane (4a) isosteres of OZ277 (1a) had good potency against *Plasmodium falciparum* in vitro, neither compound was curative in the *P. berghei* model. Similarly, trioxane 3b and tetraoxane 4b were no less potent than OZ439 (1b) in vitro, but of these, only 3b was partially curative in the *Plasmodium berghei* model.

■ PHYSICOCHEMICAL AND ADME PROPERTIES

Additional studies were conducted to explain why the ozonide heterocycle in this particular dispiro structural framework (1a and 1b) appears to have the best overall antimalarial profile.

The predicted physicochemical parameters (Table 2) indicated that OZ277 and its isosteres were more basic and

Table 2. Calculated Physicochemical Properties of Dispiro Peroxides

compd	log P/log D _{pH 7.4}	pK _a	PSA (Å ²)	H-bond donor/acceptor
1a	2.07/0.69	9.05	82.8	3/6
2a	3.21/1.83	9.05	73.6	3/5
3a	2.44/1.05	9.05	82.8	3/6
4a	2.26/0.87	9.05	92.0	3/7
1b	4.63/4.53	6.81	49.4	0/6
2b	5.66/5.56	6.82	40.2	0/5
3b	4.88/4.78	6.82	49.4	0/6
4b	4.81/4.71	6.81	58.6	0/7

less lipophilic than OZ439 and its isosteres. While the OZ277 series would be expected to have better aqueous solubility at physiological pH, the OZ439 series would likely have better permeability properties due to a higher log *D* and lower polar surface area (PSA). However, within the two peroxide series, the differences in these calculated physicochemical properties

provided little insight to account for the divergent antimalarial activities and ADME properties.

Metabolic stability studies in mouse liver microsomes indicated that with the exception of trioxane 3a, which was quite stable, the remaining OZ277 isosteres (1a, 2a, and 4a) exhibited similar metabolic stabilities (Table 3). For OZ439 and its isosteres, trioxolane 1b was marginally more metabolically stable than the remaining isosteres (2b, 3b, and 4b), and similar to that of OZ277 (1a).

Table 3. ADME Properties of Dispiro Peroxides

compd	in vitro CL _{int} (μL/min/mg)/EH ^a	in vitro blood T _{1/2} (h) ^b	in vivo AUC _{0-24h} (nM·h) ^c
1a	17/0.43	0.93	5130 ^d
2a	12/0.35	1.0	ND ^e
3a	<7/<0.2	2.3	22377
4a	14/0.37	0.91	13511
1b	14/0.38	10.4	52726
2b	28/0.55	<i>f</i>	ND
3b	38/0.63	8.6	34419
4b	31/0.58	2.6	18448

^aMouse liver microsomes. ^bMouse blood. ^c30 mg/kg oral dose to mice. ^dAUC to 8 h only as concentrations were below the limit of quantitation at 24 h. ^eND = not determined. ^fNo measurable degradation.

For the OZ277 series, the stability in mouse blood showed the same trend as seen with the microsomes, with trioxane 3a being about 2-fold more stable than 1a, 2a, and 4a (Table 3). Consistent with previous results in human and rat blood,^{11a} OZ439 (1b) was about 10-fold more stable in mouse blood compared to OZ277 (1a). No degradation was detected over the 4 h incubation period for dioxolane 2b, whereas 3b exhibited similar stability to 1b, and 4b was about 4-fold less stable compared to 1b.

In vivo, 3a had the highest area under the curve (AUC) of the OZ277 isosteres, consistent with its better metabolic stability and marginally better blood stability; this trioxane also produced the longest average survival time for the OZ277 series, despite having somewhat weaker activity against *P. falciparum* in vitro (Table 3). As shown in Figure 3, panel A, 3a was also the only compound from the OZ277 series that had plasma concentrations above about 10 ng/mL for more than 24 h. In contrast, concentrations of both 1a and 4a dropped below 1 ng/mL by the 24 h time point. Interestingly, the 2-fold increase in the AUC of 4a vs 1a did not translate into markedly superior in vivo antimalarial efficacy of the former. For the OZ439 series, the rank ordering of AUC values mirrored the in vitro blood stability data with 1b and 3b having similar AUC values, whereas 4b had an AUC that was approximately 2- to 3-fold lower. The rank order of exposure (i.e., 1b > 3b > 4b) was also consistent with the in vivo efficacy results where 1b resulted in the longest survival (>30 days), 3b resulted in an average survival of 27 days, and animals treated with 4b survived for an average of only 15 days. Each of the compounds from the OZ439 series maintained plasma concentrations above about 30 ng/mL for more than 24 h, with concentrations of 1b and 3b being above about 200 ng/mL at the 24 h time point (Figure 3, panel B).




Figure 3. Plasma concentration versus time profiles for (A) OZ277 isosteres and (B) OZ439 isosteres following oral administration of 30 mg/kg to noninfected Swiss outbred mice. Open symbols represent individual concentrations for $n = 2$ mice per time point, and filled symbols represent the average concentrations. Symbols: **1a** and **1b** (circles), **3a** and **3b** (squares), and **4a** and **4b** (triangles). For **1a**, concentrations were below the limit of quantitation of 0.5 ng/mL by the 24 h time point.

SUMMARY AND CONCLUSIONS

In summary, this work confirms that dioxolanes such as **2a** and **2b** have very weak antimalarial properties, which we attribute to their reaction with ferrous iron to form inactive diol reaction products.¹⁷ In the OZ277 series, trioxane **3a** had the best ADME profile, but this was compensated by a somewhat lower antimalarial potency, which may be due in part to subtle differences in the iron(II) and heme reaction profiles of trioxanes, trioxolanes, and tetraoxanes.^{19b,21} For the OZ439 series, there was a good correlation between antimalarial efficacy and in vivo exposure (AUC) with the rank order being trioxolane **1b** > trioxane **3b** > tetraoxane **4b**. As we have noted before,^{11a} the superior exposure profile of 8'-aryl (OZ439) vs 8'-alkyl (OZ277) ozonides translates into superior antimalarial efficacy (as evidenced by increased survival time) of the former. Our data suggest that this same trend also holds true for trioxanes (e.g., **3b** versus **3a**) and tetraoxanes (e.g., **4b** versus **4a**), which is consistent with recent data²² indicating that aryl substituted tetraoxanes may have superior antimalarial efficacies compared to alkyl substituted tetraoxanes such as RKA182.¹³ Finally, we suggest that the inferior blood stability and PK profiles of tetraoxanes **4a** and **4b** versus the corresponding trioxanes **3a** and **3b** results from the second peroxide bond of the former. For example, each of the degenerate conformers of the tetraoxanes has an equatorial peroxide bond with a more accessible LUMO^{4e,19b,21c} that would be expected to react more readily with iron(II) than the corresponding trioxanes which have only one conformer with an equatorial peroxide bond.

EXPERIMENTAL SECTION

General. Melting points are uncorrected. ¹H and ¹³C NMR spectra were recorded on a 500 MHz spectrometer. All chemical shifts are reported in parts per million (ppm) and are relative to internal (CH₃)₄Si (0 ppm) for ¹H and CDCl₃ (77.0 ppm) or DMSO-*d*₆ (39.7 ppm) for ¹³C NMR. Combustion and HPLC analysis confirmed that all target compounds possessed purities ≥95%. As indicated below, starting materials were commercially available or were prepared according to known procedures.

cis-Adamantane-2-spiro-3'-8'-[[(2'-amino-2'-methylpropyl)amino]carbonyl]methyl]-1',2'-dioxaspiro[4.5]decane *p*-Tosylate (2a**).** Step 1. To a solution of I₂ (0.254 g, 1.0

mmol) and 50% H₂O₂ (4.5 mL, 40 mmol) in MeOH^{15b} (50 mL) was added methyl 2-(4-oxocyclohexyl)acetate (**5**) (1.70 g, 10 mmol). The mixture was stirred at rt for 24 h and concentrated to a residue that was partitioned between CH₂Cl₂ (30 mL) and water (30 mL). The aqueous layer was extracted with CH₂Cl₂ (2 × 30 mL). The combined extracts were washed with water and brine, dried over MgSO₄, filtered, and concentrated to afford methyl 2-(4-hydroperoxy-4-methoxycyclohexyl)acetate as a 1:1 mixture of diastereomers (2.15 g, 99%), which was used immediately in the next step: ¹H NMR (CDCl₃) δ 0.92–2.46 (m, 11H), 3.30 (s, 1.5H), 3.34 (s, 1.5H), 3.70 (s, 3H), 7.42 (s, 0.5H), 7.52 (s, 0.5H). **Step 2.** The unpurified methyl 2-(4-hydroperoxy-4-methoxycyclohexyl)acetate (2.15 g, 9.86 mmol) in DMF (100 mL) was treated with Et₃N (4.5 mL, 32 mmol) and Et₃SiOTf (2.54 mL, 12 mmol) at 0 °C. The reaction mixture was stirred at rt for 24 h and then diluted with ice-cold hexane (100 mL) and ice-water (100 mL). The organic layer was separated, and the aqueous layer was extracted with hexane (3 × 100 mL). The extracts were combined, dried over MgSO₄, and concentrated to afford methyl 2-[4-methoxy-4-(triethylsilylperoxy)cyclohexyl]acetate (**7**) as a 1:1 mixture of diastereomers (3.02 g, 92%), which was used immediately in the next step: ¹H NMR (CDCl₃) δ 0.68–0.80 (m, 6H), 0.94–1.08 (m, 9H), 0.84–2.44 (m, 11H), 3.26 (s, 1.5H), 3.29 (s, 1.5H), 3.67 (s, 3H). **Step 3.** To a –78 °C solution of **7** (3.02 g, 9.10 mmol) in CH₂Cl₂ (50 mL) was added 2-methyleneadamantane (0.67 g, 4.53 mmol) followed by 1 M SnCl₄ in CH₂Cl₂ (10 mL, 10 mmol). The resulting mixture was stirred at –78 °C for 30 min and then kept at –30 °C overnight. The reaction mixture was allowed to warm to –3 °C and quenched with ice-water (50 mL). After separation of the organic layer, the aqueous layer was extracted with CH₂Cl₂ (2 × 50 mL). The combined extracts were washed with water (50 mL) and brine (50 mL), dried over MgSO₄, filtered, and concentrated. Purification by silica gel chromatography (0–10% ether in hexane) afforded *cis*-adamantane-2-spiro-3'-8'-[(methoxycarbonyl)methyl]-1',2'-dioxaspiro[4.5]decane (**9**) as a colorless solid (0.60 g, 40%): mp 119–120 °C; ¹H NMR (CDCl₃) δ 1.24–1.36 (m, 2H), 1.44–1.96 (m, 17H), 1.95–2.02 (m, 2H), 2.06–2.14 (m, 2H), 2.13 (s, 2H), 2.20 (d, *J* = 7.5 Hz, 2H), 3.66 (s, 3H); ¹³C NMR (CDCl₃) δ 26.44, 26.99, 29.07, 33.45, 34.93, 35.64, 36.24, 37.21, 41.17, 51.42, 55.47, 84.02, 88.71, 173.47. **Step 4.** To a solution of **9** (0.45 g, 1.35 mmol) in EtOH (20 mL) was added 15% aq KOH solution (2 mL). The resulting mixture was stirred at 60 °C for 20 h, concentrated to ~5 mL, then diluted with water (10 mL), and acidified with acetic acid (5 mL). The precipitate was collected by filtration, washed with cold water, and dried in vacuo at 40 °C to afford *cis*-adamantane-2-spiro-3'-

8'-(carboxymethyl)-1',2'-dioxaspiro[4.5]decane as a colorless solid (0.40 g, 93%): mp 184–185 °C; ^1H NMR (CDCl_3) δ 1.27–1.40 (m, 2H), 1.42–1.96 (m, 17H), 1.95–2.04 (m, 2H), 2.06–2.14 (m, 2H), 2.13 (s, 2H), 2.24 (d, J = 7.5 Hz, 2H), 11.14 (brs, 1H); ^{13}C NMR (125.7 MHz, CDCl_3) δ 26.44, 26.99, 29.00, 33.26, 33.45, 34.90, 35.64, 36.24, 37.20, 41.04, 55.46, 83.98, 88.75, 178.74. **Step 5.** A solution of *cis*-adamantane-2-spiro-3'-8'-(4'-carboxymethyl)-1',2'-dioxaspiro[4.5]decane (0.22 g, 0.69 mmol), HOSu (0.09 g, 0.78 mmol), and EDCI (0.15 g, 0.78 mmol) in DMF (10 mL) was stirred at rt for 24 h. Under ice cooling, the reaction was quenched with water (30 mL). The precipitate was collected by filtration, washed with cold water, and dried in a vacuum oven at 40 °C to afford the dioxolane active ester as a colorless solid (0.28 g, 99%): mp 188–189 °C; ^1H NMR (CDCl_3) δ 1.33–1.98 (m, 19H), 1.98–2.06 (m, 2H), 2.06–2.14 (m, 2H), 2.13 (s, 2H), 2.27 (d, J = 7.0 Hz, 2H), 2.96 (s, 4H). **Step 6.** To a solution of 1,2-diamino-2-methylpropane (0.30 g, 3.41 mmol) in CHCl_3 (10 mL) was added dropwise to the solution of the dioxolane active ester (0.28 g, 0.68 mmol) in CHCl_3 (10 mL). The resulting mixture was stirred at rt for 4 h and then quenched with water (10 mL). After separation of the organic layer, the aqueous layer was extracted with CHCl_3 (2×20 mL). The combined extracts were washed with water (2×10 mL) and brine (10 mL), dried over MgSO_4 , filtered, and concentrated. The residue was dissolved in ether (10 mL), and then the solution of *p*-toluenesulfonic acid monohydrate (0.14 g) in ether (10 mL) was added. The precipitate was collected by filtration to afford **2a** as a colorless solid (0.35 g, 90%): mp 224–225 °C; ^1H NMR (CDCl_3) δ 0.96–1.08 (m, 2H), 1.33 (s, 6H), 1.24–1.94 (m, 21H), 2.06 (s, 2H), 2.02–2.16 (m, 2H), 2.39 (s, 3H), 3.39 (d, J = 6.5 Hz, 2H), 7.25 (d, J = 9.0 Hz, 2H), 7.30 (brs, 3H), 7.66 (d, J = 9.0 Hz, 2H); ^{13}C NMR (CDCl_3) δ 21.35, 23.98, 26.47, 27.02, 28.92, 32.08, 33.21, 33.48, 34.93, 35.65, 36.24, 37.24, 46.23, 55.52, 56.07, 84.00, 88.49, 125.53, 129.44, 140.34, 141.54, 173.26. Anal. Calcd for $\text{C}_{30}\text{H}_{46}\text{N}_2\text{O}_6\text{S}$: C, 64.03; H, 8.24; N, 4.98. Found: C, 64.08; H, 7.99; N, 4.71.

***cis*-Adamantane-2-spiro-3'-8'-[4'-[2'-(4'-morpholinyl)-ethoxy]phenyl]-1',2'-dioxaspiro[4.5]decane Mesylate (2b).** **Step 1.** To a solution of **1**₂ (0.254 g, 1.0 mmol) and 30% H_2O_2 (4.5 mL, 40 mmol) in MeOH^{15b} (50 mL) was added 4-(4-acetoxyphe-nyl)-cyclohexanone (**6**) (2.32 g, 10 mmol). The mixture was stirred at rt for 24 h and concentrated to a residue that was partitioned between CH_2Cl_2 (30 mL) and water (30 mL). The aqueous layer was extracted with CH_2Cl_2 (2×30 mL). The combined extracts were washed with water and brine, dried over MgSO_4 , filtered, and concentrated to afford 4-(4-hydroperoxy-4-methoxycyclohexyl)phenyl acetate as a 3:2 mixture of diastereomers (2.8 g, 100%), which was used immediately in the next step: ^1H NMR (CDCl_3) δ 1.52–2.00 (m, 6H), 2.17–2.34 (m, 5H), 2.48–2.62 (m, 1H), 3.34 (s, 1.2H), 3.36 (s, 1.8H), 6.97–7.04 (m, 2H), 7.20–7.26 (m, 2H), 7.58 (s, 0.4H), 7.62 (s, 0.6H); ^{13}C NMR (CDCl_3) δ 21.12, 30.06, 30.30, 30.78, 31.30, 33.96, 41.28, 43.01, 48.31, 48.56, 104.90, 105.12, 121.31, 121.33, 127.75, 127.76, 143.65, 143.75, 148.86, 169.73. **Step 2.** The unpurified 4-(4-hydroperoxy-4-methoxycyclohexyl)phenyl acetate (2.8 g, 10 mmol) in DMF (100 mL) was treated with Et_3N (4.5 mL, 32 mmol) and Et_3SiOTf (2.54 mL, 12 mmol) at 0 °C. The reaction mixture was stirred at rt for 24 h and then diluted with ice-cold hexane (100 mL) and ice–water (100 mL). The organic layer was separated, and the aqueous layer was extracted with hexane (3×100 mL). The extracts were combined, dried over MgSO_4 , and concentrated. Purification by sg chromatography (0–10% ether in hexane) afforded 4-[4-methoxy-4-(triethylsilylperoxy)cyclohexyl]phenyl acetate (**8**) as a 2:3 mixture of diastereomers (2.18 g, 55%): ^1H NMR (CDCl_3) δ 0.68–0.82 (m, 6H), 0.94–1.07 (m, 9H), 1.42–1.84 (m, 6H), 2.18–2.25 (m, 1.2H), 2.29 (s, 1.8H), 2.29 (s, 1.2H), 2.31–2.38 (m, 0.8H), 2.50–2.59 (m, 1H), 3.32 (s, 1.8H), 3.34 (s, 1.2H), 6.97–7.03 (m, 2H), 7.17–7.25 (m, 2H); ^{13}C NMR (CDCl_3) δ 3.79, 3.80, 5.79, 6.57, 6.75, 6.79, 21.13, 30.21, 30.51, 31.32, 31.71, 43.01, 43.26, 48.15, 48.40, 104.20, 104.35, 121.26, 121.32, 127.68, 127.79, 143.99, 144.17, 148.82, 169.68. **Step 3.** To a –78 °C solution of **8** (8.0 g, 20 mmol) in CH_2Cl_2 (200 mL) was added 2-methyleneadamantane (3.2 g, 21.6 mmol) followed by 1 M SnCl_4 in CH_2Cl_2 (30 mL, 30 mmol). The resulting mixture was stirred at –78 °C for 30 min and then kept at –30 °C overnight. The reaction

mixture was allowed to warm up to –3 °C and quenched with ice–water (100 mL). After separation of the organic layer, the aqueous layer was extracted with CH_2Cl_2 (2×100 mL). The combined extracts were washed with water (100 mL) and brine (100 mL), dried over MgSO_4 , filtered, and concentrated. Crystallization from EtOH afforded *cis*-adamantane-2-spiro-3'-8'-(4'-acetoxyphe-nyl)-1',2'-dioxaspiro[4.5]decane (**10**) as a colorless solid (5.60 g, 70%): mp 149–150 °C; ^1H NMR (CDCl_3) δ 1.52–1.88 (m, 16H), 1.93–1.98 (m, 2H), 2.07–2.16 (m, 4H), 2.18 (s, 2H), 2.28 (s, 3H), 2.45–2.53 (m, 1H), 6.98 (d, J = 8.5 Hz, 2H), 7.19 (d, J = 8.5 Hz, 2H); ^{13}C NMR (CDCl_3) δ 21.14, 26.46, 27.02, 30.52, 33.47, 35.63, 35.67, 36.28, 37.22, 42.83, 55.61, 83.89, 88.79, 121.25, 127.75, 144.49, 148.70, 169.73. **Step 4.** To a solution of **10** (5.60 g, 14.0 mmol) in THF (80 mL) and EtOH (160 mL) was added 15% aq KOH solution (5 mL). The resulting mixture was stirred at 60 °C for 20 h. The solution was concentrated to ~5 mL, and the residue was diluted with water (10 mL) and acidified with acetic acid (5 mL). The precipitate was collected by filtration, washed with cold water, and dried in a vacuum oven at 40 °C to afford *cis*-adamantane-2-spiro-3'-8'-(4'-hydroxyphe-nyl)-1',2'-dioxaspiro[4.5]decane as a colorless solid (4.50 g, 90%): mp 210–211 °C; ^1H NMR (500 MHz, CDCl_3) δ 1.52–1.90 (m, 16H), 1.93–2.00 (m, 2H), 2.06–2.17 (m, 4H), 2.18 (s, 2H), 2.38–2.48 (m, 1H), 4.72 (brs, 1H), 6.75 (d, J = 8.5 Hz, 2H), 7.07 (d, J = 8.5 Hz, 2H); ^{13}C NMR (125.7 MHz, CDCl_3) δ 26.45, 27.01, 30.71, 33.48, 35.67, 35.72, 36.28, 37.22, 42.47, 55.63, 84.03, 88.83, 115.10, 127.83, 139.25, 153.71. **Step 5.** To a solution of *cis*-adamantane-2-spiro-3'-8'-(4'-hydroxyphe-nyl)-1',2'-dioxaspiro[4.5]decane (0.50 g, 1.40 mmol) in dry acetonitrile (80 mL) were added powdered NaOH (0.30 g, 7.50 mmol) and tetrabutylammonium hydrogen sulfate (0.06 g, 0.20 mmol). The mixture was stirred at rt for 30 min before 3-chloropropylamine hydrochloride (0.70 g, 3.75 mmol) was added. The reaction mixture was stirred at 60 °C overnight, cooled to rt, filtered, and washed with CH_2Cl_2 . After the filtrate was concentrated, the residue was dissolved in CH_2Cl_2 , washed with water and brine, dried over MgSO_4 , filtered, and concentrated. The residue was dissolved in CH_2Cl_2 (5 mL), and then a solution of methanesulfonic acid (MSA) (0.11 g) in ether (20 mL) was added. The precipitate was collected by filtration to afford **2b** as a colorless solid (0.41 g, 51%): mp 175–176 °C; ^1H NMR (CDCl_3) δ 1.52–1.87 (m, 16H), 1.92–1.97 (m, 2H), 2.07–2.16 (m, 4H), 2.18 (s, 2H), 2.40–2.50 (m, 1H), 2.83 (s, 3H), 3.02–3.12 (m, 2H), 3.50–3.57 (m, 2H), 3.62–3.70 (m, 2H), 3.97–4.04 (m, 2H), 4.06–4.16 (m, 2H), 4.44–4.51 (m, 2H), 6.82 (d, J = 8.5 Hz, 2H), 7.14 (d, J = 8.5 Hz, 2H), 11.84 (brs, 1H); ^{13}C NMR (CDCl_3) δ 26.45, 27.00, 30.63, 33.47, 35.63, 35.66, 36.28, 37.20, 39.36, 42.45, 52.88, 55.59, 56.79, 62.84, 63.86, 83.90, 88.81, 114.36, 128.02, 140.85, 155.20. Anal. Calcd for $\text{C}_{30}\text{H}_{45}\text{NO}_7\text{S}$: C, 63.92; H, 8.05; N, 2.48. Found: C, 63.69; H, 7.89; N, 2.52.

Adamantane-2-spiro-3'-9'-[[[2'-amino-2'-methylpropyl)-amino]carbonyl]methyl]-1',2',4'-trioxaspiro[5.5]undecane *p*-Tosylate (3a). **Step 1.** A solution of adamantane-2-spiro-3'-9'-(carboxymethyl)-1',2',5'-trioxaspiro[5.5]undecane (**11**)¹⁸ (0.20 g, 0.60 mmol), HOSu (0.08 g, 0.70 mmol), and EDCI (0.14 g, 0.73 mmol) in DMF (10 mL) was stirred at rt for 24 h. The reaction was quenched with water (30 mL) at 0 °C, and the precipitate was collected by filtration, washed with cold water, and dried in vacuo at 40 °C to give the trioxane active ester (0.24 g, 96%) as a colorless solid, which was used directly in the next step: ^1H NMR (CDCl_3) δ 1.00–2.10 (m, 21H), 2.48–2.64 (m, 2H), 2.76–2.90 (m, 4H), 2.95 (brs, 1H), 3.18 (brs, 1H), 3.76 (s, 2H). **Step 2.** To a solution of 1,2-diamino-2-methylpropane (0.20 g, 2.30 mmol) in CHCl_3 (5 mL) was added dropwise to the solution of the trioxane active ester (0.24 g, 0.58 mmol) in CHCl_3 (15 mL). The resulting mixture was stirred at rt for 5 h and then quenched with water (20 mL). After separation of the organic layer, the aqueous layer was extracted with CHCl_3 (2×20 mL). The combined extracts were washed with water (2×20 mL) and brine (20 mL), dried over MgSO_4 , filtered, and concentrated to afford the free base of **3a** (0.18 g) as a colorless solid. The free base of **3a** (0.18 g) was dissolved in EtOAc (20 mL) to which was added a solution of *p*-toluenesulfonic acid monohydrate (0.09 g) in EtOAc (10 mL). The solid was collected and dried in vacuo at 40 °C to afford **3a**

as a colorless solid (0.24 g, 69%): mp 166–167 °C; ^1H NMR (CDCl_3) δ 1.34 (s, 6H), 0.50–2.15 (m, 23H), 2.41 (s, 3H), 2.41 (brs, 1H), 2.90 (brs, 1H), 3.40 (s, 2H), 3.54 (s, 2H), 7.25 (d, J = 8.0 Hz, 2H), 7.33 (t, J = 7.0 Hz, 1H), 7.65 (d, J = 8.0 Hz, 2H), 7.76 (brs, 3H); ^{13}C NMR (CDCl_3): δ 21.39, 23.90, 27.12, 28.51 (br), 31.44 (br), 33.33, 33.58, 36.05 (br), 37.14, 41.97, 46.15, 56.27, 62.45, 77.58, 104.23, 125.50, 129.41, 140.34, 141.78, 172.99. Anal. Calcd for $\text{C}_{30}\text{H}_{46}\text{N}_2\text{O}_7\text{S}$: C, 62.26; H, 8.01; N, 4.84. Found: C, 62.49; H, 7.86; N, 4.77.

Adamantane-2-spiro-3'-9'-[4'-(2'-(4'-morpholinyl)ethoxy)phenyl]-1',2',4'-trioxaspiro[5.5]undecane Mesylate (3b). **Step 1.** To a solution of *n*-BuLi (2.5 M in hexane, 6.3 mL, 15.8 mmol) in dry ether (50 mL) was added methyltriphenylphosphonium bromide (5.7 g, 16.0 mmol). The resulting mixture was stirred at rt for 1 h, and then a solution of 4-(4-hydroxyphenyl)cyclohexanone (**12**) (1.5 g, 7.9 mmol) in dry ether (30 mL) was added dropwise. The mixture was then refluxed overnight. After removal of ether, the residue was partitioned between ether (30 mL) and water (30 mL). The aqueous layer was extracted with ether (2 \times 30 mL). The combined extracts were washed with water and brine, dried over MgSO_4 , filtered, and concentrated. Purification by sg chromatography (0–10% ether in hexane) afforded 4-(4-methylenecyclohexyl)phenol as a colorless solid (0.92 g, 49%): mp 89–90 °C; ^1H NMR (CDCl_3) δ 1.43–1.55 (m, 2H), 1.91–1.99 (m, 2H), 2.12–2.22 (m, 2H), 2.37–2.44 (m, 2H), 2.56–2.66 (m, 1H), 4.63 (s, 1H), 4.67 (m, 2H), 6.73–6.78 (m, 2H), 7.03–7.10 (m, 2H); ^{13}C NMR (CDCl_3) δ 35.15, 35.75, 43.24, 107.28, 115.10, 127.87, 139.28, 148.91, 153.61. **Step 2.** To a solution of 4-(4-methylenecyclohexyl)phenol (0.92 g, 4.65 mmol) in CH_2Cl_2 (10 mL) at 0 °C was added dry pyridine (5 mL) followed by acetic anhydride (2 mL). The resulting mixture was stirred at rt overnight. After removal of the solvents, the residue was partitioned between CH_2Cl_2 (30 mL) and water (30 mL). The aqueous layer was extracted with CH_2Cl_2 (2 \times 30 mL). The combined extracts were washed with 1 M HCl (2 \times 10 mL) and water (10 mL), dried over MgSO_4 , filtered, and concentrated to afford 4-(4-methylenecyclohexyl)phenyl acetate (**13**) as a colorless solid (1.06 g, 99%): ^1H NMR (CDCl_3) δ 1.43–1.60 (m, 2H), 1.91–1.99 (m, 2H), 2.12–2.22 (m, 2H), 2.29 (s, 3H), 2.37–2.44 (m, 2H), 2.56–2.66 (m, 1H), 4.68 (m, 2H), 6.99 (d, J = 8.5 Hz, 2H), 7.20 (d, J = 8.5 Hz, 2H). **Step 3.** Peracetic acid (5.7 g, 32 wt % in aq acetic acid) was mixed with CHCl_3 (25 mL), the aqueous layer was discarded, and the organic layer was dried over MgSO_4 to yield an anhydrous peracetic acid solution, which was added dropwise to an ice cold solution of **13** (1.80 g, 7.83 mmol) in CHCl_3 (15 mL) containing sodium acetate (0.5 g). The reaction mixture was then warmed to rt and stirred overnight. The reaction mixture was washed with water (3 \times 20 mL) and saturated NaHCO_3 (3 \times 20 mL), dried over MgSO_4 , filtered, and concentrated to afford 4-(1-oxaspiro[2.5]octan-6-yl)-phenyl acetate as a pale yellow oil (1.80 g, 93%): ^1H NMR (CDCl_3) δ 1.32–1.50 (m, 2H), 1.76–2.00 (m, 4H), 2.00–2.16 (m, 2H), 2.30 (s, 3H), 2.56–2.66 (m, 1H), 2.70 (s, 2H), 7.01 (d, J = 8.5 Hz, 2H), 7.24 (d, J = 8.5 Hz, 2H). **Step 4.** Anhydrous MgSO_4 (10 g) was added to a mixture of 50% H_2O_2 (10.0 g, 114 mmol) and ether (150 mL) at 0 °C. After stirring for 20 min, the mixture was filtered and the filtrate was added to a mixture of 4-(1-oxaspiro[2.5]octan-6-yl)phenyl acetate (1.80 g, 7.32 mmol) and molybdenyl acetylacetonate (228 mg, 0.7 mmol) in ether (10 mL). After the reaction mixture was stirred overnight at rt, it was washed with water (100 mL) and brine (100 mL), dried over MgSO_4 , filtered, and concentrated to afford 4-[(4-hydroperoxy-4-(hydroxymethyl)cyclohexyl)]phenyl acetate (**14**) as a pale yellow oil (1.46 g, 71%): ^1H NMR (CDCl_3) δ 1.35–2.20 (m, 8H), 2.30 (s, 3H), 2.46–2.56 (m, 1H), 3.91 (s, 2H), 7.01 (d, J = 8.5 Hz, 2H), 7.21 (d, J = 8.5 Hz, 2H), 7.63 (brs, 1H). **Step 5.** 10-Camphorsulfonic acid (CSA) (0.20 g, 0.88 mmol) was added to a mixture of **14** (1.46 g, 5.20 mmol) and 2-adamantanone (1.20 g, 8.00 mmol) in CH_2Cl_2 (50 mL). The reaction mixture was stirred overnight at rt, washed with saturated NaHCO_3 (20 mL), water (20 mL), and brine (20 mL), dried over MgSO_4 , filtered, and concentrated. The crude product was purified by crystallization from 5:1 EtOH:H₂O to afford adamantane-2-spiro-3'-9'-(4'-acetoxyphe-nyl)-1',2',4'-trioxaspiro[5.5]undecane as a colorless solid (0.46 g, 21%): mp 155–156 °C; ^1H NMR (CDCl_3) δ 1.30–2.10 (m, 20H), 2.28 (s,

3H), 2.56–2.66 (m, 1H), 2.61 (brs, 2H), 3.87 (brs, 2H), 7.00 (d, J = 8.0 Hz, 2H), 7.18 (d, J = 8.0 Hz, 2H); ^{13}C NMR (CDCl_3) δ 21.13, 27.12, 28.60 (br), 29.66 (br), 30.55 (br), 33.18 (br), 33.36, 36.00 (br), 37.14, 39.23, 42.96, 62.28, 77.65, 104.40, 121.40, 127.63, 143.37, 148.87, 169.70. **Step 6.** To a solution of adamantane-2-spiro-3'-9'-(4'-acetoxyphe-nyl)-1',2',4'-trioxaspiro[5.5]undecane (0.30 g, 0.73 mmol) in THF (10 mL) and EtOH (20 mL) was added 15% aq KOH (1 mL). The resulting mixture was stirred at 50 °C for 4 h, concentrated to ~5 mL, diluted with water (30 mL), and acidified with acetic acid (2 mL). The precipitate was collected by filtration, washed with cold water, and dried in vacuo at 40 °C to afford adamantane-2-spiro-3'-9'-(4'-hydroxyphenyl)-1',2',4'-trioxaspiro[5.5]undecane (**15**) as a colorless solid (0.24 g, 89%): mp 178–179 °C; ^1H NMR (CDCl_3) δ 1.30–2.20 (m, 20H), 2.51–2.60 (m, 1H), 2.82 (brs, 1H), 2.98 (brs, 1H), 3.89 (brs, 2H), 6.77 (d, J = 8.5 Hz, 2H), 7.06 (d, J = 8.5 Hz, 2H); ^{13}C NMR (CDCl_3) δ 27.13, 30.69 (br), 32.26 (br), 33.18, 33.36, 34.18 (br), 37.15, 42.67, 62.30, 77.75, 104.40, 115.17, 127.74, 138.19, 153.78. **Step 7.** To a solution of **15** (0.10 g, 0.27 mmol) in dry acetonitrile (20 mL) were added powered NaOH (0.20 g, 5.0 mmol) and tetrabutylammonium hydrogen sulfate (0.01 g, 0.03 mmol). The mixture was stirred at rt for 30 min before *N*-(2-chloroethyl)-morpholine hydrochloride (0.14 g, 0.75 mmol) was added. The reaction mixture was stirred at 60 °C overnight, cooled to rt, filtered, and washed with CH_2Cl_2 . After the filtrate was concentrated, the residue was dissolved in CH_2Cl_2 , washed with water and brine, dried over MgSO_4 , filtered, and concentrated. The residue (0.12 g) was dissolved in EtOAc (10 mL) to which was added a solution of MSA (0.02 g) in ether (20 mL). The precipitate was collected by filtration to afford **3b** as a colorless solid (0.13 g, 83%): mp 188–189 °C; ^1H NMR (CDCl_3) δ 1.30–2.14 (m, 20H), 2.53–2.62 (m, 1H), 2.83 (s, 3H), 2.74–2.90 (m, 1H), 2.96 (brs, 1H), 3.02–3.13 (m, 2H), 3.51–3.58 (m, 2H), 3.64–3.70 (m, 2H), 3.89 (brs, 2H), 3.98–4.05 (m, 2H), 4.06–4.16 (m, 2H), 4.47–4.53 (m, 2H), 6.84 (d, J = 8.5 Hz, 2H), 7.13 (d, J = 8.5 Hz, 2H), 11.81 (brs, 1H); ^{13}C NMR (CDCl_3) δ 27.12, 28.46 (br), 29.68 (br), 30.62 (br), 33.36, 34.77 (br), 36.01 (br), 37.13, 39.36, 42.66, 52.94, 56.86, 62.28, 62.88, 63.83, 77.65, 104.40, 114.45, 127.92, 139.70, 155.35. Anal. Calcd for $\text{C}_{30}\text{H}_{45}\text{NO}_8\text{S}$: C, 61.20; H, 7.88; N, 2.38. Found: C, 61.15; H, 7.95; N, 2.36.

Adamantane-2-spiro-3'-9'-[[(2'-amino-2'-methylpropyl)-amino]carbonyl]methyl]-1',2',4',5'-tetraoxaspiro[5.5]undecane Mesylate (4a). To a solution of adamantane-2-spiro-3'-9'-(carboxymethyl)-1',2',4',5'-tetraoxaspiro[5.5]undecane (**16**)¹³ (339 mg, 1 mmol) in CH_3CN (10 mL) and CH_2Cl_2 (10 mL) was added HOSu (127 mg, 1.1 mmol) followed by EDCI (211 mg, 1.1 mmol). After the mixture was stirred at rt for 16 h, 1,2-diamino-2-methylpropane (264 mg, 3 mmol) was added. After stirring for 2 h, the mixture was concentrated, diluted with water (50 mL), and extracted with CH_2Cl_2 (2 \times 20 mL). The combined organic layers were washed with water (30 mL) and brine (20 mL), dried over MgSO_4 , and concentrated. To a solution of the residue in EtOAc (5 mL) was added a solution of MSA (96 mg, 1 mmol) in EtOAc (5 mL). The precipitate was collected by filtration and purified by crystallization from CH_3CN to give **4a** as a colorless solid (243 mg, 48%): mp 150 °C dec; ^1H NMR (CDCl_3) δ 1.18–1.37 (m, 2H), 1.36 (s, 6H), 1.42–2.08 (m, 19H), 2.17 (d, J = 7.3 Hz, 2H), 3.00–3.26 (m, 2H), 11.15 (brs, 1H); ^{13}C NMR (CDCl_3) δ 23.93, 26.97, 26.99, 27.75 (br), 28.36 (br), 28.78 (br), 30.04 (br), 31.07 (br), 33.07, 33.11, 33.44, 34.24 (br), 36.87, 39.86, 42.63, 46.25, 56.14, 107.62, 110.37, 172.80. Anal. Calcd for $\text{C}_{23}\text{H}_{40}\text{N}_2\text{O}_8\text{S}$: C, 54.74; H, 7.99; N, 5.55. Found: C, 54.79; H, 8.01; N, 5.42.

Adamantane-2-spiro-3'-9'-[4'-(2'-(4'-morpholinyl)ethoxy)phenyl]-1',2',4',5'-tetraoxaspiro[5.5]undecane Mesylate (4b). To a solution of adamantane-2-spiro-3'-9'-(4'-hydroxyphenyl)-1',2',4',5'-tetraoxaspiro[5.5]undecane (**17**)¹⁸ (0.5 g, 1.34 mmol) in dry acetonitrile (50 mL) were added powered NaOH (0.30 g, 7.50 mmol) and tetrabutylammonium hydrogen sulfate (0.06 g, 0.20 mmol). The mixture was stirred at rt for 30 min before *N*-(2-chloroethyl) morpholine hydrochloride (0.75 g, 4.03 mmol) was added. The reaction mixture was stirred at 60 °C overnight, cooled to rt, filtered, and washed with CH_2Cl_2 . After the filtrate was

concentrated, the residue was dissolved in CH_2Cl_2 , washed with water and brine, dried over MgSO_4 , filtered, and concentrated. The residue (0.60 g) was dissolved in EtOAc (10 mL) to which was added a solution of MSA (0.11 g) in ether (20 mL). The precipitate was collected by filtration to afford **4b** as a colorless solid (0.45 g, 58%): mp 182–183 °C; ^1H NMR (CDCl_3) δ 1.56–2.10 (m, 20H), 2.53–2.63 (m, 1H), 2.83 (s, 3H), 3.01–3.13 (m, 2H), 3.16–3.36 (m, 2H), 3.48–3.58 (m, 2H), 3.67 (d, J = 12.5 Hz, 2H), 3.95–4.16 (m, 4H), 4.45–4.53 (m, 2H), 6.84 (d, J = 8.5 Hz, 2H), 7.16 (d, J = 7.0 Hz, 2H), 11.83 (brs, 1H); ^{13}C NMR (CDCl_3) δ 27.03, 29.84 (br), 31.90 (br), 33.14, 34.28 (br), 36.92, 39.36, 42.76, 52.93, 56.85, 62.87, 63.84, 107.45, 110.52, 114.43, 128.08, 139.74, 155.39. Anal. Calcd for $\text{C}_{29}\text{H}_{43}\text{NO}_9$: C, 59.88; H, 7.45; N, 2.41. Found: C, 59.97; H, 7.67; N, 2.33.

In Vitro and in Vivo Antimalarial Activity. In vitro and in vivo antimalarial activities^{9,11a,23} were measured using the chloroquine-resistant K1 and chloroquine-sensitive NF54 strains of *P. falciparum*, and *P. berghei*-infected mice, respectively. In vivo data were obtained using single 30 mg/kg oral doses (relative to the free base of each compound) of 1–4 administered in a nonsolubilizing, standard suspension vehicle (SSV) comprising 0.5% w/v carboxymethyl cellulose, 0.5% v/v benzyl alcohol, 0.4% v/v Tween 80, and 0.9% w/v sodium chloride in water, except for **3a**, which was dosed in 3% v/v ethanol and 7% v/v Tween 80 in water. Activity is defined as percent reduction in parasitemia on day 3 postinfection compared to an untreated control group. For example, a compound with an activity of 99.9% is 10-fold more active than one with an activity of 99.0% and 100-fold more active than one with an activity of 90%. Cures in the *P. berghei* model are defined as having no detectable parasites on day 30 postinfection.

Physicochemical Properties. Physicochemical properties ($\log D_{\text{pH } 7.4}$, pK_a and PSA) were calculated using ACD/Laboratories Release 9.0 software (Advanced Chemistry Laboratories, Toronto).

Metabolic Stability. Metabolic stability was assessed in vitro by incubating compounds (1 μM) at 37 °C in mouse liver microsomes (Xenotech, Lenexa, KS) suspended in 0.1 M phosphate buffer (pH 7.4) at a final protein concentration of 0.4 mg/mL. Metabolic reactions were initiated by the addition of an NADPH-regenerating system (1 mg/mL NADP, 1 mg/mL glucose-6-phosphate, 1 U/mL glucose-6-phosphate dehydrogenase) and MgCl_2 (0.67 mg/mL) and were quenched at various time points up to 60 min by the addition of ice-cold acetonitrile. Quenched samples were centrifuged, and the relative loss of parent compound was monitored by LC–MS using a Waters/Micromass Xevo TQ triple quadrupole mass spectrometer coupled to a Waters Acquity UPLC (Waters Corporation, Milford, MA). Chromatography was conducted using a Supelco Ascentis Express RP Amide column (50 \times 2.1 mm, 2.7 μm particle size, Supelco, Bellefonte, PA) operated at 40 °C and compounds were eluted with a methanol–water gradient buffered with 0.05% formic acid and delivered at a flow rate of 0.4 mL/min. Mass spectrometry was conducted in positive electrospray ionization mode with multiple-reaction monitoring using MS/MS parameters optimized for each compound. Concentration versus time data were fitted to an exponential decay function to determine the first order rate constant for substrate depletion, which was then used to calculate the degradation half-life, an in vitro intrinsic clearance value (mL/min/mg microsomal protein), and subsequently a predicted in vivo intrinsic clearance value.²⁴ Predicted in vivo CL_{int} values were converted to predicted in vivo hepatic extraction ratios (E_{H}) using the following equation: $E_{\text{H}} = \text{CL}_{\text{int}}/Q + \text{CL}_{\text{int}}$ where Q is liver blood flow, which was assumed to be 90 mL/min/kg for mice.²⁵

Blood Stability. Blood stability was assessed by spiking compounds into freshly collected mouse blood and incubating at 37 °C. At various time points over a 4 h period, aliquots of blood were removed and quenched immediately by the addition of 0.4 M potassium dichromate (10% v/v) solution followed by snap freezing. Concentrations were assessed by LC–MS using the conditions described above for the microsomal stability studies. Samples were processed by protein precipitation with acetonitrile using a 2:1 volume ratio, and concentrations were determined by comparison to

calibration standards prepared in blank mouse whole blood processed in the same manner.

Mouse Exposure Studies. Animal studies were conducted using established procedures^{9,11a} in accordance with the Australian Code of Practice for the Care and Use of Animals for Scientific Purposes, and the study protocols were reviewed and approved by the Monash Institute of Pharmaceutical Sciences Animal Ethics Committee. Exposure following oral administration to noninfected mice was assessed at a dose of 30 mg/kg using an aqueous formulation vehicle containing 0.5% (w/v) hydroxypropylmethylcellulose, 0.5% (v/v) benzyl alcohol, and 0.4% (v/v) Tween 80. Compounds were administered by oral gavage (3 mL/kg) to 6–8 week old male Swiss outbred mice. An abbreviated blood sampling protocol was used to collect samples at 1, 4, 8, and 24 h postdose (two mice at each time point) with blood sampling either by cheek-bleed or terminal cardiac puncture with a maximum of two samples per mouse. Plasma was immediately separated from erythrocytes and frozen at –20 °C prior to analysis by LC–MS. Chromatography, MS, and sample processing were as described for the metabolic stability, and concentrations were determined by comparison to a calibration curve prepared in mouse plasma. Limits of quantitation were between 0.5 and 1 ng/mL for all analytes in plasma. Area under the plasma concentration–time profiles (AUC) were approximated using WinNonLin version 5.2.1 (Pharsight Corporation, Mountain View, CA).

AUTHOR INFORMATION

Corresponding Author

*Phone: 402.559.5362. E-mail: jvenners@unmc.edu.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This investigation received financial and scientific support from Medicines for Malaria Venture (MMV) and the Nebraska Research Initiative (NRI). We thank the manuscript reviewers for their many useful comments and suggestions.

ABBREVIATIONS USED

CSA, 10-camphorsulfonic acid; EDCI, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride; HOSu, N-hydroxysuccinimide; MSA, methanesulfonic acid; SSV, standard suspending vehicle

REFERENCES

- (1) (a) Haynes, R. K. From artemisinin to new artemisinin antimalarials: biosynthesis, extraction, old and new derivatives, stereochemistry and medicinal chemistry requirements. *Curr. Topics Med. Chem.* **2006**, *6*, 509–537. (b) White, N. J. Qinghaosu (artemisinin): the price of success. *Science* **2008**, *320*, 330–334.
- (2) (a) Jefford, C. W. New developments in synthetic peroxidic drugs as artemisinin mimics. *Drug Discovery Today* **2007**, *12*, 487–95. (b) Muraleedharan, K. M.; Avery, M. A. Progress in the development of peroxide-based anti-parasitic agents. *Drug Discovery Today* **2009**, *14*, 793–803. (c) Tang, Y.; Dong, Y.; Vennerstrom, J. L. Synthetic peroxides as antimalarials. *Med. Res. Rev.* **2004**, *24*, 425–448. (d) Jefford, C. W. Synthetic peroxides as potent antimalarials. News and views. *Curr. Top. Med. Chem.* **2012**, *12*, 373–399.
- (3) Kaiser, M.; Wittlin, S.; Nehrbass-Stuedli, A.; Dong, Y.; Wang, X.; Hemphill, A.; Matile, H.; Brun, R.; Vennerstrom, J. L. Peroxide bond-dependent antiparasitic specificity of artemisinin and OZ277 (RBx11160). *Antimicrob. Agents Chemother.* **2007**, *51*, 2991–2993.
- (4) (a) Crespo, M. D.; Avery, T. D.; Hanssen, E.; Fox, E.; Robinson, T. V.; Valente, P.; Taylor, D. K.; Tilley, L. Artemisinin and a series of novel endoperoxide Antimalarials exert early effects on digestive vacuole morphology. *Antimicrob. Agents Chemother.* **2008**, *52*, 98–109. (b) Elliott, D. A.; McIntosh, M. T.; Hosgood, H. D.; Chen, S.; Zhang, G.; Baevova, P.; Joiner, K. A. Four distinct pathways of hemoglobin

- uptake in the malaria parasite *Plasmodium falciparum*. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 2463–2468. (c) Hartwig, C. L.; Rosenthal, A. S.; D'Angelo, J.; Griffin, C. E.; Posner, G. H.; Cooper, R. A. Accumulation of artemisinin trioxane derivatives within neutral lipids of *Plasmodium falciparum* malaria parasites is endoperoxide-dependent. *Biochem. Pharmacol.* **2009**, *77*, 322–336. (d) Klonis, N.; Crespo-Ortiz, M. P.; Bottova, I.; Abu-Bakar, N.; Kenny, S.; Rosenthal, P. J.; Tilley, L. Artemisinin activity against *Plasmodium falciparum* requires hemoglobin uptake and digestion. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 11405–11410. (e) Hartwig, C. L.; Lauterwasser, E. M.W.; Mahajan, S. S.; Hoke, J. M.; Cooper, R. A.; Renslo, A. R. Investigating the antimalarial action of 1,2,4-trioxolanes with fluorescent chemical probes. *J. Med. Chem.* **2011**, *54*, 8207–8213.
- (5) (a) Meshnick, S. R. Artemisinin: mechanisms of action, resistance and toxicity. *Int. J. Parasitol.* **2002**, *32*, 1655–1660. (b) O'Neill, P. M.; Posner, G. H. A medicinal chemistry perspective on artemisinin and related endoperoxides. *J. Med. Chem.* **2004**, *47*, 2945–2964.
- (6) Jefford, C. W.; Kohmoto, S.; Jaggi, D.; Timari, G.; Rossier, J. C.; Rudaz, M.; Barbuzzi, O.; Gerard, D.; Burger, U.; Kamalaprjia, P.; Mareda, J.; Bernardinelli, G.; Manzanares, I.; Canfield, C. J.; Fleck, S. L.; Robinson, B. L.; Peters, W. Synthesis, structure, and antimalarial activity of some enantiomerically pure, cis-fused cyclopenteno-1,2,4-trioxanes. *Helv. Chim. Acta* **1995**, *78*, 647–662.
- (7) Jaquet, C.; Stohler, H. R.; Chollet, J.; Peters, W. Antimalarial activity of the bicyclic peroxide Ro 42–1611 (arteeflene) in experimental models. *Trop. Med. Parasitol.* **1994**, *45*, 266–271.
- (8) Tilley, L.; Charman, S. A.; Vennerstrom, J. L. Semisynthetic artemisinin and synthetic peroxide antimalarials. In *Neglected Diseases and Drug Discovery*; Palmer, M. J., Wells, T. N. C., Eds.; Royal Society of Chemistry: Cambridge, U.K., 2011; pp 33–64.
- (9) Vennerstrom, J. L.; Arbe-Barnes, S.; Brun, R.; Charman, S. A.; Chiu, F. C.; Chollet, J.; Dong, Y.; Dorn, A.; Hunziker, D.; Matile, H.; McIntosh, K.; Padmanilayam, M.; Santo Tomas, J.; Scheurer, C.; Scoreaux, B.; Tang, Y.; Urwyler, H.; Wittlin, S.; Charman, W. N. Identification of an antimalarial synthetic trioxolane drug development candidate. *Nature* **2004**, *430*, 900–904.
- (10) (a) Gautam, A.; Ahmed, T.; Sharma, P.; Varshney, B.; Kothari, M.; Saha, N.; Roy, A.; Moehrle, J. J.; Paliwal, J. Pharmacokinetics and pharmacodynamics of artefene maleate following multiple oral doses in adult patients with *P. falciparum* malaria. *J. Clin. Pharmacol.* **2011**, *51*, 1519–1528. (b) Olliaro, P.; Wells, T. N. C. The global portfolio of new antimalarial medicines under development. *Clin. Pharmacol. Ther.* **2009**, *85*, 584–595.
- (11) (a) Charman, S. A.; Arbe-Barnes, S.; Bathurst, I. C.; Brun, R.; Campbell, M.; Charman, W. N.; Chiu, F. C.; Chollet, J.; Craft, J. C.; Creek, D. J.; Dong, Y.; Matile, H.; Maurer, M.; Morizzi, J.; Nguyen, T.; Papastogiannidis, P.; Scheurer, C.; Shackleford, D. M.; Sriraghavan, K.; Stingelin, L.; Tang, Y.; Urwyler, H.; Wang, X.; White, K. L.; Wittlin, S.; Zhou, L.; Vennerstrom, J. L. Synthetic ozonide drug candidate OZ439 offers new hope for a single-dose cure of uncomplicated malaria. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 4400–4405. (b) Moehrle, J. J.; Duparc, S.; Siethoff, C.; van Giersbergen, P. L. M.; Craft, J. C.; Arbe-Barnes, S.; Charman, S. A.; Gutierrez, M.; Wittlin, S.; Vennerstrom, J. L. First-in-man safety and pharmacokinetics of synthetic ozonide OZ439 demonstrates an improved exposure profile relative to other peroxide antimalarials. *Br. J. Clin. Pharmacol.* **2013**, *75* (2), 524–537.
- (12) Coslédan, F.; Fraise, L.; Pellet, A.; Guillou, F.; Mordmuller, B.; Kremsner, P. G.; Moreno, A.; Mazier, D.; Maffrand, J. P.; Meunier, B. Selection of a trioxaquine as an antimalarial drug candidate. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 17579–17584.
- (13) O'Neill, P. M.; Amewu, R. K.; Nixon, G. L.; ElGarah, F. B.; Mungthin, M.; Chadwick, J.; Shone, A. E.; Vivas, L.; Lander, H.; Barton, V.; Muangnoicharoen, S.; Bray, P. G.; Davies, J.; Park, B. K.; Wittlin, S.; Brun, R.; Preschel, M.; Zhang, K. S.; Ward, S. A. Identification of a 1,2,4,5-tetraoxane antimalarial drug-development candidate (RKA 182) with superior properties to the semisynthetic artemisinins. *Angew. Chem., Int. Ed.* **2010**, *49*, 5693–5697.
- (14) (a) Griesbaum, K.; Liu, X. J.; Kassiaris, A.; Scherer, M. Ozonolyses of O-alkylated ketoximes in the presence of carbonyl groups: A facile access to ozonides. *Liebigs Ann.-Recl.* **1997**, 1381–1390. (b) Tang, Y.; Dong, Y.; Karle, J. M.; DiTusa, C. A.; Vennerstrom, J. L. Synthesis of tetrasubstituted ozonides by the Griesbaum cozonolysis reaction: Diastereoselectivity and functional group transformations by post-ozonolysis reactions. *J. Org. Chem.* **2004**, *69*, 6470–6473.
- (15) (a) Dussault, P. H.; Zope, U. Hydroperoxide-mediated C-C bond formation—Synthesis of 1,2-dioxolanes from alkoxyhydroperoxides in the presence of Lewis-acids. *Tetrahedron Lett.* **1995**, *36*, 3655–3658. (b) Zmitek, K.; Zupan, M.; Stavber, S.; Iskra, J. The effect of iodine on the peroxidation of carbonyl compounds. *J. Org. Chem.* **2007**, *72*, 6534–6540.
- (16) Ramirez, A.; Woerpel, K. A. Synthesis of 1,2-dioxolanes by annulation reactions of peroxy-carbenium ions with alkenes. *Org. Lett.* **2005**, *7*, 4617–4620.
- (17) Wang, X.; Dong, Y.; Wittlin, S.; Creek, D.; Chollet, J.; Charman, S. A.; Tomas, J. S.; Scheurer, C.; Snyder, C.; Vennerstrom, J. L. Spiro- and dispiro-1,2-dioxolanes: contribution of iron(II)-mediated one-electron vs two-electron reduction to the activity of antimalarial peroxides. *J. Med. Chem.* **2007**, *50*, 5840–5847.
- (18) Wang, X. F.; Zhao, Q. J.; Vargas, M.; Dong, Y. X.; Sriraghavan, K.; Keiser, J.; Vennerstrom, J. L. The activity of dispiro peroxides against *Fasciola hepatica*. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 5320–5323.
- (19) (a) Li, Y.; Hao, H. D.; Wu, Y. K. Facile ring-opening of oxiranes by H_2O_2 catalyzed by phosphomolybdic acid. *Org. Lett.* **2009**, *11*, 2691–2694. (b) Tang, Y.; Dong, Y.; Wang, X.; Sriraghavan, K.; Wood, J. K.; Vennerstrom, J. L. Dispiro-1,2,4-trioxane analogues of a prototype dispiro-1,2,4-trioxolane: mechanistic comparators for artemisinin in the context of reaction pathways with iron(II). *J. Org. Chem.* **2005**, *70*, 5103–5110.
- (20) Erhardt, S.; Macgregor, S. A.; McCullough, K. J.; Savill, K.; Taylor, B. J. Model studies of beta-scission ring-opening reactions of cyclohexyloxy radicals: Application to thermal rearrangements of dispiro-1,2,4-trioxanes. *Org. Lett.* **2007**, *9*, 5569–5572.
- (21) (a) Bousejra-El Garah, F.; Wong, M. H. L.; Amewu, R. K.; Muangnoicharoen, S.; Maggs, J. L.; Stigliani, J. L.; Park, B. K.; Chadwick, J.; Ward, S. A.; O'Neill, P. M. Comparison of the reactivity of antimalarial 1,2,4,5-tetraoxanes with 1,2,4-trioxolanes in the presence of ferrous iron salts, heme, and ferrous iron salts/phosphatidylcholine. *J. Med. Chem.* **2011**, *54*, 6443–6455. (b) Creek, D. J.; Charman, W. N.; Chiu, F. C.; Prankerd, R. J.; Dong, Y.; Vennerstrom, J. L.; Charman, S. A. Relationship between antimalarial activity and heme alkylation for spiro- and dispiro-1,2,4-trioxolane antimalarials. *Antimicrob. Agents Chemother.* **2008**, *52*, 1291–1296. (c) Creek, D. J.; Charman, W. N.; Chiu, F. C.; Prankerd, R. J.; McCullough, K. J.; Dong, Y.; Vennerstrom, J. L.; Charman, S. A. Iron-mediated degradation kinetics of substituted dispiro-1,2,4-trioxolane antimalarials. *J. Pharm. Sci.* **2007**, *96*, 2945–2956. (d) Fernandez, I.; Robert, A. Peroxide bond strength of antimalarial drugs containing an endoperoxide cycle. Relation with biological activity. *Org. Biomol. Chem.* **2011**, *9*, 4098–4107. (e) Wang, X.; Creek, D. J.; Schiaffo, C. E.; Dong, Y.; Chollet, J.; Scheurer, C.; Wittlin, S.; Charman, S. A.; Dussault, P. H.; Wood, J. K.; Vennerstrom, J. L. Spiroadamantyl 1,2,4-trioxolane, 1,2,4-trioxane, and 1,2,4-trioxepane pairs: relationship between peroxide bond iron(II) reactivity, heme alkylation efficiency, and antimalarial activity. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 4542–4545. (f) Sabbani, S.; Stocks, P. A.; Ellis, G. L.; Davies, J.; Hedenstrom, E.; Ward, S. A.; O'Neill, P. M. Piperidine dispiro-1,2,4-trioxane analogues. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 5804–5808.
- (22) Chadwick, J.; Amewu, R. K.; Marti, F.; Bousejra-El Garah, F.; Sharma, R.; Berry, N. G.; Stocks, P. A.; Burrell-Saward, H.; Wittlin, S.; Rottmann, M.; Brun, R.; Taramelli, D.; Parapini, S.; Ward, S. A.; O'Neill, P. M. Antimalarial mannoxanes: Hybrid antimalarial drugs with outstanding oral activity profiles and a potential dual mechanism of action. *ChemMedChem* **2011**, *6*, 1357–1361.
- (23) Desjardins, R. E.; Canfield, C. J.; Haynes, J. D.; Chulay, J. D. Quantitative assessment of antimalarial activity in vitro by a

semiautomated microdilution technique. *Antimicrob. Agents Chemother.* **1979**, *16*, 710–718.

(24) Obach, R. S. Prediction of human clearance of twenty-nine drugs from hepatic microsomal intrinsic clearance data: An examination of in vitro half-life approach and nonspecific binding to microsomes. *Drug Metab. Dispos.* **1999**, *27*, 1350–1359.

(25) Davies, B.; Morris, T. Physiological parameters in laboratory animals and humans. *Pharm. Res.* **1993**, *10*, 1093–1095.