

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6876335>

Composition, Enantiomeric Distribution, and Antibacterial Activity of the Essential Oil of *Achillea ligustica* All. from Corsica

ARTICLE in JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY · AUGUST 2006

Impact Factor: 2.91 · DOI: 10.1021/jf060752y · Source: PubMed

CITATIONS

27

READS

54

5 AUTHORS, INCLUDING:



Jean-Jacques Filippi

University of Nice-Sophia Antipolis

49 PUBLICATIONS 417 CITATIONS

SEE PROFILE



Don Antoine Lanfranchi

University of Strasbourg

27 PUBLICATIONS 323 CITATIONS

SEE PROFILE



Uwe J. Meierhenrich

University of Nice-Sophia Antipolis

177 PUBLICATIONS 1,958 CITATIONS

SEE PROFILE

Composition, Enantiomeric Distribution, and Antibacterial Activity of the Essential Oil of *Achillea ligustica* All. from Corsica

JEAN-JACQUES FILIPPI,^{*,†,‡} DON-ANTOINE LANFRANCHI,[§] SOIZIC PRADO,[#]
NICOLAS BALDOVINI,[†] AND UWE J. MEIERHENRICH[†]

LCMBA, UMR CNRS 6001, Université de Nice-Sophia Antipolis, Parc Valrose,
F-06108 Nice Cedex 2, France; Bioanorganische Chemie 130, Universität Hohenheim,
Garbenstrasse 30, D-70593 Stuttgart, Germany; Laboratoire de Stéréochimie, ECPM,
25 rue Becquerel, F-67008 Strasbourg Cedex 2, France; and Unité de Génétique Moléculaire
Bactérienne, Unité de Chimie Organique, Institut Pasteur, 28 rue du Docteur Roux,
F-75724 Paris Cedex 15, France

The essential oil of *Achillea ligustica* from Corsica was investigated by gas chromatography (GC) and gas chromatography–mass spectrometry (GC-MS). A total of 82 compounds representing 94.0% of the oil were tentatively identified. The main constituents were the camphane derivatives, representing >30% (camphor, 21.3%; borneol, 6.2%; bornyl acetate, 3.5%) of the whole oil, and santolina alcohol (19.3%). The enantiomeric distribution of 8 chiral constituents was determined by GC-MS using two enantioselective stationary phases (DIME- β -CD and Lipodex-E). Racemic santolina alcohol, required for optimization of the enantioselective GC conditions, was prepared by an original two-step synthesis from 2,5-dimethylhexa-2,4-diene. The whole essential oil was tested for its antibacterial activity against a wide range of bacteria using a paper disk method. The results show a promising activity against *Streptomyces* species.

KEYWORDS: *Achillea ligustica*; essential oil composition; enantioselective analysis; antibacterial properties; santolina alcohol

INTRODUCTION

Achillea ligustica is an herbaceous Asteraceae widespread in the Mediterranean area. In Corsica, this plant is found under the vernacular name “erba santa” (holy herb), which refers to its role in the Corsican traditions. Hence, it was used in cataplasms to relieve sprains and insect bites and had also a reputation for stopping hemorrhages (1). This is apparently a nontoxic plant as it has been traditionally added to cakes consumed on Good Friday.

Three studies have been reported on the composition of the essential oil of *A. ligustica*: Maffei et al. (2) found that samples prepared from plants collected in Italy were rich in artemisia ketone and 2,7-dimethyl-4,6-octadien-2-ol, whereas Tzakou et al. (3) showed that the hydrodistillation of *A. ligustica* harvested in Greece furnished essential oils containing linalool and 1,8-cineole as main constituents. Tuberoso et al. (4) recently reported the composition of samples of *A. ligustica* collected in different

sites of Sardinia and showed that the essential oils were rich in santolina alcohol.

In the course of our studies on Mediterranean aromatic plants, we report here on the composition of Corsican *A. ligustica* essential oil, determined by means of gas chromatography (GC) and gas chromatography–mass spectrometry (GC-MS) techniques. Stereochemical configurations of the main components were measured by GC using chiral stationary phases, after oil fractionation. Additionally, the essential oil was also tested for its potential to inhibit *in vitro* the growth of various bacterial strains.

MATERIALS AND METHODS

Plant Material. The aerial parts of wild *A. ligustica* were collected at blooming stage at U Rugnicon, near Ajaccio (Corsica), in May 2004. A voucher specimen (voucher no. B-5071) was deposited in the herbarium of the Museum d'Histoire Naturelle de la ville de Nice, where the taxonomic identification of the plant material was confirmed.

Essential Oil Isolation. Fresh aerial parts (1000 g) were hydro-distilled for 5 h in a Clevenger-type apparatus. The essential oil was obtained as a pale blue liquid with a 0.4% yield (w/w).

GC and GC-MS Analyses. GC and GC-MS analyses on apolar columns were carried out using an Agilent 6890 N gas chromatograph apparatus equipped with a flame ionization detector (FID) and coupled to a quadrupole Agilent 5973 network mass selective detector working

* Address correspondence to this author at the Universität Hohenheim, Bioanorganische Chemie 130, D-70593 Stuttgart, Germany (e-mail jfilippi@uni-hohenheim.de or jfilippi@unice.fr).

[†] Université de Nice-Sophia Antipolis.

[‡] Universität Hohenheim.

[§] Laboratoire de Stéréochimie, ECPM.

[#] Institut Pasteur.

in electron impact (EI) mode at 70 eV (scanning over 35–350 amu range). The gas chromatograph was equipped with two fused silica capillary columns HP-1 (PDMS, 50 m × 0.2 mm i.d., film thickness = 0.33 μm). The analytical parameters (identical for GC and GC-MS analyses unless specified) were the following: The carrier gas was helium at a flow rate of 1 mL/min. The oven temperature was programmed from 60 to 250 °C at 2 °C/min and held isothermal for 40 min. The injector (split mode, ratio 1/100) temperature was 250 °C. The FID temperature was set at 250 °C, and in the GC-MS analyses, the temperatures of the ion source and transfer line were 170 and 280 °C, respectively.

GC-MS analyses on polar column were performed with a Hewlett-Packard 5890 chromatograph coupled with a 5970A mass spectrometer of the same company (EI mode at 70 eV, mass range of 35–400 amu), using the following conditions: fused-silica capillary column, Agilent HP-20M (50 m × 0.2 mm i.d., film thickness = 0.1 μm); injection mode, split 1:30; oven temperature, programmed from 60 to 220 °C at 2 °C/min and then held isothermal for 30 min; carrier gas, helium (0.8 mL/min); injector and transfer line temperatures, 220 and 230 °C, respectively.

The constituents of the essential oil and its fractions were identified by comparison of their mass spectral pattern and retention indices (RI) with those of pure compounds registered in commercial libraries and literature data (5–9) or a laboratory-made database built from authentic compounds.

Enantioselective GC Analysis. Enantioselective GC analyses were performed using an Agilent 6890 gas chromatograph coupled with a 5973 mass selective detector (EI mode at 70 eV, mass range of 35–400 amu). Two different stationary phases were applied for this study. Hydrodex β-6-tBDM (25 m × 0.25 mm i.d.), Macherey-Nagel, Germany, was used with the following conditions: the oven temperature was held isothermal at 40 °C for 5 min and then programmed from 40 to 90 °C at 0.5 °C/min and then from 90 to 200 °C at 2 °C/min and held isothermal at 200 °C for 30 min. The carrier gas was helium, at a flow rate of 1.5 mL/min. The injector (split mode, 1/100) and the transfer line temperatures were 220 and 230 °C, respectively. In the case of β-pinene and sabinene, which presented coelution on the above-mentioned stationary phase, enantiomer separations were performed by using an enantioselective capillary Lipodex-E column (25 m × 0.25 mm i.d.; Macherey-Nagel, Germany) mounted on a HP 5890 gas chromatograph operated under the following conditions: the oven temperature was programmed from 40 to 180 °C at 1 °C/min. The carrier gas was nitrogen with a constant column head pressure of 11.5 psi. The injector (split, 1/100) and detector temperature was set at 200 °C.

NMR. NMR spectra were recorded on a Bruker DRX 500 spectrometer using deuterated chloroform as solvent. The chemical shift values are reported in parts per million with reference to TMS.

Chemicals. Standard compounds of definite enantiomeric purity were used to assess the enantiomeric ratios measured by GC-MS. Some of them were purchased from chemical supply companies: hex-3(Z)-en-1-ol (Acros, 928-96-1), benzaldehyde (Aldrich, B1334), (–)-α-pinene (Aldrich, 305715), (+)-camphene (Aldrich, C301), (+)-sabinene (Robertet, Grasse), (–)-β-pinene (Aldrich, 112089), β-myrcene (Aldrich, M100005), α-terpinene (Aldrich, 223182), p-cymene (Aldrich, C121452), limonene (Aldrich, 18,316-4), 1,8-cineole (Aldrich, C8-060-1), santolina alcohol (Fluka, 84500), γ-terpinene (Acros, 207501000), terpinolene (Fluka, 86484), linalool (Fluka, 62140), (+)-camphor (Aldrich, 857300), isoborneol (Aldrich, I13901), (–)-borneol (Acros, 464-45-9), (–)-terpinen-4-ol (Acros, 20126-76-5), α-terpineol (Acros, 301610250), myrtenol (Aldrich, W343900), eugenol (Aldrich, E51791), β-caryophyllene (Aldrich, W225207), caryophyllene oxide (Fluka, 22076). Chemicals for synthetic use including 2,5-dimethylhexa-2,4-diene, peracetic acid (40%), and vinylmagnesium bromide (1 M THF solution) were purchased from Sigma-Aldrich.

Essential Oil Fractionation. The essential oil (2.75 g) was fractionated by flash chromatography (FC) using pentane with increasing amounts of diethyl ether (0, 5, 10, 20, and 50%) as eluent. Five fractions were obtained, and the third fraction was further fractionated by column

chromatography using 40–63 μm silica gel and 1% AgNO₃ doped silica gel, to provide 200 mg of santolina alcohol, pure enough for NMR identification.

Synthesis of Santolina Alcohol. In a 250 mL round-bottom flask placed in an ice bath was added dropwise 20 mmol of peracetic acid to a well-stirred mixture of 2,5-dimethylhexa-2,4-diene (2.16 g, 19.6 mmol) and Na₂CO₃ (8.34 g) in CH₂Cl₂ (50 mL). The reaction was stirred at room temperature until a negative KI–starch test was achieved. The reaction mixture was then washed with a saturated NaHCO₃ solution and brine. After drying, the solvent was evaporated under vacuum to provide 2.7 g of crude 2,2-dimethyl-3-(2-methylprop-1-enyl)oxirane as a pale yellow oil (the crude product contained mainly the expected epoxide; see Supporting Information Figure S4). This crude product was used directly in the next step: in a two-neck round-bottom flask placed in an ice bath and under magnetic stirring was added dropwise 20 mL of a 1 M THF solution of vinylmagnesium bromide to a well-stirred solution of crude 2,2-dimethyl-3-(2-methylprop-1-enyl)oxirane (1.3 g; ≈8 mmol) in 5 mL of anhydrous diethyl ether under nitrogen atmosphere. The reaction was allowed to stand overnight at room temperature. The reaction mixture was then quenched with a saturated solution of ammonium chloride. The aqueous layer was extracted with diethyl ether, and the organic phases were collected, washed with brine, and dried over magnesium sulfate. The solvent was evaporated under vacuum to provide a crude pale yellow oil, which was purified by column chromatography on silica gel (40–63 μm) to yield 2,5-dimethyl-3-vinylhex-4-en-2-ol (santolina alcohol): ¹H NMR (Figures S1 and S2) δ 1.15 [s, 3H; C(OH)(CH₃)₂], 1.17 [s, 3H; C(OH)(CH₃)₂], 1.65 [d, 3H, J = 1.4 Hz; CH=C(CH₃)₂], 1.75 [d, 1H, J = 1.4 Hz; CH=C(CH₃)₂], 2.98 (dd, 1H, J = 8.6, 9.5 Hz, H₃), 5.00–5.20 [m, 3H, CH=C(CH₃)₂, CH=CH₂], 5.78 (ddd, 1H, J = 8.2, 9.6, 17.8 Hz, CH=CH₂); ¹³C NMR δ 18.41, 26.37, 26.80, 27.12, 54.63, 72.75, 116.54, 122.74, 134.99, 138.07 [¹H and ¹³C NMR spectra are in good agreement with those reported by Bertus et al. (10)]; EI-MS, m/z (%) 41 (18), 43 (26), 55 (12), 59 (100), 67 (11), 77 (8), 79 (16), 81 (81), 95 (8), 96 (67).

Bacterial Strains and Culture Conditions. References strains of *Escherichia coli* (ATCC 11775), *Corynebacterium jeikeium* (ATCC 43734), and *Nocardia asteroides* (ATCC 19247) were obtained from the Collection de l'Institut Pasteur, Paris, France. *Streptomyces coelicolor* (M15), *Streptomyces avidinii* (ATCC 31267), *Streptomyces albus*, *Pseudomonas aeruginosa* (13C3104), *Enterococcus faecalis* (14C1104), and *Staphylococcus aureus* (18C2204) strains were isolated from patients and kindly provided by Philippe Mazodier and Laurent Marsollier (Institut Pasteur). *N. asteroides* and *C. jeikeium* were cultivated for 48 h in brain heart infusion medium (BHI) at 30 and 37 °C, respectively. The other bacteria species were cultivated for 48 h in Mueller Huntington's medium (MH) at 37 and at 30 °C for *Streptomyces*.

Disk Diffusion Assay. The antibacterial activity of the essential oil was evaluated using the standardized filter paper disk (6 mm non-impregnated disk; antibiotica assay disks, grade 2668, Schleicher and Schuell) diffusion method according to the Kirby–Bauer method (11). Briefly, culture suspension of the tested microorganisms (≈10⁶ CFU/mL) was spread on the solid medium plates (50 mL). Filter paper disks were impregnated with 10 μL of serial dilutions in dimethyl sulfoxide (DMSO, Sigma) of the essential oil and placed onto the solid medium plates. The diameter of inhibition was measured after 24 or 48 h of incubation at 30 or 37 °C. Ten microliters of ampicillin (1 mg/mL) (Sigma) and DMSO were used as positive and negative controls, respectively. All assays were performed in triplicate.

RESULTS AND DISCUSSION

GC-MS analyses were performed on the essential oil of *A. ligustica*, as well as on the five fractions obtained after FC of the whole oil, to further identify minor constituents and to separate some coeluting compounds. The chemical composition of the whole essential oil is reported in Table 1.

Among the main constituents, the camphane derivatives were the most abundant, representing 30.0% of the oil (camphor,

Table 1. Chemical Composition of *A. ligustica* Essential Oil

| no. | component ^a | % ^b | RI HP1 ^c | RI HP20M ^c | oil fraction | identification ^d | enantiomeric ratio ^e |
|-----|--|------------------|---------------------|-----------------------|--------------|-----------------------------|--|
| 1 | hex-3(Z)-en-1-ol | 0.1 | 833 | 1347 | oil, F5 | RI, MS, Std | |
| 2 | non-1-ene ^f | tr | 886 | | F1 | RI, MS | |
| 3 | santolinatriene ^f | 1.3 | 902 | 1027 | oil, F1 | RI, MS | |
| 4 | artemisiatriene ^f | tr | 918 | | F1 | RI, MS | |
| 5 | tricyclene ^f | 0.1 | 921 | | oil, F1 | RI, MS | |
| 6 | α -thujene ^f | tr | 923 | | oil, F1 | RI, MS | |
| 7 | benzaldehyde | tr | 927 | 1471 | oil, F4 | RI, MS, Std | |
| 8 | α -pinene | 1.1 | 931 | 1002 | oil, F1 | RI, MS, Std | (1S,5S)-(-) 41:59 (1R,5R)-(+) ^g |
| 9 | camphene | 2.5 | 944 | 1043 | oil, F1 | RI, MS, Std | (1R)-(+)- 78:22 (1S)-(-) ^g |
| 10 | sabinene | 0.3 | 965 | 1099 | oil, F1 | RI, MS, Std | (1R,5R)-(+)- 89:11 (1S,5S)-(-) ^h |
| 11 | β -pinene | 3.2 | 971 | 1083 | oil, F1 | RI, MS, Std | (1S,5S)-(-) 98:2 (1R,5R)-(+) ^h |
| 12 | β -myrcene | tr | 981 | | F1 | RI, MS, Std | |
| 13 | yomogi alcohol ^f | 1.2 | 984 | 1377 | oil, F3, F4 | RI, MS | |
| 14 | isobutyl 2-methylbutyrate ^f | tr | 988 | | F2 | RI, MS | |
| 15 | isobutyl isovalerate ^f | tr | 990 | | F2 | RI, MS | |
| 16 | α -terpinene | 0.3 | 1010 | 1153 | oil, F1 | RI, MS, Std | |
| 17 | <i>p</i> -cymene | 1.0 | 1012 | 1240 | oil, F1 | RI, MS, Std | |
| 18 | limonene | tr | 1019 | 1168 | F1 | RI, MS, Std | |
| 19 | 1,8-cineole | 1.0 | 1020 | 1176 | oil, F2 | RI, MS, Std | |
| 20 | santolina alcohol | 19.3 | 1024 | 1375 | oil, F3, F4 | RI, MS, NMR, Std | (3S)-(+)- 100:0 (3R)-(-) ^g |
| 21 | artemisia ketone ^f | 5.9 | 1044 | 1320 | oil, F2 | RI, MS | |
| 22 | γ -terpinene | 0.9 | 1049 | 1219 | oil, F1 | RI, MS, Std | |
| 23 | <i>trans</i> -sabinene hydrate ^f | 0.4 | 1053 | 1425 | oil, F3 | RI, MS | |
| 24 | artemisia alcohol ^f | 0.5 | 1069 | 1476 | oil, F3 | RI, MS | |
| 25 | terpinolene | 0.2 | 1079 | 1253 | oil, F1 | RI, MS, Std | |
| 26 | filifolone ^f | 0.2 | 1082 | 1395 | oil, F2 | RI, MS | |
| 27 | linalool | 1.4 | 1084 | 1515 | oil, F3, F4 | RI, MS, Std | |
| 28 | α -thujone ^f | 1.9 | 1086 | 1377 | oil, F2 | RI, MS | |
| 29 | 2-methylbutyl 2-methylbutyrate ^f | 0.1 | 1090 | 1260 | oil, F2 | RI, MS | |
| 30 | <i>iso</i> -amyl isovalerate ^f | 0.1 | 1093 | 1277 | oil, F2 | RI, MS | |
| 31 | β -thujone ^f | 0.3 | 1097 | 1395 | oil, F2 | RI, MS | |
| 32 | chrysanthemone ^f | 0.4 | 1099 | 1458 | oil, F2 | RI, MS | |
| 33 | 4-acetyl-1-methylcyclohex-1-ene ^f | 0.5 | 1105 | 1504 | oil, F2 | RI, MS | |
| 34 | <i>p</i> -menth-2-en-1-ol ^f | 0.1 | 1108 | | oil, F5 | RI, MS | |
| 35 | camphor | 21.3 | 1123 | 1464 | oil, F2, F3 | RI, MS, Std | (1S,4S)-(-) 40:60 (1R,4R)-(+) ^g |
| 36 | <i>trans</i> -pinocarveol ^f | 0.3 | 1124 | 1602 | oil, F4 | RI, MS | |
| 37 | camphene hydrate ^f | tr | 1129 | 1545 | F4 | RI, MS | |
| 38 | pinocarvone ^f | 0.1 | 1135 | 1514 | oil, F2 | RI, MS | |
| 39 | isoborneol | tr | 1137 | | F4 | RI, MS, Std | |
| 40 | <i>cis</i> -chrysanthemol ^f | 0.2 | 1146 | 1642 | oil, F4, F5 | RI, MS | |
| 41 | lavandulol ^f | tr ⁱ | | 1638 | F4 | RI, MS | |
| 42 | borneol | 6.2 ^j | 1149 | 1652 | oil, F4 | RI, MS, Std | (1S,2R,4S)-(-) 73:27 (1R,2S,4R)-(+) ^g |
| 43 | <i>p</i> -cymen-8-ol ^f | tr | 1157 | 1796 | oil, F5 | RI, MS | |
| 44 | terpinen-4-ol | 2.8 | 1162 | 1555 | oil, F3, F4 | RI, MS, Std | (4S)-(+)- 56:44 (4R)-(-) ^g |
| 45 | artemisyl acetate ^f | 0.2 | 1164 | 1331 | oil, F2 | RI, MS | |
| 46 | myrtenal ^f | 0.2 | 1168 | 1570 | oil, F2 | RI, MS | |
| 47 | α -terpineol | 0.6 | 1171 | 1650 | oil, F4, F5 | RI, MS, Std | |
| 48 | myrtenol | 0.3 | 1177 | 1742 | oil, F4 | RI, MS, Std | |
| 49 | <i>trans</i> -carveol ^f | 0.3 | 1196 | 1785 | oil, F4 | RI, MS | |
| 50 | <i>cis</i> -carveol ^f | 0.1 | 1207 | | oil, F4 | RI, MS | |
| 51 | carvone ^f | 0.2 | 1212 | 1773 | oil, F2 | RI, MS | |
| 52 | <i>cis</i> -chrysanthemyl acetate ^f | 0.9 | 1244 | 1533 | oil, F2 | RI, MS | |
| 53 | bornyl acetate ^f | 3.5 | 1269 | 1540 | oil, F2 | RI, MS | |
| 54 | <i>trans</i> -sabinyl acetate ^f | 0.2 ^j | 1272 | 1609 | oil, F2 | RI, MS | |
| 55 | lavandulyl acetate ^f | 0.3 ^j | | 1576 | oil, F2 | RI, MS | |
| 56 | unknown 1 ^k | 0.5 | 1285 | 1629 | oil, F2 | | |
| 57 | <i>trans</i> -carvyl acetate ^f | 0.4 | 1314 | 1695 | oil, F2 | RI, MS | |
| 58 | eugenol | 0.3 | 1326 | 2100 | oil, F4 | RI, MS, Std | |
| 59 | <i>cis</i> -carvyl acetate ^f | 0.3 | 1340 | 1727 | oil, F2 | RI, MS | |
| 60 | benzyl isovalerate ^f | 0.1 | 1362 | 1844 | oil, F2 | RI, MS | |
| 61 | α -copaene ^f | 0.1 | 1372 | 1459 | oil, F1 | RI, MS | |
| 62 | β -caryophyllene | 0.3 | 1413 | 1555 | oil, F1 | RI, MS, Std | |
| 63 | α -humulene ^f | tr ⁱ | 1442 | 1622 | F1 | RI, MS | |
| 64 | β -farnesene ^f | 0.1 ⁱ | 1444 | 1641 | oil, F1 | RI, MS | |
| 65 | alloaromadendrene ^f | 0.2 ^j | 1446 | 1559 | oil, F1 | RI, MS | |
| 66 | γ -muurolene ^f | 0.1 ⁱ | | 1648 | F1 | RI, MS | |
| 67 | <i>ar</i> -curcumene ^f | 0.2 | 1465 | 1738 | oil, F1 | RI, MS | |
| 68 | germacrene-D ^f | 3.1 | 1473 | 1667 | oil, F1 | RI, MS | |
| 69 | bicyclogermacrene ^f | 0.2 ^j | | 1687 | oil, F1 | RI, MS | |
| 70 | ledene ^f | 0.1 ⁱ | 1488 | 1653 | F1 | RI, MS | |
| 71 | α -muurolene ^f | 0.1 | 1490 | 1749 | oil, F1 | RI, MS | |
| 72 | δ -cadinene ^f | 0.2 ^j | | 1718 | oil, F1 | RI, MS | |
| 73 | unknown 2 ^{l,j} | 0.1 ⁱ | 1508 | 1770 | oil, F2 | | |
| 74 | α -calacorene ^f | 0.3 | 1524 | 1904 | oil, F1 | RI, MS | |
| 75 | palustrol ^f | 0.2 | 1555 | 1870 | oil, F2 | RI, MS | |

Table 1 (Continued)

| no. | component ^a | % ^b | RI HP1 ^c | RI HP20M ^c | oil fraction | identification ^d | enantiomeric ratio ^e |
|-----|-------------------------------------|----------------|---------------------|-----------------------|--------------|-----------------------------|---------------------------------|
| 76 | spathulenol ^f | 0.1 | 1557 | 2061 | oil, F4 | RI, MS | |
| 77 | caryophyllene oxide | 0.3 | 1563 | 1917 | oil, F2 | RI, MS, Std | |
| 78 | viridiflorol ^f | 3.2 | 1576 | 2026 | oil, F3, F4 | RI, MS | |
| 79 | ledol ^f | 0.3 | 1586 | 1968 | oil, F4 | RI, MS | |
| 80 | zingiberenol ^f | 0.2 | 1593 | 2062 | oil, F4 | RI, MS | |
| 81 | unknown 3 ^{l,m} | 0.6 | 1601 | 2096 | oil, F3, F4 | | |
| 82 | 1- <i>epi</i> -cubenol ^f | 0.8 | 1608 | 2006 | oil, F2, F3 | RI, MS | |
| 83 | β -eudesmol ^f | 0.4 | 1627 | 2162 | oil, F5 | RI, MS | |
| 84 | cadalene ^f | tr | 1645 | 2157 | F1 | RI, MS | |
| 85 | α -bisabolol ^f | 0.4 | 1662 | 2168 | oil, F4 | RI, MS | |
| | identified compounds | 82 | 94.0 | | | | |
| | monoterpene hydrocarbons | 14 | 10.9 | | | | |
| | oxygenated monoterpenes | 35 | 71.0 | | | | |
| | sesquiterpene hydrocarbons | 14 | 5.1 | | | | |
| | oxygenated sesquiterpenes | 9 | 5.9 | | | | |
| | others | 10 | 1.2 | | | | |

^a Compounds are listed in order of their elution from an HP1 column. ^b FID percentage. tr = trace (<0.1%), trace levels of unidentified compounds are not mentioned. ^c RI = retention indices as determined on HP1 and HP20M columns using the homologous series of C₅–₂₆ *n*-alkanes. ^d Method of identification: MS, by comparison of the mass spectrum with those of the computer mass libraries; RI, by comparison of RI with those from the literature; Std, by injection of an authentic sample (present in our laboratory-made database); NMR, after isolation of the compound by column chromatography and spectral analysis. ^e Determined by enantioselective gas chromatography (see Materials and Methods for conditions). ^f Compound tentatively identified according to its mass spectrum and by comparison of its retention indices with those of the literature. ^g Determined on heptakis(2,3-*O*,*O*-dimethyl-6-*O*-*tert*-butyldimethylsilyl)- β -cyclodextrin (Hydrodex β -6TBDM). ^h Determined on octakis(3-*O*-butyryl-2,6-*O*,*O*-dipentyl)- γ -cyclodextrin (Lipodex E; Figure S3). ⁱ Percentage determined after analysis on HP20M column. ^j MS spectra of unknown compounds (see Supporting Information, Figure S5, for full spectra). ^k Unknown 1, *m/z* (%): 41 (28.7), 43 (100), 67 (10.5), 69 (23.9), 79 (15.4), 81 (17.6), 91 (17.6), 109 (95.4), 134 (33.3), 151 (27.3). ^l Unknown 2, *m/z* (%): 41 (51.0), 43 (91.2), 81 (65.0), 91 (75.1), 93 (48.7), 105 (100), 106 (71.5), 119 (45.5), 147 (45.6), 220 (96.2). ^m Unknown 3, *m/z* (%): 41 (46.9), 43 (47.2), 79 (44.2), 91 (54.2), 93 (37.4), 97 (52.2), 105 (41.2), 109 (39.1), 159 (100), 177 (36.3), 220 (7.3).




Figure 1. EI-MS spectrum of santolina alcohol.

21.3%; borneol, 6.2%; bornyl acetate, 3.5%). According to the commercial spectral libraries and the literature (7, 12), the second most abundant compound (RI 1024/1375) showed a mass spectrum identical with that of 2,7-dimethyl-4(*E*),6-octadien-2-ol, previously reported by Maffei et al. in the essential oil of northern Italian *A. ligustica* (2). However, as the retention index of this compound was not present in our databases, we could not confirm its identification by this crucial parameter. Then, we isolated this product from the oil by a multistep fractionation involving the use of AgNO₃-impregnated silica gel, and thereafter we could identify it undoubtedly as santolina alcohol by means of ¹H and ¹³C NMR experiments and comparison with a commercial sample. The confusion between the mass spectral data of santolina alcohol (Figure 1) and 2,7-dimethyl-4(*E*),6-octadien-2-ol was pointed out by Näf-Müller et al. in 1981 (13).

In agreement with the literature (2, 4), the sesquiterpenic content of the oil was mostly represented by germacrene-D (3.1%) and viridiflorol (3.2%). However, neither chamazulene nor guaiazulene was detected despite the specific blue color of the hydrocarbon FC fraction. Considering the high content and the variety of proazulenilactones in *Achillea* species (14, 15),




Figure 2. Synthesis of *rac*-santolina alcohol from 2,5-dimethylhexa-2,4-diene.

other hitherto unknown azulenes can reasonably be present in the oil. As a result, *A. ligustica* from Corsica can be regarded as a potential new chemotype, completely different from the samples harvested in Greece (3), albeit closer to the northern Italian and Sardinian oils (2, 4).

To determine stereochemical information on the main chiral constituents of the essential oil, enantioselective analyses were carried out on the five FC fractions (Table 1). Enantiomer resolution was achieved by using octakis(2,6-di-*O*-pentyl-3-*O*-butyryl)- γ -cyclodextrin (Lipodex-E) and heptakis(2,3-di-*O*-methyl-6-*O*-*tert*-butyldimethylsilyl)- β -cyclodextrin (DIME- β -CD; Hydrodex β -6tBDM), which was previously reported to resolve monoterpenoids efficiently (16). Although chiral multidimensional GC is the most powerful tool for the chiral analysis of essential oils (17), the monodimensional chiral GC study of the FC fractions of the total oil is an economical alternative. The validity of the results requires a careful check of the absence of coelutions within the split peaks used for enantiomeric ratio determination (Figure S3). This check is based on the control of the nature of the constituents of the fractions on both chiral and achiral columns, as well as on the detailed examination of the mass spectral pattern of each split peak.

For all studied chiral compounds, analysis of racemic and/or enantiomerically enriched standard samples allowed us to determine the optimal chromatographic conditions required for their resolution and to attribute each peak to its corresponding enantiomer. As the commercially available santolina alcohol was enantiomerically pure, we had to prepare a racemic sample of this compound to determine the chromatographic conditions required for its chiral analysis. Racemic santolina alcohol 3 was then synthesized according to an original two-step procedure




Figure 3. Enantiomer separation of santolina alcohol on heptakis(2,3-O,O-dimethyl-6-O-*tert*-butyldimethylsilyl)- β -cyclodextrin (Hydrodex β -6tBDM; see Materials and Methods).

Table 2. Antimicrobial Activity of *A. ligustica* Essential Oil As Determined by the Disk Diffusion Assay^a

| microorganism | property | oil concentration | | | ampicillin 10 μ g/mL |
|----------------------|---------------|-------------------|-----|-----|-----------------------------|
| | | 100% | 10% | 1% | |
| <i>S. albus</i> | | 20 | 17 | 9 | 07 |
| <i>S. avidinii</i> | | 17 | 12 | 0 | 08 |
| <i>S. coelicolor</i> | | 18 | 11 | 0 | 08 |
| <i>E. faecalis</i> | Gram-positive | 8 | 0 | 0 | 29 |
| <i>S. aureus</i> | Gram-positive | 9 | 7 | 6.5 | 15 |
| <i>P. aeruginosa</i> | Gram-negative | 9 | 7 | 0 | 0 |
| <i>E. coli</i> | Gram-negative | 8 | 7 | 0 | 21 |
| <i>N. asteroides</i> | | 11.5 | 8 | 0 | nd |
| <i>C. jeikeium</i> | | 17 | 1 | 0 | 20 |

^a Values represent diameter of inhibition zone (mm) at indicated dilution (90/10) (EO/DMSO). No inhibition was observed for any of the tested strains with pure DMSO.

from commercial 2,5-dimethylhexa-2,4-diene **1**. This diene was epoxidized as described by Crandall et al. (18), and the crude oxirane **2** thus obtained (Figure S4) was used in the second step without purification, to be opened by vinylmagnesium bromide (Figure 2). *rac*-Santolina alcohol **3** was used to identify chromatographic conditions suitable for its enantiomeric resolution, and proved that the santolina alcohol contained in the *A. ligustica* essential oil was the pure dextrogyre form (Figure 3).

The antibacterial activity of the essential oil of *A. ligustica* was evaluated against Gram-positive (*E. faecalis*, *S. aureus*, *N. asteroides*, *C. jeikeium*), Gram-negative (*P. aeruginosa*, *E. coli*), and *Streptomyces* (*S. albus*, *S. coelicolor*, *S. avidinii*) bacteria. Antibacterial activity was determined by measuring the zone of growth inhibition of serial dilutions of the essential oil. Ampicillin was employed as positive control. The essential oil was found to have a significant antimicrobial activity against all of the tested bacteria. These results are in agreement with the data reported by Tuberoso et al. (4) except for the growth inhibition of *P. aeruginosa*. This variation could be explained by the different geographical origins of the samples. However, this activity was less potent than that of ampicillin except on *Streptomyces* strains. As shown in Table 2, the crude extract of essential oil and its 10^{-1} diluted solution possess a modest activity against Gram-positive and Gram-negative bacteria

strains compared to control. No activity was observed at a dilution of 10^{-2} for these strains. Nevertheless, on *Streptomyces*, the diameter of inhibition is ≈ 17 mm, which is more than that for ampicillin (8 mm).

The antibacterial activity measured for the essential oil could be attributed to the major compounds, mainly camphor (21.3%) and santolina alcohol (19.3%). Indeed, the antimicrobial activity of various essential oils has been attributed to camphor (19–21). However, Yasphe et al. (22) measured the capacity of camphor to inhibit the growth of various strains, and no inhibition was observed against *E. coli* or *P. aeruginosa*. Hence, santolina alcohol and other minor constituents contributed probably to the antibacterial properties of this essential oil. Actually, antibacterial activities against *E. coli* and *P. aeruginosa* were reported for essential oils rich in santolina alcohol (23, 24).

In conclusion, the essential oil of *A. ligustica* demonstrated an interesting selectivity as it has a modest antimicrobial activity against the tested Gram-positive and Gram-negative strains, but a very potent activity against *Streptomyces* bacteria, never described before. Further studies examining a large number of clinically isolated strains could be interesting to evaluate the therapeutic potential of this essential oil.

ACKNOWLEDGMENT

We thank Dr. Gabriel Alziar of the Museum d'histoire naturelle de la ville de Nice for the taxonomic recognition of the plant and for sharing his deep knowledge of Mediterranean flora. S.P. thanks Dr. Gilles Reyssset and Dr. Yves Janin for useful discussion and critical reading of the manuscript. The CRBIP (Centre de Ressources Biologiques de l'Institut Pasteur) is also greatly acknowledged for providing the bacterial strains.

Supporting Information Available: Figure S1, ^1H NMR spectrum santolina alcohol isolated from the oil; Figure S2, ^{13}C NMR spectrum (JMOD); Figure S3, enantiomer separation of sabinene and β -pinene (Lipodex E); Figure S4, ^1H NMR spectrum of 2,2-dimethyl-3-(2-methylprop-1-enyl)oxirane; Figure S5, EI-MS spectra of unknown compounds in *A. ligustica* essential oil. This material is available free of charge via the Internet at <http://pubs.acs.org>.

LITERATURE CITED

- (1) Simonpoli, P. In *Arburi, Arbe, Arbighiule, Savoirs Populaires sur les Plantes de Corse*; Parc Naturel Régional de la Corse: Ajaccio, Corsica, 1993.
- (2) Maffei, M.; Germano, F.; Doglia, G.; Chialva, F. Essential oils, chromosome numbers and karyotypes from Italian *Achillea* species. Part II. *J. Essent. Oil Res.* **1993**, *5*, 61–70.
- (3) Tzakou, O.; Loukis, A.; Verykokidou, E.; Roussis, V. Chemical constituents of the essential oil of *Achillea ligustica* All. from Greece. *J. Essent. Oil Res.* **1995**, *7*, 549–550.
- (4) Tuberoso, C. I. G.; Kowalczyk, A.; Coroneo, V.; Russo, M. T.; Dessi, S.; Cabras, P. Chemical composition and antioxidant, antimicrobial, antifungal activities of the essential oil of *Achillea ligustica* All. *J. Agric. Food Chem.* **2005**, *53*, 10148–10153.
- (5) Boelens Aroma Chemical Information Service (BACIS). *ESO 2000, The Complete Database of Essential Oils*; Huizen, The Netherlands, 1999.
- (6) Joulain, D.; König, W. A. *The Atlas of Spectral Data of Sesquiterpene Hydrocarbons*; E. B.-Verlag: Hamburg, Germany, 1998.
- (7) McLafferty, F. W.; Staufner, D. B. In *The Wiley NBS Registry of Mass Spectral Data*; Wiley: New York, 1989.
- (8) Joulain, D.; König, W. A.; Hochmuth, D. H. *Terpenoids and Related Constituents of Essential Oils*; MassFinder 2.1; Hamburg, Germany, 2001.
- (9) National Institute of Standards and Technology (NIST). NIST/EPA/NIH Mass Spectral Library, version 1.7, 1999.
- (10) Bertus, P.; Drouin, L.; Laroche, C.; Szymoniak, J. *Tetrahedron* **2004**, *60*, 1375–1383.
- (11) Bauer, A. W.; Kirby, W. M.; Sherris, J. C.; Turck, M. Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* **1966**, *45*, 493–496.
- (12) Eptsein, W. W.; McGee, L. R.; Poulter, C. D.; Marsh, L. L. Mass spectral data for gas chromatograph–mass spectral identification of some irregular monoterpenes. *J. Chem. Eng. Data* **1976**, *21*, 500–502.
- (13) Näf-Müller, R.; Pickenhagen, W.; Willhalm, B. New irregular monoterpenes in *Artemisia vulgaris*. *Helv. Chim. Acta* **1981**, *64*, 1424–1430.
- (14) Glasl, S.; Presser, A.; Gunbilig, D.; Werner, I.; Narantuya, S.; Haslinger, E.; Jurenitsch, J.; Kubelka, W. Highly hydroxylated guaianolides of *Achillea asiatica* and Middle European *Achillea* species. *Phytochemistry* **2001**, *58*, 1189–1194.
- (15) Ahmed, A. A.; Gáti, T.; Hussein, T. A.; Ali, A. T.; Tzakou, O. A.; Couladis, M. A.; Mabry, T. J.; Tóth, G. Ligustolide A and B, two novel sesquiterpenes with rare skeletons and three 1,10-seco-guaianolide derivatives from *Achillea ligustica*. *Tetrahedron* **2003**, *59*, 3729–3735.
- (16) Bayer, M.; Mosandl, A. Improved gas chromatographic stereo-differentiation of chiral main constituents from different essential oils using a mixture of chiral stationary phases. *Flavour Fragrance J.* **2004**, *19*, 515–517.
- (17) Tranchida, P. Q.; Dugo, P.; Dugo, G.; Mondello, L. Comprehensive two-dimensional chromatography in food analysis. *J. Chromatogr. A* **2004**, *1054*, 3–16.
- (18) Crandall, J. K.; Banks, D. B.; Colyer, R. A.; Watkins, R. J.; Arrington, J. P. A synthesis of homoallylic alcohols. *J. Org. Chem.* **1968**, *33*, 423–425.
- (19) Palic, R.; Stojanovic, G.; Tanja, N. Composition and antibacterial activity of *Achillea crithmifolia* and *Achillea nobilis* essential oils. *J. Essent. Oil Res.* **2003**, *15*, 434–437.
- (20) Cha, J. D.; Jeong, M. R.; Jeong, S. I.; Moon, S. E.; Kim, J. Y.; Kil, B. S.; Song, Y. H. Chemical composition and antimicrobial activity of the essential oils of *Artemisia scoparia* and *A. capillaris*. *Planta Med.* **2005**, *71*, 186–190.
- (21) Tirillini, B.; Velasquez, E. R.; Pellegrino, R. Chemical composition and antimicrobial activity of essential oil of *Piper angustifolium*. *Planta Med.* **1996**, *62*, 372–373.
- (22) Yasphe, Y.; Feuerstein, I.; Barel, S.; Segal, R. The antibacterial and antispasmodic activity of *Artemisia herba alba* Asso. II. Examination of essential oils from various chemotypes. *Int. J. Crude Drug Res.* **1987**, *25*, 89–96.
- (23) Yasphe, J.; Segal, R.; Breuer, A.; Erdreich-Naftali, G. Antibacterial activity of *Artemisia herba alba*. *J. Pharm. Sci.* **1979**, *68*, 924–925.
- (24) El-Shazly, A. M.; Hafez, S. S.; Wink, M. Comparative study of the essential oils and extracts of *Achillea fragrantissima* (Forssk.) Sch. Bip. and *Achillea santolina* (Asteraceae) from Egypt. *Pharmazie* **2004**, *59*, 226–230.

Received for review March 17, 2006. Revised manuscript received June 14, 2006. Accepted June 18, 2006. J.J.F. is grateful for an ATER position at the University of Nice-Sophia Antipolis (UNSA) and an enantio-selective GC-MS system funded by the Deutsche Forschungsgemeinschaft. This work was presented as an oral communication at the 36th International Symposium on Essential Oils, Sept 4–7, 2005 (Budapest, Hungary).

JF060752Y