

Published in final edited form as:


J Am Chem Soc. 2007 May 23; 129(20): 6382–6383. doi:10.1021/ja071426h.

Resonance Raman Detection of the Hydroperoxo Intermediate in the Cytochrome P450 Enzymatic Cycle

Piotr J. Mak, Ilia G. Denisov, Doreen Victoria, Thomas M. Makris, Tianjing Deng, Stephen G. Sligar, and James R. Kincaid

Contribution from the Department of Chemistry, Marquette University, Milwaukee, Wisconsin 53233 and Department of Biochemistry, Center for Biophysics and Computational Biology, and the Beckman Institute, University of Illinois, Urbana-Champaign, Illinois 61801

Abstract



The Resonance Raman spectra of the hydroperoxo complex of camphor-bound CYP101 have been obtained by cryoradiolytic reduction of the oxygenated ferrous form that had been rapidly frozen in water-glycerol frozen solution; EPR spectroscopy was employed to confirm the identity of the trapped intermediate. The $\nu(\text{O}-\text{O})$ mode, appearing at 799cm^{-1} , is observed for the first time in a peroxo-heme adduct. It is assigned unambiguously by employing isotopomeric mixtures of oxygen gas containing 50% $^{16}\text{O}^{18}\text{O}$, confirming the presence of an intact O-O fragment. The $\nu(\text{Fe}-\text{O})$ mode is observed at 559cm^{-1} (H_2O). Furthermore, both modes shift down by 3cm^{-1} , documenting the formulation as a hydroperoxo-complex, in agreement with EPR data.

The Cytochromes P450 heme-thiolate enzymes facilitate quite difficult chemical transformations through a multi-step reaction cycle that culminates in the generation of a remarkably potent oxidizing species capable of hydroxylating even inert substrates.¹⁻³ Following substrate binding to the resting state ferric enzyme, two sequential one-electron reductions bracketing the binding of molecular oxygen and a subsequent proton delivery step lead to heterolytic O-O bond cleavage and formation of a highly reactive ferryl heme species comparable to the so-called Compound I intermediate of peroxidases.⁴⁻⁶ Thus, key precursors to this critical cleavage reaction are activated heme-bound peroxo- and hydroperoxo fragments; i.e., (Protoporphyrin)Fe(III)(O_2^{2-}) or Fe(III)($\text{O}-\text{OH}^-$). A useful approach to access and study these species is to generate and trap the relatively stable oxy-ferrous P450 complex and then to subject the cryotrapped (77K) sample to radiolytic reduction using radiation from synchrotron, ^{60}Co γ -ray or ^{32}P sources.^{7,8} Enzymatic intermediates produced by

james.kincaid@mu.edu, s-sligar@uiuc.edu.

Supporting Information Available: Figure S1 shows RR spectra of non-irradiated oxy adducts, while S2 and S3 show EPR and low frequency RR spectra of cryoreduced form.

cryoradiolytic reduction of the oxygenated complex of cytochrome P450_{cam} (CYP101) have been detected by both electronic absorption and EPR spectroscopic methods.^{7,9,10} However, critical mechanistic information has been missing and it is now important to attempt to provide more detailed structural characterization of the active sites of these species.

Resonance Raman (RR) spectroscopy is an especially attractive probe of such species, effectively interrogating both the heme macrocycle structure and various iron-ligand fragments.¹¹⁻¹⁵ In fact, the feasibility of coupling this powerful spectroscopic probe with cryogenic radiolysis of heme protein samples was recently demonstrated and suggested to be a promising new approach to structurally characterize these important intermediates.^{3,14,15} The present work uses RR to provide the first direct observation of the structure sensitive internal vibrational modes of the (Fe-OOH) fragment of the hydroperoxo-ferric intermediate of the CYP101 enzyme.

Samples of the oxygenated form of CYP101 were prepared by bubbling dioxygen gas through 30% glycerol/buffer solutions of the (ferrous) enzyme contained in 5 mm NMR tubes. The tube was shaken for several seconds to ensure efficient mixing and quickly frozen in liquid nitrogen. The RR spectra of these trapped O₂ adducts are entirely consistent with those published previously,¹⁶⁻¹⁸ with some slight shifts to higher frequency being due to low temperature effects. The spectrum of the ¹⁶O₂-ligated form, shown in trace A of Figure 1, exhibits the strong $\nu(^{16}\text{O}-^{16}\text{O})$ band at 1139 cm⁻¹, as expected.¹⁶⁻¹⁸ Trace B shows the difference spectrum obtained by subtraction of the spectra of the ¹⁸O₂ and ¹⁶O₂ samples, while trace C shows the difference spectrum observed for the ¹⁶O₂/H₂O and ¹⁶O₂/D₂O samples, the latter confirming the lack of a shift for these dioxygen ligated forms in D₂O buffers, also as expected.¹⁶⁻¹⁸ The above RR spectral data confirm the integrity of the samples of O₂ adducts. For reasons described below, it was decided to include samples prepared from so-called “scrambled” mixtures of O₂ isotopomers; i.e., [1:2:1] mixtures of [¹⁶O₂:¹⁶O¹⁸O:¹⁸O₂]. These mixtures were prepared as described elsewhere.¹⁹ Thus, trace D exhibits a spectral pattern expected for that of the ¹⁶O₂-ligated species subtracted from that of the adduct formed from the scrambled isotopic mixture, the negative peak in the region of the $\nu(^{16}\text{O}-^{16}\text{O})$ resulting from combination of positive and negative $\nu(^{16}\text{O}-^{16}\text{O})$ features in a 1:4 ratio; i.e., the scrambled mixture contains only 25% ¹⁶O₂. Similarly, trace E exhibits the expected pattern for the difference spectrum indicated.

These samples also exhibit the features expected for the $\nu(\text{Fe}-^{16}\text{O})$ and $\nu(\text{Fe}-^{18}\text{O})$ modes of the Fe-O₂ fragments (Supporting Information, Figure S1), showing a positive feature at 546 cm⁻¹ for the former and at 515 cm⁻¹ for the latter, frequencies near those previously observed for these adducts in solution at 4° C.^{17,18} It is also noted that these lowered frequencies, relative to corresponding values observed near 570 cm⁻¹ for hemoglobin and myoglobin, are also consistent with expectations based on consideration of the trans-axial ligand effects, as has been carefully considered by Babcock and coworkers.²⁰

These frozen oxygenated samples were then irradiated with a ⁶⁰Co γ -source to produce the reduced hydroperoxo- forms; the EPR spectra confirmed the conversion with approximately 60% yield (Figure S2). The RR spectra were acquired using the 442 nm excitation line, a wavelength in closer resonance with the Soret band of the hydroperoxo form.⁹ We note that attempts to acquire the RR spectra of the reduced (i.e., hydroperoxo-) form with the 413 nm excitation yielded only the spectral lines of residual O₂ adducts.

The RR spectra acquired for the γ irradiated samples are shown in traces F through J. In trace F it is seen that there are multiple resonance enhanced heme modes in this region; in fact, casual inspection of the absolute spectra in this spectral region provides no clear indication of modes associated with the Fe-O-O fragment of a peroxo formulation. However, as can be seen in

traces G through J, upon subtraction of the overlapping heme macrocycle modes, the $^{16}\text{O}/^{18}\text{O}$ difference features attributable to such a fragment are clearly evident; i.e., the 40 cm^{-1} shift indicated by the positive and negative components shown in trace G is entirely consistent with that expected for the $\nu(\text{O-O})$ mode of an Fe-O-O fragment of a peroxo/hydroperoxo formulation.²¹⁻²⁴ However, bands observed near this frequency region, which also exhibit comparable $^{16}\text{O}/^{18}\text{O}$ shifts, have also been observed for Fe(IV)=O fragments in Compound I and Compound II derivatives of various heme proteins and, without additional experimental data, there might be lingering ambiguity regarding the nature of the species observed in the present experiments.¹¹⁻¹³

A conclusive set of experiments is realized, however, by employing the "scrambled" isotopomeric mixture mentioned earlier; i.e., the 1:2:1 mixture of ($^{16}\text{O}_2$: $^{16}\text{O}^{18}\text{O}$: $^{18}\text{O}_2$). Thus, the revealing patterns observed in traces I and J for the (scrambled- $^{16}\text{O}_2$) and (scrambled - $^{18}\text{O}_2$) difference traces, respectively, confirm the fact that the oxygen isotope- sensitive modes represent species with intact O-O fragments; i.e., these difference traces are consistent only with an Fe-O-O⁻ fragment and not an Fe=O fragment, for which only a two-band difference pattern would be observed. Furthermore, comparison of difference traces G and H show that both the $\nu(^{16}\text{O}-^{16}\text{O})$ and the $\nu(^{18}\text{O}-^{18}\text{O})$ modes, observed at 799 and 759 cm^{-1} in H_2O solutions, shift down to 796 and 755 cm^{-1} , respectively, in deuterated glycerol buffers. These shifts are quite comparable to those observed for other M-OOH(D) fragments, supporting the formulation of this intermediate as a hydroperoxo- species.²¹⁻²⁴

As is shown in Supporting Information (Figure S3), the low frequency spectra also exhibit new $^{16}\text{O}/^{18}\text{O}$ isotopic sensitive bands at $559/532\text{ cm}^{-1}$ that are most reasonably assigned to the $\nu(\text{Fe-O})$ modes of a Fe-OOH(D) fragment, based on a 3 cm^{-1} shift in D_2O and comparisons with other such species.^{14,15} Thus, the present results indicate that the $\nu(\text{Fe-O})$ bond is stronger for the hydroperoxo species relative to the non-reduced oxy complex, based on a shift of 13 cm^{-1} to higher frequency. This shift is somewhat smaller, however, than the $\sim 45\text{ cm}^{-1}$ shift observed for hydroperoxo-myoglobin, relative to oxyMyoglobin.¹⁴

This observation of a trapped hydroperoxo form is quite consistent with behavior expected for the native Cytochrome P450_{cam} enzyme, which has previously been shown to proceed directly at 77K to the hydroperoxo- form, with rather small fractions of the unprotonated species (Fe-OO⁻) being present in the preparations.⁷ Similarly, recent crystallographic studies of cryotrapped CPO Compound III also were interpreted to lead to radiation induced conversion to the hydroperoxo form, with no evidence for a trapped peroxo-ligated species.²⁵ Future studies with mutant proteins that restrict proton delivery to the Fe-O-O fragment and may facilitate systematic comparisons between oxygenated, peroxo- and hydroperoxo forms, are planned. Finally, the successful stabilization and structural characterization of these immediate precursors to the reactive products formed from O-O bond cleavage may possibly lead to an effective strategy to structurally characterize fleeting intermediates that appear later in the cycle.¹⁻⁶

Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This work was supported by grants from the National Institutes of Health (DK35153 to J.R.K. and a Merit Award, R37GM31756, to S.G.S.). J.R.K. acknowledges support from the Pfleischinger Habermann Fund, Marquette University. We gratefully appreciate the help provided by Dr. John Bentley, Notre Dame Radiation Laboratory (Notre Dame University, IN), a facility of the U.S. Department of Energy, Office of Basic Energy Sciences.

References

1. Ortiz de Montellano Paul, R., editor. Cytochrome P450: Structure, Mechanism, and Biochemistry. 3rd ed.. Kluwer Academic/ Plenum Publishers; New York: 2005.
2. Shaik S, Kumar D, de Visser SP, Altun A, Thiel W. Chem. Rev 2005;105:2279–2328. [PubMed: 15941215]
3. Denisov IG, Makris TM, Sligar SG, Schlichting I. Chem. Rev 2005;105:2253–2277. [PubMed: 15941214]
4. Loew GH, Harris DL. Chem. Rev 2000;100:407–420. [PubMed: 11749241]
5. Zheng J, Wang D, Thiel W, Shaik S. J. Am. Chem. Soc 2006;128:13204–13215. [PubMed: 17017800]
6. Schlichting I, Berendzen J, Chu K, Stock AM, Maves SA, Benson DE, Sweet RM, Ringe D, Petsko GA, Sligar SG. Science 2000;287:1615–1622. [PubMed: 10698731]
7. Davydov R, Makris TM, Kofman V, Werst DE, Sligar SG, Hoffman BM. J. Am. Chem. Soc 2001;123:1403–1415. [PubMed: 11456714]
8. Denisov IG, Makris TM, Sligar SG. Methods in Enzymology 2002;357:103–115. [PubMed: 12424902]
9. Sligar SG, Makris TM, Denisov IG. Biochem. Biophys. Res. Commun 2005;338:346–354. [PubMed: 16139790]
10. Makris, TM.; Denisov, IG.; Schlichting, I.; Sligar, SG. Cytochrome P450: Structure, Function, Genetics. 3rd ed.. Ortiz de Montellano, PR., editor. Kluwer Academic/Plenum Publishers; New York: 2004. p. 149-182.
11. Kitagawa T, Mizutani Y. Coord. Chem. Rev 1994;135/136:685–735.
12. Terner J, Palaniappan V, Gold A, Weiss R, Fitzgerald MM, Sullivan AM, Hosten CM. J. Inorg. Biochem 2006;100:480–501. [PubMed: 16513173]
13. Kincaid, JR. The Porphyrin Handbook. Kadish, KM.; Smith, KM.; Guillard, R., editors. 7. Academic Press; New York: 2000. p. 225-291.
14. Ibrahim M, Denisov IG, Makris TM, Kincaid JR, Sligar SG. J. Am. Chem. Soc 2003;125:13714–13718. [PubMed: 14599210]
15. Ibrahim M, Kincaid JR. J. Porphyrins Phthalocyanines 2004:215–225.
16. Bangcharoenpaupong O, Rizos AK, Champion PM, Jollie D, Sligar S. J. Biol. Chem 1986;261:8089–8092. [PubMed: 3722145]
17. Hu S, Schneider AJ, Kincaid JR. J. Am. Chem. Soc 1991;113:4815–4822.
18. MacDonald IDG, Sligar SG, Christian JF, Unno M, Champion PM. J. Am. Chem. Soc 1999;121:376–380.
19. Proniewicz LP, Nakamoto K, Kincaid JR. J. Am. Chem. Soc 1988;110:4541–4545.
20. Oertling WA, Kean RT, Wever R, Babcock GT. Inorg. Chem 1990;29:2633–2645.
21. Ho RYN, Roelfes G, Feringa BL, Que L Jr. J. Am. Chem. Soc 1999;121:264–265.
22. Chen P, Fujisawa K, Solomon EI. J. Am. Chem. Soc 2000;122:10177–10193.
23. Lehnert N, Neese F, Ho RYN, Que L, Solomon EI. J. Am. Chem. Soc 2002;124:10810–10822. [PubMed: 12207537]
24. Rajani C, Kincaid JR, Petering DH. J. Am. Chem. Soc 2004;126:3829–3836. [PubMed: 15038737]
25. Kuhnel K, Derat E, Terner J, Shaik S, Schlichting I. Proc. Natl. Acad. Sci. USA 2007;104:99–104. [PubMed: 17190816]

**Figure 1.**

Left panel: RR spectrum of $^{16}\text{O}_2$ CYP101 in 30% glycerol / buffer before irradiation (A); $^{16}\text{O}_2$ - $^{18}\text{O}_2$ in glycerol/buffer (B); $^{16}\text{O}_2$ - $^{18}\text{O}_2$ in deuterated glycerol/buffer (C); Scrambled- $^{16}\text{O}_2$ in glycerol/buffer (D); Scrambled- $^{18}\text{O}_2$ in deuterated glycerol/buffer (E). Right panel: RR spectrum of $^{16}\text{O}_2$ CYP101 in 30% glycerol / buffer after irradiation (F); $^{16}\text{O}_2$ - $^{18}\text{O}_2$ in glycerol/buffer (G); $^{16}\text{O}_2$ - $^{18}\text{O}_2$ in deuterated glycerol/buffer (H); Scrambled- $^{16}\text{O}_2$ in glycerol/buffer (I); Scrambled- $^{18}\text{O}_2$ in deuterated glycerol/buffer (J). Dashed line in trace A is for the sample with scrambled oxygen.