

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/256200424>

Time Course Production of Urolithins from Ellagic Acid by Human Gut Microbiota

ARTICLE *in* JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY · AUGUST 2013

Impact Factor: 2.91 · DOI: 10.1021/jf402498b · Source: PubMed

CITATIONS

22

READS

109

5 AUTHORS, INCLUDING:



Rocío García Villalba

Spanish National Research Council

48 PUBLICATIONS 712 CITATIONS

SEE PROFILE



Juan Carlos Espín

Spanish National Research Council

174 PUBLICATIONS 8,060 CITATIONS

SEE PROFILE



María V Selma

Spanish National Research Council

75 PUBLICATIONS 2,047 CITATIONS

SEE PROFILE



Francisco A Tomás-Barberán

Spanish National Research Council

377 PUBLICATIONS 16,072 CITATIONS

SEE PROFILE

1 Time Course Production of Urolithins from Ellagic Acid by Human
2 Gut Microbiota3 Rocío García-Villalba, David Beltrán, Juan Carlos Espín, María Victoria Selma,*
4 and Francisco A. Tomás-Barberán

5 Research Group on Quality, Safety and Bioactivity of Plant Foods, CEBAS-CSIC, Espinardo, Murcia 30100, Spain

6 **ABSTRACT:** Ellagic acid (EA) is converted to urolithins by gut microbiota. Urolithins have beneficial biological effects in
7 humans, but differences in urolithin production capacity among individuals have been shown. Therefore, the identification of the
8 urolithin production pathways and the microorganisms implicated is of high interest. EA was incubated with gut microbiota from
9 two volunteers able to produce urolithins but with different in vivo urolithin profiles (urolithin A and isourolithin A producers).
10 The metabolic capabilities observed in vivo were retained in vitro. Both individuals showed a much higher abundance of
11 *Clostridium leptum* group of *Firmicutes* phylum than *Bacteroides/Prevotella*. EA was either dissolved in DMSO or suspended in
12 water. DMSO increased EA solubility but decreased urolithin production rate due to a delay in growth of some microbial groups,
13 principally, *Clostridium coccoides*. This allowed the detection of catabolic intermediates [urolithins M-5, M-6, M-7, C, and
14 2,3,8,10-tetrahydroxy urolithin (urolithin E)]. Bacteria from *C. coccoides* group (or genera co-occurring in vivo with this group)
15 seem to be involved in production of different urolithins.16 **KEYWORDS:** ellagitannins, colon microbiota, metabolism, dibenzopyranones, polyphenols, human health

17 ■ INTRODUCTION

18 Ellagitannins and ellagic acid (EA) are plant secondary
19 metabolites that have relevant antioxidant activities in vitro,
20 potential cardiovascular protection, and anticarcinogenic and
21 anti-inflammatory effects.^{1–3} These phytochemicals are relevant
22 constituents in different foods including pomegranates, berries
23 (strawberry, raspberry, blackberry, camu–camu, etc.), nuts
24 (walnuts, acorns, chestnuts, etc.), muscadine grapes, oak-aged
25 wines, and medicinal plants and tisanes (geranium, oak leaves,
26 etc.). They are not absorbed in the gut and are metabolized in
27 vivo by the gut microbiota to produce a series of metabolites
28 known as urolithins.^{4,5} Therefore, urolithins have been
29 suggested as biomarkers of intake for strawberries, raspberries,
30 pomegranates, walnuts, and oak-aged red wines.⁶ We have
31 previously suggested that urolithins, that are much better
32 absorbed than the original ellagitannins, could be responsible
33 for the systemic biological effects of ellagitannins.^{2,5} Additional
34 evidence supports this suggestion as urolithins and their
35 glucuronide and sulfate metabolites have shown relevant effects
36 at the concentrations found in vivo.^{7–9} Urolithins share a
37 nucleus of a dibenzo-pyran-one, with different hydroxyl
38 substitutions. The first reported urolithins were urolithin A
39 and urolithin B that were found in kidney stones of sheep.¹⁰ In
40 a recent study, it has been demonstrated that other mammals
41 also produce urolithins from ellagitannins, although they have
42 different structures and different hydroxylation patterns on the
43 urolithin nucleus.¹¹ Thus, isourolithin A has been found as the
44 main in vivo metabolite in beef cattle after oak leaf intake,¹² and
45 a series of tetrahydroxy and trihydroxy urolithin metabolites
46 were produced by rats after *Geranium* extracts intake.¹³47 Recently, it has been reported that some humans are able to
48 produce isourolithin A as the main in vivo metabolite instead of
49 urolithin A after the intake of strawberries¹⁴ and raspberries.¹⁵
50 Furthermore, marked person-to-person differences have been51 observed in the level of in vivo urolithin production following
52 consumption of ellagitannin-containing food. In vitro anaerobic
53 metabolism of EA to urolithin A in human fecal suspensions
54 has also been reported.¹⁶ More recently, in vitro anaerobic
55 incubation of EA with human fecal suspensions demonstrated
56 conversion to some other urolithins, urolithins B, C, and
57 isourolithin A.¹⁷ However, neither comparison between the in
58 vitro and the in vivo urolithin patterns of these volunteers nor
59 their endogenous microbiota has been analyzed in any of these
60 studies.61 In the present study, we evaluated the in vitro time course
62 production of urolithins from EA by human fecal microbiota
63 from two volunteers who have urolithin A and isourolithin A
64 urinary excretion patterns. Detailed in vitro and in vivo
65 experiments were performed to identify the intermediary and
66 final catabolites of EA conversion by the human gut microbiota
67 to elucidate interindividual urolithin production routes and to
68 determine if EA metabolism could be linked to the microbiota
69 composition of these individuals.

70 ■ MATERIALS AND METHODS

71 **Chemicals.** EA and 6,7-dihydroxycoumarin were obtained from
72 Sigma-Aldrich (St. Louis, MO, USA). Urolithins A and B were
73 chemically synthesized by Villapharma SL (Parque Tecnológico de
74 Fuente Álamo, Murcia, Spain). Urolithins C (3,7,8-tetrahydroxy-6H-
75 dibenzo[*b,d*]pyran-6-one, Uro-C; > 95% purity) and D (2,3,7,8-
76 tetrahydroxy-6H-dibenzo[*b,d*]pyran-6-one, Uro-D; > 95% purity) were
77 purchased from Dalton Pharma Services (Toronto, Canada).
78 Methanol and acetonitrile were purchased from Romil (Barcelona, 78
Spain) and ethyl acetate and dimethyl sulfoxide from Labscan (Dublin, 79

Received: June 6, 2013

Revised: August 28, 2013

Accepted: August 28, 2013

Ireland). Formic acid and hydrochloric acid was obtained from Panreac (Barcelona, Spain). Milli-Q system (Millipore Corp., Bedford, MA) ultrapure water was used throughout this experiment. Nutrient broth (NB) was from Oxoid (Basingstoke, Hampshire, UK). L-Cysteine hydrochloride was from (Panreac Química, Barcelona, Spain). Dimethyl sulfoxide (DMSO) was from (Scharlab, Barcelona, Spain). All chemicals and reagents used in the preparation of buffers, macromineral, micromineral, and reducing solutions were obtained from Sigma-Aldrich, Panreac, and Scharlab.

Collection of Human Fecal Samples. These were donated by a healthy female (age 31) and a healthy male (age 36) in the Centro de Edafología y Biología Aplicada del Segura (CEBAS-CSIC, Murcia, Spain). These two volunteers were identified as urolithin A and isourolithin A producers, respectively, in a previous study in which urine samples were analyzed after the intake of strawberries and strawberry jam,¹⁸ although these metabolic groups were not reported. To conduct fermentation experiments in duplicate, each volunteer donated two fecal samples. They were stored at 4 °C and were further processed within 1 h of donation, as no differences in microbial concentrations were observed in fresh samples before and after processing samples for fermentation experiments. Anoxic conditions were maintained by using the AnaerobeGen compact system (Oxoid).

Fermentation Medium. This was prepared as described by Jaganath et al.¹⁹ with some modifications. Briefly, 2 g of tryptone, 2 g of glucose, 1 g of maltose and 2 g of yeast extract were mixed in 400 mL of distilled water and 100 μ L of micromineral solutions (consisting of 13.2 g of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 10 g of $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 1 g of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 8 g of $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, and distilled water up to 100 mL), 200 mL of buffer solution (2 g of NH_4CO_3 , 17.5 g Na_2CO_3 , and distilled water up to 500 mL), 200 mL of macromineral solution (2.85 g of Na_2HPO_4 , 3.1 g KH_2PO_4 , 0.3 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, and distilled water up to 500 mL), 1 mL of 1% (w/v) of resazurin solution (a redox indicator) and 0.5 mg/L of Vit K₁, and 5 mg/L and 625 mg/L hemein. The medium was adjusted to pH 7 using HCl, dispensed into the fermentation vessels, and autoclaved at 121 °C for 15 min and allowed to cool under oxygen-free nitrogen (OFN) to remove oxygen.

Conversion Experiments of EA into Urolithins with Human Fecal Cultures. Preparation of fecal suspensions and subsequent culturing experiments were conducted under anoxic conditions in an anaerobic chamber (Don Whitley Scientific Limited, Shipley, UK) with an atmosphere consisting of $\text{N}_2/\text{H}_2/\text{CO}_2$ (80:10:10) at 37 °C. Aliquots of fecal samples (10 g) were diluted 1/10 (w/v) in NB supplemented with 0.05% L-cysteine hydrochloride and homogenized by stomacher in filter bags. Aliquots of filtered fecal suspensions (2 mL) were inoculated into 200 mL of prerduced fermentation medium containing EA at 30 μ M with and without 1% DMSO. DMSO was used in order to increase EA solubility, and EA dissolved in water instead of DMSO was used as control. Duplicate cultures were prepared in parallel from each fecal suspension. In addition, controls were used; ones without fecal suspension and others without EA. Samples (6 mL) were collected at appropriate time intervals during 8 day incubation at 37 °C and stored at –20 °C until further HPLC analysis. Similarly, 3 mL were collected every 2 h during 2 day incubation at 37 °C and stored at –20 °C until further qPCR analysis.

Sample Cleanup for LC Analyses. Feces and urine of the two volunteers were extracted and analyzed. Urine samples were defrosted, vortexed, and centrifuged at 14000g for 10 min at 4 °C. The supernatant was filtered through a 0.45 μ m PVDF filter and analyzed by HPLC-DAD-IT MS. When the intensity was not enough to distinguish the metabolites, urine samples were concentrated using a Sep-Pak reverse phase C-18 extraction cartridge (Waters Millipore). The cartridges were previously activated with 10 mL of MeOH and 10 mL of water. Then 25 mL of urine acidified with 250 μ L of formic acid (1%) were passed through the cartridge that was then dried with air. The metabolites remaining in the cartridge were eluted with 3 mL of MeOH/ H_2O (50:50, v/v). Samples were analyzed after filtration through 0.45 μ m PVDF filter. Feces samples (1 g) were defrosted and homogenized with 10 mL of MeOH/DMSO/ H_2O (40:40:20) with 0.1% HCl using an Ultra-Turrax for 1 min at 24000 rpm. The mixture

was centrifuged at 5000g for 10 min at room temperature and the supernatant filtered through a 0.45 μ m PVDF filter before analysis.

Samples (5 mL) obtained in fermentation experiments with human faecal suspensions were extracted with 5 mL of ethyl acetate acidified with 1.5% formic acid. The mixture was vortexed for 2 min and centrifuged at 3500g for 10 min. The organic phase was separated and evaporated under reduced pressure until dryness. The dry samples were then redissolved in 250 μ L of methanol and filtered through a 0.45 μ m PVDF filter. Then 5 μ L of 100 μ g/mL of internal standard (6,7-dihydroxycoumarin) was added to 50 μ L of sample prior to the injection onto a column for LC-UV/vis and LC-MS analysis under the conditions described below.

LC-UV/Vis and LC-MS/MS Analyses. The analyses were performed using an Agilent 1100 HPLC system equipped with a photodiode array detector and an ion-trap mass spectrometer detector in series (Agilent Technologies, Waldbronn, Germany). Chromatographic separation was carried out on a reverse phase LiChroCART C-18 column (Merck, Darmstadt, Germany) (250 mm \times 4 mm, 4.5 μ m particle size) using water with 1% formic acid (A) and acetonitrile (B) as the mobile phases. The gradient profile was: 0–20 min, 5–30% B, 20–30 min, 30–55% B, 30–38 min, and 55–90% B, and this percentage was maintained for 2 min and then came back to the initial conditions. A volume of 10 μ L of sample was injected onto the column operating at room temperature and a flow rate of 1 mL/min. UV chromatograms were recorded at 280, 360, and 305 nm.

The HPLC system was coupled in series to an ion trap mass spectrometer (IT) equipped with an electrospray interface (ESI). Nitrogen was used as drying gas with flow of 11 L/min and temperature of 350 °C and nebulizing gas at pressure of 65 psi. The capillary voltage was set at 4 kV. Mass scan (MS) and daughter (MS-MS) spectra were recorded in negative mode in the range of m/z 100–700 with target mass of 300. Maximum accumulation time of ion trap and the number of MS repetitions to obtain the MS average spectra were set at 200 ms and 3, respectively. Compound stability was set at 75%.

Identification of all metabolites was carried out by direct comparison (UV spectra and MS) with pure standards and confirmed by their spectral properties and molecular mass. Calibration curves were obtained for EA, urolithin A, urolithin B, and urolithin C with good linearity ($R^2 > 0.998$). EA was quantified at 360 nm and urolithins at 305 nm. The limits of detection (LODs) were determined based on a signal-to-noise ratio (S/N) of 3 and of 10 for the limit of quantification (LOQ). EA, urolithin A, and urolithin B showed LODs of 0.5 μ M and LOQs of 1.67 μ M and urolithin C an LOD of 0.2 μ M and an LOQ of 0.67 μ M. Repeatability was evaluated by injecting 20 μ M of a mixture of standards four times in the same day (intraday repeatability) and in four different days (interday repeatability). The results expressed as the relative standard deviation (RSD) of peak area were $\leq 5\%$ for intraday repeatability and $\leq 8\%$ for interday repeatability. The recovery of the compounds was calculated, spiking the medium in the presence of inactivated bacteria with a standard solution of EA, urolithin A, and urolithin B in DMSO at a final concentration of 20 μ M. Recoveries of 75%, 90%, and 83% were obtained for EA, urolithin A, and urolithin B, respectively. Isourolithin A was quantified at 305 nm with the urolithin A calibration curve and urolithins M-5, M-6, and M-7 at 360 nm with the EA calibration curve.

Fecal Dry Weight Determination and DNA Extraction. To determine fecal moisture content, approximately 0.5 g (wet weight) of each fecal specimen was placed in a vacuum dryer for 3 d and reweighed. Percent fecal dry weight was calculated. Total DNA was extracted from human fecal samples using a commercial DNA extraction kit (QIAampR DNA Stool Mini Kit, Qiagen Inc., Valencia, CA). Although this kit did not contain beads, an additional step of vigorous shaking using the FastPrep Instrument was carried out. Briefly, 10 mg of fresh fecal samples ($25.3 \pm 8.9\%$ dry matter) were added to 2 mL tubes containing specialized beads (MP Biomedicals, LLC, Ohio, USA) and homogenized in the lysing matrix using the FastPrep Instrument for 30 s at a speed setting of 5.5. The following steps were carried out attending protocol supplied with the kit: briefly, incubation temperatures (95 °C, 5 min and 70 °C, 10 min), cell lysis

Table 1. Primer and Probe Sequences Used for Detection of Main Bacterial Groups in Human Feces by Quantitative Real-Time Polymerase Chain Reaction

target organism groups and species ²⁴	primers and probes	sequence 5'–3'	refs
<i>Clostridium leptum</i> group	F_Clept 09	CCT TCC GTG CCG SAG TTA	24
<i>Clostridium leptum</i>	R_Clept 08	GAA TTA AAC CAC ATA CTC CAC TGC TT	
<i>Faecalibacterium prausnitzii</i>	P_Clep 01 ^a	6FAM-CAC AATAAG TAA TCC ACC	
<i>Ruminococcus albus</i>			
<i>Bifidobacterium</i> genus	F_Bifid 09c	CGG GTG AGT AAT GCG TGA CC	24
<i>Bifidobacterium adolescentis</i>	R_Bifid 06	TGA TAG GAC GCG ACC CCA	
<i>Bifidobacterium breve</i>	P_Bifid ^a	6FAM-CTC CTG GAA ACG GGT G	
<i>Bifidobacterium infantis</i>			
<i>Clostridium coccooides</i> group	F_Ccoc 07	GAC GCC GCG TGA AGG A	24
<i>Clostridium coccooides</i>	R_Ccoc 14	AGC CCC AGC CTT TCA CAT C	
<i>Ruminococcus gnavus</i>	P_Erec482 ^a	VIC-CGG TAC CTG ACT AAG AAG	25
<i>Ruminococcus hansenii</i>			
<i>Eubacterium rectale</i>			
<i>Bacteroides/Prevotella</i> group	F_Bacter 11	CCT WCG ATG GAT AGG GGT T	24
<i>Bacteroides fragilis</i>	R_Bacter 08	CAC GCT ACT TGG CTG GTT CAG	
<i>Bacteroides ovatus</i>	P_Bac303 ^a	VIC-AAG GTC CCC CAC ATT G	26
<i>Bacteroides thetaiotaomicron</i>			
<i>Bacteroides uniformis</i>			
<i>Bacteroides vulgatus</i>			
<i>Bacteroides caccae</i>			
<i>Bacteroides eggerthii</i>			
<i>Prevotella oralis</i>			
<i>Prevotella buccae</i>			
<i>Prevotella albensis</i>			
<i>Escherichia coli</i> species	E.coli F	CAT GCC GCG TGT ATG AAG AA	27
	E.coli R	CGG GTA ACG TCA ATG AGC AAA	
<i>Lactobacillus/Leuconostoc/Pediococcus</i> group	F_Lacto 05	AGC AGT AGG GAA TCT TCC A	24
<i>Lactobacillus acidophilus</i>	R_Lacto 04	CGC CAC TGG TGT TCY TCC ATA TA	
<i>Lactobacillus casei</i>			
<i>Lactobacillus paracasei</i>			
<i>Lactobacillus delbrueckii</i>			
<i>Lactobacillus fermentum</i>			
<i>Lactobacillus johnsonii</i>			
<i>Lactobacillus plantarum</i>			
<i>Lactobacillus rhamnosus</i>			
<i>Lactobacillus helvetic</i>			
<i>Pediococcus inopinatus</i>			
<i>Pediococcus parvulus</i>			
<i>Pediococcus cellicola</i>			
<i>Pediococcus acidilactici</i>			
<i>Leuconostoc mesenteroides</i>			
<i>Leuconostoc pseudomesenteroides</i>			

^aProbe sequences.

and homogenization (centrifugation at 14,000 rpm, 1 min), adsorption of inhibitors (InhibitEx tablet), and approximate time to completion (60–80 min).

Real-Time qPCR. DNA from dominant groups of fecal bacteria was quantified with real-time qPCR and primers as summarized in Table 1. Real-time qPCR was performed using an ABI 7500 sequence detection system. Amplification and detection were carried out in 96-well plates with TaqMan Universal PCR 2× master mix (Applied-Biosystems) or with SYBR-Green PCR 2× master mix (Applied-Biosystems). Each reaction was run in duplicate in a final volume of 25 μL with 0.2 mM final concentration of each primer, 0.25 mM final

concentration of each probe, and 10 mL of appropriate dilutions of DNA samples. Probes were designed with molecular-groove binding nonfluorescence quencher (MGBNFQ). Amplifications were carried out using the following ramping profile: 1 cycle at 95 °C for 10 min, followed by 40 cycles of 95 °C for 30 s and 60 °C for 1 min. For SYBR-Green amplifications, the following dissociation protocol (a hold at 95 °C for 15 s, a hold at 60 °C for 20 s, and a slow ramp for 20 min from 60 to 95 °C) was followed after the last cycle in order to check the expected amplification products.

When PCR was performed on unknown fecal samples, the cycle threshold of each sample was compared with a standard curve 240

241 (performed in triplicate) made by diluting genomic DNA (6-fold serial
242 dilutions of *Bifidobacterium longum* DSM 20088T for *Bifidobacterium*
243 genus, *Escherichia coli* CECT 515T for *E. coli* species, *Clostridium*
244 *leptum* DSM 753T and *Clostridium coccooides* DSM 935T for *C. leptum*
245 and *C. coccooides* groups, *Bacteroides ovatus* DSM 1896T for *Bacteroides*/
246 *Prevotella* group and *Lactobacillus plantarum* CECT 748T for
247 *Lactobacillus/Leuconostoc/Pediococcus* group). Selected bacteria strains
248 were cultured anaerobically on selective broth as recommended by
249 DSMZ. For each culture, the total number of bacteria, in terms of
250 CFU, was determined by plating. Aliquots of 1 mL of culture were
251 used for DNA extraction by DNeasy blood and tissue kit (Quiagen)
252 following the manufacturer's instructions.

253 **Statistical Analyses and Data Modeling.** Fecal samples were
254 analyzed in triplicate, and the quantitative estimates represent mean
255 values \pm standard error. Data were subjected to statistical analysis of
256 variance (ANOVA). Normality of variances was tested by the
257 Shapiro–Wilk test before determining the ANOVA. Statistical analyses
258 were performed using SPSS V.14 for Windows.

259 Microbial growth curves were fitted using the function of Baranyi et
260 al.²⁰ to estimate the main growth parameters (maximum specific
261 growth rate, lag time of microorganisms before the onset of growth,
262 and estimated correlation coefficient that indicates the goodness of fit
263 of the parameters derived from experimental data). Only growth
264 curves with at least 10 data points were used for modeling, as
265 suggested by the authors.

266 ■ RESULTS

267 **In Vivo Metabolism of EA and Ellagitannins by Human**
268 **Gut Microbiota.** Two volunteers who had different routes of
269 EA metabolism in vivo were selected for the present study.
270 Volunteer 1 produced urolithin A, and volunteer 2 produced
271 isourulithin A. We had previously identified these two urolithin
272 metabolic profile groups in urine samples from different
273 volunteers after the intake of walnuts, strawberries, and
274 raspberries (unpublished data). The analyses of fecal samples
275 from both volunteers after the intake of walnuts (0.6 g/kg body
276 weight per day for 5 days) showed that volunteer 1 had a
277 chromatographic profile characterized by urolithin A that was
278 identified by its MS spectrum, its characteristic UV spectrum,¹¹
279 and by chromatographic comparisons with an authentic
280 standard, while volunteer 2 produced isourulithin A as a main
281 metabolite (Figure 1). The UV spectra of urolithin A and
282 isourulithin A allow a clear and fast identification of both
283 metabolites.¹¹ In addition, other minor metabolites were also
284 detected in fecal samples of both volunteers [urolithin M-6 (7)
285 and urolithin C (8)] (Table 2). When the urine of the same
286 volunteers was analyzed, their metabolic profiles were
287 consistent with those found in feces, and volunteer 1 excreted
288 mainly urolithin A glucuronide (1), while volunteer 2 excreted
289 two isomers of isourulithin A glucuronide (2, 3) instead
290 (Figure 2). Trace amounts of urolithin B-glucuronide (10)
291 were also detected in the urine of volunteer 2 (Figure 2B).

292 **Bacterial Populations of Human Fecal Samples Used**
293 **in This Study.** Fecal bacteria densities of the two volunteers,
294 who donated feces for the investigation of the metabolic fate of
295 EA in vivo, were also analyzed. Interindividual differences in
296 fecal microbiota were analyzed by qRT-PCR using primers and
297 probes (Table 1) that target the main bacterial groups:
298 *Bifidobacterium* genus; *C. leptum* group (*C. leptum*, *Faecalibacte-*
299 *rium prausnitzii*, and some species of *Ruminococcus* genus); *C.*
300 *coccooides* group (*C. coccooides*, *Eubacterium rectale* spp., and some
301 species of *Ruminococcus* genus); *E. coli* species; *Bacteroides*/
302 *Prevotella* group; *Lactobacillus/Leuconostoc/Pediococcus* group
303 (Table 1). All bacteria results were presented as the mean of
304 the log CFU per g feces. Fecal dry weight (%) was 31.7 and




Figure 1. HPLC-UV analysis at 305 nm of fecal samples of volunteer 1 (A) and volunteer 2 (B) after the intake of walnuts (0.6 g/kg body weight per day for 5 days). (7) urolithin M-6; (8) urolithin C; (11) isourulithin A; (12) urolithin A.

19.0% in volunteers 1 and 2. To overcome the fact that fecal samples contained different water, a correction factor ($19/31.7 = 0.6$) was applied to microbial data of volunteer 1.

Fecal sample analyses from both urolithin A (volunteer 1) and isourulithin A (volunteer 2) producers were analyzed (Figure 3). This study showed that one of the most highly represented bacterial groups, in human stools of both volunteers, was the *C. leptum* group of the *Firmicutes* phylum. No significant differences in levels of *C. leptum* group ($7\text{--}34 \times 10^7$ CFU/g) and *C. coccooides* group ($(5\text{--}8) \times 10^6$ CFU/g) of the *Firmicutes* phylum as well as the *Bacteroides/Prevotella* group ($5\text{--}12 \times 10^5$ CFU/g) of the *Bacteroidetes* phylum were observed between fecal samples from volunteers 1 and 2. However, concentrations of other bacterial groups in the volunteer 1 fecal sample were noticeably different from those of the volunteer 2 fecal sample ($P \leq 0.01$) (Figure 3). Thus, the microbiota from volunteer 1 showed a higher concentration of *Bifidobacterium* genus (6×10^8 CFU/g) of the *Actinobacteria* phylum when compared with that of volunteer 2 (1×10^6 log CFU/g) (Figure 3A). Furthermore, in volunteer 1, the fecal sample contained a higher concentration of *E. coli* species (1×10^6 CFU/g) of the *Proteobacteria* phylum than fecal sample of volunteer 2 (8×10^4 CFU/g). In contrast, a lower abundance of *Lactobacillus/Leuconostoc/Pediococcus* group of the *Firmicutes* phylum was found in volunteer 1 (2×10^4 CFU/g) which were 2.7 log units lower than counts for fecal sample of volunteer 2 (1×10^7 CFU/g) (Figure 3). These studies showed that the

Table 2. Metabolites identified in urine (U), Feces (F), Fermentation Medium Incubated with Human Fecal Suspensions in the Presence of Ellagic Acid (FM) or in the Presence or Ellagic Acid Dissolved in 1% DMSO (FM-DMSO) (sh, Shoulder)^a

no.	comps	volunteers	origins	R _t (min)	[M – H] [–]	λ _{max}
1	urolithin A glucuronide	1	U	15.75	403	246, 278, 303, 350
2	isourolithin A glucuronide	2	U	15.52	403	255, 299sh, 325
3	isourolithin A glucuronide	2	U	16.14	403	255, 299sh, 325
4	urolithin M-5	1	FM and FM-DMSO	14.65	275	261, 292sh,352
		2	FM-DMSO			
5	urolithin E	1	FM-DMSO	16.55	259	252, 276sh,365
6	ellagic acid	1 and 2	FM and FM-DMSO	17.62	301	254, 300sh, 366
7	urolithin M-6	1 and 2	F and FM and FM-DMSO	17.94	259	259, 291sh, 350
8	urolithin C	1	F and FM	18.83	243	255, 304, 349
		2	F and FM and FM-DMSO			
9	urolithin M-7	1	FM and FM-DMSO	20.49	243	250, 280sh, 367
		2	FM-DMSO			
10	urolithin B glucuronide	2	U	20.56	387	249, 273, 297, 326
11	isourolithin A	2	F and FM and FM-DMSO	23.06	227	256, 302, 329
12	urolithin A	1	F and FM and FM-DMSO	24.11	227	246, 280, 305, 356
		2	FM and FM-DMSO			
13	urolithin B	2	FM	28.21	211	249, 276, 301, 333

^aInternal standard (6,7-dihydroxycoumarin), [M – H][–] 177; retention time, 11.52 min.




Figure 2. HPLC-UV analysis at 305 nm of urine samples of volunteer 1 (A) and volunteer 2 (B) after the intake of walnuts (0.6 g/kg body weight per day for 5 days). (2) urolithin A glucuronide; (3) isourolithin A glucuronide; (10) urolithin B glucuronide.




Figure 3. Quantities of dominant groups of bacteria in fecal samples of volunteer 1 (A) and volunteer 2 (B) as determined by real-time polymerase chain reaction. The mean (±SD) counts from triplicate determinations (three DNA samples isolated from the same feces) were used to quantify the bacterial groups. (Bif) *Bifidobacterium* genus; (C. lep) *Clostridium leptum* group; (C. coc/Eu) *Clostridium coccoides* group; (E. coli) *E. coli* species; (Bac/Prev) *Bacteroides/Prevotella* group; (Lac/Leu/Pedio) *Lactobacillus/Leuconostoc/Pediococcus* group. **Significant differences ($P \leq 0.01$) in bacteria levels between volunteer 1 (A) and volunteer 2 (B). An estimation of the total amount of *Firmicutes* was obtained by adding bacterial values obtained from *C. coccoides*, *C. leptum*, and *Lactobacillus/Leuconostoc/Pediococcus*.

Bifidobacterium genus of the *Firmicutes* and *Actinobacteria* phyla were the major representative in volunteer 1.

In Vitro Catabolism of EA by Human Gut Microbiota.

The production of urolithins by the gut microbiota from the two volunteers was followed in vitro in order to study the time course of the production of the two microbial metabolite patterns found in vivo and identify potential intermediate catabolites in the route from EA to urolithin A or isourolithin A (Figure 4). The first metabolites observed when EA was incubated with the volunteer 1 (urolithin A producer) fecal microbiota were urolithin M-5 (pentahydroxy-urolithin) and urolithin M-6 (tetrahydroxy-urolithin), which peaked at 12 h

most highly represented bacterial group in volunteer 2 was the *C. leptum* of the *Firmicutes* phylum, while *C. leptum* and




Figure 4. In vitro conversion of EA during batch fermentation with human feces from two volunteers. Human fecal cultures consisted of (A) volunteer 1 fecal suspension in fermentation medium with 30 μM EA; (B) volunteer 2 fecal suspension in fermentation medium with 30 μM EA; (C) volunteer 1 fecal suspension in fermentation medium with 30 μM EA dissolved with 1% DMSO; (D) volunteer 2 fecal suspension in fermentation medium with 30 μM EA dissolved with 1% DMSO. Ellagic acid (red line with red solid diamonds); urolithin M-5 (blue line with blue solid circles); urolithin E (black lines with black solid squares); urolithin M-6 (magenta lines with magenta Xs); urolithin M-7 (cyan lines with cyan solid stars); urolithin C (green lines with green open squares); urolithin A (purple lines with purple open circles); isourolithin A (brown lines with brown solid triangles); urolithin B (black lines with black solid circles).

with a maximum concentration of 1 and 2.5 μM, respectively (Figure 4A). Then, urolithin C (trihydroxy-urolithin) started to be accumulated reaching 9.4 μM at 14 h. At that moment, a small amount of another trihydroxy-urolithin (urolithin M-7) started to be produced with a maximum of 0.29 μM at 16 h. Urolithin A was detected after 14 h incubation, and a maximum of 19.6 μM was reached at 42 h, then a plateau was maintained (Figure 4A). A large proportion of the EA supplied (30 μM) was finally converted into urolithin A (21 μM), although some EA remained unmetabolized in the medium (1.3 μM).

In the case of volunteer 2 (isourolithin A producer), no urolithin M-5 was detected and both urolithin C and urolithin M-6 appeared with a maximum after 5 h (7.5 and 3.5 μM, respectively) (Figure 4B). Then isourolithin A started to be produced, reaching a maximum of 10 μM after 24 h. Isourolithin A then decreased, and urolithin B increased to reach 6.5 μM after 60 h and then reached a plateau. At the same

time as urolithin B was produced, some urolithin A was also observed, although it reached concentrations much lower than those found for volunteer 1 (3 μM). The maximum EA conversion to urolithins was lower in the case of volunteer 2 fecal fermentation (15 μM) (Figure 4B) than in the case of volunteer 1 fecal fermentation (21 μM) (Figure 4A) and some EA remained unmetabolized in the medium (2 μM).

The urolithin production from EA when this was dissolved in DMSO and added to the fermentation medium (1%) to increase its solubility, and potentially its bacterial accessibility, was also evaluated. The EA catabolism was very different from that observed when EA was added to the fermentation medium without adding any organic solvent, and significant changes were observed in growth parameters of some bacterial groups analyzed as described in the section below. In the case of volunteer 1, the main in vitro metabolite produced in the presence of DMSO was urolithin M-5 (pentahydroxy-urolithin)

instead of urolithin A, which started to be detected after 12 h, with a maximum concentration of 9 μM at 40 h incubation. Urolithin M-6, urolithin M-7, and urolithin A were also observed but not before 16, 36, and 42 h incubation, respectively, and in much smaller maximal concentrations (0.3, 1.3, and 0.5 μM , respectively) (Figure 4C) than in the absence of DMSO (2.5, 0.3, and 19.6 μM , respectively) (Figure 4A). Urolithin C was not produced in the presence of DMSO (Figure 4C). Despite the smaller and slower urolithin production in presence of the DMSO, a large proportion of EA supplied was metabolized in the medium (24 μM) (Figure 4C). A relevant unidentified tetrahydroxy urolithin was started to be produced after 36 h of incubation, and this was named as urolithin E (Figure 4C). This had the same mass spectrum as urolithin M-6. To identify the structure of urolithin E, this was compared with the other two available tetrahydroxy-urolithins, urolithin D and urolithin M-6. All of them had the same MS spectrum but differed in the UV spectrum (Figure 5). Urolithin M-6 had a characteristic UV spectrum, and urolithin D had a similar spectrum to urolithin A. Urolithin E had a different UV spectrum, with a significant band I at 360 nm, a characteristic of those urolithins without a hydroxyl at 9-position (Figure 5).¹¹ Therefore, urolithin E was tentatively identified as 3,4,8,10-tetrahydroxy-urolithin, a new urolithin metabolite. NMR analysis of this urolithin was not possible due to the small amount produced.

In the case of volunteer 2, when EA added was dissolved in DMSO, this started to be converted to urolithin M-5, urolithin M-6, and urolithin C with a maximum concentration of 2, 6, and 7.5 μM after 7.5, 10, and 25 h, respectively. Then urolithin A started to be produced, reaching a maximum (22 μM) after 40 h (Figure 4D). In this case, only a small concentration of isourolithin A was detected (1 μM). Urolithin M-7 was also observed and reached maximal concentrations of 3 μM after 50 h. Most EA was metabolized, and only low concentrations (1 μM) remained unmetabolized in the medium (Figure 4D).

Changes in Human Gut Microbiota in Vitro. Changes in the fecal microbiota of volunteers 1 and 2 were followed during the in vitro metabolism of EA in order to elucidate which microbial groups could be implicated in the production of urolithins. The growth characteristics of the main bacterial groups, determined by q-PCR, during the conversion of EA in vitro batch fermentation are shown in Table 3. In the absence of DMSO, the main differences between fecal microbiota development of volunteers 1 and 2 were detected in the *Bifidobacterium* genus. In fact, the bifidobacteria of volunteer 2 were not able to grow in the in vitro conditions despite urolithins being produced. *Lactobacillus/Leuconostoc/Pediococcus* group also grew slower in volunteer 2 than in volunteer 1, but a higher level of bacteria was obtained at the end of the growth (N_{max}) in volunteer 2 (5.0 log CFU/mL) than in volunteer 1 culture (4.4 log CFU/mL). In contrast, *C. coccoides* group reached a higher level in volunteer 1 (N_{max} : 6.1 log CFU/mL) than in volunteer 2 culture (N_{max} : 5.0 log CFU/mL) (Table 3). In the same way, urolithin concentrations were higher in volunteer 1 than in volunteer 2 culture.

In the presence of DMSO, a delay in the growth of *C. coccoides* group was observed in both volunteer fecal fermentations. Thus, the lag period was extended in 9.7 and 22 h in the case of volunteers 1 and 2, respectively (Table 3). For human fecal suspensions of volunteer 1, the extension of the lag phase in 9.7 h of *C. coccoides* group in the presence of DMSO was accompanied by a delay of 10 h in the beginning of




Figure 5. UV spectra of the three tetrahydroxy urolithins ($[M - H]^-$ m/z 259): urolithin D, urolithin E, and urolithin M-6.

urolithin production. A similar tendency was observed in fecal suspension of volunteer 2, where the presence of DMSO lengthens both lag period (24.6 h) and time needed to achieve the maximum urolithin concentration. In contrast, the *Bacteroides/Prevotella* group achieved higher levels in the absence of DMSO (N_{max} : 7.2 log CFU/mL) than when DMSO was present (N_{max} : 6.8 log CFU/mL) in the case of volunteer 2 (Table 3).

DISCUSSION

The studies carried out so far with human biological fluids (urine and plasma) have allowed the identification of urolithin A, urolithin B, urolithin C, and isourolithin A.⁵ Other metabolites such as urolithins M-5, M-6, and M-7 had been reported in rat fecal samples and other animal materials.^{11–13}

Table 3. Growth Characteristics of Main Bacterial Groups Determined by q-PCR during Conversion of Ellagic Acid in Vitro Batch Fermentation with Human Fecal Suspensions^a

microbial groups	volunteer	fermentation medium + EA	lag time (h)	Gr _{max} (log CFU/h)	N _{max} (log CFU/mL)	correlation coefficient (R ²)
<i>Clostridium coccoides</i> group	1	without DMSO	0	0.12	6.08	0.99
		with DMSO	9.66** ^a	0.05** ^a	5.87	0.99
	2	without DMSO	2.59	0.10	5.04	0.99
		with DMSO	24.60** ^a	0.14	5.08	0.99
<i>Bifidobacterium</i> genus	1	without DMSO	0	0.15	7.91	0.92
		with DMSO	0	0.10	7.93	0.83
	2	without DMSO	no growth	no growth	no growth	
		with DMSO	no growth	no growth	no growth	
<i>Clostridium leptum</i> group	1	without DMSO	0	0.07	6.89	0.95
		with DMSO	0	0.04** ^a	6.60** ^a	0.87
	2	without DMSO	0	0.11	7.08	0.99
		with DMSO	0	0.09	7.12	0.97
<i>Lactobacillus/Leuconostoc/Pediococcus</i>	1	without DMSO	0	0.16	4.40	0.99
		with DMSO	0	0.18	4.34	0.99
	2	without DMSO	0	0.02	5.00	0.99
		with DMSO	0	0.05** ^a	4.96	0.99
<i>Bacteroides/Prevotella</i> group	1	without DMSO	0	0.12	6.40	0.96
		with DMSO	0	0.24** ^a	6.41	0.98
	2	without DMSO	0	0.21	7.17	0.99
		with DMSO	0	0.21	6.81** ^a	0.99
<i>E. coli</i> species	1	without DMSO	0	0.45	8.22	0.98
		with DMSO	0	0.40	8.30	0.99
	2	without DMSO	0	0.31	7.37	0.99
		with DMSO	0	0.27	7.83** ^a	0.99

^aFecal cultures consisted of volunteer 1 and 2 fecal suspensions in fermentation medium with 30 μ M ellagic acid (EA) dissolved or not with 1% DMSO. Significant differences in growth without and with DMSO addition. * ($P \leq 0.05$), ** ($P \leq 0.05$).

However, this is the first time that urolithin M-5, urolithin M-6, urolithin M-7, urolithin C, and the new identified metabolite urolithin E are reported to be produced by human gut microbiota. Urolithin M-6 (tetrahydroxy-urolithin) was detected with a maximum concentration between those of urolithin M-5 and two trihydroxy-urolithins (urolithin C and urolithin M-7) as could be expected for the catabolic sequence (Figure 6). This shows that these metabolites occur in vivo in human gut, although they can be quickly metabolized into urolithin A, which is the main metabolite found in plasma and urine. Analysis of human feces after the intake of ellagitannins and EA should be carried out in order to evaluate the occurrence of these urolithin metabolites in the gut conditions. Degradation of EA by human gut microbiota leads to similar urolithin metabolites in vitro (fermentation medium containing EA at 30 μ M) than in vivo (after consumption of 0.6 g walnuts/kg body weight per day for 5 days). Thus, both urolithin A and isourolithin A producers in vivo (volunteers 1 and 2) led to urolithin A and isourolithin A in vitro, respectively (Figure 4A,B). Despite the fact that urolithin metabolic profiles found in vivo were consistent with those found in vitro, further studies should be done in order to study the time course of the production of the two microbial metabolite patterns when EA containing foods are used in place of EA. This would elucidate the influence of other food constituents such as fiber on urolithin production rate.

In the human fecal culture of volunteer 1 (in vivo urolithin A producer), EA was 62% converted to urolithins at 36 h, while in volunteer 2 (in vivo isourolithin A producer) the conversion efficiency to urolithins was much lower (46% at 36 h). Taking into account the higher EA conversion efficiency of volunteer 1 and the fact that of the *C. coccoides* group achieved 1 log higher concentration in fecal fermentation sample of volunteer 1 than in volunteer 2, this suggests that this microbial group or other in vivo co-occurring bacteria could be involved in EA catabolism. In contrast, the lack of growth of *Bifidobacterium* in volunteer 2 could indicate that this microbial group is not involved in urolithin production.

Because of solubility problems of EA, this was alternatively dissolved in DMSO and incorporated into the medium at 1% v:v. In human fecal cultures where DMSO was added, different EA conversion was observed (28% and 74% in the case of volunteer 1 and 2, respectively). In the case of volunteer 1, the EA catabolism did not reach the whole transformation to urolithin A, and different intermediates were obtained. Urolithin M-5 was observed as the first step in urolithin production from EA to continue with urolithin M-6 and then urolithin M-7 to end with urolithin A but in much lower concentration than in the absence of DMSO. Furthermore, the addition of DMSO prevented the conversion of urolithin M-6 into urolithin C being one trihydroxy-urolithin (urolithin M-7), the only precursor found for the final catabolite (urolithin A)




Figure 6. Structures of the main urolithin metabolites detected and suggested catabolic routes of formation by gut microbiota.

(Figure 6). Therefore, the use of DMSO in fecal cultures offered the possibility of studying other metabolic intermediates between EA and urolithin A, such as urolithin E, another tetrahydroxy-urolithin precursor of urolithin M-7 (Figure 6). Despite the smaller urolithin production in presence of the DMSO, most of the EA was metabolized in the medium (24 μ M). This indicates that DMSO delayed urolithin producing bacteria development, allowing the action of other bacteria able to metabolize EA to simple organic acids such as phenyl acetic acid and phenylpropionic acid. However, further studies should be done in order to elucidate if higher abundance of *Bacteroidetes* and less abundance of *Firmicutes* such as *C. coccoides* are correlated with a lower urolithin production in vivo.

In the case of human fecal culture of volunteer 2, it seems that the catabolic route of EA to urolithins was modified by the effect of DMSO. In fact, much lower concentration of isourolithin A and much higher concentration of urolithin A were detected in the presence of DMSO (Figure 4D). This suggests that the microbial strains responsible for isourolithin A production in volunteer 2 grew with difficulty under these in vitro conditions. Furthermore, urolithin B was not detected, which suggests isourolithin A and not urolithin A is its only precursor and confirms this theory that was first proposed in cattle after the administration of oak leaves.¹² In addition, this is also in agreement with a previous in vitro study where urolithin B was only detected in those human fermentation cultures where isourolithin A were detected.¹⁷ A negative effect of DMSO on the growth of some bacterial groups was also observed, particularly on *C. coccoides* bacteria. A similar delay on growth of this bacterial group and the time needed to achieve the maximum urolithin concentration was observed, suggesting that this bacterial group could be involved in the EA metabolism of both volunteers.

In the literature, nothing is found about bacterial species or groups responsible for urolithin production from ellagitannins. In the present study, *C. coccoides* group (*C. coccoides* spp., *E. rectale* spp., and some species of *Ruminococcus* genus) from the *Firmicutes* phylum, or genera co-occurring in vivo with this group, are proposed as the microorganisms involved in urolithins production. *Clostridium* and *Eubacterium* genera are a common element of the metabolism of several phenolic compounds such as flavonoids.²¹ Given the phylogenetic association between these two genera, the association of them in the phenolic transformations is not surprising. The fecal samples corresponding to in vivo urolithin A and isourolithin A producers showed a higher abundance of either *C. coccoides* or *C. leptum* than that of the *Bacteroides/Prevotella* group. Recently, three robust clusters, referred to as “enterotypes”, which are not nation or continent specific have been identified. Assignment of an individual microbiome into a given enterotype is based upon the relative enrichment of that microbiome in one of three genera: *Bacteroides* (enterotype 1), *Prevotella* (enterotype 2), or *Ruminococcus* (enterotype 3).²² Further studies should be done in order to elucidate existent relations between human urolithins production capacity and relative abundance of bacteria either in the different phyla or in the different enterotypes.

Over the past decade, there has been enormous public and scientific interest in urolithins because of their proposed health-promoting effects, which, nonetheless, are far from being fully understood.²³ Future studies should take the newly identified urolithin metabolites into account when investigating EA absorption and metabolism as well as their physiologic effects in humans. Isolation and identification of bacteria strains implicated in urolithins production also require further investigation.

AUTHOR INFORMATION

Corresponding Author

*Phone: +34 968 396102. Fax: +34 968 396213. E-mail: mvselma@cebas.csic.es

Funding

This work has been supported by the projects CSD2007-00063 (Fun-C-Food; Consolider Ingenio 2010), CICYT AGL2011-22447, and BACCHUS (FP7 European Commission Grant Agreement, 312090). R.G.V. is holders of a JAE-DOC grant from CSIC (Spain) cofinanced by the European Social Fund (ESF)

Notes

The authors declare no competing financial interest.

REFERENCES

- (1) Gil, M. I.; Tomás-Barberán, F. A.; Hess-Pierce, B.; Holcroft, D. M.; Kader, A. A. Antioxidant activity of pomegranate juice and its relationship with phenolic composition and processing. *J. Agric. Food Chem.* **2000**, *48*, 4581–4589.
- (2) Larrosa, M.; García-Conesa, M. T.; Espín, J. C.; Tomás-Barberán, F. A. Ellagitannins, ellagic acid and vascular health. *Mol. Aspects Med.* **2010**, *31*, 513–539.
- (3) Landete, J. M. Ellagitannins, ellagic acid and their derived metabolites. A review about source, metabolism, function and health. *Food Res. Int.* **2011**, *44*, 1150–1160.
- (4) Cerdá, B.; Llorach, R.; Cerón, J. J.; Espín, J. C.; Tomás-Barberán, F. A. Evaluation of the bioavailability and metabolism in the rat of punicalagin, and antioxidant polyphenol of pomegranate juice. *Eur. J. Nutr.* **2003**, *42*, 18–28.
- (5) Cerdá, B.; Espín, J. C.; Parra, S.; Martínez, P.; Tomás-Barberán, F. A. The potent in vitro antioxidant ellagitannins from pomegranate juice are metabolized into bioavailable but poor antioxidant hydroxy-6H-dibenzopyran-6-one derivatives by the colonic microflora in healthy humans. *Eur. J. Nutr.* **2004**, *43*, 205–220.
- (6) Cerdá, B.; Tomás-Barberán, F. A.; Espín, J. C. Metabolism of antioxidant and chemopreventive ellagitannins from strawberries, raspberries, walnuts and oak-aged wine in humans: identification of biomarkers and individual variability. *J. Agric. Food Chem.* **2005**, *53*, 227–235.
- (7) Giménez, J. A.; González-Sarriás, A.; Larrosa, M.; Tomás-Barberán, F. A.; Espín, J. C.; García-Conesa, M. T. Ellagitannin metabolites, urolithin A glucuronide and its aglycone urolithin A, ameliorate TNF- α induced inflammation and associated molecular markers in human aortic endothelial cells. *Mol. Nutr. Food Res.* **2012**, *56*, 784–796.
- (8) Giménez, J. A.; Larrosa, M.; González-Sarriás, A.; Tomás-Barberán, F. A.; Espín, J. C.; García-Conesa, M. T. Intestinal ellagitannin metabolites urolithins and ellagic acid ameliorate cytokine induced inflammation and associated molecular markers in human colon fibroblasts. *J. Agric. Food Chem.* **2012**, *60*, 8866–76.
- (9) Seeram, N. P.; Schulman, R. N.; Heber, D. *Pomegranates. Ancient Roots to Modern Medicine*; CRC Press: Boca Raton, 2006; 244pp.
- (10) Pope, G. S. Isolation of two benzocoumarins from 'clover stone', a type of renal calculus found in sheep. *Biochem. J.* **1964**, *93*, 474–477.
- (11) González-Barrio, R.; Truchado, P.; Ito, H.; Espín, J. C.; Tomás-Barberán, F. A. UV and MS identification of urolithins and nasutins, the bioavailable metabolites of ellagitannins and ellagic acid in different mammals. *J. Agric. Food Chem.* **2011**, *59*, 1152–1162.
- (12) González-Barrio, R.; Truchado, P.; García-Villalba, R.; Hervás, G.; Frutos, P.; Espín, J. C.; Tomás-Barberán, F. A. Metabolism of oak leaf ellagitannins and urolithin production in beef cattle. *J. Agric. Food Chem.* **2012**, *60*, 3068–3077.
- (13) Ito, H.; Iguchi, A.; Hatano, T. Identification of urinary and intestinal bacterial metabolites of ellagitannin geraniin in rats. *J. Agric. Food Chem.* **2008**, *56*, 393–400.
- (14) Tulipani, S.; Urpi-Sarda, M.; García-Villalba, R.; Rabassa, M.; Lopez-Uriarte, P.; Bulló, M.; Jáuregui, O.; Tomás-Barberán, F. A.; Salas-Salvadó, J.; Espín, J. C.; Andrés-Lacueva, C. Urolithins are the main urinary microbial-derived phenolic metabolites discriminating a moderate consumption of nuts in free-living subjects with diagnosed metabolic syndrome. *J. Agric. Food Chem.* **2012**, *60*, 8930–8940.
- (15) González-Barrio, R.; Borges, G.; Mullen, W.; Crozier, A. Bioavailability of anthocyanins and ellagitannins following consumption of raspberries by healthy humans and subjects with an ileostomy. *J. Agric. Food Chem.* **2010**, *58*, 3933–3939.
- (16) Cerdá, B.; Periago, P.; Espín, J. C.; Tomás-Barberán, F. A. Identification of urolithin A as a metabolite produced by human colon microflora from ellagic acid and related compounds. *J. Agric. Food Chem.* **2005**, *53*, 5571–5576.
- (17) González-Barrio, R.; Edwards, C. A.; Crozier, A. Colonic catabolism of ellagitannins, ellagic acid, and raspberry anthocyanins: in vivo and in vitro studies. *Drug Metab. Dispos.* **2011**, *39*, 1680–1688.
- (18) Truchado, P.; Larrosa, M.; García-Conesa, M. T.; Cerdá, B.; Vidal-Guevara, M. L.; Tomás-Barberán, F. A.; Espín, J. C. Strawberry processing does not affect the production of urinary excretion of urolithins, ellagic acid metabolites, in humans. *J. Agric. Food Chem.* **2012**, *60*, 5749–5754.
- (19) Jaganath, I. B.; Mullen, W.; Lean, M. E. J.; Edwards, C. A.; Crozier, A. In vitro catabolism of rutin by human fecal bacteria and antioxidant capacity of its metabolites. *Free Radical Biol. Med.* **2009**, *47*, 1180–1189.
- (20) Baranyi, J.; Roberts, T. A.; McClure, P. A non-autonomous different equation to model bacterial growth. *Food Microbiol.* **1993**, *10*, 43–59.
- (21) Selma, M. V.; Espín, J. C.; Tomás-Barberán, F. A. Interaction between phenolics and gut microbiota: role in human health. *J. Agric. Food Chem.* **2009**, *57*, 6485–6501.
- (22) Arumugam, M.; Raes, J.; Pelletier, E.; Le Paslier, D.; Yamada, T.; Mende, D. R.; Fernandes, G. R.; Tap, J.; Bruls, T.; Batto, J. M. MetaHIT Consortium et al. enterotypes of the human gut microbiome. *Nature* **2011**, *473*, 174–180.
- (23) Espín, J. C.; Larrosa, M.; García-Conesa, M. T.; Tomás-Barberán, F. A. Biological significance of urolithins, the gut microbial ellagic acid-derived metabolites: the evidence so far. *Evidence-Based Complementary Altern. Med.* **2013**, *2013*, article ID 270418, DOI doi.org/10.1155/2013/270418
- (24) Furet, J. P.; Firmesse, O.; Gourmelon, M.; Bridonneau, C.; Tap, J.; Mondot, S.; Corthier, G. Comparative assessment of human and farm animal faecal microbiota using real-time quantitative PCR. *FEMS Microbiol. Ecol.* **2009**, *68*, 351–362.
- (25) Franks, A. H.; Harmsen, H. J.; Raangs, G. C.; Jansen, G. J.; Schut, F.; Welling, G. W. Variations of bacterial populations in human feces measured by fluorescent in situ hybridization with group-specific 16S rRNA-targeted oligonucleotide probes. *Appl. Environ. Microbiol.* **1998**, *64*, 3336–3345.
- (26) Manz, W.; Amann, R.; Ludwig, W.; Vancanneyt, M.; Schleifer, K. H. Application of a suite of 16S rRNA-specific oligonucleotide probes designed to investigate bacteria of the phylum *Cytophaga-Flavobacter-Bacteroides* in the natural environment. *Microbiology* **1996**, *142*, 1097–1106.
- (27) Huijsdens, X. W.; Linkens, R. K.; Mak, M.; Neuwissen, S. G.; Vanderbroucke-Grauls, C. M.; Savelkoul, P. H. Quantification of bacteria adherent to gastrointestinal mucosa by real-time PCR. *J. Clin. Microbiol.* **2002**, *40*, 4423–4427.