

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231538175>

# Potentiometric Study of the Coordination Tendency of 1,2-Bis(4-benzylpiperidine)glyoxime Toward Some Transition Metal Ions

ARTICLE in JOURNAL OF CHEMICAL & ENGINEERING DATA · NOVEMBER 2010

Impact Factor: 2.04 · DOI: 10.1021/jc1003803

---

CITATIONS

5

---

READS

12

3 AUTHORS, INCLUDING:



Faisal Obaidi

Al-Mustansiriya University

5 PUBLICATIONS 14 CITATIONS

SEE PROFILE



Hayati Sari

Gaziosmanpasa University

44 PUBLICATIONS 365 CITATIONS

SEE PROFILE

# Potentiometric Study of the Coordination Tendency of 1,2-Bis(4-benzylpiperidine)glyoxime Toward Some Transition Metal Ions

Faisal N. Al-Obaidi,<sup>\*,†</sup> Hayati Sari,<sup>‡</sup> and Mustafa Macit<sup>§</sup>

The University of Mustansiriyyah, Department of Chemistry, College of Science, Baghdad, Iraq, Gaziosmanpasa University, Science and Arts Faculty, Chemistry Department, 60250 Tokat, Turkey, and Ondokuzmayıs University, Department of Chemistry, 55139 Samsun, Turkey

A potentiometric study has been carried out to reveal the coordination properties of 1,2-bis(4-benzylpiperidine)glyoxime (BPG), via its reaction with certain transition metal ions,  $\text{Ni}^{2+}$ ,  $\text{Cu}^{2+}$ , and  $\text{Zn}^{2+}$ , and to determine the stability constants of the complexes formed. The experimental conditions were arranged to achieve all of the measurements and coordination in aqueous solution (mixed ethyl alcohol (10 %) and water (90 %)) at  $(25 \pm 0.1)^\circ\text{C}$  and an ionic background of  $0.1 \text{ mol}\cdot\text{dm}^{-3}$  NaCl. The overall stability constants  $\log \beta_n$  values of all species formed in solution together with the dissociation constants of the ligand were calculated by use of the SUPERQUAD computer program. The dissociation constants of the ligand BPG are 2.830, 6.066, 6.966, and 9.510. The chemical species present in the solution under our experimental conditions were demonstrated by the use of speciation diagrams.

## Introduction

The *vic*-dioxime ligands and their complexes have been the subject of detailed research for many years.<sup>1–3</sup> They are of immense significance in the field of coordination chemistry, and their complexes with transition metal ions are of considerable importance.<sup>4–10</sup> Their applications encompass an array of areas, from using them as catalysts in chemical processes<sup>2,11</sup> as well as their usage in analytical chemistry<sup>12</sup> and biochemistry.<sup>13</sup> They also have other applications in the field of medicine,<sup>14</sup> bioorganics, and electrochemistry.<sup>15,16</sup> The similarity between the *vic*-dioxime complexes and vitamin B<sub>12</sub> is an important issue which attracts much research.<sup>17</sup> *vic*-Dioximes may be of interest with regards to the photophysical and photochemical properties exhibited by some of their complexes with certain metal ions.<sup>15,18–20</sup>


The coordination between *vic*-dioxime ligands and metal ions is essentially a *N,N*-coordinate square planar containing a conjugated NCCN system;<sup>18,21–23</sup> however, other structures are also feasible.<sup>16</sup> The square planar structure forms a five-membered ring and causes the dissociation of the hydroxyl group, leaving one or both of the oxime groups in the form of an oximate, ending up with a rigid structure. Interhydrogen bonding will certainly operate in the case where aqueous solutions are used in experimental work. This would enhance the whole structure of the metal complex as a result of electronic attraction.

The aim of the present research is to assess the coordination capability of the 1,2-bis(4-benzylpiperidine)glyoxime (BPG) ligand (Scheme 1) by evaluation of the dissociation constants of the ligand and the stability of its metal complexes.

## Experimental Section

**Chemicals.** NaOH, KOH, NaCl, HCl, ethanol, 4-benzylpiperidine  $\text{CuCl}_2\cdot 2\text{H}_2\text{O}$ ,  $\text{ZnCl}_2\cdot 7\text{H}_2\text{O}$ , and  $\text{NiCl}_2\cdot 6\text{H}_2\text{O}$  were

Scheme 1



purchased from Merck and potassium hydrogen phthalate (KHP) from Fluka. All reagents were of analytical quality and were used without further purification. For the solutions,  $\text{CO}_2$ -free double-distilled deionized water was obtained with an aqua-MAX-Ultra water purification system (Young Lin Inst.); the resistivity was  $18.2 \text{ M}\Omega\cdot\text{cm}^{-1}$ .

**Synthesis of BPG.** BPG was synthesized as in the literature.<sup>24</sup> A solution of *anti*-dichloroglyoxime<sup>25</sup> (20 mmol: 3.14 g) in absolute ethanol (30 mL) at  $-10^\circ\text{C}$  was added to a solution of 4-benzylpiperidine (40 mmol: 7.01 g) in absolute ethanol (30 mL) dropwise over 1 h. The reaction mixture was stirred for 4 h at  $-10^\circ\text{C}$ . After adjusting the pH of the mixture to 6 to 7 with 1 % KOH solution at  $0^\circ\text{C}$ , the mixture was stirred for an additional 4 h. The mixture was filtered, and the filtrate was dried in a vacuum. The crude product was recrystallized from absolute ethanol. Yield, 58 %. Elemental analysis. Calculated (%): C, 71.86; H, 7.87; N, 12.89; Found (%): C, 71.95; H, 8.12; N, 12.98. <sup>1</sup>H NMR ( $\text{DMSO}-d_6$ ):  $\delta = 11.3$  (2H, s), 7.26–7.30 (10H, m), 2.90–3.00 (8H, m), 2.52–2.54 (4H, s), 1.86–1.90 (2H, m), 1.35–1.55 (8H, m) ppm. IR (KBr): [3240 (O–H), 1625 (C=N), 960 (N–O)]  $\text{cm}^{-1}$ .

**Titration Procedure.** Stock solutions ( $2.00\cdot 10^{-3} \text{ mol}\cdot\text{dm}^{-3}$ ) of the metal ions were prepared for  $\text{CuCl}_2\cdot 2\text{H}_2\text{O}$ ,  $\text{ZnCl}_2\cdot 7\text{H}_2\text{O}$ , and  $\text{NiCl}_2\cdot 6\text{H}_2\text{O}$  at  $25^\circ\text{C}$  and standardized with ethylenediaminetetraacetic acid (EDTA).<sup>26</sup> Other diluted solutions can be obtained simply by volumetric dilution. KHP was prepared as a buffer solution (0.05 M) and used for the calibration of the

\* Corresponding author. E-mail: fobaidi@uomustansiriyyah.edu.iq.

<sup>†</sup> The University of Mustansiriyyah.

<sup>‡</sup> Gaziosmanpasa University. E-mail: hsari@gop.edu.tr.

<sup>§</sup> Ondokuzmayıs University. E-mail: mmacit@omu.edu.tr.

**Table 1. Summary of the Experimental Parameters for the Potentiometric Stability Constants Measurements<sup>a</sup>**

system	BPG, with H <sup>+</sup> , Cu <sup>2+</sup> , Ni <sup>2+</sup> , and Zn <sup>2+</sup> in water
solution composition	[L] range/mol·dm <sup>-3</sup> : 0.001–0.002; [M] range/mol·dm <sup>-3</sup> : 0.001; ionic strength/mol·dm <sup>-3</sup> : 0.1 electrolyte NaCl
experimental method	pH-metric titration in range of pH 3 to 11; log β <sub>00-1</sub> – 13.78
T/°C	25.0
n <sub>tot</sub> <sup>b</sup>	250
n <sub>tit</sub> <sup>c</sup>	3
method of calculation	SUPERQUAD
titration system	Molspin

<sup>a</sup> M: Metal ion, L: ligand, β: overall stability constant. <sup>b</sup> Number of titration points per titration. <sup>c</sup> Number of titrations per metal ligand system.

combination pH electrode according to the method of Molspin.<sup>27</sup> The NaOH concentration (0.025 mol·dm<sup>-3</sup>) was standardized with primer standard KHP solution by a pH-metric titration. The HCl stock solutions used for obtaining pH 3.0 were prepared from concentrated HCl, and its concentration was determined with standard NaOH. All potentiometric titrations were carried out on solutions in 100 mL double-walled glass vessels using a Molspin automatic titration system, which interfaces to a PC, with a 10 cm<sup>3</sup> syringe and a SenTix 20 pH combination electrode (WTW, Weilheim), and the temperature was controlled at (25 ± 0.1) °C by circulating water from a constant-temperature bath. The cell was equipped with a magnetic stirrer and a tightly fitting cap which contained three holes for the combined electrode, nitrogen gas, and automatic buret. The pH was measured between pH 3.0 to 11.0 and under a nitrogen atmosphere. The ligand concentrations varied in the range (1.00·10<sup>-4</sup> to 2.00·10<sup>-4</sup>) mol·dm<sup>-3</sup>. The summary of the titration conditions is given in Table 1. The pH-metric data were used to find the dissociation and stability constants using the SUPERQUAD computer program.<sup>28</sup> The standard deviations (σ values) computed by SUPERQUAD refer to random errors.

## Results and Discussion

**Dissociation Constants of BPG.** The deprotonation equilibrium is as seen in the following equations (charges are omitted for simplicity).



and the deprotonation constants ( $K_n$ ) are given as;

$$K_n = [\text{LH}_{n-1}][\text{H}]/[\text{LH}_n]$$

Table 2 contains dissociation constant ( $\text{p}K_a$ ) values of the ligand under investigation. Four  $\text{p}K_a$  values are obtained for this ligand; two of them are associated with the oxime nitrogen atoms, and the other two are associated with hydroxyl groups. Comparing the  $\text{p}K_a$  values of the BPG ligand with that of 1,2-bis(dimethyl)glyoxime (DMG) shows to what extent the benzylpiperidine moiety in BPG affects the basicity. The  $\text{p}K_a$  of the oxime nitrogen atoms for BPG are 2.83 and 6.066, while that of dimethylglyoxime are 6.011 and 8.968.<sup>29</sup> Other values of  $\text{p}K_a$  of similar ligands are given in Table 2.<sup>30,31</sup> So, the benzylpiperidine moiety in the ligand gives significant electron withdrawing on the coordination center of the BPG. The sp<sup>3</sup> nitrogen of the piperidine might be the main cause of the decrease in basicity of the oxime nitrogen in BPG; as with its relatively high electronegativity, it tends to draw electron density


**Table 2.  $\text{p}K_a$  Values for BPG together with  $\text{p}K_a$  Values for Other Similar *vic*-Oximes, 1,2-Bis(4-benzylpiperazine)glyoxime (BBPGH2), 1,2-Bis(4-methylpiperazine)glyoxime (BMPGH2), 1-Benzyl-4-piperazineglyoxime (BPGO), and 1-Methyl-4-piperazineglyoxime (MPGO), Which Have Been Taken from Literature<sup>30,31</sup> ( $I = 0.1$  mol·dm<sup>-3</sup> NaCl, at (25 ± 0.1) °C)**


ligand	species	$\text{p}K_a$
BPG	LH <sub>4</sub>	2.830 ± 0.034
	LH <sub>3</sub>	6.066 ± 0.019
	LH <sub>2</sub>	6.966 ± 0.020
	LH	9.510 ± 0.013
BBPGH2	LH <sub>4</sub>	3.46
	LH <sub>3</sub>	5.89
	LH <sub>2</sub>	6.77
	LH	9.76
BMPGH2	LH <sub>4</sub>	2.91
	LH <sub>3</sub>	6.79
	LH <sub>2</sub>	7.97
	LH	10.00
BPGO	LH <sub>3</sub>	3.187
	LH <sub>2</sub>	7.035
	LH	9.790
MPGO	LH <sub>3</sub>	3.007
	LH <sub>2</sub>	7.620
	LH	9.564

away from the oxime group. This actually results in a reduction in the electron availability on the oxime nitrogen atoms and at the same time increases the dissociation of the hydroxyl group. In other words, the withdrawal property of benzylpiperidine has an impact on the hydrolysis of the hydroxyl group in the ligand, which would facilitate the release of protons. Protonation of the nitrogen atoms in *vic*-oximes, in acidic media, is expected to have very little influence on the acidity of the hydroxyl group, but coordination with metal ions will have an appreciable influence and ease the dissociation of the hydroxyl group.

$\text{p}K_a$  values for other similar *vic*-dioximes from the literature rationalize and support data obtained in this work.<sup>30</sup> The above results are in good agreement with our expectations on the extent of basicity of this ligand. The basicity of the ligand does not necessarily reflect the right coordination power of the ligand; in other words, the highly basic ligand does not always form the most stable complexes. One of the major factors that affects stability is steric hindrance, which sometimes even precludes coordination.

The titration curves of the ligand demonstrated in Figure 1 illustrate the extent of the acidity of the ligand. All measurements in this study were achieved in mixed water and ethanol (see the Experimental Section). Dissociation constants would certainly vary with the percentage of constituents of mixed solvents, which finally depends on electrostatic and nonelectrostatic factors.<sup>32–34</sup> All ligand species of BPG, which are expected to be formed under our experimental conditions, are

**Figure 1.** Titration curve of the BPG with 0.025 mol·dm<sup>-3</sup> NaOH at 25 °C in aqueous NaCl ( $I = 0.100$  mol·dm<sup>-3</sup> NaCl).




**Figure 2.** Species distribution diagram of BPG as a function of  $-\log[H]^+$ . L % = percentage concentration of species,  $I = 0.1 \text{ mol} \cdot \text{dm}^{-3} \text{ NaCl}$ , at  $(25 \pm 0.1)^\circ\text{C}$ .

presented in the distribution diagrams illustrated in Figure 2. The ligand species, which occurred in solution under our experimental conditions, are  $\text{LH}_4$ ,  $\text{LH}_3$ ,  $\text{LH}_2$ ,  $\text{LH}$ , and  $\text{L}$ . The experimental parameters and conditions used here are the same as those used to determine the stability constants.

The distribution diagram of the ligand assists the determination of the experimental conditions where the coordination with metal ions would take place, especially in terms of pH.

The bidentate nature of the ligand imposes a *cis*-structure on the ligand upon coordination; this actually designates the coordination power of the ligands and indicates its affinity toward metal ions. As with the *trans*-form the ligand cannot act as a bidentate; however, this form exhibits less steric strain.

**Stability Constants of Complexes.** The overall stability constants ( $\beta_{pqr}$ ) are defined as follows (charges are omitted for simplicity).




$$\beta_{pqr} = [\text{M}_p\text{L}_q\text{H}_r]/[\text{M}]^p[\text{L}]^q[\text{H}]^r$$

where M is the metal ion, L is the ligand, H is the proton, and  $p$ ,  $q$ , and  $r$  are the respective stoichiometric coefficients.

The mode of coordination of *vic*-dioxime has been mentioned in the Introduction, where the *N,N*-donor atom is engaged to form a five-membered chelating ring. Several possible species would be formulated in the solution depending on pH. In other words, the pH function regulates the subsistence of plausible species in solution.

Figure 3 shows all of the possible species in the Ni–BPG system; there are six types of Ni–BPG species which are tracked through our computer program as they formed in substantial and measurable amounts. The complexes traced at pH ranges between 4 and 6.5 are  $\text{NiH}_4\text{L}$  and  $\text{NiH}_3\text{L}$ . The dominant species in the solution at a pH between 7.5 and 8.5 is  $\text{NiH}_2\text{L}$ , and it comprises at pH 8 about 80 % of all of the species. Other species such as  $\text{NiHL}$  and  $\text{NiL}$  start to exist in significant amounts at a pH above 8.5. The species  $\text{NiH}_{-2}\text{L}$  starts to exist at pH values above 9.5.

There is a potential chance for the hydrolysis of the hydrated Ni(II) ion to take place in the pH range where the measurements were taken,<sup>35</sup> which gives rise to hydrolyzed species in solution. The formation of this type of partially hydrolyzed species not only depends on the pH but on the availability of water molecules on the solvated metal ion.<sup>36</sup>



**Figure 3.** Species distribution diagrams for the Ni–BPG system,  $I = 0.1 \text{ M NaCl}$ ,  $(25 \pm 0.1)^\circ\text{C}$ .

Potentiometric measurements showed that the existence of monocomplexes only, under the experimental conditions used, despite the ligand-to-metal ratio and the extent of the pH range, should allow the possibility of the existence of bis-complexes. The main cause for the preclusion of the bis-BPG complexes is steric hindrance, which arises from the fact that the BPG ligand is quite a bulky molecule and there will not be enough room in the coordination sphere of the metal ion for a second ligand to be accommodated either in a square planar pattern or any other constitution. So the tendency of *vic*-dioxime toward bis-complex formation will be precluded by steric requirements. In fact, steric hindrance might be raised even among the coordinate ligand itself; as upon coordination of the ligand, the two bulky moieties of 4-benzylpiperidine would not really be far away from each other. However, according to our data, it showed that all types of mono complexes formed sustain good stability within certain experimental conditions. The coordinated ligand may withstand the impact of the steric effect through its high flexibility, as the two moieties of 4-benzylpiperidine, which cause the steric effect, could arrange themselves in a way so that they are away from each other. So as long as there is a single covalent bond connected between the attachments, there will be some bending via this type of bond. This distortion in structure is expected to influence the whole complex.

The mono binary Ni(II)–BPG system exists in several forms, as has been mentioned above, and each single species originates from the preceding one as a result of proton liberation. So, there is a deprotonation process persisting throughout the intact coordination course. The deprotonation, which includes the dissociation of hydroxyl groups, occurs as a result of the formation of the coordinate bond, which causes electron withdrawal toward the metal ion and therefore lessens the dissociation of the hydroxyl groups. In other words, the coordination of the ligand with the metal ion will intensify the dissociation of the hydroxyl group or in fact any other acidic hydrogen in the system due to an increase in the electron charge flow toward the metal ion, as a result of coordinate bond formation.

Our results show that the species  $\text{NiH}_4\text{L}$ , with the stability constant value of 47.481, is the most stable of all of the complexes formed. The species  $\text{NiH}_4\text{L}$  forms no intrahydrogen bonding, as the ligand kept all hydrogen atoms intact; this is because both hydroxyl groups are quite far away from each other and they are unlikely to be linked by intrahydrogen bonding, but interhydrogen bonding is significantly feasible as long as an aqueous solution is used. The two hydroxyl groups will






**Table 3.** Stability Constant Data for the Complexation of  $\text{Cu}^{2+}$ ,  $\text{Ni}^{2+}$ , and  $\text{Zn}^{2+}$  with BPG at 25 °C in Aqueous NaCl ( $I = 0.100 \text{ mol} \cdot \text{dm}^{-3}$ );  $\beta_{pqr} = [\text{M}_p\text{L}_q\text{H}_r]/[\text{M}][\text{L}]^q[\text{H}]^r$ 

	$pqr^*$	Ni(II)	Cu(II)	Zn(II)
M(II)-BPG complexes	101	$16.345 \pm 0.026$	$14.554 \pm 0.051$	$15.40 \pm 0.020$
	111	$25.640 \pm 0.014$	$21.410 \pm 0.047$	
	121	$34.484 \pm 0.027$	$28.168 \pm 0.024$	$30.913 \pm 0.015$
	131	$41.448 \pm 0.043$	$34.405 \pm 0.023$	$38.301 \pm 0.019$
	141	$47.481 \pm 0.044$	$39.989 \pm 0.027$	$42.539 \pm 0.036$
	1-11		$6.904 \pm 0.049$	
	1-21	$-2.679 \pm 0.023$	$-2.766 \pm 0.114$	$-0.404 \pm 0.022$
	1-31		$-12.001 \pm 0.083$	$-10.215 \pm 0.043$

engage in such bonding via either their hydrogen or oxygen atoms. This type of bonding would expect to have no or very little influence on the stability of the complex; it is expected that the interhydrogen bonding is reasonably strong due to an increase in the acidity of the hydroxyl group through the course of the coordination; this of course would increase the species solubility in the solution. Accordingly no precipitate has been noticed, even when neutral species were formed. This case was justified by the fact that the expected concentrations of the complexes were relatively small.

The high tendency of the Ni(II) ion toward *vic*-dioxime is actually the main cause of why the nickel(II) complexes are more stable than the corresponding copper(II) complexes (see Table 3).<sup>37</sup> This of course does not comply with the Irving–William series.<sup>38</sup> In this case, the highest stability constants of the Ni(II) complexes over the Cu(II) complexes might be due to the possible formation of metal–metal bonding, as this type of bonding is confirmed in solid *vic*-dioxime complexes.<sup>39</sup>

Figure 4 shows the similarity in the distribution of Cu(II)–BPG species, with that of the Ni(II)–BPG system, but with eight species being tracked by the computer program. The species  $\text{CuH}_4\text{L}$  formed almost in the same pH range as  $\text{NiH}_4\text{L}$ , but with different distribution percentages at any pH. The percentage composition of  $\text{CuH}_4\text{L}$  at pH 5.5 is about 50 % out of the total concentration of Cu(II) ion, while the percentage of  $\text{NiH}_4\text{L}$  at the same pH is about 80 %. This does not necessarily imply the extent of the stability of the Ni(II) complex. The other species is  $\text{CuH}_3\text{L}$ , which is formed in considerable amounts after exceeding pH 5 up to pH 6.5 and with a maximum percentage occurrence of 50 % at a pH of 5.75. This species is expected to persist within the low acidic area so it still can hold an extra proton. Increasing the basicity of the solution caused deprotonation with the existence of other species such as  $\text{CuH}_2\text{L}$  and  $\text{CuHL}$  within a pH range from just above 6 up to 7.5 followed by  $\text{CuL}$  which formed within the pH range 6.75 to just above

**Figure 4.** Species distribution diagrams for Cu–BPG systems,  $I = 0.1 \text{ M NaCl}$ ,  $(25 \pm 0.1) ^\circ\text{C}$ .**Figure 5.** Species distribution diagrams for Zn–BPG systems,  $I = 0.1 \text{ M NaCl}$ .**Figure 6.** Structure of the  $\text{ZnH}_3\text{L}$  species.

8, where more loss of protons occurred. Increasing the pH will result in the formation of other hydrolyzed species;  $\text{CuH}_4\text{L}$  starts to form at approximately a pH of 7.5, and the maximum percentage of its composition will be about 85 % at a pH of 8.5. The other hydrolyzed species  $\text{CuH}_2\text{L}$  formed in the pH range between 9 to 10 and was found in smaller quantities when compared with other species. Its maximum concentration was at pH 9.5 with a percentage of just less than 20 %. After the pH exceeds the value of 9.5, the species  $\text{CuH}_3\text{L}$  starts to form in significant amounts, and after going beyond pH 10, this last species will be dominant in the solution.

In the M–BPH system, the moment the ligand was added to the metal ion, a certain species will come out as a result of reaction. So, the range of pH where the variety of species will start to form depends on the individual metal ion. The initial pH value of the diluted copper solution in general is less than the initial pH for the diluted nickel solution; therefore, we expected that copper complexes formed in pH will be less than that of nickel complexes. This probably may not hold for bis or tris-complexes and may depend on the structural shape of the bis or tris complexes.

Zinc(II)–BPG complexes seem to span over a wider range of pH values starting from pH 3.5 up to pH 11.5. Almost analogous species to that of the Ni(II) and Cu(II) systems are predicted to exist in solution. Stability constants of all Zn(II)–BPG species are higher than the corresponding Cu(II)–BPG species, which might suggest that the Zn(II) complexes have a tendency to adapt a tetrahedral structure. The extra stability is basically due to an increase in entropy as a result of alteration from an octahedral structure of  $[\text{Zn}(\text{H}_2\text{O})_6]^{2+}$  to a tetrahedral with a structure of  $[\text{Zn}(\text{BPG})]^{2+}$ . On the other hand, the stability of all Ni(II)–BPG complexes are higher than the corresponding Zn(II)–BPG complexes; this confirms the stability magnitude, caused by the assumed Ni–Ni bond in solution,<sup>39</sup> as mentioned previously. Also, the stability of Ni(II) complexes, in many cases, could be higher than the corresponding Zn(II) complexes without the establishment of metal–metal bonding.

The dominant species between pH 4.25 to pH 7.25 are  $\text{ZnH}_4\text{L}$ ,  $\text{ZnH}_3\text{L}$ , and  $\text{ZnH}_2\text{L}$ , with a higher percentage (more than 90

%) of  $\text{ZnH}_4\text{L}$  at a pH of less than 3.5. The species  $\text{ZnH}_3\text{L}$  forms about 90 % of the total Zn ion concentration at pH 6.5, while the highest percentage of the  $\text{ZnH}_2\text{L}$  species would be at 7.25 at about 45 %. The increase in deprotonation is quite gradual and increases with the increase of pH. The  $\text{ZnHL}$  species formed in the neutral to low basic area with a maximum formation of 40 % at pH 8. All other deprotonated species  $\text{ZnH}_2\text{L}$  and  $\text{ZnH}_3\text{L}$  are formed in the pH range between 8.25 and 10.5. Figure 5 shows the distribution diagrams for these species. The existence of the deprotonated species  $\text{ZnH}_3\text{L}$  occurs at a high pH (see Figure 5) as a result of the dissociation of the two hydroxyls on the oxime groups. The aquated  $\text{Zn(II)}$  ion at such a high pH tends to be hydrolyzed and exists in a form of hydroxy species, which will attach to the deprotonated BPG ligand as demonstrated in Figure 6.

## Literature Cited

- (1) Chaudhuri, P. Homo- and Hetero-polymetallic Exchange Coupled Metal-Oximates. *Coord. Chem. Rev.* **2003**, *243*, 143–190.
- (2) Taqui, K.; Ramachandraiah, G.; Kumar, S. Mononuclear bis(dimethylglyoximate) ruthenium(III) complexes with different appended axial groups: highly efficient catalysts for water oxidation. *J. Mol. Catal.* **1990**, *58*, 199–203.
- (3) Scrauzer, G. N.; Windgassen, J. On Hydroxyalkylcobaloximes and the Mechanism of Cobamide-Dependent Diol Dehydrase. *J. Am. Chem. Soc.* **1967**, *89*, 143–147.
- (4) Pahor, N. B.; Creos-Garlatti, R.; Geremia, S.; Randaccio, L.; Tauzher, G.; Zangrando, E. Syntheses, rate constants, and X-ray structures of alkylrhodoximes with  $\sigma$ -donating alkyl groups methyl, ethyl isopropyl. A comparison with the analogous alkylcobaloximes, a vitamin B12 model. *Inorg. Chem.* **1990**, *29*, 3437–3441.
- (5) Polson, S. M.; Hansen, L.; Marzilli, L. G. The First X-ray Structural Evidence Demonstrating Thiolate Coordination in an Organocobalt B12 Model Complex: Implications for Methionine Synthase. *Inorg. Chem.* **1997**, *36*, 307–313.
- (6) Pahor, N. B.; Forcolin, M.; Marzilli, L. G.; Randaccio, L.; Summers, M. F.; Toscano, P. J. Organocobalt  $\text{B}_{12}$  models: axial ligand effects on the structural and coordination chemistry of cobaloximes. *Coord. Chem. Rev.* **1985**, *63*, 1–125.
- (7) Yari, A.; Kakanejadifard, A. Spectrophotometric and theoretical studies on complexation of a newly synthesized vic-dioxime derivative with nickel(II) in dimethylformamide. *J. Coord. Chem.* **2007**, *60*, 1121–1132.
- (8) Pang, I.; Stynes, D. V. Kinetics of Reversible Binding of Carbon Monoxide and Benzyl Isocyanide to Ferrous Dimethylglyoxime Complexes. *Inorg. Chem.* **1977**, *16*, 590–594.
- (9) Pomposo, F.; Stynes, D. V. Iron Benzoquinone Dioxime Complexes. Electronic Effects on Axial Ligation to Ferrous Dioximes. *Inorg. Chem.* **1983**, *22*, 569–571.
- (10) Bakac, A.; Brynildson, M. E.; Espenson, J. H. Characterization of the Structure, Properties and Reactivity of a Cobalt(II) Macrocyclic Complex. *Inorg. Chem.* **1986**, *25*, 4108–4114.
- (11) Kijima, M.; Miyamori, K.; Nakamura, T.; Sato, T. Reaction of Organocobaloxime with Thiol under Irradiation. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 2549–2554.
- (12) Ghaedi, M.; Asadpour, E.; Vafaie, A. Simultaneous Preconcentration and Determination of Copper, Nickel, Cobalt, Lead, and Iron Content Using a Surfactant-Coated Alumina. *Bull. Chem. Soc. Jpn.* **2006**, *79*, 432–436.
- (13) Mehta, B. H.; Nagarkoti, B. Synthesis and characterization of some transition metal complexes of 2-naphthol oxime. *Asian J. Chem.* **2002**, *14*, 103–106.
- (14) Hall, I. H.; Bastow, K. F.; Warren, A. E.; Barnes, C. R.; Bouet, G. M. Cytotoxicity of cobalt complexes of furan oximes in murine and human tissue-cultured cell lines. *Appl. Organomet. Chem.* **1999**, *13*, 819–828.
- (15) Coskun, A.; Yilmaz, F.; Akgemci, E. G. Synthesis, characterization and electrochemical investigation of a novel vic-dioxime ligand and its some transition metal complex. *J. Inclusion Phenom. Macrocyclic Chem.* **2008**, *60*, 393–400.
- (16) Kilic, A.; Tas, E.; Gumgum, B.; Yilmaz, I. Synthesis, spectral characterization and electrochemical properties of new vic-dioxime complexes bearing carboxylate. *Transition Met. Chem.* **2006**, *31*, 645–652.
- (17) Finke, R. G.; Schiraldi, D. A.; Mayer, B. J. Towards the unification of coenzyme  $\text{B}_{12}$ -dependent diol dehydratase stereochemical and model studies: The bound radical mechanism. *Coord. Chem. Rev.* **1984**, *54*, 1–22.
- (18) Das, A. K.; Peng, S. M.; Bhattacharya, S. Chemistry of ruthenium with some dioxime ligands. Syntheses, structures and reactivities. *Polyhedron* **2001**, *20*, 327–335.
- (19) Kurtoglu, M.; Dagdelen, M.; Toroglu, S. Synthesis and biological activity of novel (E,E)-vic-dioximes. *Transition Met. Chem.* **2006**, *31*, 382–388.
- (20) Laranjeira, M. C. M.; Marusak, R. A.; Lappin, A. G. Driving force effects in proton coupled electron transfer. *Inorg. Chim. Acta* **2000**, *300*, 186–190.
- (21) Serin, S. New vic-dioxime transition metal complexes. *Transition Met. Chem.* **2001**, *26*, 300–306.
- (22) Gürsoy, S.; Cihan, A.; Koçak, M. B.; Bekaroğlu, Ö. Synthesis and Complexation of a Novel Soluble. *Monatsh. Chem.* **2001**, *132*, 967–972.
- (23) Kurtoglu, M.; Purtaş, F.; Toroglu, S. A new vic-dioxime ligand containing two azo substituents and its mononuclear nickel(II), cobalt(II), and copper(II) complexes; synthesis, characterization, spectroscopic and biological studies. *Transition Met. Chem.* **2008**, *33*, 705–710.
- (24) Steinkopf, W.; Jurgens, B. Zur Kenntnis aliphatischer Nitrokorper Uber Hydroxams. *J. Prakt. Chem.* **1911**, *83*, 453–470.
- (25) Brinzinger, H.; Titzmann, R. Notiz über einige halogenierte aliphatische oxime. *Chem. Ber.* **1952**, *85*, 344–345.
- (26) Jeffery, G. H.; Bassett, J.; Mendham, J.; Denney, R. C. *Vogel's Textbook of Quantitative Chemical Analysis*, 5th ed.; Longman: London, 1989.
- (27) Pettit, L. D. *Molspin Software for Molspin pH Meter*; Sourby Farm: Timple, Otley, U.K., 1992.
- (28) Gans, P.; Sabatini, A.; Vacca, A. SUPERQUAD: an improved general program for computation of formation constants from potentiometric data. *J. Chem. Soc., Dalton Trans.* **1985**, 1195–2000.
- (29) Al-Obaidi, F. N.; Macit, M.; Sari, H. The Potentiometric Study of Dissociation Constants of 1,2-Bis(dimethylglyoxime) and 1,2-Bis-(dichloro)glyoxime and Stability Constants of their Copper, Nickel and Zinc Complexes. *Int. J. Pure Appl. Chem.* **2010**, *5*, 87–93.
- (30) Sari, H.; Can, M.; Macit, M. Potentiometric and Theoretical Studies of Stability Constants of Glyoxime Derivatives and their Nickel, Copper, Cobalt and Zinc Complexes. *Acta Chim. Slov.* **2005**, *52*, 317–322.
- (31) Can, M.; Sari, H.; Macit, M. Potentiometric study of the new synthesized 1-Benzyl-4-Piperazineglyoxime and 1-methyl-4-piperazineglyoxime and their divalent metal complexes. *Acta Chim. Slov.* **2003**, *50*, 1–14.
- (32) Abraham, M. H.; Liszi, J. Calculations on ionic solvation-V. The calculation of partition coefficients of ions. *J. Inorg. Nucl. Chem.* **1981**, *43*, 143–151.
- (33) Feakins, D.; O'Neill, R. D.; Waghorne, W. E. Relative solvent transport numbers for ion constituents in dioxane + water and DMSO + water systems. *J. Chem. Soc., Faraday Trans.* **1983**, *79*, 2289–2298.
- (34) Latha, M. P.; Rao, V. M.; Rao, T. S.; Rao, G. N. Speciation of Ternary Complexes of Some Essential Metal Ions with L-Glutamic Acid and L-Methionine in 1,2-Propanediol–Water Mixtures. *Acta Chim. Slov.* **2007**, *54*, 160–165.
- (35) Huheey, J. E. *Inorganic Chemistry, Principles of Structure and Reactivity*, 3rd ed.; Harper International: Cambridge, U.K., 1983.
- (36) Burgess, J. *Metal Ions in Solution*; Ellis Horwood Limited, John Wiley & Sons: Chichester, U.K., 1978.
- (37) Gazda, D. B.; Fritz, J. S.; Porter, M. D. Determination of nickel(II) as the nickel dimethylglyoxime complex using colorimetric solid phase extraction. *Anal. Chim. Acta* **2004**, *508*, 53–59.
- (38) Irving, H.; Williams, R. J. P. The stability of transition-metal complexes. *J. Chem. Soc.* **1953**, 3192–3210.
- (39) Banks, C. V.; Barnum, D. W. Intermolecular Metal-Metal Bonds and Absorption Spectra of Some Nickel(II) and Palladium(II) Complexes of vic-Dioximes. *J. Am. Chem. Soc.* **1958**, *80*, 4767–4772.

Received for review April 19, 2010. Accepted November 6, 2010.

JE1003803