

J Am Chem Soc. Author manuscript; available in PMC 2014 July 14.

Published in final edited form as:

J Am Chem Soc. 2002 March 6; 124(9): 1862–1863.

# A Rationally Designed Genotoxin that Selectively Destroys Estrogen Receptor-Positive Breast Cancer Cells

Kaushik Mitra, John C. Marquis, Shawn M. Hillier, Peter T. Rye, Beatriz Zayas, Annie S. Lee, John M. Essigmann, and Robert G. Croy

Department of Chemistry and Division of Bioengineering and Environmental Health, Massachusetts Institute of Technology, Cambridge, MA 02139

The persistence of genetic damage produced by alkylating agents<sup>1</sup> as well as the antagonism of essential biochemical processes such as transcription can have lethal consequences for malignant cells.<sup>2</sup> Both mechanisms have been identified in studies to uncover the reasons for the efficacy of cisplatin in the treatment of several cancers.<sup>2a,3</sup> We describe a synthetic strategy to create bifunctional molecules that produce DNA adducts capable of binding the estrogen receptor (ER), which is aberrantly expressed in many breast cancer cells.<sup>4</sup> It is speculated that DNA adducts that form complexes with the ER will be poorly repaired in these cells because they are camouflaged from detection by DNA repair enzymes. Consequently, the DNA lesions persist. Furthermore, the DNA adducts would be expected to act as "molecular decoys" capable of displacing the ER from its natural targets and antagonizing its role in malignant growth. In healthy cells, where the abundance of the ER is minimal, no such ER-DNA adduct complexes will be present, and the cell should survive.<sup>5</sup>

In this report we describe the design and synthesis of compound 1, a bifunctional agent that can form covalent DNA adducts capable of binding the ER with high affinity and specificity. We show that 1 has selective toxicity toward ER+ breast cancer cells compared to ER- cells *in vitro*.

Compound 1 consists of a bis-chloroethyl aniline mustard as the DNA alkylating unit tethered to estradiol, the natural ligand for the ER. The site of substitution of estradiol in 1 was based on reports that relatively large alkyl groups can be attached at the 7α position with retention of high affinity for the ER<sup>6</sup>. The synthetic strategy for 1 is shown in Scheme 1. Compound 7, a key compound in the synthesis, was prepared by a modification of a published strategy. Briefly, 3 was functionalized with a 6-carbon chain at the 7-position in α-stereochemistry to provide the alkenyl steroid 4. Efficient reduction of the 6-oxo group in 4 was achieved with Et<sub>3</sub>SiH/BF<sub>3</sub>.Et<sub>2</sub>O; however, this treatment also caused the loss of 3,17-tetrahydropyranoxy (THP) groups producing diol 5. The 3,17-OHs of 5 were reprotected with THP groups to afford 6, followed by oxidation of the alkene at the terminus of the linker to provide alcohol 7. Steroid alcohol 7 was converted to bromide 8, which was subsequently allowed to react with a protected ethanolamine to give 9. Compound 9 was desilylated with tetrabutylammonium fluoride (TBAF) and converted to a carbonate

intermediate with *p*-nitrophenyl chloroformate. The carbonate was coupled to (N,N-bis-2-chloroethylaminophenyl)-propylamine that was prepared from chlorambucil via the Curtius reaction.<sup>8</sup> Deprotection of the product in HCl/methanol furnished **1**.

The affinity of **1** for the ER was first determined. Using a competitive binding assay<sup>9</sup> with  $[^3H]$ -17 $\beta$  estradiol, compound **1** was found to have a relative binding affinity (RBA) for the calf uterine ER of 30; RBA of estradiol=100.

Next, the ability of 1 to modify DNA covalently was investigated. Plasmid DNA was incubated with  $100 \, \mu M \, [^{14}C]$ - $1^{10}$  at 37 °C for up to 6 h. After unbound 1 was removed by phenol-CHCl $_3$  extraction and ethanol precipitation, the radioactivity associated with DNA was measured. The amount of radioactivity bound to DNA increased at a constant rate over the 6-h period indicating the formation of covalently bound 1 (see Supporting Information). Based on previous studies on DNA alkylation by nitrogen mustards,  $^{11}$  it is likely that covalent adducts of 1 are formed primarily at the N7 atom of guanines. To investigate the identity of the covalent adducts, 1-modified DNA was treated with 0.1N HCl, the major product from the digested DNA was isolated by reversed phase HPLC and analyzed by full scan electrospray mass spectrometry to yield a prominent molecular ion signal at  $M+H^+/z$  813.5051. This mass is consistent with a chemical structure in which one ethylene arm of the mustard of 1 is attached to guanine and the other arm contains a -OH substituted for the Cl atom.  $^{12}$ 

The affinity of DNA adducts of 1 for the ER was investigated using an Electrophoretic Mobility Shift Assay (EMSA). Substrates for this assay were prepared by reaction of 5'[<sup>32</sup>P]-labeled self-complementary oligonucleotide 5'-d(ATTATTGGCCAATAAT) with 1 for 4 hr at 37 °C. To assess quantitatively the level of covalent modification under these conditions, the DNA was treated with piperidine (1M, 90 °C, 1 h) and the products were separated on a denaturing 20% polyacrylamide gel (Figure 1A). This analysis found that 50% of the 16-mer oligo was cleaved, forming two products identical to those produced by the Maxam-Gilbert sequencing reaction for guanine (data not shown). EMSA was performed by combining the modified [32P]-labeled 16-mer with a 30 kD fragment of the ER containing the ligand binding site for estradiol (ER-LBD). <sup>13</sup> Analysis on a 4% nondenaturing polyacrylamide gel (0.5X TBE) revealed that approximately 50% of the 16mer formed a slowly migrating band (Figure 1B; lane 2). The amount of retarded band corresponded closely with the level of modification of the 16-mer as revealed by piperidine treatment. Furthermore, no slowly migrating band was seen with unmodified 16-mer (Figure 1B, lane 1) or with 16-mer modified with the nitrogen mustard chlorambucil (see Supporting Information). To examine the identity of the complex, increasing amounts of 17β-estradiol were added to the mixture containing the ER-LBD and 1-modified 16-mer. As shown in Figure 1B (lanes 3–8), addition of 17β-estradiol resulted in disruption of ER-LBD-DNA complex. These in vitro results indicated that ER-LBD selectively forms a complex with 1modified oligo, and raises the strong possibility of formation of similar complexes in cells.

Clonal survival assays were performed in ER+ and ER- breast cancer cell lines to determine if ER status affected the sensitivity of cells to the toxicity of 1. To reveal clearly the effect of DNA adducts, cells were exposed to 1 for 2 h after which fresh drug-free medium was added

to the cell cultures. This procedure minimized the possibility that unreacted 1 could function as a receptor antagonist and thereby inhibit growth or survival. Under these conditions 1 was found to possess a significantly lower  $EC_{50}$  in the ER+ cell line MCF-7 ( $EC_{50}$ =3.5 uM) as compared to the ER- cell line MDA-MB231 ( $EC_{50}$ =9.2 uM (Figure 2). In contrast to these results, chlorambucil under the same conditions did not show any difference in  $EC_{50}$  between these two cell lines (Figure 2).

The fact that ER status affected the  $EC_{50}$  of  ${\bf 1}$ , but not of chlorambucil, is consistent with the role of the ER as an effector of selective toxicity of  ${\bf 1}$ . Covalent  ${\bf 1}$ -guanine adducts have been identified in DNA isolated from treated cells (unpublished). Therefore,  ${\bf 1}$  is stable to cell culture conditions and can form DNA adducts in cells identical to those formed *in vitro*. We are currently investigating the role of  ${\bf 1}$ -DNA adducts *in vivo* to determine if they are less efficiently repaired in ER+ cells and to evaluate whether the association of the ER with DNA adducts of  ${\bf 1}$  is directly responsible for the greater sensitivity of MCF-7 cells.

## **Supplementary Material**

Refer to Web version on PubMed Central for supplementary material.

## **Acknowledgments**

We thank John Wishnok and W. Sara Gardinar-Stillwell for helpful discussions. We are grateful to the National Cancer Institute (1-RO1-CA77743-01) for support of this work, to the Susan G. Komen Foundation for a fellowship to KM (PDF00-000679), the National Institutes of Health for support of the NMR facility (1S10RR13886-01), and the BEH Mass Spectrometry Laboratory for use of their instruments.

### References

- (a) Dolan EM. Adv Drug Delivery Rev. 1997; 26:105–118.(b) Alaoui-Jamali M, Loubaba BB, Robyn S, Tapiero H, Batist G. Cancer Chemother Pharmacol. 1994; 34:153–158. [PubMed: 8194166] (c) Frankfurt OS, Seckinger D, Sugarbaker EV. Anticancer Res. 1993; 13:947–952. [PubMed: 8352564] (d) Frankfurt OS. Int J Cancer. 1991; 48:916–923. [PubMed: 1907257] (e) Ross WE, Ewig RAG, Kohn KW. Cancer Res. 1978; 38:1502–1506. [PubMed: 565677] (f) Hansson J, Lewenshon R, Ringborg U, Nilsson B. Cancer Res. 1987; 47:2631–2637. [PubMed: 3567896] (g) De Silva IU, McHugh PJ, Clingen PH, Hartley JA. Mol Cellular Biol. 2000; 20:7980–7990. [PubMed: 11027268] (h) Hansson J, Keyse SM, Lindahl T, Wood RD. Cancer Res. 1991; 51:3384–3390. [PubMed: 2054778]
- Examples include the "hijacking" of transcription factors by cisplatin DNA lesions (Zhai X, Beckmann H, Jantzen HM, Essigmann JM. Biochemistry. 1998; 37:16307–16315. [PubMed: 9819223] ) and hormonal antagonists (Lerner LJ, Jordan VC. Cancer Res. 1990; 50:4177–4189. [PubMed: 2194650] ).
- (a) Chu G. J Biol Chem. 1994; 269:787–790. [PubMed: 8288625] (b) Toney JH, Donahue BA, Kellett PJ, Bruhn SL, Essigmann JM, Lippard SJ. Proc Natl Acad Sci USA. 1989; 86:8328–8332. [PubMed: 2530581] (c) Brown SJ, Kellett PJ, Lippard SJ. Science. 261:603–605. [PubMed: 8342024] (d) Trimmer EE, Zamble DB, Lippard SJ, Essigmann JM. Biochemistry. 1998; 37:352–362. [PubMed: 9425057] (e) Huang JC, Zamble DB, Reardon JT, Lippard SJ, Sancar A. Proc Natl Acad Sci USA. 1994; 91:10394–10398. [PubMed: 7937961] (f) Jamieson ER, Lippard SJ. Chem Rev. 1999; 99:2467–2498. [PubMed: 11749487] (g) McANulty MM, Lippard SJ. Mutat Res. 1996; 362:75–86. [PubMed: 8538651] (h) Szymkowski DE, Yarema K, Essigmann JM, Lippard SJ, Wood RD. Proc Natl Acad Sci USA. 1992; 89:10772–10776. [PubMed: 1438274]
- 4. Fern M, Borg X, Johansson U, Norgren A, Olsson H, Ryd∂n S, Sellberg G. Acta Oncologica. 1990; 29:129–135. [PubMed: 2334566]

5. The feasability of this strategy has been demonstrated in a preliminary report.Rink SM, Yarema KJ, Solomon MS, Paige LA, Tadayoni-Rebek BM, Essigmann JM, Croy RG. Proc Natl Acad Sci USA. 1996; 93:15063–15068. [PubMed: 8986764]

- (a) Bucourt R, Vignau M, Torelli V, Richard-Foy H, Geynet C, Secco-Millet C, Redeuilh G, Baulieu EE. J Biol Chem. 1978; 253:8221–8227. [PubMed: 556354] (b) Bowler J, Lilley TJ, Pittman JD, Wakeling AE. Steroids. 1989; 54:71–99. [PubMed: 2815158] (c) DaSilva JN, van Lier JE. J Med Chem. 1990; 33:430–434. [PubMed: 2296033]
- 7. (a) Skaddan MB, Wuest FR, Katzenellenbogen JA. J Org Chem. 1999; 64:8108–8121. [PubMed: 11674724] (b) Tedesco R, Fiaschi R, Napolitano E. Synthesis. 1995:1493–1495.
- 8. Valu KK, Gourdie TA, Boritzki TJ, Gravatt GL, Baguley BC, Wilson WR, Wakelin LP, Woodgate PD, Denny WA. J Med Chem. 1990; 33:3014–3019. [PubMed: 2231599]
- 9. Korenman SG. Endocrinology. 1970; 87:1119-1123.
- 10. [<sup>14</sup>C]-**1** was obtained by reacting the carbonate intermediate with 4-(*N*,*N*-bis-2-chloroethylaminophenyl)-[1-<sup>14</sup>C] propylamine in step xii in Scheme 1; additional details in Supporting Information.
- (a) Kohn KW, Hartley JA, Mattes WB. Nucleic Acids Res. 1987; 15:10531–10549. [PubMed: 3697095] (b) Povirk LF, Shuker DE. Mutation Res. 1994; 318:205–226. [PubMed: 7527485] (c) Osborne MD, Wilman DEV, Lawley PD. Chem Res Toxicol. 1995; 8:316–320. [PubMed: 7766817]
- 12. DNA modified with  $d_4$ -1 (deuterium labeled 1) and analyzed using the same conditions for hydrolysis, HPLC and mass spectrometry revealed a molecular ion signal at M+H<sup>+</sup>/z 817.5. This result is consistent with  $d_4$ -1 attached to guanine as described. The preparation of  $d_4$ -1 is described in Supporting Information.
- 13. The 30 kD ER-LBD was obtained by expression of a gene fragment containing amino acids 282 to 595 of the human estrogen receptor in *Sf*21 cells using the baculovirus expression system (see Supporting Information) studies, gel shift experiments and toxicity experiments (18 pages). This information is available free of charge via the internet at <a href="http://www.pubs.acs.org">http://www.pubs.acs.org</a>.





Figure 1. A) Piperidine treatment of self-complementary deoxy oligonucleotide 5′-d(AATATTGGCCAATATT) treated with 1. Lane 1: untreated oligonucleotide, Lane 2: oligonucleotide + 200  $\mu$ M 1. B) Retarded mobility of oligonucleotide-1 in the presence of ER-LBD illustrated by EMSA, and disappearance of the retarded band by competition with estradiol. Lane 1: untreated oligonucleotide + ER– LBD, Lane 2: oligonucleotide modified by 200  $\mu$ M 1 + ER-LBD, Lanes 3–8: modified oligonucleotide + 10–300 nM estradiol.



**Figure 2.** Survival of MCF-7 (ER+) (+,0) and MD-MB231 (ER-) (&,#) cells treated with **1** (closed symbols) or chlorambucil (open symbols).

#### Scheme 1.

Conditions: i) DHP, pyridinium *p*-toluenesulfonate, reflux, 92%; ii) *n*-BuLi, KO*t*Bu, B(OMe)<sub>3</sub>, H<sub>2</sub>O<sub>2</sub>; iii) PCC, 59%; iv) 6-iodohexene, KO*t*Bu, Et<sub>3</sub>B, 47%; v) CH<sub>3</sub>COCl/MeOH, Et<sub>3</sub>SiH, BF<sub>3</sub>.Et<sub>2</sub>O, 74%; vi) DHP, pyridinium *p*-toluenesulfonate, 77%; vii) BH<sub>3</sub>.THF, KOH/H<sub>2</sub>O<sub>2</sub>, 66%; viii) methane sulfonyl chloride, LiBr, 86%; ix) Ph<sub>2</sub>P(O)NH-CH<sub>2</sub>-CH<sub>2</sub>-OtBDMS, NaH, cat. tetra-n-butylammonium bromide, 73%; x) TBAF, 73%; xi) *p*-nitrophenylchloroformate, DIEA; xii) [4-(3-aminopropyl)-phenyl]-bis-(2-chloroethyl)amine, DIEA; xiii) H+, 60%