pubs.acs.org/jcim

Comparative Analysis of Threshold and Tessellation Methods for Determining Protein Contacts


Jeremy Esque,**,†,† Christophe Oguey,† and Alexandre G. de Brevern‡

[†]LPTM, CNRS UMR 8089, Université de Cergy Pontoise, 2 av. Adolphe Chauvin, 95302 Cergy-Pontoise, France [‡]INSERM UMR-S 665, Dynamique des Structures et Interactions des Macromolécules Biologiques (DSIMB),

Université Paris Diderot, Paris 7, INTS, 6, rue Alexandre Cabanel, 75739 Paris Cedex 15, France



ABSTRACT: The 3D structure of a protein is the main physical support of a protein's biological function; 3D protein folds are primarily maintained through interactions between amino acids. Inter-residue contacts are essential for the stability of protein folds. Therefore, many methodologies in the fields of structure analysis, structure prediction, and structure—function relationships are based on residue contacts. The present study provides a comparative analysis of two approaches for determining contacts: the classical distance-threshold method and an application of Laguerre, or weighted Voronoi tessellation.



First, we examined mean contact distributions and their dependence on residue volumes, accessibility and hydrophobicity. In general, the different methods gave concordant results, although the method based on $C\alpha$ distances showed significant discrepancies with the all-atom tessellation method. We also analyzed preferential contacts between all amino acid species and studied the influence of protein chain length, the proximity of the residues along the sequence, and the secondary structure environment. Interestingly, the discrepancies between methods were occasionally large enough to substantially change the relative preferences of some contacts. Finally, a case study on disulfide bridges demonstrated the importance of the structural environment in determining contacts from tessellation. In conclusion, the tessellation method is more accurate because of its fine adaptation to local protein topology, with far-reaching implications for most contact-based prediction methods of protein folding.

■ INTRODUCTION

Three-dimensional protein structures are the physical supports of biological functions. Atomic interactions are essential for protein folding and for stabilizing the protein folds that make up three-dimensional structures. All amino acids share a common backbone, and their side-chains determine their physicochemical specificities. ^{1,2} Interactions between amino acids consist in forces (or energies), whereas contacts between amino acids describe the spatial proximity of residues. Contacts are defined by using the spatial coordinates of structure, whereas forces are most often indirectly inferred from their effects on structure, motion, chemical activity or any kind of response.

Inter-residue interactions can be classified into two main groups, those involving covalent (stable and strong) bonds and those involving weak bonds. A typical example of a covalent bond is the disulfide bridge that links two cysteines, which may be located far apart in the protein sequence, and that thereby stabilizes the structure.³ Weaker noncovalent forces, such as hydrogen bonds (H-bonds), van der Waals interactions, or hydrophobic effects, are also closely and commonly involved in folding and stabilizing protein structures. For example, the protein core is mainly maintained through nonpolar interactions; ⁴ some hydrophobic

units are thought to be potential nucleation sites during protein folding. Hydrogen bonds involve various donor groups, for example, N–H or O–H, and acceptor groups, for example, N or O, C–H or the π -system. Hydrogen bonds are responsible for the formation of repetitive secondary structural elements as 3_{10} -, α -, π -helices, β -sheets, and many turns. These types of bonds therefore involve short-range interactions and/or contacts along the sequence in both α -helices and β -turns, and longer range interactions in β -sheets. Secondary structures have been widely analyzed and used for predicting three-dimensional protein structures.

The term "contacts" covers many types of interactions, as mentioned above. Since contacts describe spatial proximity, the corresponding interactions are mostly local, for example, distance constraints caused by steric or electrostatic effects. Protein contacts are widely used to detect protein domains or protein subunits, ^{5,9} for example, the DDOMAIN, ¹⁰ PUU, ¹¹ DOMAK, ¹² 3Dee, ¹³ DIAL, or Protein Peeling ^{14,15} software programs. Information on protein contacts have proven to be useful for research and its

Received: May 17, 2010 Published: January 12, 2011



applications on protein folding and stability mechanisms, \$5,16-21 the development of inter-residue potentials, \$22,23 the identification of amino acid side-chain clusters with structural and/or functional roles, \$24-26 or the analysis of the intrinsic disorder of proteins. In particular, two interesting lines of research deserve to be highlighted. First, the relative frequency of noncovalent interactions has been used to define extracellular or intracellular proteins. Second, a good description of local protein structures, called structural alphabets, \$1,28,29 shows that these local protein structures, namely protein blocks, 30,31 are characterized by specific contact patterns. Second

In the past few years, much research has been dedicated to predicting inter-residue contacts. $^{33-40}$ Accordingly, 3D structures can be recovered from contact maps. $^{33-40}$ Due to their major importance, contact prediction methods have been the focus of recent meetings of Critical Assessment of Techniques for Protein Structure Prediction (CASP) experiments. $^{41-47}$ In spite of progress, contact map prediction and folding prediction remain major challenges. Interestingly, the prediction of β -sheets is the hardest case to solve. Long-range interactions and contacts are the most difficult to predict of all 3D protein features; they are also the main reason behind the failure of protein fold predictions. 19

The usual approach for defining contacts is based on a distance threshold, τ , between C α atoms (or pseudoatoms). In this study, we assess an alternative, tessellation approach based on Laguerre and Voronoi diagrams. Both diagrams partition space into convex polyhedra, one around each atom or residue, depending on the scale of interest. The polyhedral faces separating two contiguous polyhedra define contacts in a parameter-free manner. $^{49-51}$

Voronoi tessellations have been used to investigate a variety of protein properties, for example, protein—protein interactions, ⁵² standard volumes of residues. ⁵³ In a previous paper, we presented the usefulness of tessellation methods in protein structure analysis and particularly in the analysis of residue volumes. ⁵⁴

The present work presents a comparative analysis of contacts defined using the usual distance—threshold approach and those defined based on Laguerre tessellation. This study is therefore a direct continuation of previous research on protein contacts and tessellations. The distance method depends on a threshold τ defined more or less arbitrarily, while the tessellation approach does not rely on any metrical bound. In a previous study, the general differences of contact distributions were investigated, but the differences in residue pair assignments were not analyzed. Moreover, space partitioning was considered only at the residue scale. Here, the Laguerre tessellation was built at the atomic scale and the distance method was examined at both the atomic and residual ($C\alpha$ atom) scale. In tessellation methods, a realistic solvent must be added around proteins to account for exposed residues.

Contacts were evaluated on an updated, nonredundant databank of protein structures. Systematic analyses relating contacts to relative residue accessibility, protein size and proximity along the sequence or secondary structures were performed. Comparisons of the results of the different methods revealed a number of discrepancies depending on the method and the scale of the analyses. Scale differences caused greater discrepancies than methods themselves. Being the most frequent contacts, cysteine-cysteine contacts were examined in more detail.

In summary, this study highlights the usefulness of Laguerre tessellation in defining protein contacts and their potential applications, for example, reconstructing protein 3D structures from contact maps, predicting contacts, defining a mean force potential or refining structure models from the distance constraints using nuclear magnetic resonance data.

■ MATERIALS AND METHODS

Data Set. A nonredundant globular protein databank was built. It contained 818 polypeptide chains representing 187 433 residues. The protein data set was generated by the PISCES database 56,57 from files in the Protein Databank (PDB). The selected proteins had a high resolution (better than 2.5 Å); and only proteins sharing less than 25% of sequence identity were used. To ensure an unbiased study, no missing atoms or residues along the chain were allowed: all proteins were complete. All protein structures were treated using GROMACS 4.0.5 software and relaxed to near equilibrium through a short molecular dynamics run. During the simulation, the protein was frozen, that is, constraints were applied to limit protein movement. Further details on molecular dynamics runs are given below.

Addition of Water Molecules. The addition of water molecules was performed using GROMACS 4.0.5 software. $^{59-62}$ Each simulation was done under an OPLS-AA force field⁶³ with the TIP 4P water model. 64 The structure was immersed in a periodic water box neutralized with Na⁺ or Cl⁻ counterions. Each system was energy-minimized with a steepest-descent algorithm for 1000 steps. During the following steps, temperature and pressure were maintained constant at 300 K and 1 bar using the Berendsen algorithm.⁶⁵ The coupling time constants were $\tau_t = 0.1$ ps and $\tau_{\rm p}=0.5$ ps for temperature and pressure, respectively. An integration step of 2 fs was chosen and bond length was constrained using the LINCS algorithm. 66 A cutoff of 1.4 nm was used for nonbonded interactions in association with the generalized-reaction-field algorithm⁶⁷ for long-range electrostatic interactions using a dielectric constant. 65 For this study, the protocol is slightly different from our previous paper; 54 however no significant effect on the contact statistics was observed when replacing the old databank⁵⁴ by the one of this study and changing the energy relaxation procedure.

Tessellations for Proteins. A tessellation is a partition of space by a collection of polyhedra filling that space without overlaps or gaps. Laguerre tessellation is based on a set of sites, each defined by a point and a weight. In our case, sites are defined by atomic positions and weights (see below) of the system comprising the protein and the solvent. In the Laguerre tessellation, each polyhedron is convex and most often surrounds a single site. The shape of these polyhedra depends on the weights and mutual positions of neighboring sites. The Voronoi partition is a special case of the Laguerre tessellation where all the weights are equal. Further details on Laguerre diagrams and its dual (Delaunay tessellation) can be found in the literature. S4,68

Laguerre Weights. The Laguerre weights were set to $w = r^2$, where the atomic van der Waals radius r takes the default values in GROMACS, ⁶¹ that is, r = 1.5 for C, 1.05 for O, 0.4 for H, 1.1 for N, 1.6 for S (in Å). This simple relation was sufficient and optimal for our purposes. Dimensionally, the weight w of a site must be a length squared. The optimal value of the (dimensionless) proportionality constant between w and r^2 has previously been shown to be 1. ⁵⁴ This value minimizes the weighted sum of the residue volume variances.

Contact Definitions. Contacts are classically defined using a distance threshold, τ . This distance-threshold method can be considered at two scales. At the first coarse-grained scale, only

one point is retained for each residue and two residues are in contact if their Cas are separated by a distance of less than 8 Å. The contact numbers generated by this criterion are called $C\alpha$ contact numbers (CCN). At a finer atomic scale, two residues are in contact if they share a pair of atoms, one in each residue (not including H atoms), within 4.5 Å of each other. The corresponding count is the atomic contact number (ACN). In Laguerre and Voronoi tessellations, a contact between two residues occurs whenever two atoms (one of each residue) are separated by a common face in the tessellation. These contacts will be noted Laguerre contacts (LC) and Voronoi contacts (VC), and the corresponding counts LCN and VCN. Because proteins are polymeric chains, the immediate neighbors of any residue are systematically present in its spatial surrounding. In all subsequent analyses, these neighbors are discarded from the contact counts. 1,69 More precisely, all the neighbors at position $\pm 1, ..., \pm D/2$ in the sequence are excluded from the statistics. The parameter D was set to 6.1

Relative Frequencies. The preferential contacts sorted according to the amino acid species are specified by relative frequencies. The *relative frequency* of amino acid j in contact with amino acid i, rf_{ij} (also denoted $rf(i \rightarrow j)$), is the frequency of j as a neighbor of i normalized to its own frequency f_j^{DB} . In statistical terms, rf_{ij} is the proportion of j in the set of contacts of i over the proportion of j in the databank:

$$\mathrm{rf}_{ij} = \frac{f(j|i)}{f_j^{\mathrm{DB}}} = \frac{\#\mathrm{contacts}(ij)}{\#\mathrm{contacts}(i)} \cdot \frac{\#\mathrm{res}}{\#\mathrm{res}(j)} \tag{1}$$

with f(j|i)=(#contacts(ij))/(#contacts(i)) the frequency of amino acid j in contact with amino acid i, that is, the ratio between the number of contacts between i and j (#contacts(ij)), and the total number of contacts for amino acid i (#contacts(i)); f_j^{DB} =(#res(j))/(#res) is the frequency at which amino acid j occurs in the protein databank (i.e., the number of residues of amino acid species j, #res(j), over the total number of residues in the databank, #res).

Relative frequencies depend on the method used to determine the contacts. As for contact numbers (CN), CRF denotes relative frequencies obtained by the coarse-grained $C\alpha$ method, ARF by the all-atom distance method, LRF by the Laguerre tessellation method, and VRF by the Voronoi tessellation method.

To check the influence of different criteria, such as secondary structure or protein size, on contact numbers or relative frequencies, the databank was divided into subsets according to criteria defined in the analysis undertaken (see Results and Discussion). In each case, the differences were defined as $\mathrm{drf} = \mathrm{rf}_c - \mathrm{rf}$, the relative frequency evaluated on the specific subset c (rf_c) minus its counterpart evaluated over the entire databank (rf). All the investigated methods yielded specific differences dCRF, dARF, dLRF, and dVRF.

Software. The Laguerre or Voronoi tessellations were computed using VLDP (Voronoi Laguerre Delaunay Protein), a computer program developed at the Theoretical Physics and Modeling Laboratory (*Laboratoire Physique Théorique et Modélisation*, Cergy, France). The program builds a Delaunay tessellation and its Laguerre dual by incremental insertion of any set of sites. The surface accessibility of residues was evaluated using NACCESS (version 2.1.1). The secondary structures were assigned using DSSP software (version 2000, CMBI), according to three classes: α -helices (α , 3. α) and α -helices), β -strands (β -sheets) and coils (β -bridges, turns, bends, and coils). The molecular pictures were created using PyMol software.

Definition of Buried Residues. A residue was considered buried if its accessible surface area (ASA, given by NACCESS, ⁷⁰ probe radius of 1.4 Å) and polyhedral interface area (PIA, deduced from the Laguerre or Voronoi tessellation) were both evaluated at zero. PIA is defined as the residue surface area in contact with solvent, divided by its total surface area (facing solvent or other residues).

■ RESULTS AND DISCUSSION

Relationship between Residue Volumes and Contacts. Protein folds are maintained by atomic interactions between their residues. The amount of space occupied by each residue and their contacts both contribute to the proper conformation of the protein's structure. Residue occupancy is usually computed using the van der Waals volumes of the residue's atoms. An alternative method is to evaluate the volume as the sum of its atomic Laguerre polyhedra. Figure 1 shows the correlation between the average Laguerre volumes of residues and the mean contact numbers defined by Laguerre tessellation (LCN).

Considering all residues (exposed and buried, Figure 1a), the amino acid species are scattered around the least-squares regression line. The quality of the linear relationship is acceptable, with a Pearson correlation coefficient (PCC) value of 0.70 (ideal value would be 1). The regression line separates hydrophobic residues (aliphatic, aromatic), found above the line, from hydrophilic residues (polar, charged), found below the line. Thus, LCN reflects the hydrophobicity of residues, or their tendency to be buried. For instance, the hydrophobic character of cysteine (C) is because this residue is involved in disulfide bridges that occur mainly deep within the protein 3D structure. Unlike cysteine, lysine (K) is often found on the protein's surface and is thus located below the regression line.

When only buried residues are plotted, points are aligned close to the regression line, corroborated by a high PCC value close to 0.96 (Figure 1b). In the protein core, the residue assembly conforms to the packing of condensed matter, ⁷⁵ the contact number increases with residue size. Following the Lewis law, CN is proportional to surface area. ^{76,77}

The same analysis was performed on accessible (ASA >25%, Figure 1c) and nonaccessible residues (ASA <25%). Interestingly, these two subsets had high PCC values close to 0.95 (see Supporting Informaion 1). Taken separately, each subset of either accessible or entirely buried residues followed a linear relationship with very good fits, although the linear equations had different coefficients in each case (Figure 1). This explains why the data set incorporating both types of residues had a lower PCC (see Figure 1a).

For all three data sets, ACN and VCN showed results similar to LCN, that is, PCC values were close to 0.9 for accessible and nonaccessible residues separately and ranged from 0.6 to 0.7 when both types of residues were considered together (see Supporting Informaion 1). However, the C α distance method showed a different and peculiar pattern. Here, the set of accessible residues followed a linear relationship between CCN and van der Waals volumes, despite a relatively low PCC = 0.73. On the other hand, the buried residues showed a negative correlation (PCC = -0.76). As a result, when both types of residues were considered together, the correlation decreased to nearly zero (that is, PCC = -0.03, see Supporting Informaion 1). Hence, in the protein core, CCN overestimates the number of contacts of small residues and, conversely, underestimates contacts of large residues. This is clearly an artifact of the constant distance-threshold criterion.




Figure 1. Plot of mean residue contact number (LCN) against mean Laguerre volume (\mathring{A}^3). Mean values are taken over all residues (a), restricted to buried residues (ASA = PIA = 0) (b), or restricted to exposed residues (ASA > 25%) (c). Similar plots were obtained for Voronoi tessellations (not shown). Linear least-squares regression lines are indicated (dashed lines): (a) $y = 0.03 \,\mathring{A}^{-3} \, x + 1.13$; (b) $y = 0.05 \,\mathring{A}^{-3} \, x + 3.94$; (c) $y = 0.03 \,\mathring{A}^{-3} \, x + 1.97$.

Relationship between Residue Accessibility and Contact Number. Accessibility quantifies the exposure of residues to the solvent and this exposure is statistically related to the residue's proximity to the protein surface. The residue contact number reflects its environment and obviously depends on accessibility. Samanta et al. noted an exponential relationship between residue contact number and accessible surface area (ASA). Samanta's protocol is different from ours in that two residues have as many contacts as atomic pairs in contact, whereas, in our case, amino acid i is in contact with amino acid j if at least one atomic pair is in contact. This enumeration of contacts is more similar to the coarse-grained method and is more straightforward.

Figure 2 displays variations of ASA (or LPIA) according to LCN for two typical and similarly sized residues: Lysine (K) and Methionine (M). However, K is more hydrophilic than M.² Thus, K was found more often in the low LCN region reflecting characteristics of surface residues, whereas M had a higher propensity for larger LCN corresponding to the protein core (see distributions at the bottom of Figure 2). These profiles confirm some observations made by Samanta et al. The slope of the regression line is steeper for hydrophobic than for hydrophilic residues. The M profile reaches the asymptote at an LCN value of around 11 (Figure 2b), while K does not reach the asymptote before an LCN value of 16 (Figure 2a).




Figure 2. Mean accessible surface area (ASA) and mean Laguerre polyhedral interface area (LPIA) with respect to the Laguerre contact number (LCN). Two typical residues are illustrated: (a) lysine and (b) methionine. The lines are fits to the following function: y = a(1 - erf(bx)). Lower panels for both species give the residue population with respect to LCN to show its influence on variation in accessibility.

As functions of LCN, two patterns were distinguished from the ASA and LPIA variations: (i) a decreasing linear relationship for

low contact numbers, depending on residue hydrophobicity and (ii) an asymptote close to zero for high contact numbers. However, the LPIA curve was always below the ASA curve for low contact numbers, with the opposite relationship for high LCN values. For low LCN values, the dominance of ASA can be explained by the different normalization conventions: PIA is limited to 100% but ASA is not. 54 For high LCN values, where accessibility is low, the difference is mainly due to the probe radius parameter used in NACCESS, 1.4 Å. As noted in ref 54, this value (close to the average van der Waals radius of water) is fairly large compared to surface sinuosities. Consequently, ASA often equaled 0 even when water-residue contacts occurred in the tessellation, meaning that LPIA was not equal to zero. LPIA is more sensitive than ASA in detecting small areas of exposure to the solvent. Similar conclusions hold for the comparison of ASA with VPIA. The other contact methods led to similar observations; CCN and ACN showed similar curves with respect to variation in ASA. These trends were verified using other residues: the same analysis performed on arginine (R) and phenylalanine (F) gave similar results. Finally inter-residue contacts do not depend only on their volume, but also on their hydrophilicity and thus their accessi-

Mean Contact Numbers Deduced from Distance-Threshold and Tessellation Methods. As discussed above, the usual approach for predicting contacts or defining potentials of mean force is to set a distance threshold, τ . The literature contains a range of values for this parameter, depending on the data and scale (atom or residue) of interest. For contacts based on Cα, the cutoff distances used are typically 8, 10 or 12 Å.^{69,80,81} If all the atoms in an amino acid are considered and if contacts are defined in terms of minimal atom—atom distances, threshold values are lower: $\tau = 4$,^{79,82} 5.5,⁸³ and 4.5 Å.⁴⁹ The advantage of the Laguerre or Voronoi tessellation methods is that they do not need any threshold parameter. Space filling defines the neighborhoods, and thus adapts to the local geometry of residue packing.

Table 1 gives the mean residue contact number calculated using the four contact methods (CCN, ACN, LCN and VCN). The overall mean CCN and ACN values were very similar (4.7–4.8), as were those for LCN and VCN (5.6–5.7). On average, the tessellation methods resulted in 0.8–0.9 more contacts per residue than the distance methods. The overall averages can be rendered equal by adjusting the threshold to nearly 5 Å for the all-atom method. But we kept the value 4.5 Å, more standard in the literature. Some specific mean CCN and ACN values showed large discrepancies, while this was not the case for LCN and VCN. For instance, ACN and CCN differed by 2.71 for tryptophan (W), 1.61 for tyrosine (Y) and 1.82 for phenylalanine (F), all defined as aromatic residues (see Table 1).

Figure 3 displays the relative differences between the tessellation method (LCN) and the distance-threshold methods (CCN or ACN). For small residues, LCN values are smaller than CCN, but larger for large residues; a fairly linear progression interpolates between these extremes. For small residues, the $C\alpha$ method includes not only the nearest neighbors but also a few higher order ones, leading to a large overestimate. Similarly, the threshold approach misses some immediate neighbors for large residues, leading to a large underestimate. The effect of a fixed threshold value, independent of the residue size, can be clearly seen

In addition, other particularities can be highlighted. M and K are two equally sized residues, but they showed strong discrepancies. More hydrophilic than M, K showed a small difference

Table 1. Mean Residue Contact Number Calculated Using a Panel of Four Contact Methods^a

AA	N	CCN	σ	ACN	σ	VCN	σ	LCN	σ
A	15160	5.3	3.0	4.1	2.2	4.9	2.6	4.9	2.6
С	2266	6.6	2.8	5.7	2.1	6.8	2.5	6.9	2.5
D	11123	3.5	2.7	3.2	2.3	3.5	2.6	3.6	2.6
E	13189	3.3	2.4	3.3	2.3	3.6	2.6	3.7	2.6
F	7626	5.5	2.6	7.3	2.8	8.7	3.4	8.8	3.4
G	13336	4.9	3.3	3.1	2.0	3.7	2.4	3.8	2.4
Н	4212	4.6	2.8	4.9	2.8	5.7	3.2	5.8	3.2
I	11087	6.0	2.7	6.6	2.5	8.0	3.0	8.1	3.0
K	11252	3.7	2.4	3.6	2.3	4.0	2.6	4.1	2.7
L	17818	5.4	2.6	6.3	2.5	7.7	3.1	7.7	3.1
M	3449	5.4	2.7	6.4	2.8	7.6	3.4	7.7	3.4
N	8135	4.1	2.8	3.7	2.5	4.2	2.9	4.2	2.9
P	8596	4.3	2.9	3.6	2.4	4.4	2.9	4.5	2.9
Q	7035	3.9	2.6	3.9	2.5	4.4	2.9	4.5	2.9
R	9382	4.2	2.6	4.9	2.9	5.6	3.3	5.6	3.3
S	10869	4.5	3.0	3.5	2.3	4.1	2.7	4.2	2.7
T	10044	4.9	2.9	4.2	2.4	5.0	2.9	5.1	2.9
V	13601	6.2	2.8	5.8	2.4	7.1	2.9	7.2	2.9
W	2605	5.3	2.5	8.0	3.0	9.3	3.5	9.4	3.5
Y	6648	5.4	2.7	7.0	3.0	8.1	3.5	8.2	3.5
$\sum N$	187433								
avg(CN)		4.8		4.7		5.5		5.6	

^a The mean residue contact numbers and the corresponding standard deviations σ were computed for the distance-threshold methods (CCN, ACN) and the tessellation methods (VCN, LCN). N, residue count; $\sum N$, total residue count in the databank; avg(CN), global weighted average mean residue contact numbers.

between LCN and CCN, whereas this difference was great for M. The same pattern can be observed for R and F. Thus, physicochemical properties are also involved in setting contacts. K or R, are more often exposed and localized near the protein surface (mean ASA of K $\approx 53\%^{54}$). Hence, their environment is less dense and only incompletely filled by neighboring residues. The direct consequence is a decrease in the mean contact number. Finally, this comparison revealed a relationship between mean contact number and the propensity to be at or close to the surface. This conclusion can be visualized in Figure 2.

At the finer, atomic scale, discrepancies between LCN and ACN did not depend on residue volumes (see Figure 3, green bars). Both methods take residue size into account through the number of atoms composing each residue. In this case, differences arose due to residue shape and physicochemical properties. A group of small and/or hydrophobic residues (G, A, C, P, V, L, I, M) and some small polar residues (S and T) showed the greatest relative discrepancies, followed by the aromatic residues (H, F, Y, W), and finally by a group of polar or charged residues (D, N, E, Q, K, R). The discrepancies between LCN and ACN in the first two groups were comparable, whereas those of the third, polar-charged group were significantly smaller. Differences between LCN and ACN were always positive, indicating that some contacts found by the tessellation method cover a distance larger than 4.5 Å. Thus, the tessellation method (performed at atomic resolution) generally gives higher contact counts than the atomic threshold method and the observed discrepancies are partly correlated to residue hydrophobicity. As already mentioned, the global average




Figure 3. Discrepancies in relative mean contact number between the Laguerre and distance methods. The bar heights indicate the percent of relative differences (LCN-ACN)/LCN or (LCN-CCN)/LCN for each residue; residues are ordered according to increasing volume. ⁵⁴

difference can be reduced to zero by increasing the threshold to 5 Å; however the (dis)agreement was very poorly quantified by the PCC values. Indeed, the computations on a sample of thresholds ranging from 4.5 to 7.0 Å showed a constant PCC value close to 0.99, indicating that the mean LCN and ACN remained linearly related.

Analysis of Global Relative Frequencies. The relative frequencies (rf) give information on favored and unfavored contacts observed between residue pairs. Figure 4 shows the correlation of rf computed from Laguerre tessellation (LRF) with rf for the threshold approach at both scales, $C\alpha$ (CRF) and all atoms (ARF). The frequency of cysteine—cysteine contacts was always high because of the special nature of disulfide bridges: $CRF[C \rightarrow C] = 6.45$, $ARF[C \rightarrow C] = 5.00$, $VRF[C \rightarrow C] = 6.50$, and $LRF[C \rightarrow C] = 6.47$. Therefore, the corresponding points were isolated and are only shown in an inset in Figure 4. The points in the (CRF, LRF) plot are scattered around the linear regression line and the correlation is indeed moderate (PCC = 0.85, see Figure 4a), whereas a sharper linear correlation is observed between ARF and LRF (PCC = 0.98, see Figure 4b).

Laguerre and Voronoi tessellations gave highly similar results (see Supporting Information 2a). Overall, 207 pairs of amino acids (\sim 50%) had the same LRF and VRF up to the third decimal. The greatest differences occurred for [C \rightarrow C] and [Q \rightarrow H], with values of -0.05 and 0.04, respectively. Hence, the subsequent analyses focused on Laguerre tessellation.

The details of the rf discrepancies are displayed as matrices in Figure 5, with a resolution threshold of 0.2. For any pair of amino acids, the contact tendency is simply the rf value compared to the value 1. Contacts tend to be either overrepresented, rf > 1, or underrepresented, rf < 1 (see also Supporting Information 2b). Three kinds of rf changes can be distinguished: (1) positive enhancement: relative contact frequency determined by the Laguerre tessellation (LRF) is significantly increased compared to the distance-threshold method, the trend does not change (overrepresented or underrepresented); (2) negative enhancement: LRF is significantly decreased compared to the distance-threshold method, again without change in trends; (3) tendency inversion: over-represented contacts of one approach are found to be under-represented in another. The differences will be stated as




Figure 4. Correlation between relative frequencies given by tessellation and distance-threshold methods. LRF is plotted against (a) CRF and (b) ARF. The lines correspond to least-squares fit: (a) f(x) = 1.4x - 0.3 and (b) f(x) = 1.6x - 0.6. The insets show the complete data including the isolated cysteine—cysteine pair.




Figure 5. Discrepancies in relative frequency between the tessellation and distance-threshold methods. The rf differences are given as matrices indexed by the amino acid species (a) LRF-CRF and (b) LRF-ARF. The color key of frequency differences is provided.

variations of LRF with respect to ARF or CRF, considered as the reference. Among the 148 changes (37% of the matrix entries) found in the LRF-CRF matrix (see Figure 5a), 87 were positive enhancements (higher LRF values), 24 were negative enhancements and 37 were inversions (28 negative and 9 positive changes, representing nearly 10% of all the contacts.). Reductions (LRF lower) mainly occur with small residues (A, G, S). The contacts of aromatic residues (W, Y, F) and of some aliphatic/hydrophobic residues (M, L, I) were enhanced by LRF. These results corroborate those found for mean contacts (see above); the Ca distance method overestimated the contacts of small residues and, conversely, underestimated those of bulky residues, such as aromatic residues. As expected, the negative inversions (LRF < CRF) were observed mainly for contacts involving A and G residues (see Table 2). Interestingly, the positive inversions (LRF > CRF) were mainly observed for contacts involving arginine (R): LRF-CRF = 0.64 for $[D \rightarrow R]$ and 0.64 for $[E \rightarrow R]$. The enhanced LRF is well explained by the electronic attraction between the positively charged R and negatively charged D (or E), but the distances involved in those contacts are sometimes too large to be included in CRF.

Figure 5b displays the differences between LRFs and ARFs. Among the 84 changes (21% of all contacts), 65 are positive enhancements compared to 19 negative enhancements (lower

Table 2. Inversion Cases in the Comparison of LRF with CRF for the Whole Databank^a

amino acid pairs	LRF	CRF	LRF-CRF
$[A \rightarrow G]$	0.73	1.05	-0.32
$[C \rightarrow G]$	0.73	1.04	-0.31
$[D \rightarrow A]$	0.77	1.02	-0.25
$[D \rightarrow C]$	0.93	1.19	-0.26
$[D \rightarrow G]$	0.77	1.17	-0.40
$[D \rightarrow S]$	0.89	1.09	-0.20
$[E \rightarrow A]$	0.78	1.04	-0.26
$[F \rightarrow A]$	0.82	1.02	-0.20
$[G \rightarrow G]$	0.93	1.30	-0.37
$[H \rightarrow A]$	0.77	1.01	-0.24
$[H \rightarrow G]$	0.72	1.10	-0.38
$[I \rightarrow A]$	0.90	1.15	-0.25
$[L \rightarrow A]$	0.93	1.23	-0.30
$[M \rightarrow A]$	0.92	1.15	-0.23
$[M \rightarrow G]$	0.65	1.03	-0.38
$[N \rightarrow G]$	0.78	1.10	-0.32
$[P \rightarrow A]$	0.83	1.06	-0.23
$[P \rightarrow G]$	0.77	1.17	-0.40
$[P \rightarrow S]$	0.83	1.03	-0.20
$[Q \rightarrow A]$	0.84	1.06	-0.22
$[R \rightarrow A]$	0.78	1.02	-0.24
$[R \rightarrow G]$	0.77	1.05	-0.28
$[S \rightarrow G]$	0.80	1.17	-0.37
$[T \rightarrow G]$	0.74	1.07	-0.33
$[V \rightarrow A]$	0.97	1.20	-0.23
$[W \rightarrow A]$	0.80	1.00	-0.20
$[W \rightarrow G]$	0.65	1.03	-0.38
$[Y \rightarrow A]$	0.79	1.00	-0.21
$[D \rightarrow W]$	1.64	0.97	0.67
$[E \rightarrow F]$	1.17	0.97	0.20
$[G \rightarrow R]$	1.15	0.91	0.24
$[I \rightarrow W]$	1.49	0.93	0.56
$[N \rightarrow R]$	1.15	0.86	0.29
$[P \rightarrow R]$	1.19	0.90	0.29
$[Q \rightarrow R]$	1.21	0.98	0.23
$[R \rightarrow R]$	1.12	0.84	0.28
$[S \rightarrow R]$	1.12	0.87	0.25
0 D 1 1 C . 1 1 1	. 1	(IDE CDE): 4	

^a Boldface indicates the largest values (LRF or CRF) in the comparison.

LRF values). No inversions were observed. As expected, the LRF vs ARF differences were weaker than those between LRF and CRF, both in number and amplitude. With the exception of cysteine—cysteine contacts, which showed a difference of 1.47, the largest differences ranged from -0.24 to 0.49 for LRF vs ARF against -0.40 to 0.78 for LRF vs CRF. The positive changes mainly involved the contacts of aromatic residues, particularly W, but also Y, F, and aliphatic residues as L, M, or I.

The contacts between cysteines depart greatly from the other pairs of contacts. First, their relative frequency differences were LRF-CRF [C \rightarrow C] = 0.02 and LRF-ARF [C \rightarrow C] = 1.47. This result can be partly attributed to the fact that disulfide bonds can form between cysteines, whose C α distances range from 4.2 to 7.5 Å. The tessellation method is able to find contacts between

two cysteines separated by more than 4.5 Å. Thus, their contact numbers are equivalent to those found by the C α method (with threshold of 8 Å), whereas the all-atom threshold at 4.5 Å fails to detect some of them.

The details of the relative frequencies reveal compensations in the contributions to the mean residue contact numbers. For instance for D, the relative difference LCN vs CCN was 0.42 (see Figure 3) and is the sum of negative (A, C, G, S) and positive differences (H, K, R, W, Y) from Figure 5a. In other words, the mean contact counts sometimes even out sharper discrepancies revealed only when the contacts are sorted according to the species of both partners, as in relative frequencies.

To obtain a more accurate view, contacts were sorted (see Figure 6) to determine which contacts were shared or different between methods. Three categories were distinguished: (1) contacts found in both the Laguerre and threshold methods, (2) contacts specific to the Laguerre tessellation method, and (3) contacts specific to the distance-threshold method. In the bar graph given in Figure 6, the height of each bar represents the total contact count for the corresponding amino acid; the hatched section of the bar represents the contacts found by both methods. The nonhatched part of the bar corresponds to contacts found by only one of the two methods. In Figure 6a, of all the contacts found by the Cα threshold method, a proportion ranging from 16 to 34% depending on the amino acid, were only found by the Cα threshold method. The remainder, ranging from 66 to 84% of CCN, was observed for both (Cα and Laguerre) methods. Common contacts represented 47 to 84% of LCN, the contacts exclusively found for Laguerre tessellations, represented from 16 to 53% of LCN.

Comparison of LCN with ACN (see Figure 6b) shows that the Laguerre-specific contacts ranged from 12 to 20%, compared to 0.4 to 3.2% of ACN for the contacts specific to the all-atom distance method. These results demonstrate that both methods at the atomic scale (Laguerre tessellation and all-atom distance) share a larger set of common contacts than the Laguerre tessellation with the $C\alpha$ threshold method.

Analysis of Relative Frequencies According to Protein Size. The protein fold depends on the length of the protein chain; protein size may therefore act on (un)favored contacts. We defined four classes of protein size (L being the number of residues in the protein chain), L < 150, L = 150-250, L = 251-400, and L > 400, as proposed by Brocchieri et al.; ⁸⁵ and we examined the differences (dLRFs = LRFs - LRF), where LRFs is the Laguerre relative frequency calculated over the subset of proteins belonging to class s (size) whereas LRF is calculated over the entire databank (see Materials and Methods section).

To discern significant changes due to the contact method, we focused on the sets of amino acid pairs that satisfied the following criteria: (1) dLRF < 0.2 and dCRF (or dARF) > 0.2; (2) dLRF > 0.2 and dCRF (or dARF) < 0.2; (3) dLRF < -0.2 and dCRF (or dARF) < -0.2 and dCRF (or dARF) < -0.2. Only the most striking changes are listed in Table 3; the selected amino acid pairs and the corresponding values of dLRF, dCRF, and dARF are given for each protein size class.

Comparing dLRF and dCRF (or dARF), the greatest number of discrepancies was observed for small proteins (L < 150). On average, small proteins had a larger conformational variety, with a smaller proportion of well-characterized secondary structures; therefore, more discrepancies may be expected in this class. For small proteins, dCRF differed from dLRF mainly for contacts involving bulky residues, such as methionine (M) or tryptophan




Figure 6. Contacts found in the Laguerre and distance-threshold methods. The total contact numbers of each amino acid species, computed over the whole databank, are displayed as bar graphs for (a) the Laguerre vs Cα distance methods and (b) the Laguerre vs all-atom distance methods. The hatched portion of the bars represents the contacts common to both methods. Conversely, for each residue, the remaining percentages give the proportion of contacts found exclusively by each method (solid-colored bars without hatching).

(W), but also with hydrophobic residues, such as cysteine (C), glycine (G), histidine (H), isoleucine (I), and valine (V). The greatest discrepancy was observed for $[M \to M]$, with a dLRF value of 0.46 compared to a dCRF valued of -0.05. Regarding dARF, the differences with dLRF were found for contacts with aromatic residues (F and W) and hydrophobic amino acids (C, G, H, I, S, and T). The greatest discrepancy, observed for a dLRF value of -0.3 and a dARF value of 0.0, was for the $[H \to W]$ contact. Interestingly, for proteins including 150-250 amino acids, the selected changes involved contacts with only three main amino acid species (C, M, and W), comparing dLRF with either dCRF or dARF. In the third protein size class (251-400),

four amino acids were affected by changes in contact definition: C, H, I, and M. Finally, in the last class, cysteine (C) was involved in five of the seven recorded changes.

Globally, a linear relation was found between the drf's (such as dLRF and dCRF or with dARF, see Supporting Information 3). In order of increasing size, the (dLRF, dCRF) PCC for the four protein classes were 0.91, 0.82, 0.98, and 0.89, respectively. For (dLRF, dARF), the PCC were 0.91, 0.76, 0.95, and 0.88, respectively.

Analysis of rf According to Distance along Sequence. Among the possible types of interactions, two major types may be distinguished: short- and long-range interactions. 86,87 Long-range

Table 3. Influence of Protein Size on Relative Frequencies^a

			_
	<150		
amino acid pairs	dLRF	dCRF	dARF
$[C \rightarrow H]$	-0.22	-0.10	-0.27
$[C \rightarrow I]$	-0.27	-0.26	-0.12
$[C \rightarrow P]$	0.05	0.07	0.22
$[C \rightarrow V]$	-0.23	-0.19	-0.20
$[D \rightarrow G]$	-0.10	-0.21	-0.07
$[D \rightarrow H]$	-0.27	-0.10	-0.26
$[F \rightarrow C]$	0.22	0.31	0.11
$[F \rightarrow F]$	0.29	0.21	0.16
$[G \rightarrow G]$	-0.25	-0.37	-0.19
$[H \rightarrow K]$	0.10	0.13	0.29
$[H \rightarrow T]$	-0.20	-0.22	-0.09
$[H \rightarrow W]$	0.30	0.24	0.00
$[K \rightarrow C]$	0.20	0.18	0.12
$[M \rightarrow F]$	0.12	0.16	0.31
$[M \rightarrow I]$	0.06	0.24	-0.01
$[M \rightarrow M]$	0.46	-0.05	0.25
$[M \rightarrow W]$	-0.20	0.12	0.06
$[N \rightarrow S]$	-0.19	-0.21	-0.01
$[Q \rightarrow C]$	0.13	0.09	0.29
$[R \rightarrow W]$	-0.16	-0.04	-0.21
$[S \rightarrow C]$	0.38	0.41	0.19
$[S \rightarrow H]$	-0.17	-0.21	-0.20
$[S \rightarrow N]$	-0.23	-0.20	-0.02
$[T \rightarrow C]$	0.14	0.22	0.19
$[T \rightarrow H]$	-0.28	-0.23	-0.15
$[W \rightarrow M]$	-0.24	-0.01	-0.19
$[W \rightarrow T]$	-0.10	-0.22	0.01
$[W \rightarrow W]$	-0.24	-0.08	-0.36
$[Y \rightarrow I]$	0.16	0.09	0.20
	150-250		
amino acid pairs	dLRF	dCRF	dARF
$[C \rightarrow W]$	-0.16	-0.23	-0.51
$[H \rightarrow C]$	0.17	0.24	0.14
$[K \rightarrow W]$	0.20	0.12	0.13
$[M \rightarrow C]$	0.04	0.21	-0.13
$[M \rightarrow M]$	-0.02	-0.18	-0.25
$[M \rightarrow C]$	-0.09	-0.26	-0.24
$[W \rightarrow W]$	0.14	0.18	0.27
	251 400		
amino acid pairs	251-400 dLRF	dCRF	dARF
$[C \rightarrow H]$	0.15	0.09	0.20
$[C \rightarrow I]$	0.21	0.21	0.16
$[H \rightarrow C]$	0.14	0.08	0.22
$[I \rightarrow C]$	0.18	0.20	0.15
$[M \rightarrow M]$	0.11	0.30	0.19
amino acid pairs	>400 dLRF	dCRF	dARF
$[C \rightarrow M]$	0.28	0.09	0.22
$[C \rightarrow R]$	0.20	0.19	0.20

Table 3. Continued

$[H \rightarrow S]$	0.09	0.20	0.03
$[M \rightarrow C]$	0.14	0.00	0.25
$[Q \rightarrow C]$	-0.11	-0.21	-0.07
$[S \rightarrow C]$	-0.21	-0.16	-0.12
$[Y \rightarrow W]$	-0.17	-0.11	-0.25

^a Four classes of protein size are considered: <150, 150–250, 251–400, >400 (residue number). dLRF = LRFs – LRF is the difference between the average restricted to proteins of the specified size and the global average, computed using the Laguerre method. dCRF and dARF are defined analogously for the distance methods. The listed amino acid pairs were selected according to one of the criteria 1, 2, 3, or 4 in Analysis of Relative Frequencies According to Protein Size section.

interactions are essential for the onset and prediction of protein folding. 19,40 To investigate the dependence on separation along the sequence, we defined three zones of distance along the protein sequence: near (5-20 residues), far (21-50 residues)and very far (>50 residues) as previously proposed. 85 As above, a threshold of 0.2 (or -0.2) was chosen to characterize significant changes between dLRF and dCRF (or dARF). Table 4 summarizes those differences for all three contact methods. The comparison of dLRF with dCRF (or dARF) shows that the main discrepancies occurred at very far contacts. A simple hypothesis would relate this pattern to the fact that very far contacts preferentially involve β -sheets and some loops. We found small residues (C, G, P), some charged (E, K), and aromatic residues (F, W and Y), were involved in the differences between methods. Among the large discrepancies, the following amino acid pairs were the most interesting ones: $[D \rightarrow W]$, $[E \rightarrow R]$, and $[W \rightarrow W]$. They all showed a dLRF value of >0.2 whereas the dCRF value was <0.1. Regarding dARF, the Laguerre contact excess dLRF mainly differed from dARF for contacts involving hydrophobic residues (F, H, M, Y and W) and Cysteine (C). The greatest discrepancy was for $[W \rightarrow W]$ pairs, with a dLRF value of 0.26 and a dARF value equal to -0.05.

For the other two distance zones, the selected partners were slightly different. For *near* contacts along the sequence, the discrepancies between dLRF and dCRF involved contacts with small, aromatic and hydrophobic residues (C, T, V, W, and Y). The comparison of dLRF with dARF showed discrepancies for contacts mainly involving C, but also G, V, I. For the *far* contacts, the set of residues involved in discrepancies was more heterogeneous, for example, P, C, W, Q, M, and H for differences between dLRF and dARF; V, P, N, C, E, V, W, G, and K for differences between dLRF and dCRF.

The analysis of rf as a function of distance along a sequence shows that the contacts or interactions between residues very distant along the sequence are difficult to determine and are more likely to result in discrepancies between the methods. As expected because of its size, its physicochemical properties, and its implication in various interactions, ⁸² W is often involved in the strongest discrepancies at any distance.

As in the previous section, the overall discrepancies can be summarized through correlation coefficients. For *near* contacts, the (dCRF, dLRF), the PCC value equaled 0.85 and 0.61 for dARF vs dLRF. For *far* contacts, PCC values were 0.73 for dLRF vs dCRF and 0.64 for dLRF vs dARF, respectively. Finally, for *very far* contacts, the PCC was 0.57 for dLRF vs dCRF and 0.78 for dLRF vs dARF. Therefore, the dLRF vs dCRF correlation decreased with increasing contact distance, whereas the dLRF vs

Table 4. Relative Frequency Modulation According to Distance along the Sequence a

tance along the Sequence ^a						
	near (5-20 residues)					
amino acid pairs	dLRF	dCRF	dARF			
$[A \rightarrow A]$	-0.17	-0.16	-0.26			
$[A \rightarrow C]$	0.16	0.23	0.07			
$[A \rightarrow I]$	0.20	0.08	-0.01			
$[A \to V]$	0.20	0.15	0.00			
$[C \rightarrow M]$	-0.18	-0.20	-0.14			
$[C \rightarrow T]$	0.13	0.20	0.08			
$[E \rightarrow C]$	0.20	0.20	0.07			
$[E \rightarrow E]$	-0.22	-0.20	-0.14			
$[E \rightarrow G]$	0.08	0.00	0.20			
$[E \rightarrow T]$	0.06	0.05	0.22			
$[E \rightarrow V]$	0.23	0.25	0.11			
$[E \rightarrow M]$	0.17	0.25	0.11			
$[E \rightarrow Y]$	0.16	0.23	0.13			
$[F \rightarrow C]$	0.18	0.27	0.09			
$[G \rightarrow C]$						
$[G \rightarrow C]$	0.14	0.24	0.37			
$[H \rightarrow C]$	-0.12	-0.16 0.28	-0.22			
$[K \rightarrow C]$	0.20		0.11			
$[K \rightarrow C]$ $[K \rightarrow W]$	0.20	0.25	0.08 0.16			
	0.21	0.22				
$[M \rightarrow C]$	0.07	0.00	-0.21			
$[M \rightarrow M]$	-0.10	-0.25	-0.32			
$[N \rightarrow C]$	0.09	0.22	0.24			
$[b \rightarrow C]$	0.30	0.31	0.18			
$[P \to Y]$	0.10	0.22	0.06			
$[Q \rightarrow V]$	0.16	0.22	0.05			
$[Q \rightarrow W]$	0.15	0.29	0.13			
$[Q \to Y]$	0.16	0.29	0.09			
$[R \rightarrow V]$	0.17	0.20	0.05			
$[T \rightarrow C]$	0.18	0.46	0.19			
$[W \rightarrow C]$	0.20	-0.01	0.21			
$[W \rightarrow W]$	-0.05	0.29	0.01			
amino acid pairs	far (21–50 residues) dLRF	dCRF	dARF			
•			und			
$[C \rightarrow H]$	-0.02	-0.03	-0.33			
$[D \rightarrow C]$	0.23	0.29	0.10			
$[D \rightarrow G]$	0.09	0.22	0.09			
$[D \to G]$	-0.19	-0.22	-0.10			
$[E \rightarrow E]$	-0.22	-0.23	-0.15			
$[E \rightarrow G]$	0.08	0.21	0.03			
$[E \rightarrow K]$	-0.07	-0.23	-0.10			
$[E \rightarrow Q]$	-0.21	-0.23	-0.16			
$[H \rightarrow M]$	0.07	0.06	-0.20			
$[H \rightarrow W]$	-0.14	-0.23	-0.08			
$[I \rightarrow \Lambda]$	0.17	0.26	0.11			
$[K \to E]$	-0.08	-0.21	-0.03			
$[K \rightarrow C]$	0.13	0.26	0.24			
$[M \rightarrow C]$	0.20	0.15	0.35			
$[M \rightarrow W]$	0.01	-0.20	-0.23			
$[N \rightarrow C]$	0.24	0.16	0.24			

			ARTICLE
Table 4. Continued			
$[N \rightarrow N]$	0.12	0.23	0.00
$[P \rightarrow P]$	0.22	0.12	0.04
$[P \rightarrow W]$	-0.09	-0.10	-0.24
$[Q \rightarrow C]$	0.20	0.36	0.12
$[Q \rightarrow Q]$	-0.22	-0.32	-0.14
$[V \rightarrow V]$	0.16	0.21	0.13
$[M \rightarrow C]$	0.00	0.18	0.21
[Y C]	0.11	0.30	0.23
v	ery far (>50 residue	s)	
amino acid pairs	dLRF	dCRF	dARF
$[A \rightarrow F]$	0.24	0.07	0.08
$[A \rightarrow W]$	0.32	0.04	0.22
$[A \rightarrow Y]$	0.20	0.02	0.12
$[C \rightarrow F]$	0.23	0.00	0.15
$[D \rightarrow H]$	0.34	0.19	0.11
$[D \rightarrow R]$	0.32	-0.02	0.05
$[D \rightarrow W]$	0.22	-0.11	0.13
$[D \rightarrow X]$	0.32	-0.02	0.15
$[E \rightarrow H]$	0.26	0.07	0.04
$[E \to K]$	-0.10	-0.25	-0.20
$[E \rightarrow b]$	0.19	0.26	0.14
$[E \rightarrow K]$	0.31	-0.15	0.04
$[E \rightarrow M]$	0.27	-0.02	0.22
$[E \rightarrow A]$	0.24	0.01	0.09
$[G \rightarrow C]$	-0.11	-0.12	-0.21
$[G \rightarrow M]$	0.13	0.05	0.20
$[G \rightarrow W]$	0.30	0.05	0.17
$[H \rightarrow C]$	-0.31	-0.25	-0.07
$[H \rightarrow G]$	0.01	0.20	0.04
$[H \rightarrow M]$	0.35	0.02	-0.05
$[I \to E]$	0.20	-0.01	-0.01
$[I \rightarrow W]$	0.23	-0.01	0.06

[a w]	0.30	0.03	0.17
$[H \rightarrow C]$	-0.31	-0.25	-0.07
$[H \rightarrow G]$	0.01	0.20	0.04
$[H \rightarrow W]$	0.35	0.02	-0.05
$[I \rightarrow F]$	0.20	-0.01	-0.01
$[I \rightarrow W]$	0.23	-0.01	0.06
$[K \rightarrow E]$	-0.14	-0.25	-0.21
$[K \rightarrow G]$	0.06	0.20	0.01
$[K \rightarrow K]$	-0.10	-0.20	-0.10
$[L \rightarrow W]$	0.20	0.04	0.15
$[N \rightarrow G]$	0.00	0.20	0.01
$[N \rightarrow Y]$	0.25	0.05	0.08
$[P \rightarrow W]$	0.28	0.16	0.33
$[P \rightarrow Y]$	0.20	0.00	0.10
$[Q \rightarrow E]$	-0.15	-0.20	-0.15
$[Q \rightarrow G]$	0.03	0.22	0.02
$[R \rightarrow E]$	-0.05	-0.22	-0.10
$[R \rightarrow G]$	0.03	0.23	-0.01
$[R \rightarrow P]$	0.19	0.26	0.16
$[R \rightarrow W]$	0.23	0.13	0.16
$[S \rightarrow C]$	-0.24	-0.28	-0.11
$[S \rightarrow M]$	0.21	0.12	0.10
$[S \rightarrow P]$	0.16	0.21	0.17
$[T \rightarrow W]$	0.28	0.03	0.06
$[W \rightarrow C]$	-0.20	0.03	-0.17
$[M \rightarrow B]$	0.12	0.31	0.06
$[M \rightarrow Q]$	-0.08	-0.20	-0.07

Table 4. Continued

$[W \rightarrow W]$	0.26	-0.17	-0.054
$[Y \rightarrow C]$	-0.21	-0.19	0.00
$[Y \rightarrow W]$	0.26	0.09	0.07

 $[^]a$ Excess dLRF, dCRF and dARF induced by specifying the distance between residue are compared for three distance zones: near (5–20 residues), far (21–50 residues), and very far (>50 residues). Only the cases with one absolute value >0.2 are presented.

Table 5. Contact Differences for α -Helices and β -Sheets^a

	dI	LRF	dO	CRF	d.	ARF
amino acid pairs	helix	sheet	helix	sheet	helix	sheet
$[H \rightarrow E]$	-0.23	0.33	-0.17	0.29	-0.18	0.19
$[I \rightarrow A]$	-0.22	0.21	-0.16	0.17	-0.13	0.14
$[L \rightarrow A]$	-0.22	0.24	-0.21	0.19	-0.16	0.14
$[P \rightarrow M]$	-0.11	0.18	-0.06	0.1	-0.24	0.26
$[T \rightarrow Q]$	-0.22	0.23	-0.15	0.19	-0.15	0.22
$[V \rightarrow A]$	-0.2	0.28	-0.17	0.26	-0.1	0.21
$[W \rightarrow Q]$	-0.25	0.25	-0.24	0.22	-0.21	0.33
$[C \rightarrow F]$	0.38	-0.42	0.2	-0.44	0.18	-0.31
$[C \rightarrow H]$	-0.01	-0.12	0.1	-0.04	0.21	-0.2
$[C \rightarrow V]$	0.24	-0.46	0.2	-0.44	0.13	-0.35
$[F \rightarrow F]$	0.21	-0.62	0.03	-0.35	0.06	-0.44
$[I \rightarrow C]$	0.1	-0.25	0.2	-0.3	0.01	-0.18
$[\Gamma \rightarrow C]$	0.16	-0.17	0.2	-0.21	0.06	-0.1
$[M \rightarrow C]$	0.26	-0.29	0.31	-0.21	0.01	-0.15
$[M \rightarrow F]$	0.2	-0.51	0.13	-0.34	0.09	-0.39
$[M \rightarrow V]$	0.22	-0.49	0.14	-0.45	0.15	-0.38
$[W \rightarrow F]$	0.22	-0.57	0.04	-0.29	0.04	-0.35

^a The table gives the relative frequency changes (drf) resulting from the secondary structure environment of the residues. Only the amino acid pairs showing contrasting changes are displayed, that is, both (helix and sheet) drf of absolute value >0.2 but of opposite sign, in at least one of the methods. The pairs satisfying this criterion are displayed in bold.

dARF correlation increased. In some cases, because of its special position in Figure 4, the cysteine—cysteine interaction had a strong influence on the PCC values. For example, without taking $[C \rightarrow C]$ into account, PCC values for dCRF vs dLRF were 0.73 (near), 0.73 (far) and 0.47 (very far). These values for dARF vs dLRF were 0.44 (near), 0.59 (far), and 0.68 (very far). Except the far contacts for dCRF, the absence of $[C \rightarrow C]$ decreased the PCC values. Finally, a difference in contact distributions between secondary structures analyzed in each distance zone may also account for some PCC variations. For instance, Laguerre tessellation and the all-atom threshold method counted 26.4% and 28.0% of near contact frequency between two β-strands, respectively; whereas the Cα method resulted in 32.1%.

Analysis of rf According to Secondary Structures. The secondary structure elements (SSEs) are local protein structures, known to be involved in the stability of protein 3D folds. The residue interactions and contacts observed in SSEs differ depending on their environment, thus a significant dependence on secondary structure may be expected. Indeed, α -helices are primarily maintained by short-range interactions, while β -sheets mainly involve long-range interactions. For specificity, the analysis was

Table 6. Counts of Cysteine—Cysteine Contacts and Disulfide Bridges^a

	threshold	Voronoi	Laguerre
Cysh	1754	1523	1502
Cyss	512	743	764

^a The numbers of free cysteines (Cysh) (no S—S contact with other cysteine) and of half-cystines (Cyss) are indicated as provided by the three contact methods. "Threshold" stands for the all-atom threshold method. The half-cystines are defined either by a distance shorter than 2.1 Å in the threshold method or, in the tessellation method, by a face shared by two sulfurs (in this case, the contact may be covalent or not).




Figure 7. Laguerre face area vs bond distance in tessellation for contacts between cysteine sulfurs. Each point represents a Laguerre contact between the S atoms of two cysteines.

performed on the residues showing specific rf changes, that is, important opposing changes in the two repetitive structures. As above, drf is the difference between the relative frequency calculated on a subset of residues (e.g., both i and j in α -helices) and its counterpart evaluated on the whole databank.

We only considered a limited number of representative cases. In Table 5, the amino acid pairs were selected as follows: $\{drf(\alpha-helices)>0.2$ and $drf(\beta-sheet)<-0.2\}$ or $\{drf(\alpha-helices)<-0.2$ and $drf(\beta-sheet)>0.2\}$ in at least one of the methods. In a majority of cases, the criterion was fulfilled by the Laguerre tessellation (dLRF). The values of the other methods (C α and all-atom threshold methods) were often close, for example, they had similar patterns. One exception was for [C \rightarrow H] contacts in helices where the dLRF value was 0.01 while the dCRF value was -0.1 and the dARF, -0.21. Small residues, such as A or C, often appeared in the selected pairs. Cysteines are well-known for maintaining protein structures by forming disulfide bridges, mainly in β -sheets.

As a general conclusion, discrepancies between the Laguerre tessellation and the distance-threshold methods ($C\alpha$ method or all-atom) are modulated by both residue proximity along the sequence and their secondary structure.

Specific Case Example: Disulfide Bridges. Disulfide bonds contribute to protein tertiary or quaternary structure by forming relatively strong covalent bonds between cysteines, which can be either quite distant in the amino acid sequence or even members of different peptide chains (quaternary case). However, the question




Figure 8. Four typical configurations of cysteine pairs in contact. The cysteines are shown as balls. The blue polygon of area A, is the Laguerre face between the Cysteine sulfurs, distance d apart. (a) 1lpb, 95 d = 2.04 Å, A = 7.71 Å 2 ; (b) 1pl3, 96 d = 2.04 Å, A = 12.80 Å 2 ; (c) 2bm5, 97 d = 5.025 Å, A = 0.09 Å 2 ; (d) 1b25, 98 d = 5.57 Å, A = 7.35 Å 2 . Views made with PyMol.

of disulfide bridges is still a challenge.84,88-92 A collection of criteria has been proposed to identify cystine, based either on the distance between two sulfurs of less than 2.3 Å³ or on the distance between cysteine Cas ranging from 4.2 to 7.5 Å. 84,92 A cystine is formed by the oxidation of two cysteine residues which covalently link and make a disulfide bond. A half-cystine is a cysteine involved in a disulfide bridge with another cysteine to form a cystine. In our analysis, cystines were located using either a distance between two sulfurs lower than 2.1 Å, or the occurrence of a Laguerre (or Voronoi) face separating the sulfur polyhedra of two distinct cysteines. The free cysteines and half-cystines were enumerated in each method and the results are given in Table 6. The contact criterion depended on the method. The threshold of 2.1 Å ensures that all the cysteine (C) contacts found by the distance-threshold method form covalently bonded cystines. All these contacts were also found by the tessellation methods. Indeed, covalent bonds imply that distances are short enough to be detected by all the considered methods. The tessellations included additional C-C contacts over distances greater than 2.1 Å still labeled as half-cystines even if those bonds are almost certainly not covalent. Thus the number of half-cystines detected by the Laguerre or Voronoi methods was higher than by the distance-threshold method (512 for the threshold method, 743 for Voronoi and 764 for Laguerre). Moreover, the odd number of half-cystines produced using the Voronoi method indicates that some contacts involve more than two cysteines. The C-C contact counts, detailed in Supporting Information 4, confirm that some proteins, such as the transferase (PDB code 1d0q⁹³) or the Vhs domain of Tom1 protein (PDB code 1elk⁹⁴), have three cysteines that are in contact.

The tessellation methods do not account for either the physical nature of the interactions or the absolute distance, so there is no guarantee that the associated pairs of cysteines are covalently bound. Nevertheless, these contacts reflect spatial closeness. More insight can be obtained from the correlation between (1) the distance between cysteine sulfurs and (2) the area of the corresponding face in Laguerre tessellation, displayed in Figure 7. The data were split into two distinct clusters, clearly separated along the distance spectrum: one sharply centered on a mean distance of 2 Å, certainly involving the covalently bound Cystines, and the other, scattered at values greater than 3.2 Å, corresponding to noncovalent contacts. The gap between 2.1 and 3.2 Å leaves no ambiguity in qualifying these contacts. While the covalent distance is fixed to nearly 2 Å, the corresponding face area spreads over quite a broad interval ranging from 8 to 13 Å². For distances greater than 3 Å, the distance and area showed a negative correlation and a middle range PCC value of -0.72, similar to normal kinds of contacts.

Figure 8 represents the molecular configurations of the four main cases observed: covalent distance ($<2.1\,\text{\AA}$) and a bottom range area (see Figure 8a), covalent distance and top range area (see Figure 8b), normal distance ($>3\,\text{\AA}$) and small area (see Figure 8c), normal distance and larger area (see Figure 8d). Figure 8c and d demonstrates the importance of the orientation of the cysteines on the Laguerre face area. When both cysteines are parallel, the contact tends to be small, with a small face area (see Figure 8c), while the area increases when the cysteine sulfurs face each other as in Figure 8d.

In summary, the distance method is very effective in selecting only the covalently linked cystines, while the tessellation method, not limited by any threshold, detects the relative proximity of sulfurs even in absence of any tight bond interaction. Moreover, distance is not the only factor; the conformation around the cysteines also plays a role in the contacts found by tessellation methods. Therefore, tessellations, especially the Laguerre tessellation with well-tuned weights, may even provide deeper insight into the geometry of the contacts.

■ CONCLUSION

Currently, the knowledge of protein folding still poses a challenge for fully understanding the functionality of proteins and predicting their structure. Exploring the interactions and contacts between residues is a key step to furthering our knowledge in this area. Here, we proposed a detailed analysis of the contacts which can be specified by geometrical criteria, whereas interactions rely on forces or energies. We carried out a comparative analysis of two contact definitions: distance-threshold methods and tessellation methods. The distance-threshold method is useful and realistic when the contacts surrounding a residue are specified by a particular distance range. This type of method does not need any solvent around the protein, which may save computer memory and run time. The tessellation method provides a more realistic representation of the local ordering in the structure; the contacts deduced from tessellation essentially consist of a complete list of neighbors in the first layer around any residue. The method is flexible and adapts itself to density inhomogeneities. However, this tessellation approach needs the presence of solvent if accessible residues are to be incorporated in the analysis. The Voronoi tessellation method does not depend on any parameter, but it is known to even out local inhomogeneities, which may lead to some undesirable bias. 54 At the coarse-grained and atomic scales, the Laguerre (or weighted Voronoi) tessellation method provides the most precise account of space occupation by the constituent atoms, residues or molecular units. However, it relies on a set of weights that need tuning, 51,54 even though the simple formula $w = r^2$, in terms of the van der Waals radius r, was found to be optimal at the atomic scale. Regarding contacts, Laguerre and Voronoi partitions give very similar results, with about 99% of common contacts (see Supporthing Information 5). The few cases of discrepancies mainly involve residues at the protein surface.

Much more significant are the discrepancies found in comparing the tessellation and distance-threshold methods. On average, these differences compensate each other, an indication that the threshold has been set to an appropriate intermediate value. However, the discrepancies become more and more visible when the contacts are differentiated by amino acid species, or even by pairs of species as in the relative frequencies.

ASSOCIATED CONTENT

Supporting Information. PCC values for the correlation between residue volumes and mean contacts, relative frequencies from tessellations; differences between Laguerre and Voronoi data (2a) and Laguerre LRF values (2b), (dLR, dCRF) correlations vs protein size, a list of proteins is also provided including cysteine—cysteine contact counts, and finally, the contacts shared by the Laguerre and Voronoi methods are displayed in the same way as Figure 6. This material is available free of charge via the Internet at http://pubs.acs.org/.

AUTHOR INFORMATION

Corresponding Author

*E-mail: jeremy.esque@univ-paris-diderot.fr.

ACKNOWLEDGMENT

A PhD grant to E.J. from the French Ministry of Higher Education and Research is acknowledged. We thank Xabier Oyharçabal for his contribution to the early development of VLDP and all the developers of the freely available software, which greatly facilitated our work (cited in Materials & Methods section). This work was supported by grants from the French Ministry of Research, Paris Diderot University Paris 7, French National Center for Scientific Research (CNRS), French National Institute for Blood Transfusion (INTS) and French National Institute for Health and Medical Research (INSERM). The authors thank Ms. C. Engel-Gautie, a native English speaker and professional scientific editor, who has improved our manuscript thanks to her corrections.

■ REFERENCES

- (1) Faure, G.; Bornot, A.; de Brevern, A. G. Protein contacts, interresidue interactions and side-chain modelling. *Biochimie* **2008**, *90*, 626–639.
- (2) Taylor, W. R. The classification of amino acid conservation. *J. Theor. Biol.* **1986**, *119*, 205–218.
- (3) Bhattacharyya, R.; Pal, D.; Chakrabarti, P. Disulfide bonds, their stereospecific environment and conservation in protein structures. *Protein Eng. Des. Sel.* **2004**, *17*, 795–808.
- (4) Rose, G. D.; Wolfenden, R. Hydrogen bonding, hydrophobicity, packing, and protein folding. *Annu. Rev. Biophys. Biomol. Struct.* **1993**, 22, 381–415.
- (5) Tsai, C. J.; Nussinov, R. Hydrophobic folding units derived from dissimilar monomer structures and their interactions. *Protein Sci.* **1997**, *6*, 24–42.
- (6) Brandl, M.; Weiss, M. S.; Jabs, A.; Sühnel, J.; Hilgenfeld, R. $C-H\cdots\pi$ -interactions in proteins. *J. Mol. Biol.* **2001**, 307, 357–377.
- (7) Myers, J. K.; Pace, C. N. Hydrogen bonding stabilizes globular proteins. *Biophys. J.* **1996**, *71*, 2033–2039.
- (8) Weiss, M. S.; Brandl, M.; Sühnel, J.; Pal, D.; Hilgenfeld, R. More hydrogen bonds for the (structural) biologist. *Trends Biochem. Sci.* **2001**, 26, 521–523.
- (9) Wetlaufer, D. B. Nucleation, rapid folding, and globular intrachain regions in proteins. *Proc. Natl. Acad. Sci. U.S.A.* 1973, 70, 697–701.
- (10) Zhou, H.; Xue, B.; Zhou, Y. DDOMAIN: Dividing structures into domains using a normalized domain-domain interaction profile. *Protein Sci.* **2007**, *16*, 947–955.
- (11) Holm, L.; Sander, C. Parser for protein folding units. *Proteins* 1994, 19, 256–268.
- (12) Siddiqui, A. S.; Barton, G. J. Continuous and discontinuous domains: an algorithm for the automatic generation of reliable protein domain definitions. *Protein Sci.* **1995**, *4*, 872–884.
- (13) Siddiqui, A. S.; Dengler, U.; Barton, G. J. 3Dee: A database of protein structural domains. *Bioinformatics* **2001**, *17*, 200–201.
- (14) Gelly, J. C.; Etchebest, C.; Hazout, S.; de Brevern, A. G. Protein Peeling 2: A web server to convert protein structures into series of protein units. *Nucleic Acids Res.* **2006**, *34*, W75–W78.
- (15) Gelly, J. C.; de Brevern, A. G.; Hazout, S. "Protein Peeling": An approach for splitting a 3D protein structure into compact fragments. *Bioinformatics* **2006**, 22, 129–133.
- (16) Dedmon, M. M.; Lindorff-Larsen, K.; Christodoulou, J.; Vendruscolo, M.; Dobson, C. M. Mapping long-range interactions in alphasynuclein using spin-label NMR and ensemble molecular dynamics simulations. *J. Am. Chem. Soc.* **2005**, *127*, 476–477.
- (17) Dudev, M.; Lim, C. Discovering structural motifs using a structural alphabet: application to magnesium-binding sites. *BMC Bioinformatics* **2007**, *8*, 106.

- (18) Fiser, A.; Dosztányi, Z.; Simon, I. The role of long-range interactions in defining the secondary structure of proteins is overestimated. *Comput. Appl. Biosci.* **1997**, *13*, 297–301.
- (19) Gromiha, M. M.; Selvaraj, S. Importance of long-range interactions in protein folding. *Biophys. Chem.* **1999**, *77*, 49–68.
- (20) Ku, S.-Y.; Hu, Y.-J. Protein structure search and local structure characterization. *BMC Bioinformatics* **2008**, *9*, 349.
- (21) Sander, O.; Sommer, I.; Lengauer, T. Local protein structure prediction using discriminative models. *BMC Bioinformatics* **2006**, *7*, 14.
- (22) Bahar, I.; Jernigan, R. L. Inter-residue potentials in globular proteins and the dominance of highly specific hydrophilic interactions at close separation. *J. Mol. Biol.* **1997**, *266*, 195–214.
- (23) Miyazawa, S.; Jernigan, R. L. Residue-residue potentials with a favorable contact pair term and an unfavorable high packing density term, for simulation and threading. *J. Mol. Biol.* **1996**, 256, 623–644.
- (24) Dosztanyi, Z.; Fiser, A.; Simon, I. Stabilization centers in proteins: Identification, characterization and predictions. *J. Mol. Biol.* **1997**, 272, 597–612.
- (25) Kannan, N.; Selvaraj, S.; Gromiha, M. M.; Vishveshwara, S. Clusters in alpha/beta barrel proteins: implications for protein structure, function, and folding: a graph theoretical approach. *Proteins* **2001**, *43*, 103–112.
- (26) Kannan, N.; Vishveshwara, S. Identification of side-chain clusters in protein structures by a graph spectral method. *J. Mol. Biol.* **1999**, 292, 441–464.
- (27) Gugolya, Z.; Dosztányi, Z.; Simon, I. Interresidue interactions in protein classes. *Proteins* **1997**, 27, 360–366.
- (28) de Brevern, A. G. New assessment of a structural alphabet. *In Silico Biol.* **2005**, *5*, 283–289.
- (29) Etchebest, C.; Benros, C.; Hazout, S.; de Brevern, A. G. A structural alphabet for local protein structures: improved prediction methods. *Proteins* **2005**, *59*, 810–827.
- (30) de Brevern, A. G.; Etchebest, C.; Hazout, S. Bayesian probabilistic approach for predicting backbone structures in terms of protein blocks. *Proteins* **2000**, *41*, 271–287.
- (31) Joseph, A.; Agarwal, G.; Mahajan, S.; Gelly, J.-C.; Swapna, L.; Offmann, B.; Cadet, F.; Bornot, A.; Tyagi, M.; Valadié, H.; Schneider, B.; Etchebest, C.; Srinivasan, N.; de Brevern, A. A short survey on protein blocks. *Biophys. Rev.* **2010**, *2*, 137–145.
- (32) Faure, G.; Bornot, A.; de Brevern, A. G. Analysis of protein contacts into Protein Units. *Biochimie* **2009**, *91*, 876–887.
- (33) Bonneau, R.; Ruczinski, I.; Tsai, J.; Baker, D. Contact order and ab initio protein structure prediction. *Protein Sci.* 2002, 11, 1937–1944.
- (34) Gront, D.; Kolinski, A. A new approach to prediction of short-range conformational propensities in proteins. *Bioinformatics* **2005**, *21*, 981–987.
- (35) Hamilton, N.; Burrage, K.; Ragan, M. A.; Huber, T. Protein contact prediction using patterns of correlation. *Proteins* **2004**, *56*, 679–684.
- (36) Hamilton, N.; Huber, T. An introduction to protein contact prediction. *Methods Mol. Biol.* **2008**, *453*, 87–104.
- (37) Olmea, O.; Valencia, A. Improving contact predictions by the combination of correlated mutations and other sources of sequence information. *Fold Des.* **1997**, *2*, S25–S32.
- (38) Pollastri, G.; Baldi, P. Prediction of contact maps by GIOHMMs and recurrent neural networks using lateral propagation from all four cardinal corners. *Bioinformatics* **2002**, *18*, S62–S70.
- (39) Pollastri, G.; Baldi, P.; Fariselli, P.; Casadio, R. Improved prediction of the number of residue contacts in proteins by recurrent neural networks. *Bioinformatics* **2001**, *17*, S234–S242.
- (40) Punta, M.; Rost, B. PROFcon: Novel prediction of long-range contacts. *Bioinformatics* **2005**, *21*, 2960–2968.
- (41) Moult, J.; Fidelis, K.; Kryshtafovych, A.; Rost, B.; Hubbard, T.; Tramontano, A. Critical assessment of methods of protein structure prediction—Round VII. *Proteins* **2007**, *69*, 3–9.
- (42) Moult, J.; Fidelis, K.; Kryshtafovych, A.; Rost, B.; Tramontano, A. Critical assessment of methods of protein structure prediction—Round VIII. *Proteins* **2009**, *77*, 1–4.

- (43) Moult, J.; Fidelis, K.; Rost, B.; Hubbard, T.; Tramontano, A. Critical assessment of methods of protein structure prediction (CASP)—Round 6. *Proteins* **2005**, *61*, 3–7.
- (44) Moult, J.; Fidelis, K.; Zemla, A.; Hubbard, T. Critical assessment of methods of protein structure prediction (CASP): Round IV. *Proteins* **2001**, No. Suppl 5, 2–7.
- (45) Moult, J.; Fidelis, K.; Zemla, A.; Hubbard, T. Critical assessment of methods of protein structure prediction (CASP)—Round V. *Proteins* **2003**, 53, 334–339.
- (46) Moult, J.; Hubbard, T.; Bryant, S. H.; Fidelis, K.; Pedersen, J. T. Critical assessment of methods of protein structure prediction (CASP): Round II. *Proteins* 1997, No. Suppl 1, 2–6.
- (47) Moult, J.; Hubbard, T.; Fidelis, K.; Pedersen, J. T. Critical assessment of methods of protein structure prediction (CASP): Round III. *Proteins* **1999**, No. Suppl 3, 2–6.
- (48) Gromiha, M. M.; Thangakani, A. M.; Selvaraj, S. FOLD-RATE: Prediction of protein folding rates from amino acid sequence. *Nucleic Acids Res.* **2006**, *34*, W70–W74.
- (49) McConkey, B. J.; Sobolev, V.; Edelman, M. Quantification of protein surfaces, volumes and atom—atom contacts using a constrained Voronoi procedure. *Bioinformatics* **2002**, *18*, 1365–1373.
- (50) Sadoc, J. F. Spectral properties of contact matrix: Application to proteins. *Eur. Phys. J. E: Soft Matter Biol. Phys.* **2005**, *18*, 321–333.
- (51) Sadoc, J. F.; Jullien, R.; Rivier, N. The Laguerre polyhedral decomposition: application to protein folds. *EPJ B* **2003**, *33*, 355–363.
- (52) Janin, J.; Chothia, C. The structure of protein—protein recognition sites. *J. Biol. Chem.* **1990**, 265, 16027–16030.
- (53) Harpaz, Y.; Gerstein, M.; Chothia, C. Volume changes on protein folding. *Structure* **1994**, *2*, 641–9.
- (54) Esque, J.; Oguey, C.; de Brevern, A. G. A novel evaluation of residue and protein volumes by means of Laguerre tessellation. *J. Chem. Inf. Model.* **2010**, *50*, 947–960.
- (55) da Silveira, C. H.; Pires, D. E. V.; Minardi, R. C.; Ribeiro, C.; Veloso, C. J. M.; Lopes, J. C. D.; Meira, W., Jr.; Neshich, G.; Ramos, C. H. I.; Habesch, R.; Santoro, M. M. Protein cutoff scanning: A comparative analysis of cutoff dependent and cutoff free methods for prospecting contacts in proteins. *Proteins* **2009**, *74*, 727–743.
- (56) Wang, G.; Dunbrack, R. L., Jr. PISCES: A protein sequence culling server. *Bioinformatics* **2003**, *19*, 1589–1591.
- (57) Wang, G.; Dunbrack, R. L., Jr. PISCES: Recent improvements to a PDB sequence culling server. *Nucleic Acids Res.* **2005**, 33, W94–W98.
- (58) Berman, H. M.; Westbrook, J.; Feng, Z.; Gilliland, G.; Bhat, T. N.; Weissig, H.; Shindyalov, I. N.; Bourne, P. E. The Protein Data Bank. *Nucleic Acids Res.* **2000**, 28, 235–242.
- (59) Berendsen, H. J. C.; van der Spoel, D.; van Drunen, R. GROMACS, A message-passing parallel molecular dynamics implementation. *Comput. Phys. Commun.* **1995**, *91*, 43–56.
- (60) Hess, B.; Kutzner, C.; van der Spoel, D.; Lindahl, E. GRO-MACS 4: Algorithms for highly efficient, load-balanced, and scalable molecular simulation. *J. Chem. Theory Comput.* **2008**, *4*, 435–447.
- (61) Lindahl, E.; Hess, B.; van der Spoel, D. GROMACS 3.0: A package for molecular simulation and trajectory analysis. *J. Mol. Model.* **2001**, *7*, 306–317.
- (62) Van Der Spoel, D.; Lindahl, E.; Hess, B.; Groenhof, G.; Mark, A. E.; Berendsen, H. J. C. GROMACS: Fast, flexible, and free. *J. Comput. Chem.* **2005**, *26*, 1701–1718.
- (63) Jorgensen, W. L.; Maxwell, D. S.; Tirado-Rives, J. Development and testing of the OPLS all-atom force field on conformational energetics and properties of organic liquids. *J. Am. Chem. Soc.* **1996**, *118*, 11225–11236.
- (64) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **1983**, *79*, 926–935.
- (65) Berendsen, H. J. C.; Postma, J. P. M.; van Gunsteren, W. F.; Dinola, A.; Haak, J. R. Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* **1984**, *81*, 3684–3690.

- (66) Hess, B.; Bekker, H.; Berendsen, H. J. C.; Fraaije, J. G. E. M. LINCS, A linear constraint solver for molecular simulations. *J. Comput. Chem.* **1997**, *18*, 1463–1472.
- (67) Tironi, I. G.; Sperb, R.; Smith, P. E.; van Gunsteren, W. F. A generalized reaction field method for molecular dynamics simulations. *J. Chem. Phys.* **1995**, *102*, 5451–5459.
- (68) Sugihara, K. Laguerre Voronoi diagram on the sphere. *Journal for Geometry and Graphics* **2002**, *6*, 69–81.
- (69) Porto, M.; Bastolla, U.; Roman, H. E.; Vendruscolo, M. Reconstruction of protein structures from a vectorial representation. *Phys. Rev. Lett.* **2004**, 92, 218101.
- (70) Hubbard, S. J.; Thornton, J. M. *NACCESS*, version 2.1.1; Dept of Biochemistry and Molecular Biology, University College: London, 1993.
- (71) Kabsch, W.; Sander, C. Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* **1983**, *22*, 2577–2637.
- (72) Delano, W. L. *The PyMOL Molecular Graphics System*; version 1.2r1; DeLano Scientific: San Carlos, CA, 2002; http://www.pymol.org.
- (73) Janin, J. Surface and inside volumes in globular proteins. *Nature* **1979**, 277, 491–492.
- (74) Rose, G. D.; Geselowitz, A. R.; Lesser, G. J.; Lee, R. H.; Zehfus, M. H. Hydrophobicity of amino acid residues in globular proteins. *Science* **1985**, 229, 834–838.
- (75) Soyer, A.; Chomilier, J.; Mornon, J. P.; Jullien, R.; Sadoc, J. F. Voronoi tessellation reveals the condensed matter character of folded proteins. *Phys. Rev. Lett.* **2000**, *85*, 3532–3535.
- (76) Lewis, F. T. The correlation between cell division and the shapes and sizes of prismatic cells in the epidermis of cucumis. *Anat. Rec.* **1928**, *38*, *341–376*.
- (77) Lewis, F. T. A comparison between the mosaic of polygons in a film of artificial emulsion and the pattern of simple epithelium in surface view (cucumber epidermis and human amnion). *Anat. Rec.* **1931**, *50*, 235–265.
- (78) Samanta, U.; Bahadur, R. P.; Chakrabarti, P. Quantifying the accessible surface area of protein residues in their local environment. *Protein Eng.* **2002**, *15*, 659–667.
- (79) Samanta, U.; Pal, D.; Chakrabarti, P. Environment of tryptophan side chains in proteins. *Proteins* **2000**, *38*, 288–300.
- (80) Cheng, J.; Baldi, P. Improved residue contact prediction using support vector machines and a large feature set. *BMC Bioinformatics* **2007**, *8*, 113.
- (81) Vassura, M.; Margara, L.; Di Lena, P.; Medri, F.; Fariselli, P.; Casadio, R. Reconstruction of 3D structures from protein contact maps. *IEEE/ACM Trans. Comput. Biol. Bioinform.* **2008**, *5*, 357–367.
- (82) Samanta, U.; Pal, D.; Chakrabarti, P. Packing of aromatic rings against tryptophan residues in proteins. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* 1999, 55, 1421–1427.
- (83) Meurisse, R.; Brasseur, R.; Thomas, A. Aromatic side-chain interactions in proteins. Near- and far-sequence His-X pairs. *Biochim. Biophys. Acta, Proteins Proteomics* **2003**, *1649*, 85–96.
- (84) Czaplewski, C.; Oldziej, S.; Liwo, A.; Scheraga, H. A. Prediction of the structures of proteins with the UNRES force field, including dynamic formation and breaking of disulfide bonds. *Protein Eng. Des. Sel.* **2004**, *17*, 29–36.
- (85) Brocchieri, L.; Karlin, S. How are close residues of protein structures distributed in primary sequence? *Proc. Natl. Acad. Sci. U.S.A.* **1995**, *92*, 12136–12140.
- (86) Go, N.; Taketomi, H. Respective roles of short- and long-range interactions in protein folding. *Proc. Natl. Acad. Sci. U.S.A.* **1978**, 75, 559–563.
- (87) Tanaka, S.; Scheraga, H. A. Model of protein folding: inclusion of short-, medium-, and long-range interactions. *Proc. Natl. Acad. Sci. U.S.A.* **1975**, 72, 3802–3806.
- (88) Chen, Y.-C.; Lin, Y.-S.; Lin, C.-J.; Hwang, J.-K. Prediction of the bonding states of cysteines using the support vector machines based on multiple feature vectors and cysteine state sequences. *Proteins* **2004**, *55*, 1036–1042.

- (89) Martelli, P. L.; Fariselli, P.; Malaguti, L.; Casadio, R. Prediction of the disulfide bonding state of cysteines in proteins with hidden neural networks. *Protein Eng.* **2002**, *15*, 951–953.
- (90) Mucchielli-Giorgi, M. H.; Hazout, S.; Tufféry, P. Predicting the disulfide bonding state of cysteines using protein descriptors. *Proteins* **2002**, *46*, 243–249.
- (91) Song, J.; Yuan, Z.; Tan, H.; Huber, T.; Burrage, K. Predicting disulfide connectivity from protein sequence using multiple sequence feature vectors and secondary structure. *Bioinformatics* **2007**, *23*, 3147–3154.
- (92) Dani, V. S.; Ramakrishnan, C.; Varadarajan, R. MODIP revisited: Re-evaluation and refinement of an automated procedure for modeling of disulfide bonds in proteins. *Protein Eng.* **2003**, *16*, 187–193.
- (93) Wada, T.; Yamazaki, T.; Kyogoku, Y. The structure and the characteristic DNA binding property of the C-terminal domain of the RNA polymerase alpha subunit from *Thermus thermophilus*. *J. Biol. Chem.* **2000**, 275, 16057–16063.
- (94) Misra, S.; Beach, B. M.; Hurley, J. H. Structure of the VHS domain of human Tom1 (target of myb 1): insights into interactions with proteins and membranes. *Biochemistry* **2000**, *39*, 11282–11290.
- (95) Egloff, M. P.; Marguet, F.; Buono, G.; Verger, R.; Cambillau, C.; van Tilbeurgh, H. The 2.46 A resolution structure of the pancreatic lipase-colipase complex inhibited by a C11 alkyl phosphonate. *Biochemistry* **1995**, *34*, 2751–2762.
- (96) Rotsaert, F. A.; Hallberg, B. M.; de Vries, S.; Moenne-Loccoz, P.; Divne, C.; Renganathan, V.; Gold, M. H. Biophysical and structural analysis of a novel heme B iron ligation in the flavocytochrome cellobiose dehydrogenase. *J. Biol. Chem.* **2003**, 278, 33224–33231.
- (97) Hedge, P. J.; Spratt, B. G. Resistance to beta-lactam antibiotics by re-modelling the active site of an *E. coli* penicillin-binding protein. *Nature* **1985**, *318*, 478–480.
- (98) Hu, Y.; Faham, S.; Roy, R.; Adams, M. W.; Rees, D. C. Formaldehyde ferredoxin oxidoreductase from *Pyrococcus furiosus*: the 1.85 A resolution crystal structure and its mechanistic implications. *J. Mol. Biol.* **1999**, 286, 899–914.