

Silage Fermentation and *in Vitro* Degradation of Monosaccharide Constituents of Wheat Harvested at Two Stages of Maturity

Daniel Ben-Ghedalia,* Amos Kabala, Joshua Miron, and Edith Yosef

The Metabolic Unit, Institute of Animal Science, The Volcani Center, ARO, Bet Dagan 50250, Israel

Wheat plants (*Triticum aestivum*) were harvested at the bloom and soft-dough stages and ensiled in laboratory silos, and the effect of stage of maturity on the recovery of soluble and cell wall components and on the *in vitro* digestibility of monosaccharide residues was examined. A total of 8–9% neutral detergent fiber (NDF) monosaccharides residues was solubilized and partly utilized during ensilage. Glucose and galactose were the most fermentable sugars among the ND soluble monosaccharides. Fructose was equally fermented in both silages, but the higher recovery of reducing sugars in the soft-dough silage as compared with the bloom silage (44.1 vs 26.8) indicates the possible participation of starch in the fermentation process of the former. Generally, the silages were less digestible than the source materials, but the difference was larger in the bloom treatment. Thus, a gap of 8.1 percentage units in total NDF monosaccharide digestibility (*D*) between the source materials was reduced to 5.5 units between the silages. The decline in the *D* values of the monosaccharide residues of the whole plant following ensilage was 11.5 and 6.0 percentage units for the bloom and soft-dough stages, respectively. The *D* of total monosaccharide residues of whole plant material was equal within the source materials but was 5.3 percentage units higher in the soft-dough silage than in the bloom one. Digestibility measurement at the monosaccharide level of source materials and silages offers a sharp tool for determining the optimal stage for harvesting wheat for silage.

Keywords: Wheat silage; stage of maturity; monosaccharide digestibility; monosaccharide fermentation

INTRODUCTION

The optimal stage for harvesting wheat for silage is a major issue in semiarid areas in which wheat is grown as a winter crop. It has been shown recently (Ben-Ghedalia et al., 1995) that the decrease in neutral detergent fiber (NDF) monosaccharide digestibility between the bloom and soft-dough stages is compensated by an increase of 36% in the yield of digestible dry matter. Nevertheless, the concern of getting a forage that is a combination of "straw and starch" is raised occasionally as the argument in favor of harvesting wheat forage at the bloom stage of maturity (Arieli and Adin, 1994). Starch accumulated in the spike provides an abundant source of fermentable carbohydrates for successfully ensiling the soft-dough plants. However, the entire "arsenal" of silage fermentable carbohydrates found in wheat harvested at the bloom stage has not been fully discovered yet. For instance, the more fragile cell walls (CW) and labile CW components found in the younger wheat plant may participate in the fermentation process, and if they do, is their contribution important or necessary? There are numerous studies on the patterns of silage fermentation of wheat harvested at different stages of maturity (Ashbell et al., 1983; Ashbell and Kashanchi, 1987; Weinberg et al., 1991); none of this information is related, however, to the fate of CW constituents. Moreover, the role of CW monosaccharide components in the process of silage fermentation has been investigated in grasses (Morrison, 1979) and in lucerne (Ben-Ghedalia and Yosef, 1989a,b; Campbell et al., 1990), but relevant information on wheat does not exist yet.

Chemical and *in vitro* investigation of fresh plant material (Ben-Ghedalia et al., 1995) is important for following the changes occurring in the wheat plant during maturation. Plant material, however, is modified during ensilage. To obtain information required for determining the optimal stage of harvesting, both source materials and silages should be evaluated.

The objectives of this experiment were to study the effect of maturation on the *in vitro* degradability of carbohydrates in wheat silages and in the respective source materials and to assess the recovery of soluble and CW monosaccharide components following the ensilage of wheat harvested at bloom and soft-dough stages of maturity.

MATERIALS AND METHODS

Plant Material and Ensiling Procedure. Winter wheat (*Triticum aestivum* var. Ariel) plants were harvested (10 cm above ground) from a commercial field at bloom and after 19 days at the soft-dough stage. Shortly after harvest, plants were transferred to the laboratory and chopped into pieces 1–2 cm in length, and a portion representing the source material was kept at –20 °C until analyzed. The second portion of the chopped soft-dough material, containing 35% dry matter (DM), was ensiled directly in hermetically sealed 2-L laboratory glass silos, whereas the bloom stage material, containing 25% DM, was sun wilted to reach 35% DM prior to ensiling. Both materials were ensiled, four replicates per treatment, for 60 days at 21–23 °C. After this period, the silos were opened for assessing the recovery of fructose, total reducing sugars, neutral detergent soluble (NDS) monosaccharides, and NDF monosaccharide components and to determine the profile of fermentation products and the *in vitro* digestibility of monosaccharide constituents.

Analytical Procedures and *in Vitro* Determination. DM and organic matter (OM) contents were determined according to AOAC methods (Association of Official Analytical

* Author to whom correspondence should be addressed (fax 972-3-9604023).

Table 1. Concentration (Grams per 100 g of DM) of Monosaccharide Residues and Their Distribution (Percent) in Neutral Detergent Fiber (NDF) and Neutral Detergent Solubles (NDS) of Wheat Plants Harvested at Two Stages of Maturity

| monosaccharide | bloom stage | | | soft-dough stage | | |
|----------------|-------------|-----------------|------|------------------|-----------------|------|
| | composition | distribution in | | composition | distribution in | |
| | | NDF | NDS | | NDF | NDS |
| glucose | 31.4 | 83.3 | 16.7 | 34.6 | 64.3 | 35.7 |
| xylose | 14.1 | 100 | 0 | 11.8 | 100 | 0 |
| arabinose | 2.36 | 95.3 | 4.70 | 2.69 | 92.1 | 7.91 |
| uronic acids | 2.87 | 60.1 | 39.9 | 2.40 | 59.7 | 40.3 |
| galactose | 0.94 | 49.9 | 50.1 | 0.79 | 62.3 | 37.7 |
| mannose | 0.44 | 38.0 | 62.0 | 0.48 | 41.0 | 59.0 |

Table 2. Composition (Grams per 100 g of DM) and pH of Wheat Source Materials (SM) and Silages Harvested at Bloom and Soft-Dough Stages

| component | bloom | | soft-dough | |
|----------------------------|-------|--------|------------|--------|
| | SM | silage | SM | silage |
| pH | | 4.47 | | 4.24 |
| lactic acid | | 3.18 | | 4.28 |
| acetic acid | | 2.77 | | 3.16 |
| isobutyric acid | | 0.12 | | 0.24 |
| butyric acid | | 0.20 | | 0 |
| fructose | 11.7 | 5.64 | 14.6 | 6.49 |
| reducing sugars | 12.7 | 3.84 | 10.3 | 5.20 |
| NDF ^a | 58.4 | 62.3 | 50.9 | 53.5 |
| NDF glucose | 26.2 | 28.5 | 22.3 | 24.5 |
| hemicellulose ^b | 18.7 | 18.3 | 16.4 | 16.5 |
| lignin | 5.86 | 7.12 | 5.66 | 6.56 |
| silica | 3.32 | 3.82 | 3.68 | 4.64 |

^a Neutral detergent fiber. ^b Based on monosaccharide analysis.

Chemists, 1984). Silage samples were extracted with distilled water at 0 °C; this extract was used for pH, volatile fatty acids (VFA) (Ben-Ghedalia and Yosef, 1989c), lactic acid (Pryce, 1969), and reducing sugars (Miller, 1959) determination.

Another portion of the silages and the corresponding source materials was freeze-dried, ground to 1 mm, and used for measuring and preparing NDF according to the method of Van Soest et al. (1991), employing the amylase procedure. Permanganate-lignin and silica were determined according to the method of Van Soest et al. (1991).

To extract fructose, 5 g DM of freeze-dried and ground samples of source materials and silages underwent extraction with 100 mL of 0.1 N HCl for 1 h. Total fructose in the extracts was determined according to the procedure of Boratynski (1984).

Monosaccharide components of the freeze-dried source materials and silages and their NDF preparations were determined after hydrolysis with 24 N H₂SO₄ for 1 h at 21 °C followed by 1 N H₂SO₄ for 5 h at 100 °C, as described by Ben-Ghedalia and Miron (1984). The free sugars released were converted to alditol acetates and determined by GLC (Blakeney et al., 1983). Uronic acids in the hydrolysates were determined colorimetrically (Blumenkrantz and Asboe-Hansen, 1973).

In vitro digestibility of source materials and silages was determined (in four tubes = replicates) according to the two-stage fermentation technique of Tilley and Terry (1963). Residual DM, NDF, and monosaccharide components remaining in each fermentation tube were determined as described above.

Results were analyzed statistically by using a completely randomized design (Little and Hills, 1978) consisting of two stages of maturity (treatments), with four replicates (silos) per treatment.

RESULTS AND DISCUSSION

Silage Fermentation. The monosaccharide composition of the source materials is shown in Table 1. The major change between the bloom and soft-dough stages was expressed in a doubling of the proportion of NDS

Table 3. Recovery after 60 days of Fermentation (Percent of Pre-ensilage Component)^a of Neutral Detergent Fiber (NDF) Monosaccharide Components in Wheat Silages Harvested at Two Stages of Maturity

| monosaccharide | bloom | soft-dough | SEM |
|---------------------------|-------------------|-------------------|------|
| glucose | 95.9 | 95.0 | 1.41 |
| xylose | 87.0 | 88.7 | 1.54 |
| arabinose | 76.0 | 80.9 | 1.90 |
| uronic acids | 89.6 | 90.4 | 1.18 |
| galactose | 70.6 ^a | 77.2 ^b | 1.65 |
| mannose | 59.4 ^a | 89.5 ^b | 1.47 |
| total NDF monosaccharides | 91.4 | 91.7 | 1.31 |

^a At day = 0, each component was regarded as 100%. Values in the same line marked by different superscripts are significantly different, *P* < 0.05.

glucose due to starch accumulation in the spikes. Table 2 presents the composition of silages and source materials. Theoretically, starch may participate in the process of silage fermentation (Henderson, 1993). However, the high concentrations of fructose (11.7 and 14.6%) and reducing sugars (10.6 and 12.7%) in the wheat plants prove that, in this study, there was no shortage in readily fermentable carbohydrates. Both silages were good; pH values of 4.47 and 4.24 for the bloom and soft-dough silages reflect the concentrations of lactic acid, 3.18 and 4.28%, respectively. Along with the decrease in the concentration of the fermentable carbohydrates, there was an increase in the content of the fibrous materials, in both silages. This point is worth mentioning since it is probably associated with changes in digestibility following ensilage.

The participation of NDF monosaccharide constituents in the process of silage fermentation is shown in Table 3. The general view is that NDF monosaccharides contribute very little to that process. This is, however, based on very sparse information. For instance, 100% recovery of NDF glucose and xylose residues after silage fermentation was found with lucerne (Ben-Ghedalia and Yosef, 1989a,b), whereas Morrison (1979) reported losses of 5% for cellulose and 10–20% for hemicellulose in grasses. Table 3 is the first to show the relevant data on wheat. The general picture is that NDF monosaccharide residues are being solubilized and, most likely, partly utilized during the ensilage of wheat. The extent of solubilization was 4–5, 11–13, 19–24, and 10% for NDF glucose, xylose, arabinose, and uronic acids, respectively. A total of 8–9% NDF monosaccharides were solubilized.

The recovery of NDS monosaccharides, fructose, and total reducing sugars is shown in Table 4. Glucose and galactose were the most fermentable sugars among the NDS monosaccharides, having the major and significant contribution in the bloom wheat silages. Since during ensilage there is a transfer of monosaccharides from NDF to NDS (Table 3), it is difficult to assess precisely the contribution of NDS sugars to the fermentation

Table 4. Recovery after 60 days of Fermentation (Percent of Pre-ensilage Component)^a of Neutral Detergent Soluble Monosaccharides (NDS), Fructose, and Total Reducing Sugars in Wheat Silages Harvested at Two Stages of Maturity

| component | bloom | soft-dough | SEM |
|---------------------------|-------------------|-------------------|------|
| glucose | 16.4 ^a | 56.2 ^b | 5.41 |
| xylose | 147 | 320 | 59.0 |
| arabinose | 183 | 152 | 15.5 |
| uronic acids | 86.3 | 87.5 | 2.92 |
| galactose | 35.2 ^a | 76.4 ^b | 3.99 |
| mannose | 98.7 ^a | 594 ^b | 27.6 |
| total NDS monosaccharides | 36.6 ^a | 78.6 ^b | 6.08 |
| fructose | 42.7 | 38.9 | 2.19 |
| total reducing sugars | 26.8 ^a | 44.1 ^b | 3.06 |

^a At day = 0, each component was regarded as 100%. Values in the same line marked by different superscripts are significantly different, $P < 0.05$.

process. Soluble xylose and arabinose can participate in silage fermentation. Chamberlain (1988) had shown that 50–70% of added xylose was fermented during ryegrass fermentation, and in our previous studies (Ben-Ghedalia and Yosef, 1989a,b) up to 80% of the indigenous ND arabinose found in lucerne was fermented during ensilage. The high recoveries of these ND sugars in the present study (Table 4) indicate that their role in ensilage was very limited, which is not surprising in view of the high concentrations of fructose and NDS glucose found in the source materials (Tables 1 and 2). Lactic acid bacteria are devoid of pectinases (Ben-Ghedalia et al., 1993), and therefore galacturonic acid, the major building block of the rhamnogalacturonans, is not consumed during ensilage (Ben-Ghedalia and Yosef, 1989a; Ben-Ghedalia et al., 1991, 1993). Mannose is usually a minor monosaccharide component found largely in the NDS and may be consumed during silage fermentation, particularly when there is a shortage in the preferable substrates (Ben-Ghedalia and Yosef, 1989a,b). The significantly very high recovery of mannose in the soft-dough silage (600%, due to an increase of its content from 0.48 to 2.20%) was probably a result of microbial biosynthesis. Fructose was equally fermented in both silages; however, the significantly different recoveries of total reducing sugars found in the silages of this study, and the mannose phenomenon mentioned above, hint at the possibility of different fermentation patterns. In this context, the possible participation of starch in the ensilage of the soft-dough plants cannot be excluded.

In Vitro Digestibility. Carbohydrate digestibility was determined recently in fresh wheat plants harvested at the bloom and soft-dough stages (Ben-Ghedalia et al., 1995). Wheat is usually preserved as silage; however, in view of the changes occurring in the carbohydrate fractions during ensilage (Tables 3 and 4), it is important to compare the digestibility (D) of these fractions between source materials and silages and between the silages themselves. This is shown for the NDF monosaccharide residues in Table 5. Generally, silages were less digestible than the source materials, but the difference was larger in the bloom than in the soft-dough treatment. Except for NDF galactose, the decline in the digestibility of all the other NDF monosaccharides following ensilage was greater in the bloom stage than in the soft-dough stage. The decline in total NDF monosaccharide D was 5.1 and 2.5 percentage units in the bloom and soft-dough stages, respectively. Thus, a gap of 8.1 D percentage units in total NDF

Table 5. In Vitro Digestibility (Percent)^a of NDF Monosaccharides of Wheat Source Materials (SM) and Silages Harvested at Two Stages of Maturity

| component | bloom | | soft-dough | | SEM |
|---------------------------|-------------------|-------------------|-------------------|-------------------|------|
| | SM | silage | SM | silage | |
| glucose | 66.4 ^a | 62.4 ^b | 58.2 ^c | 55.6 ^d | 0.54 |
| xylose | 63.4 ^a | 56.9 ^b | 53.0 ^c | 51.8 ^c | 0.61 |
| arabinose | 78.8 ^a | 71.6 ^c | 77.0 ^b | 72.4 ^c | 0.35 |
| uronic acids | 67.1 ^a | 62.9 ^b | 61.3 ^b | 58.3 ^c | 0.53 |
| galactose | 85.8 ^a | 78.1 ^c | 79.8 ^b | 70.8 ^d | 0.34 |
| mannose | 86.2 ^a | 77.2 ^c | 84.7 ^b | 85.0 ^b | 0.24 |
| total NDF monosaccharides | 66.4 ^a | 61.3 ^b | 58.3 ^c | 55.8 ^d | 0.55 |
| NDF | 63.3 ^a | 57.8 ^b | 56.3 ^b | 50.8 ^c | 0.62 |

^a Values in the same line marked by different superscripts are significantly different, $P < 0.05$.

Table 6. In Vitro Digestibility (Percent)^a of Whole Plant Monosaccharide Residues of Source Materials (SM) and Silages Harvested at Two Stages of Maturity

| component | bloom | | soft-dough | | SEM |
|-----------------------|-------------------|-------------------|-------------------|--------------------|------|
| | SM | silage | SM | silage | |
| glucose | 73.1 ^a | 60.4 ^c | 74.9 ^a | 65.7 ^b | 0.60 |
| xylose | 65.5 ^a | 55.3 ^c | 58.8 ^b | 57.1 ^{bc} | 0.82 |
| arabinose | 79.1 ^a | 71.4 ^c | 79.6 ^a | 75.5 ^b | 0.79 |
| uronic acids | 75.2 ^a | 69.7 ^b | 73.0 ^a | 67.6 ^b | 0.91 |
| galactose | 82.2 ^a | 62.6 ^d | 75.3 ^b | 67.2 ^c | 0.82 |
| mannose | 87.7 ^b | 89.6 ^b | 90.6 ^b | 97.2 ^a | 0.97 |
| total monosaccharides | 71.7 ^a | 60.2 ^c | 71.6 ^a | 65.5 ^b | 0.60 |
| organic matter | 69.2 ^a | 59.0 ^c | 70.5 ^a | 65.6 ^b | 0.51 |

^a Values in the same line marked by different superscripts are significantly different, $P < 0.05$.

monosaccharide D between the source materials was reduced to 5.5 D percentage units between the silages. This picture is not reflected in the NDF fraction, which comprises also the phenolic component. Therefore, in this study as in many others, NDF monosaccharide digestibility data are better measurement for differentiating treatment effect.

The D values of the monosaccharide residues in whole plant material is shown in Table 6. The presence of starch in the soft-dough source material compensated for the decline in NDF glucose D due to maturation. The decline in D of total monosaccharide residues of the whole plant following ensilage was 11.5 and 6 percentage units for the bloom and soft-dough stages, respectively. Thus, while the D of total monosaccharide residues was equal within the source materials, it was higher by 5.3 percentage units in the soft-dough silage than in the bloom one. Accordingly, the D of the individual monosaccharide components except the uronic acids was higher in the soft-dough silage than in the bloom silage. It was found in a recent study that between the bloom and soft-dough stages there is a DM yield increase of 36% (Ben-Ghedalia et al., 1995). In the present study, carbohydrate D of the soft-dough silage was 9% higher than that of the bloom silage; thus, by harvesting wheat for silage at that stage, the yield gain in terms of digestible energy would be more than 45%.

For reaching the right conclusion regarding the optimal stage of harvesting wheat for silage, it is recommended that both source materials and silages be evaluated at the monosaccharide level.

LITERATURE CITED

- Arieli, A.; Adin, G. Effect of wheat silage maturity on digestion and milk yield in dairy cows. *J. Dairy Sci.* **1994**, *77*, 237–243.

- Ashbell, G.; Kashanchi, Y. In silo losses from wheat ensiled in bunker silos in a subtropical climate. *J. Sci. Food Agric.* **1987**, *40*, 95–103.
- Ashbell, G.; Theune, H. H.; Sklan, D. Changes in the amino acid compounds of the whole crop wheat during ensiling and after fermentation. *J. Sci. Food Agric.* **1983**, *34*, 321–327.
- Association of Official Analytical Chemists. *Official Methods of Analysis*, 14th ed.; AOAC: Washington, DC, 1984.
- Ben-Ghedalia, D.; Miron, J. The digestion of total and cell wall monosaccharides of alfalfa by sheep. *J. Nutr.* **1984**, *114*, 880–887.
- Ben-Ghedalia, D.; Yosef, E. SO₂-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation. *Anim. Feed Sci. Technol.* **1989a**, *22*, 255–262.
- Ben-Ghedalia, D.; Yosef, E. Ozonated cotton stalks as a silage additive: the participation of pectin and cell wall monosaccharides in lucerne silage fermentation. *J. Anim. Physiol. Anim. Nutr.* **1989b**, *61*, 105–110.
- Ben-Ghedalia, D.; Yosef, E. SO₂-Treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation. *Anim. Feed Sci. Technol.* **1989c**, *22*, 247–254.
- Ben-Ghedalia, D.; Miron, J.; Yosef, E. Fermentation of alfalfa silages with ozonated cotton stalks added. *J. Agric. Food Chem.* **1991**, *39*, 2002–2005.
- Ben-Ghedalia, D.; Yosef, E.; Miron, J. Pectin fermentation and utilization by natural microflora during ryegrass ensilage. *Anim. Feed Sci. Technol.* **1993**, *41*, 113–119.
- Ben-Ghedalia, D.; Kabala, A.; Miron, J. Composition and *in vitro* digestibility of carbohydrates of wheat plants harvested at bloom and soft-dough stages. *J. Sci. Food Agric.* **1995**, *68*, 111–116.
- Blakeney, A. B.; Harris, P. J.; Henry, R. J.; Stone, B. A. A simple and rapid preparation of alditol acetates for monosaccharides analysis. *Carbohydr. Res.* **1983**, *113*, 291–299.
- Blumenkrantz, N.; Asboe-Hansen, G. New method for quantitative determination of uronic acids. *Anal. Biochem.* **1973**, *54*, 484–489.
- Boratynski, J. Colorimetric method for the determination of ketoses using phenol-acetone-boric acid reagent (PABR). *Anal. Biochem.* **1984**, *137*, 528–532.
- Campbell, C.; Taylor, K.; Matsouka, S.; Marshall, S.; Buchanan-Smith, J. G. Inoculants and enzymes as additives for lucerne silage with measurements of changes in structural carbohydrates and pectin during the ensiling period. *Proc. 9th Silage Conf.*, Newcastle upon Tyne, Sept 1990; pp 14–15.
- Chamberlain, D. G. Effect of added glucose and xylose on the fermentation of perennial ryegrass silage inoculated with *Lactobacillus plantarum*. *J. Sci. Food Agric.* **1988**, *46*, 129–138.
- Henderson, N. Silage additives. *Anim. Feed Sci. Technol.* **1993**, *45*, 35–56.
- Little, T. M.; Hills, F. J. In *Agricultural Experimentation Design and Analysis*; Wiley: New York, 1978, pp 48–75.
- Miller, G. L. Use of dinitrosalicylic acid reagent for determination of reducing sugars. *Anal. Chem.* **1959**, *31*, 426–428.
- Morrison, I. M. Changes in the cell wall components of laboratory silages and the effect of various additives of these changes. *J. Agric. Sci. Camb.* **1979**, *93*, 581–586.
- Pryce, J. D. A. Modification of the Barker-Summerson method for the determination of lactic acid. *Analyst* **1969**, *94*, 1151–1152.
- Tilley, J. M.; Terry, R. M. A two stage technique for the *in vitro* digestion of forage crops. *J. Br. Grassl. Soc.* **1963**, *18*, 104–111.
- Van-Soest, P. J.; Robertson, J. B.; Lewis, B. A. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* **1991**, *74*, 3583–3597.
- Weinberg, Z. G.; Ashbell, G.; Hen, Y.; Harduf, Z. Ensiling whole wheat for ruminant feeding at different stages of maturity. *Anim. Feed Sci. Technol.* **1991**, *32*, 313–320.

Received for review March 7, 1995. Accepted June 12, 1995.*
JF950137M

* Abstract published in *Advance ACS Abstracts*, August 1, 1995.