

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231435303>

# Synthesis and Spectroscopy of Tricyclo(3.3.3.0<sup>3,7</sup>)undec-3(7)-ene: Confirmation of Computational Predictions Regarding the Effects of Pyramidalization on Alkene Ionization Energies...

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · MAY 1993

Impact Factor: 12.11 · DOI: 10.1021/ja00062a074

---

CITATIONS

26

---

READS

10

8 AUTHORS, INCLUDING:



**Knut R Asmis**

University of Leipzig

103 PUBLICATIONS 2,424 CITATIONS

SEE PROFILE

# Synthesis and Spectroscopy of Tricyclo[3.3.3.0<sup>3,7</sup>]undec-3(7)-ene: Confirmation of Computational Predictions Regarding the Effects of Pyramidalization on Alkene Ionization Energies and Electron Affinities

Joseph M. Smith, David A. Hrovat, and Weston Thatcher Borden\*

Department of Chemistry  
University of Washington  
Seattle, Washington 98195


Michael Allan,\* Knut R. Asmis, Christophe Bulliard, Edwin Haselbach, and Urs C. Meier

Institut de Chimie Physique de l'Université  
CH-1700 Fribourg, Switzerland

Received February 5, 1993

Revised Manuscript Received March 22, 1993


We have reported the syntheses of the  $n = 1^1$  and  $2^2$  members of a homologous series of pyramidalized alkenes (**1**),<sup>3</sup> their spectroscopic study in matrix isolation,<sup>4,5</sup> and some of their chemistry.<sup>6</sup> The generation of a benzo derivative of **1**,  $n = 2$ ,<sup>7</sup> and a bis-ethano derivative of the  $n = 0$  member of this series have also been described.<sup>8</sup> In this communication we report the synthesis and some spectroscopy of **1**,  $n = 3$ .



Previous attempts to prepare diol **2**, a promising precursor of **1**,  $n = 3$ , have been unsuccessful. Double ring expansion of bicyclo[3.2.2]nonane-6,8-dione,<sup>9</sup> followed by transannular reductive ring closure,<sup>10</sup> failed when the ring expansion proceeded with the undesired regiochemistry.<sup>11</sup> Efforts to prepare **2** from diiodide **3** by a ring closure, analogous to the one that proved successful in the synthesis of the 10-selena derivative of **1**,  $n = 3$ ,<sup>12</sup> foundered when both dithiane and malonate ester monoadducts of **3** failed to undergo cyclization under a wide variety of conditions.<sup>13</sup>

A third route to **2**, which incorporates features of the two unsuccessful pathways, has resulted in its synthesis.<sup>14</sup> As shown in Scheme I, we have formally inserted a methylene group between

Scheme I<sup>a</sup>



<sup>a</sup> (a) Na, TMSCl, toluene, reflux, 67%; (b) CH<sub>3</sub>OH, reflux, 84%; (c) acetyl chloride, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, 82%; (d) SmI<sub>2</sub>,<sup>14</sup> THF, 91%; (e) N<sub>2</sub>CHCO<sub>2</sub>Et, Et<sub>3</sub>OBf<sub>4</sub>,<sup>15</sup> CH<sub>2</sub>Cl<sub>2</sub>, 45%; (f) 5% NaOH, 98%; (g) *p*-dioxane, reflux, 82%; (h) N<sub>2</sub>H<sub>4</sub>, NaOH, diethylene glycol, 185 °C, 78%; (i) 20% AcOH, 85 °C, 81%; (j) CH<sub>3</sub>Li, THF, then MsCl, 55%.

the two carboxylic carbons of diacid **4**<sup>12</sup> by performing an acyloin ring closure on dimethyl ester **5**, reductively removing<sup>15</sup> the hydroxyl group from **7**, and then ring expanding<sup>16</sup> ketone **9**. As in Leonard's synthesis of tricyclo[3.3.3]undecane (manxane),<sup>17</sup> the regiochemistry of the ring expansion step is irrelevant, since **12a** and **12b** both undergo Wolf-Kishner reduction to **13**. Hydrolysis of **13** affords diol **2** in 9.6% overall yield from **4**.

Diol **2** was converted to dimesylate **14**, which was reduced to **1**,  $n = 3$ , using either sodium naphthalide<sup>12,18</sup> or, more conveniently, sodium amalgam in ether. The latter method allows ether solutions of pure olefin to be obtained simply by filtering the reaction mixture. Unlike its lower homologues,<sup>1-8</sup> **1**,  $n = 3$ , is stable to dimerization at room temperature; but it reacts very rapidly with atmospheric oxygen.<sup>19</sup>

Careful exclusion of air from samples of the olefin allowed it to be characterized spectroscopically. A very weak band was observed in the IR at 1615 cm<sup>-1</sup>, which corresponds to the strongest band in the Raman spectrum and is assigned to the C=C stretch in the olefin. An RHF/3-21G<sup>20</sup> vibrational analysis<sup>21</sup> gives a frequency of 1851 cm<sup>-1</sup> for this mode. After scaling by 0.873,<sup>22</sup> in order to account for the effects of anharmonicity and electron correlation,<sup>24</sup> the C=C stretching frequency in **1**,  $n = 3$ , is predicted to occur at 1615 cm<sup>-1</sup>.

The C=C stretching frequency found in **1**,  $n = 3$ , is 10 cm<sup>-1</sup> lower than that in its 10-selena analog.<sup>12</sup> The lower C=C stretching frequency in and greater reactivity toward O<sub>2</sub> of the

(1) Renzoni, G. E.; Yin, T.-K.; Borden, W. T. *J. Am. Chem. Soc.* **1986**, *108*, 7121.

(2) Renzoni, G. E.; Yin, T.-K.; Miyake, F.; Borden, W. T. *Tetrahedron* **1986**, *42*, 1581.

(3) Review of pyramidalized alkenes: Borden, W. T. *Chem. Rev.* **1989**, *89*, 1095.

(4) Radziszewski, J. G.; Yin, T.-K.; Renzoni, G. E.; Hrovat, D. A.; Borden, W. T.; Michl, J. *J. Am. Chem. Soc.* **1993**, *115*, 1454.

(5) Radziszewski, J. G.; Yin, T.-K.; Miyake, F.; Renzoni, G. E.; Borden, W. T.; Michl, J. *J. Am. Chem. Soc.*, **1986**, *108*, 3544.

(6) Yin, T.-K.; Radziszewski, J. G.; Renzoni, G. E.; Downing, J. W.; Michl, J.; Borden, W. T. *J. Am. Chem. Soc.* **1987**, *109*, 820.

(7) Greenhouse, R.; Borden, W. T.; Hirotsu, K.; Clardy, J. *J. Am. Chem. Soc.* **1977**, *99*, 1664.

(8) Branan, B. M.; Paquette, L. A.; Hrovat, D. A.; Borden, W. T. *J. Am. Chem. Soc.* **1992**, *114*, 774.

(9) Wood, G.; Woo, E. P. *Can. J. Chem.* **1968**, *46*, 3173.

(10) Borden, W. T.; Ravindranathan, T. *J. Org. Chem.* **1971**, *36*, 4125.

(11) Greenhouse, R.; Borden, W. T.; Ravindranathan, T.; Hirotsu, K.; Clardy, J. *J. Am. Chem. Soc.* **1977**, *99*, 6955.

(12) Hrovat, D. A.; Miyake, F.; Trammell, G.; Gilbert, K. E.; Mitchell, J.; Clardy, J.; Borden, W. T. *J. Am. Chem. Soc.* **1987**, *109*, 5524.

(13) Smith, J. M. Ph.D. Thesis, University of Washington, 1993.

(14) Experimental details as well as spectroscopic and analytical data for **1**,  $n = 3$ , **2**, and **5-14** are available as supplementary material.

(15) Molander, G. A.; Hahn, G. *J. Org. Chem.* **1986**, *51*, 1135.

(16) Mock, W. L.; Hartman, M. E. *J. Org. Chem.* **1977**, *42*, 459, 466.

(17) Coll, J. C.; Leonard, N. J. *J. Am. Chem. Soc.* **1970**, *92*, 6685. Coll, J. C.; Crist, D. R.; Barrio, M. G.; Leonard, N. J. *J. Am. Chem. Soc.* **1972**, *94*, 7092. See also: Parker, W.; Doyle, M. *Tetrahedron Lett.* **1970**, 3619.

(18) Carnahan, J. C.; Closson, W. D. *Tetrahedron Lett.* **1972**, 3447.

(19) The three products formed—epoxide, allylic alcohol, and diketone—are those expected from exposure of this pyramidalized alkene to oxygen. See, for example: Wiberg, K. B.; Maturro, M. G.; Okarma, P. J.; Jason, M. E. *J. Am. Chem. Soc.* **1984**, *106*, 2194. Bartlett, P. D.; Banavali, R. *J. Org. Chem.* **1991**, *56*, 6043 and references cited therein.

(20) Binkley, J. S.; Pople, J. A.; Hehre, W. J. *J. Am. Chem. Soc.* **1980**, *102*, 939.

(21) Performed, using Gaussian 90: Frisch, M. J.; Head-Gordon, M.; Trucks, G. W.; Foresman, J. B.; Schlegel, H. B.; Raghavachari, K.; Robb, M.; Binkley, J. S.; Gonzalez, C.; Defrees, D. J.; Fox, D. J.; Whiteside, R. A.; Seeger, R.; Melius, C. F.; Baker, J.; Martin, R. L.; Kahn, L. R.; Stewart, J. P.; Topiol, S.; and Pople, J. A.; Gaussian, Inc., Pittsburgh, PA, 1990.

(22) This scaling factor is the average of those required to fit the RHF/3-21G C=C stretching frequencies for **1**,  $n = 2$  (1811 cm<sup>-1</sup>), and bicyclo[3.3.0]oct-1(5)-ene (1898 cm<sup>-1</sup>) to the experimental values (respectively, 1557<sup>23</sup> and 1675<sup>23,1685</sup> cm<sup>-1</sup>).

(23) Warner, P.; La Rose, R.; Schleis, T. *Tetrahedron Lett.* **1974**, 1409.

(24) Hehre, W. J.; Random, L.; Schleyer, P. v. R.; Pople, J. A. *Ab Initio Molecular Orbital Theory*; Wiley: New York, 1986; pp 226–261.

hydrocarbon is consistent with the fact that the doubly bonded carbon atoms in it are calculated to have larger pyramidalization angles ( $\phi = 25.0^\circ$  and  $25.2^\circ$ )<sup>25</sup> than those found in a salt of the selenium analog.<sup>12</sup>

Both the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of **1**,  $n = 3$ , are temperature dependent. The identity of the protons and carbons whose resonances broaden and coalesce indicates that the dynamic process responsible for these spectral changes is flipping of the trimethylene bridge. From the frequency separations at low temperatures and the temperatures,  $T_c$ , at which coalescence was observed in the resonances for the *exo* and *endo* allylic protons and the allylic and olefinic carbons, values of  $\Delta G^\ddagger = 14.6 - 15.0 \pm 0.2$  kcal/mol for bridge flipping were obtained at  $T_c = 304 - 328 \pm 3$  K.<sup>26</sup> In very good agreement with these NMR results, MM2 calculations<sup>27</sup> predict that trimethylene bridge flipping should require 14.4 kcal/mol.

The average chemical shift of  $\delta$  157.28 for the olefinic carbons in **1**,  $n = 3$ , is larger than that of  $\delta$  150.74 in the 10-selena derivative<sup>12</sup> and that of  $\delta$  146.0<sup>28</sup> in bicyclo[3.3.0]oct-1(5)-ene (BCO),<sup>29</sup> the unbridged reference olefin. The monotonic change in these  $^{13}\text{C}$  chemical shifts with increasing pyramidalization indicates that pyramidalization shifts the resonances for olefinic carbons to lower fields.<sup>30</sup>

Olefin **1**,  $n = 3$ , exhibits a UV absorption with  $\lambda_{\text{max}} = 217 \pm 5$  nm in pentane. The more highly pyramidalized olefin, **1**,  $n = 2$ , has a UV absorption at even longer wavelength,  $\lambda_{\text{max}} = 245 \pm 15$  nm.<sup>5</sup> In contrast, BCO does not show a UV maximum above 200 nm in solution. We find the maximum for the  $\pi \rightarrow \pi^*$  singlet excitation in the electron energy loss (EEL) spectrum of BCO to be at 6.54 eV, which is close to the 6.61 eV reported for tetramethylethylene (TME)<sup>31</sup> and corresponds to a wavelength of 190 nm.

The experimental finding of a red shift in  $\lambda_{\text{max}}$  with increasing olefin pyramidalization is consistent with the results of *ab initio* calculations.<sup>25,32</sup> However, the calculations also predict that lowering the energy of the LUMO, rather than raising that of the HOMO, plays the dominant role in decreasing the HOMO-LUMO energy separation.<sup>33</sup> For example, RHF/3-21G calculations on **1**,  $n = 3$ , find that the HOMO energy is 0.25 eV higher than that of BCO; whereas the LUMO energy of **1**,  $n = 3$ , is computed to be 0.79 eV lower than that of BCO.<sup>25</sup> The predicted changes in the HOMO and LUMO energies appear to be rather insensitive to the basis set used.<sup>34</sup>

HOMO and LUMO energies can be related, respectively, to ionization energies (IEs) and electron affinities (EAs) via Koopmans' theorem.<sup>35</sup> We have used photoelectron (PE)<sup>36</sup> spectroscopy to measure the IEs of **1**,  $n = 3$ , and BCO and employed electron transmission (ET)<sup>37</sup> spectroscopy to measure their EAs. These spectra<sup>14</sup> confirm the prediction that the effect

of pyramidalization on lowering the energy of the LUMO is much greater than its effect on raising the energy of the HOMO.<sup>38</sup>

The PE spectrum of BCO gives an adiabatic IE of 8.11 eV, which is close to that of 8.27 eV reported for TME.<sup>41</sup> The adiabatic IE of **1**,  $n = 3$ , is 7.80 eV, which is lower than that of BCO by 0.31 eV. This difference between the measured IEs is in very good agreement with the difference between the calculated HOMO energies.<sup>25,34,42</sup>

The ET spectrum of BCO gives a vertical EA of -2.44 eV, which again is close to the EA of -2.27 reported for TME.<sup>43</sup> However, the ET spectrum of **1**,  $n = 3$ , gives an EA of -1.70 eV. The 0.74 eV greater electron affinity of **1**,  $n = 3$ , is also in very good agreement with the difference between the calculated LUMO energies.<sup>25,34</sup>

This good agreement suggests that the calculations might also be able to predict which members of this series of pyramidalized olefins will form bound radical anions in the gas phase.<sup>44</sup> After scaling the computed difference of 3.50 eV between the LUMO energies of BCO and **1**,  $n = 0$ ,<sup>25</sup> by the difference of 0.74 eV between the experimental EAs of BCO and **1**,  $n = 3$ , divided by the difference of 0.79 eV between their calculated LUMO energies,<sup>25</sup> our experimental value of EA = -2.44 eV for BCO indicates that **1**,  $n = 0$ , should have a vertical EA that is positive by 0.84 eV (0.26 eV if RHF/6-31G\*//TCSCF/6-31G\* LUMO energies<sup>34</sup> are used). Therefore, this most highly pyramidalized ( $\phi = 61.2^\circ$ )<sup>25</sup> member of the homologous series of alkenes **1** is predicted to form a bound radical anion.<sup>46</sup> The chemistry observed for a bis-ethano derivative of **1**,  $n = 0$ ,<sup>8</sup> is consistent with the prediction of a LUMO of very low energy in this alkene.

**Acknowledgment.** We thank the National Science Foundation for support of this work and Professor John W. Macklin of the UW Chemistry Department for his help in obtaining the Raman spectrum of the title olefin. The work done in Switzerland is part of Project 20-34071.92 of the Schweizerischer Nationalfonds zur Förderung der wissenschaftlichen Forschung.

**Supplementary Material Available:** Synthetic details as well as spectroscopic and analytical data for **1**,  $n = 3$ , **2**, and **5-14** and figures containing photoelectron spectra and electron transmission spectra of **1** and BCO (21 pages). Ordering information is given on any current masthead page.

(37) Reviews: Jordan, K. D.; Burrow, P. D. *Chem. Rev.* **1987**, *87*, 557. Allan, M. J. *Electron Spectrosc. Relat. Phenom.* **1989**, *48*, 219. The latter review contains a description of the spectrometer used in these experiments.

(38) A similar prediction, regarding the effect of alkyne bending,<sup>32</sup> has also been confirmed by IE<sup>39</sup> and EA<sup>40</sup> measurements.

(39) Batich, E.; Ermer, O.; Heilbronner, E.; Wiseman, J. R. *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 312. Schmidt, H.; Schweig, A.; Krebs, A. *Tetrahedron Lett.* **1974**, 3237.

(40) Ng, L.; Jordan, K. D.; Krebs, A.; Rüger, W. *J. Am. Chem. Soc.* **1982**, *104*, 7414.

(41) Moscatel, P.; Grosjean, D.; Mouvier, G.; Dubois, J. *J. Electron Spectrosc. Relat. Phenom.* **1973**, *2*, 225. Mintz, D. M.; Kupperman, A. *J. Chem. Phys.* **1979**, *71*, 3499.

(42) A preliminary PE spectrum of **1**,  $n = 2$ , obtained by pyrolyzing the  $\beta$ -lactone precursor<sup>2</sup> and passing the pyrolysate directly into the inlet of the spectrometer, gives an adiabatic IE for **1**,  $n = 2$ , which, as predicted,<sup>25,34</sup> is lower than that of **1**,  $n = 3$ , by less than 0.1 eV.

(43) Jordan, K. D.; Burrow, P. D. *J. Am. Chem. Soc.* **1980**, *102*, 6882.

(44) Extrapolation of the EAs measured for cyclic alkynes allowed Ng et al. to predict that benzyne radical anion should be bound,<sup>40</sup> a prediction that was subsequently verified experimentally.<sup>45</sup>

(45) Leopold, D. G.; Miller, A. E. S.; Lineberger, W. C. *J. Am. Chem. Soc.* **1986**, *108*, 1379.

(46) Other scaling factors that have been used to fit calculated LUMO energies to observed EAs<sup>47</sup> predict a vertical EA for **1**,  $n = 0$ , that is positive,<sup>47a</sup> very close to zero,<sup>47b,c</sup> or slightly negative.<sup>47c</sup> However, even if the vertical EA were slightly negative, geometry relaxation in the radical anion could cause the adiabatic EA to be positive.

(47) (a) Chou, P. K.; Kass, S. R. *J. Am. Chem. Soc.* **1991**, *113*, 697. (b) Chen, D.; Gallup, G. A. *J. Chem. Phys.* **1990**, *93*, 8893. (c) Staley, S. W.; Strnad, J. T. *Chem. Phys. Lett.* **1992**, *200*, 527.

(25) Hrovat, D. A.; Borden, W. T. *J. Am. Chem. Soc.* **1988**, *110*, 4710.

(26) Taking the rate constant at coalescence to be  $(\nu_A - \nu_B)\pi/(2)^{1/2}$ ; Gutowsky, H. S.; Holm, C. H. *J. Chem. Phys.* **1956**, *25*, 1228.

(27) Allinger, N. L. *J. Am. Chem. Soc.* **1977**, *99*, 8127.

(28) Becker, K. B. *Helv. Chim. Acta* **1977**, *60*, 68.

(29) Corey, E. J.; Block, E. J. *Org. Chem.* **1969**, *34*, 1233. Paquette, L. A.; Hauser, R. W. *J. Am. Chem. Soc.* **1969**, *91*, 3870. Borden, W. T.; Ravindranathan, T. *J. Org. Chem.* **1971**, *36*, 4125. Borden, W. T.; Concannon, P. W.; Phillips, D. I. *Tetrahedron Lett.* **1973**, 3161.

(30) Alkyne bending has the same effect on the chemical shifts of acetylenic carbons: Krebs, A.; Kimling, H. *Justus Liebigs Ann. Chem.* **1974**, 2074.

(31) Merer, A. J.; Mulliken, R. S. *Chem. Rev.* **1969**, *69*, 639.

(32) Strozier, R. W.; Caramella, P.; Houk, K. N. *J. Am. Chem. Soc.* **1979**, *101*, 1340.

(33) The larger magnitude of the calculated change in the LUMO energy can be easily understood by considering the changes in both the overlap and the hybridization of the AOs of which these MOs are comprised.<sup>25</sup>

(34) RHF/6-31G\*//TCSCF/6-31G\* calculations put the HOMO energies of **1**,  $n = 3-0$  respectively 0.30, 0.41, 0.57, and 0.60 eV above the HOMO energy of BCO and the LUMO energies 0.88, 1.53, 2.24, and 3.21 eV below the LUMO energy of BCO.

(35) Koopmans, T. *Physica* **1933**, *1*, 104.

(36) Measured with a modified Turner-type spectrometer: Dressler, R.; Neuhaus, L.; Allan, M. J. *Electron Spectrosc. Relat. Phenom.* **1983**, *31*, 181.