



## Calorie restriction and reversal of Type 2 diabetes

Roy Taylor

**To cite this article:** Roy Taylor (2016): Calorie restriction and reversal of Type 2 diabetes, Expert Review of Endocrinology & Metabolism, DOI: [10.1080/17446651.2016.1239525](https://doi.org/10.1080/17446651.2016.1239525)

**To link to this article:** <http://dx.doi.org/10.1080/17446651.2016.1239525>



Accepted author version posted online: 21 Sep 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

**Publisher:** Taylor & Francis

**Journal:** *Expert Review of Endocrinology & Metabolism*

**DOI:** 10.1080/17446651.2016.1239525

## **Calorie restriction and reversal of Type 2 diabetes**

**Roy Taylor**

**Magnetic Resonance Centre, Institute for Cellular Medicine, Newcastle University, Newcastle upon Tyne, UK**

Word count main text: 3979

Word count abstract: 117

Figures: 1

Correspondence to:

Professor Roy Taylor,

Newcastle Magnetic Resonance Centre,

Campus for Ageing and Vitality,

Newcastle upon Tyne NE4 5PL

United Kingdom

Telephone: +44 (0)191 222 1152

Fax: +44 (0)191 222 1151

Email: [Roy.Taylor@ncl.ac.uk](mailto:Roy.Taylor@ncl.ac.uk)



## Abstract

**Introduction.** Type 2 diabetes a major global health problems and has been believed to be a lifelong condition with inevitable worsening. Steadily increasing numbers of drugs appeared to be required to achieve even modest control. Early type 2 diabetes has now been shown to be reversed by substantial weight loss and this has allowed temporal tracking of the underlying pathophysiological changes.

**Areas covered.** In early type 2 diabetes, negative calorie balance decreases liver fat within days, and allows return of normal control of hepatic glucose production. Over 8 weeks, the negative calorie balance allows raised levels of intra-pancreatic fat and simultaneously first phase insulin secretion to normalise. These findings are consistent with the 2008 Twin Cycle Hypothesis of the aetiology and pathogenesis of type 2 diabetes. Individuals develop type 2 diabetes when they exceed their personal fat threshold for safe storage of fat and there is no difference in pathophysiology between those with BMI above or below 30 kg/m<sup>2</sup>.

**Expert Commentary.** Type 2 diabetes can now be understood as a state of excess fat in liver and pancreas, and remains reversible for at least 10 years in most individuals.

## 1. Introduction

The pathogenesis of type 2 diabetes has been regarded as involving two separate processes bringing about insulin resistance and relative beta cell failure [1-3]. However, the rapid changes in population prevalence of the condition depending upon food surplus or shortage [4, 5] suggests either that both processes are affected by nutrient supply or that a single pathway is involved. By tracking the pathophysiological changes which occur during the transition from type 2 diabetes back to normal metabolic control it has been possible to test the predictions of the 2008 Twin Cycle Hypothesis [6]. This review discusses these findings in relation to established knowledge.

## 2. Contribution of the individual metabolic organs to type 2 diabetes

### 2.1. Liver

Inside the liver, all triglyceride is stored inside hepatocytes. The sensitivity of the liver to insulin is strongly related to the triglyceride content [7]. As fasting glucose concentration is principally determined by the rate of glucose production from the liver, improving insulin sensitivity of the liver decreases fasting plasma glucose [8, 9].

Excess accumulation of liver fat occurs when the total daily calorie intake exceeds expenditure day after day, and year after year. If more calories are ingested than metabolised in any 24 hour period then: a) any excess fat is stored subcutaneously, in visceral fat or in the liver; b) any excess carbohydrate cannot be stored as such once the glycogen depots are full and it has to be turned into fat by *de novo* lipogenesis. This process only happens in the liver in humans, and triglyceride synthesised by this route is particularly likely to be stored in hepatocytes rather than transported for storage in subcutaneous adipose tissue where it is metabolically inert. As *de novo* lipogenesis is stimulated by insulin, those people who are relatively insulin resistant in muscle - and who therefore have a raised plasma insulin level - are especially likely to accumulate fat in the liver. This could explain the reason why muscle insulin resistance is a very early signal of risk for type 2 diabetes [2, 10].

The practical application of this was elegantly shown by overfeeding sucrose to overweight volunteers for three weeks. The result was a sharp increase in liver triglyceride content [11]. This was associated with a 30% rise in serum ALT, reflecting the associated metabolic stress on the hepatocytes. A larger scale demonstration of the end result of this process was achieved by Sattar and colleagues [12]. During the 15 year follow up of the West of Scotland Coronary Prevention Study group of 6595 men, some developed diabetes. Stored plasma samples were analysed to allow retrospective reconstruction of the pattern of change in liver enzymes in those who were about to develop diabetes compared with the majority who did not develop diabetes. The diagnosis of type 2 diabetes was preceded by a steady rise in serum ALT over at least 18 months. As the mean level was well within the normal range it would be unlikely to be detected in any one individual. Clearly, liver fat builds up during the pre-diabetic phase of type 2 diabetes.

The recent studies of reversal of type 2 diabetes to non-diabetic glucose control show that the first impact of a hypocaloric diet is to decrease liver fat stores. In the Counterpoint study, a diet of 600-800 kcal/day achieved a 30% decrease in liver triglyceride in the first 7 days, bringing it down to the level of weight matched controls [13]. Simultaneously, fasting plasma glucose fell from 9.2 to 5.9mmol/l. A 7 day very low calorie diet was observed to improve fasting plasma glucose with no effect on first phase insulin secretion, and change in hepatic insulin sensitivity was not measured [14]. This effect of sudden negative calorie balance on fasting plasma glucose is seen not only after a purely dietary intervention, but also after the food restriction imposed by bariatric surgery [15].

## *2.2. Muscle*

Lower sensitivity of skeletal muscle to insulin is widely regarded as the earliest feature predicting onset of type 2 diabetes [10]. It prevents muscle from contributing to immediate post-meal storage of glucose as glycogen and hence post-prandial glucose levels rise considerably [16, 17]. However, it is important to recognise that 'insulin resistance' is not a distinct condition but a continuously distributed parameter in any population. The distribution of whole body (largely muscle) insulin sensitivity in type 2 diabetes is entirely within the range of the non-diabetic population but merely towards the lower range [18]. Separation of the distinct contributions of muscle and liver have

shown that early improvement in control of fasting plasma glucose is related only to improvement in liver insulin sensitivity [8, 13]. It has recently been demonstrated that the resumption of normal or near normal diurnal blood glucose control does not require improvement in muscle insulin sensitivity [13, 19]. Although this may at first appear surprising, it is supported by a wide range of earlier observations. Mice totally lacking in skeletal muscle insulin receptors do not develop diabetes [20]. Humans who have the PPP1R3A genetic variant of muscle glycogen synthase cannot store glycogen in muscle after meals but can have normal glucose tolerance [21]. Many normoglycaemic individuals maintain normal blood glucose with a degree of muscle insulin resistance identical to those who develop type 2 diabetes [18].

Examination of individuals at the extremes of the range of insulin sensitivity led to the concept that there might be an underlying primary defect in mitochondrial function in skeletal muscle which predisposed to type 2 diabetes [22]. However, no defect is present in early type 2 diabetes but can be shown to be directly related to ambient plasma glucose concentration [23]. The rates of mitochondrial ATP production can be modified by increasing or decreasing plasma fatty acid concentration [24, 25]. Also, the onset of insulin stimulation of mitochondrial ATP synthesis is slow, gradually increasing over hours and quite distinct from the almost instantaneous onset of insulin's metabolic effects [26]. It appears that there are secondary effects on mitochondrial function of hyperglycaemia and excess fatty acids, but no evidence for a primary mitochondrial defect underlying common type 2 diabetes.

The physiologic importance of muscle insulin resistance is likely to operate over a time period of many years. The presence of longstanding muscle insulin resistance will not of itself cause blood glucose levels to rise but the raised plasma insulin levels will expedite accumulation of liver fat by stimulation of *de novo* lipogenesis [10].

### 2.3. Beta cell

*In vitro* study of islets exposed to a sudden increase in fatty acid concentration demonstrates the interaction between glucose and excess fatty acid supply [27]. When islets from ZDF rats genetically predisposed to develop diabetes are exposed to even a low concentration of palmitic acid for 48 hours, the beta cells no longer secrete insulin normally in response to glucose. However, islets from animals not genetically predisposed to glucose can withstand the excess fat supply and retain glucose mediated

insulin secretion [27]. The elegant studies from Clarke's group have extended knowledge about effects of specific fatty acid upon islet and INS-1 cells and have shown it to be reversible [28-30]. The relevance to human islets has been demonstrated in that exposure to a low concentration of saturated fatty acids brings about avid uptake of fat and loss of glucose mediated insulin secretion [31]. These studies demonstrate potential mechanisms, even though the effect of excess supply of fatty acids *in vivo* is more subtle and over very long time periods.

It is particularly noteworthy that diabetes in the ZDF model is completely preventable by restriction of food intake [32]. These observations together illustrate very well the interaction between genetic susceptibility and environmental factors in the aetiology of type 2 diabetes. In the ZDF rat, a rapid increase in pancreatic fat precedes the onset of hyperglycaemia [27]. This is supported by the observations using a precise method of measuring intra-pancreatic fat in people with type 2 diabetes [13, 19]. However, it must be emphasized that the pancreas is typically one third smaller in type 2 diabetes, with highly irregular borders [33] and that if 'off the shelf' programs to determine pancreas fat are used then falsely high values will be obtained due to visceral fat being included as being within the organ, given its highly irregular shape and small volume in type 2 diabetes.

Post-mortem studies appear to show that beta cell number at diagnosis of type 2 diabetes is around 50%, and lower in long duration diabetes [1, 34]. This appears to agree with the finding that beta cell function is decreased to around 50% of normal at the time of diagnosis and decreases thereafter [3]. However, lineage tracing of beta-cells in mice during ageing and multiple pregnancies has shown that apparent loss of beta-cell mass in this animal model of diabetes is due to dedifferentiation rather than cell death [35]. Loss of insulin staining as a consequence of loss of the specialised function to produce insulin was demonstrated in parallel with induction of genes not expressed in normal adult beta-cells. This has now been confirmed in human islets prepared but not used for islet transplantation [36, 37]. Hence, the impression that type 2 diabetes is characterised by irreversible beta cell death has been shown to be flawed in that the beta cells are not 50% dead, but simply not identifiable by insulin immunostaining. Up to a certain point, de-differentiation is reversible allowing full recovery of the



specialised function to produce insulin. The corollary is that the beta cell defect can reasonably be expected to be reversible in vivo at least during the early years of type 2 diabetes.

### **3. Insight into the potential reversibility of type 2 diabetes**

The first hint that type 2 diabetes was a fully reversible syndrome came from bariatric surgery over half a century ago, and drawn to widespread attention by Walter Pories [38]. A randomised prospective study demonstrated the superiority over intensive medical therapy for type 2 diabetes [39]. Remission of diabetes was shown to be related to the degree of weight loss rather than allocation to either group. This was achieved in 73% of the surgical group and 13% of the medical group as surgery was simply more effective in achieving and maintaining weight loss [40]. Type 2 diabetes can be reversed by applying a surgical procedure which diminishes total body fat load.

The normalisation of glucose in type 2 diabetes within days after bariatric surgery, before substantial weight loss [15], led to the widespread belief that surgery itself brought about specific changes mediated via incretin hormone secretion [41, 42]. This reasoning overlooked the major change which follows bariatric surgery: an acute, profound decrease in calorie intake. Typically those undergoing bariatric surgery have a mean body weight around 150 kg [15] and would therefore require daily intake of at least 13.4 MJ/day (3,200 kcal/day) for weight maintenance [43]. It is known that hypocaloric dieting will mobilise fat first from the liver and other ectopic sites and only then from visceral and subcutaneous fat [44]. This process has been studied in detail during more moderate calorie restriction in type 2 diabetes over approximately 8 weeks [8]. Fasting plasma glucose was shown to be improved because of an 81% decrease in liver fat content and normalisation of hepatic insulin sensitivity with no change in the insulin resistance of muscle. The Counterpoint and Counterbalance studies have both shown that more vigorous calorie restriction brings about more rapid fall in liver fat and in fasting plasma glucose [13, 19]. The temporal association is highly suggestive of a causal relationship. If this is maintained then intra-pancreatic fat levels fall and in step with this, insulin secretion rates return to normal. It can now be said to be established that in the first 10 years after diagnosis, type 2 diabetes can be reversed

to non-diabetic glucose control in the majority of people who achieve 15% or more decrease in body weight by any means.

#### **4. Synthesizing recent information**

In 2008, information from study of: liver fat [7, 9]; moderate calorie restriction in type 2 diabetes [8]; and the immediate effect of bariatric surgery [15] led to the Twin Cycle Hypothesis (Figure 1)[6]. This postulated that the accumulation of fat in liver and secondarily in the pancreas would initiate self-reinforcing cycles which would interact to bring about eventual failure of beta cell function and onset of type 2 diabetes. The purpose of formulating the hypothesis was to permit rigorous testing of its predictions.

If the rapid changes in metabolism following bariatric surgery [45] were indeed a consequence of sudden change in calorie balance, then both defects in insulin secretion and hepatic insulin sensitivity should disappear by inducing sudden negative calorie balance. To test the hypothesis a group with type 2 diabetes within 4 years of diagnosis were studied before and during a 600 kcal/day diet [13]. Within 7 days, liver fat decreased by 30% and became similar to that of the matched control group, and hepatic insulin sensitivity normalised. The close association between liver fat content and hepatic glucose production had previously been established [8, 9, 46]. Plasma glucose normalised by day 7 of the diet, but the changes in post-meal blood glucose control were more gradual. Pancreas fat content decreased gradually over 8 weeks to become the same as in the control group, in step with the increase in both first phase and total insulin secretion. Fat content in the islets was not directly measured although it is known that islets take up fat avidly [31] and that islet fat content closely reflects total pancreatic fat content in animal models [47]. In the Counterpoint study, beta cell function was tested by a gold standard method using a stepped glucose infusion with subsequent arginine bolus. The typical very poor glucose-induced initial rapid peak of insulin secretion (the first phase insulin response) was confirmed at baseline in the study. This first phase response increased gradually over 8 weeks of the very low calorie diet to become indistinguishable from that of age and weight matched non-diabetic control subjects. Although a cause and effect relationship between raised intra-

organ fat levels and metabolic effect has not yet been proven, the time course data following dietary intervention study are highly suggestive of a causal link [13].

Further work has defined for how long after diagnosis type 2 diabetes remains reversible [19]. In the Counterbalance study, the mean weight loss of just under 15% represented weight loss of between 8 and 22kg. Whereas 87% of the short-duration group (<4 years) achieved non-diabetic fasting plasma glucose levels immediately after acute weight loss, only 50% of the long-duration group (8-23 years) did so. In a recent audit of bariatric surgery metabolic outcomes, HbA1c of <43mmol/mol (6.1%) was achieved by 62% and 26% respectively in those with duration of diabetes <4 or >8 years respectively [48], reflecting earlier observations [49]. As the duration of diabetes increases, it appears that beta-cells pass a point of no return and become unable to return to fully differentiated endocrine lineages perhaps with apoptosis. Whether this loss of the paracrine effect of the trophic properties of insulin is directly associated with the steadily progressive loss of pancreatic volume and increasing irregularity of the organ remains to be established [33, 50].

Although the Twin Cycle Hypothesis primarily explains the aetiology and underlying pathophysiology of type 2 diabetes, it indicates that reversal of diabetes should be long term if weight remains steady. Previously it has been shown that diabetic individuals post-bariatric surgery remain non-diabetic for up to 10 years unless substantial weight gain occurs [51]. Following dietary weight loss, 6 months of follow up with no weight regain showed that blood glucose control remained normal. Case reports and case series confirm the longer term normal metabolism following dietary weight loss provided weight remains steady [52, 53].

## **5. Understanding the apparent heterogeneity of type 2 diabetes**

It is frequently stated that non-obese people with T2DM have less insulin resistance but a greater beta cell defect than those who are overweight or obese [1, 54-56]. However, insulin resistance increases as a function of increasing BMI, whether or not an individual has normal blood glucose control [57]. When direct comparisons is made between people matched for BMI to determine the impact of insulin resistance on

the development of T2DM, it can be seen that people with T2DM have modestly greater insulin resistance at any level of BMI [58]. When individuals with T2DM are compared with BMI matched late-onset autoimmune diabetes the same is seen [59]. The apparent enigma of the usually higher fasting plasma insulin levels seen in obese individuals with T2DM, compared with their non-diabetic counterparts, is explained when the greater insulin resistance of obesity itself is taken into account [60, 61]. This must not be confused with the fall in fasting plasma insulin levels observed with increasing duration of type 2 diabetes, as this reflects apparently terminal failure of beta cells [19]. Finally, the post-prandial insulin response to test meals is similar in non-obese and obese people with T2DM [62]. Separately, the first phase insulin response to an intravenous glucose challenge is almost absent in T2DM, whatever the BMI, and in impaired glucose tolerance no effect of BMI has been demonstrated on either first or second phase insulin secretion [62].

The concept that non-obese people with T2DM have lesser degrees of insulin resistance and greater beta cell impairment has been extrapolated to therapeutic decisions. The ADA/EASD guidelines make the point that “Common practice has favoured metformin in heavier patients” [63], and non-obese patients are assumed to be able to respond less well to the glucose lowering effect of metformin. This assumption has been examined and disproven: the improvement in HbA<sub>1c</sub> between matched non-obese and obese groups with T2DM given metformin was almost identical [64, 65].

By assembling this information together with information on the distribution of BMI for the UKPDS, it has been possible to put forward the Personal Fat Threshold concept [66]. This allows understanding that type 2 diabetes only develops if an individual accumulates more fat than can be stored in the metabolically inert subcutaneous depots. The capacity of this depot varies considerably and is probably genetically determined, perhaps modified by early life environment. If however this very individual capacity is exceeded then the fat will accumulate in liver, then pancreas. Whereas 72% of people with BMI over 40kg/m<sup>2</sup> have no diabetes [5], a substantial proportion of people presenting with type 2 diabetes may have a normal BMI (36% in the UKPDS cohort [67]). Dramatically, the Nurses’ Health Study shows that developing a BMI between 23-25 is associated with a 4-fold increased risk of type 2 diabetes compared with having a BMI of less than 22kg/m<sup>2</sup> [68]. In contrast to the population

risk of developing type 2 diabetes, BMI is not necessarily a good guide to management of an individual patient.

The Personal Fat Threshold concept is also reflected in the recent data from the Counterbalance study. It may be thought that fat removed from liver and pancreas by short term hypocaloric dieting might gradually be replaced by the remaining excess from subcutaneous and visceral depots during subsequent isocaloric food intake. If so, follow up would be expected to reveal re-accumulation of intra-pancreatic fat with or without decline in beta-cell function. This has been examined during six months of weight stability after acute weight loss in people with type 2 diabetes [19]. There was no accumulation of fat in either pancreas or liver and first phase insulin response remained normal even though the weight loss of just under 15 kg left half of the group in the obese category, with the rest overweight [19].

## **6. Understanding the time course of type 2 diabetes**

The earliest risk factor for development of type 2 diabetes is believed to be low insulin sensitivity in skeletal muscle. There is no fixed cut off, but those people who will develop diabetes have insulin sensitivity mainly in the lowest population quartile [18]. In pre-diabetic individuals, raised plasma insulin levels compensate and allow normal plasma glucose control. However, as the process of *de novo* lipogenesis is stimulated by higher insulin levels [69] this sets the scene for hepatic fat accumulation as predicted by the Twin Cycle Hypothesis. Excess fat deposition in the liver is present before the onset of classical type 2 diabetes [12, 70-72]. In established type 2 diabetes, liver fat is supra-normal [71]. When ultrasound is used rather than magnetic resonance methods, only more severe degrees of steatosis are detected, and the prevalence of fatty liver is likely to be underestimated. One ultrasound study reported 70% of people with type 2 diabetes as having fatty liver [72]. Nonetheless, even using ultrasound, the prognostic power to predict onset of type 2 diabetes just by presence of hepatic steatosis is impressive. A large study of individuals with normal glucose tolerance at baseline showed a very low 8 year incidence of type 2 diabetes if fatty liver had been excluded at baseline whereas if present the hazard ratio for diabetes was 5.5 (range 3.6-8.5) [70]. The temporal progression from weight gain to raised liver enzymes, onwards to

hypertriglyceridemia and then glucose intolerance is described in the legend to Figure 1 [73]. This has been discussed in a recent review of the aetiology of type 2 diabetes [74].

In obese young people, decreased beta cell function predicts deterioration of glucose tolerance [75, 76]. Also, the rate of decline in glucose tolerance in first-degree relatives of type 2 diabetic individuals is a function of loss of beta-cell competence, whilst insulin sensitivity changes little [77]. In populations with high incidence of type 2 diabetes transition from hyperinsulinaemic normal glucose tolerance to overt diabetes involves a large, rapid rise in glucose levels with a relatively small further loss of acute  $\beta$ -cell competence [78]. Recently, the Whitehall II study has shown in a large population followed prospectively that those people developing diabetes exhibit a sudden rise in fasting glucose as beta cell function deteriorates [79]. This shows the pancreas to respond to an increasing plasma glucose with a normal, brisk insulin response largely in the two years before the detection of diabetes, even though fasting plasma glucose may have been at the upper border of normal for several years. It had previously been assumed that a linear rise in fasting plasma glucose and gradual beta cell decompensation would occur [80].

The Counterbalance study provides some of the clearest information to date on the change to the later, less reversible phase of type 2 diabetes [19]. At baseline, those individuals who could not achieve long term fasting plasma glucose levels of under 7mmol/l by substantial weight loss were characterised not only by low fasting plasma insulin levels but also by strikingly poor first phase insulin responses. Confirming that severe beta cell failure had occurred, the degree of hepatic steatosis had decreased. Taken together, these observations suggest that eventually, and most commonly after 10 years of type 2 diabetes, the beta cells pass beyond a state of reversible de-differentiation to a relatively irreversible loss of ability to make insulin.

It has been accepted for some time that the beta cell dysfunction of established diabetes progresses in an inevitable manner from the time of diagnosis [3, 77, 81] even though insulin resistance can be modified at least to some extent. However, it is now clear that the beta cell defect, and not solely hepatic insulin resistance, may be entirely reversed by weight loss at least early in the course of type 2 diabetes [13, 82]. The low insulin sensitivity of muscle tissue does not change materially either during the onset of

diabetes or during any subsequent reversal. Overall, the recent information on the inhibitory effects of excess fat on beta cell function and gene expression permits a new understanding of the time course of type 2 diabetes, both during the pre-diabetic phase and in the years after diagnosis.

## **7. Expert Commentary and Five Year View**

Once the methodology to achieve precise quantitation of intra-pancreatic fat is more widely applied, studies in different ethnic groups can be launched. This will allow an approach to the genetic determinants of intra-pancreatic fat accumulation. The process of supplying fatty acids to the pancreas will require detailed study. Dietary calorie restraint allows correction of all the metabolic features of type 2 diabetes, and it can be considered that pharmacological means could mimic some aspects of this metabolic normalisation.

## **8. Key Issues**

- The predictions of the Twin Cycle Hypothesis have been borne out by recent in vivo studies on reversing type 2 diabetes
- In the first 10 years of type 2 diabetes for most people: Negative calorie balance rapidly normalises liver fat content, hepatic glucose production and fasting plasma glucose; and if the negative calorie balance is sustained, intra-pancreatic fat content and insulin secretion also normalize
- After around 10 years of type 2 diabetes, beta cells enter a phase of failure which is not reversible by decrease in fat supply
- For any one individual, type 2 diabetes develops some years after a Personal fat Threshold has been exceeded. If the total fat load of the body is decreased below this constitutionally determined threshold then the condition may be entirely reversible

## **Acknowledgements**

The author is grateful to the noble Newcastle volunteers, who over several decades of research into type 2 diabetes have given time, effort and blood in the name of research. The work would not have been possible without the extraordinary hard work and skill of many research fellows and especially Peter Carey, Balasubramanian Ravikumar, Ee Lin Lim, Sarah Steven, Mavin Macauley, Carl Peters and Ahmad Al-Mrabeh.



## Funding

No funding to declare.

## Declaration of interest

*The authors have no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.*

## References

Papers of special note have been highlighted as:

\* of interest

\*\* of considerable interest

- [1] Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC (2003) Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. *Diabetes* 52: 102-110
- [2] Lillioja S, Mott DM, Howard BV, et al. (1988) Impaired glucose tolerance as a disorder of insulin action: Longitudinal and cross sectional studies in Pima Indians. *New England Journal of Medicine* 318: 1217-1225
- [3] Rudenski AS, Hadden DR, Atkinson AB, et al. (1988) Natural history of pancreatic islet B-cell function in type 2 diabetes mellitus studied over six years by homeostasis model assessment. *Diabetic Medicine* 5: 36-41
- [4] Franco M, Bilal U, Ordunez P, et al. (2013) Population-wide weight loss and regain in relation to diabetes burden and cardiovascular mortality in Cuba 1980-2010: repeated cross sectional surveys and ecological comparison of secular trends. *Brit Med J* 346: f1515
- [5] Gregg EW, Cheng YJ, Narayan KM, Thompson TJ, Williamson DF (2007) The relative contributions of different levels of overweight and obesity to the increased prevalence of diabetes in the United States: 1976-2004. *Prev Med* 45: 348-352
- [6] Taylor R (2008) Pathogenesis of Type 2 diabetes: Tracing the reverse route from cure to cause. *Diabetologia* 51: 1781-1789
- [7] Seppala-Lindroos A, Vehkavaara S, Hakkinen AM, et al. (2002) Fat accumulation in the liver is associated with defects in insulin suppression of glucose production and serum free fatty acids independent of obesity in normal men. *The Journal of Clinical Endocrinology and Metabolism* 87: 3023-3028
- [8] Petersen KF, Dufour S, Befroy D, Lehrke M, Hendler RE, Shulman GI (2005) Reversal of nonalcoholic hepatic steatosis, hepatic insulin resistance, and hyperglycemia by moderate weight reduction in patients with type 2 diabetes. *Diabetes* 54: 603-608
- [9] Ravikumar B, Gerrard J, Dalla Man C, et al. (2008) Pioglitazone decreases fasting and postprandial endogenous glucose production in proportion to decrease in hepatic triglyceride content. *Diabetes* 57: 2288-2295
- [10] Petersen KF, Dufour S, Savage DB, et al. (2007) The role of skeletal muscle insulin resistance in the pathogenesis of the metabolic syndrome. *Proc Natl Acad Sci U S A* 104: 12587-12594
- [11] Sevastianova K, Santos A, Kotronen A, et al. (2012) Effect of short-term carbohydrate overfeeding and long-term weight loss on liver fat in overweight humans. *American Journal of Clinical Nutrition* 96: 427-434



- [12] Sattar N, McConnachie A, Ford I, et al. (2007) Serial metabolic measurements and conversion to type 2 diabetes in the west of Scotland coronary prevention study: specific elevations in alanine aminotransferase and triglycerides suggest hepatic fat accumulation as a potential contributing factor. *Diabetes* 56: 984-991
- [13] Lim EL, Hollingsworth KG, Aribisala BS, Chen MJ, Mathers JC, Taylor R (2011) Reversal of type 2 diabetes: normalisation of beta cell function in association with decreased pancreas and liver triacylglycerol. *Diabetologia* 54: 2506-2514
- [14] Malandrucco I, Pasqualetti P, Giordani I, et al. (2012) Very-low-calorie diet: a quick therapeutic tool to improve beta cell function in morbidly obese patients with type 2 diabetes. *Am J Clin Nutr* 95: 609-613
- [15] Guidone C, Manco M, Valera-Mora E, et al. (2006) Mechanisms of recovery from type 2 diabetes after malabsorptive bariatric surgery. *Diabetes* 55: 2025-2031
- [16] Carey PE, Halliday J, Snaar JEM, Morris PG, Taylor R (2003) Direct assessment of muscle glycogen storage after mixed meals in normal and type 2 diabetic subjects. *American Journal of Physiology* 284: E286-294
- [17] Macauley M, Smith FE, Thelwall PE, Hollingsworth KG, Taylor R (2015) Diurnal variation in skeletal muscle and liver glycogen in humans with normal health and Type 2 diabetes. *Clin Sci (Lond)* 128: 707-713
- [18] Taylor R (2012) Insulin resistance and type 2 diabetes. *Diabetes* 61: 778-779
- [19] Steven S, Hollingsworth KG, Al-Mrabeh A, et al. (2016) Very Low Calorie Diet and 6 Months of Weight Stability in Type 2 Diabetes: Pathophysiological Changes in Responders and Nonresponders. *Diabetes Care* 39: 158-165
- [20] Bruning JC, Michael MD, Winnay JN, et al. (1998) A muscle-specific insulin receptor knockout exhibits features of the metabolic syndrome of NIDDM without altering glucose tolerance. *Mol Cell* 2: 559-569
- [21] Savage DB, Zhai L, Ravikumar B, et al. (2008) A prevalent variant in PPP1R3A impairs glycogen synthesis and reduces muscle glycogen content in humans and mice. *PLoS Med* 5: e27
- [22] Petersen KF, Dufour S, Befroy D, Garcia R, Shulman GI (2004) Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *The New England journal of medicine* 350: 664-671
- [23] Schrauwen-Hinderling VB, Kooi ME, Hesselink MK, et al. (2007) Impaired in vivo mitochondrial function but similar intramyocellular lipid content in patients with type 2 diabetes mellitus and BMI-matched control subjects. *Diabetologia* 50: 113-120
- [24] Brehm A, Krssak M, Schmid AI, Nowotny P, Waldhausl W, Roden M (2006) Increased lipid availability impairs insulin-stimulated ATP synthesis in human skeletal muscle. *Diabetes* 55: 136-140
- [25] Lim EL, Hollingsworth KG, Smith FE, Thelwall PE, Taylor R (2011) Inhibition of lipolysis in Type 2 diabetes normalizes glucose disposal without change in muscle glycogen synthesis rates. *Clin Sci (Lond)* 121: 169-177
- [26] Lim EL, Hollingsworth KG, Thelwall PE, Taylor R (2010) Measuring the acute effect of insulin infusion on ATP turnover rate in human skeletal muscle using phosphorus-31 magnetic resonance saturation transfer spectroscopy. *NMR Biomed* 23: 952-957
- [27] Lee Y, Hirose H, Ohneda M, Johnson JH, McGarry JD, Unger RH (1994) B-Cell lipotoxicity in the pathogenesis of non-insulin-dependent diabetes mellitus of obese rats: impairment in adipocyte-B-Cell relationships. *Proceedings of the National Academy of Science of USA* 91: 10878-10882
- [28] Moffitt JH, Fielding BA, Evershed R, Berstan R, Currie JM, Clark A (2005) Adverse physicochemical properties of tripalmitin in beta cells lead to morphological changes and lipotoxicity in vitro. *Diabetologia* 48: 1819-1829
- [29] Pinnick K, Neville M, Clark A, Fielding B (2010) Reversibility of metabolic and morphological changes associated with chronic exposure of pancreatic islet beta-cells to fatty acids. *Journal of cellular biochemistry* 109: 683-692

- [30] Pinnick KE, Collins SC, Londos C, Gauguier D, Clark A, Fielding BA (2008) Pancreatic ectopic fat is characterized by adipocyte infiltration and altered lipid composition. *Obesity* 16: 522-530
- [31] Lalloyer F, Vandewalle B, Percevault F, et al. (2006) Peroxisome proliferator-activated receptor alpha improves pancreatic adaptation to insulin resistance in obese mice and reduces lipotoxicity in human islets. *Diabetes* 55: 1605-1613
- [32] Ohneda M, Inman LR, Unger RH (1995) Caloric restriction in obese pre-diabetic rats prevents beta-cell depletion, loss of beta-cell GLUT 2 and glucose incompetence. *Diabetologia* 38: 173-179
- [33] Macauley M, Percival K, Thelwall PE, Hollingsworth KG, Taylor R (2015) Altered volume, morphology and composition of the pancreas in type 2 diabetes. *PLoS One* 10: e0126825
- [34] Rahier J, Guiot Y, Goebbels RM, Sempoux C, Henquin JC (2008) Pancreatic beta-cell mass in European subjects with type 2 diabetes. *Diabetes, obesity & metabolism* 10 Suppl 4: 32-42
- [35] Talchai C, Xuan S, Lin Hua V, Sussel L, Accili D Pancreatic  $\beta$ -Cell Dedifferentiation as a Mechanism of Diabetic  $\beta$ -Cell Failure. *Cell* 150: 1223-1234
- [36] White MG, Marshall HL, Rigby R, et al. (2013) Expression of mesenchymal and alpha-cell phenotypic markers in islet beta-cells in recently diagnosed diabetes. *Diabetes Care* 36: 3818-3820
- [37] Cinti F, Bouchi R, Kim-Muller JY, et al. (2016) Evidence of beta-Cell Dedifferentiation in Human Type 2 Diabetes. *The Journal of Clinical Endocrinology and Metabolism* 101: 1044-1054
- [38] Pories WJ, MacDonald KG, Jr., Morgan EJ, et al. (1992) Surgical treatment of obesity and its effect on diabetes: 10-y follow-up. *Am J Clin Nutr* 55: 582S-585S
- [39] Dixon JB, O'Brien PE, Playfair J, et al. (2008) Adjustable gastric banding and conventional therapy for type 2 diabetes: a randomized controlled trial. *JAMA : the journal of the American Medical Association* 299: 316-323
- [40] Ferrannini E, Mingrone G (2009) Impact of different bariatric surgical procedures on insulin action and beta-cell function in type 2 diabetes. *Diabetes Care* 32: 514-520
- [41] Rubino F, Forgione A, Cummings DE, et al. (2006) The mechanism of diabetes control after gastrointestinal bypass surgery reveals a role of the proximal small intestine in the pathophysiology of type 2 diabetes. *Ann Surg* 244: 741-749
- [42] Holst JJ (2011) Postprandial insulin secretion after gastric bypass surgery: the role of glucagon-like peptide 1. *Diabetes* 60: 2203-2205
- [43] Gibney ER, Murgatroyd P, Wright A, Jebb S, Elia M (2003) Measurement of total energy expenditure in grossly obese women: comparison of the bicarbonate-urea method with whole-body calorimetry and free-living doubly labelled water. *Int J Obes Relat Metab Disord* 27: 641-647
- [44] Colles SL, Dixon JB, Marks P, Strauss BJ, O'Brien PE (2006) Preoperative weight loss with a very-low-energy diet: quantitation of changes in liver and abdominal fat by serial imaging. *Am J Clin Nutr* 84: 304-311
- [45] Hadden DR (2001) Obstetric and diabetic care for pregnancy in diabetic women: 10 years outcome analysis, 1985-1995. *Diabetic Medicine* 18: 546-553
- [46] Gastaldelli A, Cusi K, Pettiti M, et al. (2007) Relationship between hepatic/visceral fat and hepatic insulin resistance in nondiabetic and type 2 diabetic subjects. *Gastroenterology* 133: 496-506
- [47] Lee Y, Lingvay I, Szczepaniak LS, Ravazzola M, Orci L, Unger RH (2010) Pancreatic steatosis: harbinger of type 2 diabetes in obese rodents. *Int J Obes (Lond)* 34: 396-400
- [48] Steven S, Carey PE, Small PK, Taylor R (2015) Reversal of Type 2 diabetes after bariatric surgery is determined by the degree of achieved weight loss in both short- and long-duration diabetes. *Diabetic medicine : a journal of the British Diabetic Association* 32: 47-53
- [49] Hall TC, Pellen MG, Sedman PC, Jain PK (2010) Preoperative factors predicting remission of type 2 diabetes mellitus after Roux-en-Y gastric bypass surgery for obesity. *Obes Surg* 20: 1245-1250
- [50] Al-Mrabeh A, Hollingsworth KG, Steven S, Taylor R (2016) Morphology of the pancreas in type 2 diabetes: Effect of weight loss with or without normalisation of insulin secretory capacity *Diabetologia* 59: 1753-1759
- [51] Sjostrom L, Lindroos AK, Peltonen M, et al. (2004) Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery. *The New England journal of medicine* 351: 2683-2693

- [52] Peters C, Steven S, Taylor R (2015) Reversal of type 2 diabetes by weight loss despite presence of macro-and micro-vascular complications. In: Draznin B (ed) Diabetes Case Studies: Real Problems, Practical Solutions. American Diabetes Association, Alexandria, VA
- [53] Steven S, Lim E, Taylor R (2013) Population response to information on reversibility of type 2 diabetes. *Diabetic Medicine* 30: e135-138
- [54] Vaag A, Lund SS (2007) Non-obese patients with type 2 diabetes and prediabetic subjects: distinct phenotypes requiring special diabetes treatment and (or) prevention? *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme* 32: 912-920
- [55] Ostenson CG, Efendic S (2007) Islet gene expression and function in type 2 diabetes; studies in the Goto-Kakizaki rat and humans. *Diabetes, obesity & metabolism* 9 Suppl 2: 180-186
- [56] Turner RC, Holman RR, Matthews D, Hockaday TD, Peto J (1979) Insulin deficiency and insulin resistance interaction in diabetes: estimation of their relative contribution by feedback analysis from basal plasma insulin and glucose concentrations. *Metabolism: Clinical and Experimental* 28: 1086-1096
- [57] Hinnouho GM, Czernichow S, Dugravot A, Batty GD, Kivimaki M, Singh-Manoux A (2013) Metabolically healthy obesity and risk of mortality: does the definition of metabolic health matter? *Diabetes Care* 36: 2294-2300
- [58] Hollenbeck CB, Chen Y-DI, Reaven GM (1984) A comparison of the relative effects of obesity and non-insulin dependent diabetes mellitus on in vivo insulin stimulated glucose utilization. *Diabetes* 33: 622-626
- [59] Juhl CB, Bradley U, Holst JJ, et al. (2014) Similar weight-adjusted insulin secretion and insulin sensitivity in short-duration late autoimmune diabetes of adulthood (LADA) and Type 2 diabetes: Action LADA 8. *Diabetic medicine : a journal of the British Diabetic Association* 31: 941-945
- [60] McCarthy ST, Harris E, Turner RC (1977) Glucose control of basal insulin secretion in diabetes. *Diabetologia* 13: 93-97
- [61] Turner RC, McCarthy ST, Holman RR, Harris E (1976) Beta-cell function improved by supplementing basal insulin secretion in mild diabetes. *British Medical Journal* 1: 1252-1254
- [62] Reaven GM, Chen YD, Hollenbeck CB, Sheu WH, Ostrega D, Polonsky KS (1993) Plasma insulin, C-peptide, and proinsulin concentrations in obese and nonobese individuals with varying degrees of glucose tolerance. *The Journal of clinical endocrinology and metabolism* 76: 44-48
- [63] Inzucchi SE, Bergenstal RM, Buse JB, et al. (2012) Management of hyperglycaemia in type 2 diabetes: a patient-centered approach. Position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* 55: 1577-1596
- [64] Bi Y, Tong GY, Yang HJ, et al. (2013) The beneficial effect of metformin on beta-cell function in non-obese Chinese subjects with newly diagnosed type 2 diabetes. *Diabetes/Metabolism Research and Reviews* 29: 664-672
- [65] Ito H, Ishida H, Takeuchi Y, et al. (2010) Long-term effect of metformin on blood glucose control in non-obese patients with type 2 diabetes mellitus. *Nutrition & Metabolism* 7: 83
- [66] Taylor R, Holman R (2015) Normal weight individuals who develop Type 2 diabetes: the personal fat threshold. *Clinical Science* 128: 405-410
- [67] UKPDS (1991) UK Prospective Diabetes Study (UKPDS). VIII. Study design, progress and performance. *Diabetologia* 34: 877-890
- [68] Hu FB, Manson JE, Stampfer MJ, et al. (2001) Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *The New England journal of Medicine* 345: 790-797
- [69] Schwarz JM, Linfoot P, Dare D, Aghajanian K (2003) Hepatic de novo lipogenesis in normoinsulinemic and hyperinsulinemic subjects consuming high-fat, low-carbohydrate and low-fat, high-carbohydrate isoenergetic diets. *Am J Clin Nutr* 77: 43-50
- [70] Shibata M, Kihara Y, Taguchi M, Tashiro M, Otsuki M (2007) Nonalcoholic fatty liver disease is a risk factor for type 2 diabetes in middle-aged Japanese men. *Diabetes Care* 30: 2940-2944
- [71] Perseghin G, Lattuada G, De Cobelli F, et al. (2007) Habitual physical activity is associated with intrahepatic fat content in humans. *Diabetes Care* 30: 683-688

- [72] Targher G, Bertolini L, Padovani R, et al. (2007) Prevalence of nonalcoholic fatty liver disease and its association with cardiovascular disease among type 2 diabetic patients. *Diabetes Care* 30: 1212-1218
- [73] Suzuki A, Angulo P, Lymp J, et al. (2005) Chronological development of elevated aminotransferases in a nonalcoholic population. *Hepatology* (Baltimore, Md 41: 64-71
- [74] Taylor R (2013) Type 2 diabetes: etiology and reversibility. *Diabetes Care* 36: 1047-1055
- [75] Giannini C, Weiss R, Cali A, et al. (2012) Evidence for early defects in insulin sensitivity and secretion before the onset of glucose dysregulation in obese youths: a longitudinal study. *Diabetes* 61: 606-614
- [76] Cali AM, Man CD, Cobelli C, et al. (2009) Primary defects in beta-cell function further exacerbated by worsening of insulin resistance mark the development of impaired glucose tolerance in obese adolescents. *Diabetes Care* 32: 456-461
- [77] Festa A, Williams K, D'Agostino R, Jr., Wagenknecht LE, Haffner SM (2006) The natural course of beta-cell function in nondiabetic and diabetic individuals: the Insulin Resistance Atherosclerosis Study. *Diabetes* 55: 1114-1120
- [78] Ferrannini E, Nannipieri M, Williams K, Gonzales C, Haffner SM, Stern MP (2004) Mode of onset of type 2 diabetes from normal or impaired glucose tolerance. *Diabetes* 53: 160-165
- [79] Tabak AG, Jokela M, Akbaraly TN, Brunner EJ, Kivimaki M, Witte DR (2009) Trajectories of glycaemia, insulin sensitivity, and insulin secretion before diagnosis of type 2 diabetes: an analysis from the Whitehall II study. *Lancet* 373: 2215-2221
- [80] Hanley AJ, Williams K, Festa A, et al. (2004) Elevations in markers of liver injury and risk of type 2 diabetes: the insulin resistance atherosclerosis study. *Diabetes* 53: 2623-2632
- [81] Matthews DR, Cull CA, Stratton IM, Holman RR, Turner RC (1998) UKPDS 26: Sulphonylurea failure in non-insulin-dependent diabetic patients over six years. UK Prospective Diabetes Study (UKPDS) Group. *Diabetic medicine : a journal of the British Diabetic Association* 15: 297-303
- [82] Camastra S, Manco M, Mari A, et al. (2007) Beta-cell function in severely obese type 2 diabetic patients: long-term effects of bariatric surgery. *Diabetes Care* 30: 1002-1004
- [83] Steven S, Hollingsworth KG, Small P, et al. (2016) Weight loss decreases excess pancreatic triacylglycerol specifically in type 2 diabetes. *Diabetes Care* 39: 158-165

## Reference Annotations

- [7] This study demonstrated clearly the link between liver fat content and insulin sensitivity of the liver.
- [8] This work demonstrates that the fatty liver of type 2 diabetes would respond to moderate calorie restriction with improvement in fasting plasma glucose.
- [11] Overfeeding carbohydrate produces a rapid increase in liver fat content.
- [6] This summarized the information which allowed postulation of the Twin Cycle Hypothesis
- [13] The Counterpoint study tested the basic predictions of the Twin Cycle Hypothesis and confirmed the underlying postulates.
- [33] This paper reports the abnormal gross morphology of the pancreas in type 2 diabetes and shows why measurement of pancreatic fat content is so technically demanding.

[19] The Counterbalance study showed that type 2 diabetes becomes less likely to be reversed by substantial weight loss after around 10 years after diagnosis, and also that the return to non-diabetic glucose control is durable on an isocaloric diet.

[83] This study shows that equivalent weight loss in people with normal glucose tolerance or with type 2 diabetes produces decrease in pancreatic fat content only in type 2 diabetes.

### **Financial Disclosures and Acknowledgements**

This work received no external financial support. RT is grateful to the noble Newcastle volunteers, who over several decades of research into type 2 diabetes have given time, effort and blood in the name of research. The work would not have been possible without the extraordinary hard work and skill of many research fellows and especially Peter Carey, Balasubramanian Ravikumar, Ee Lin Lim, Sarah Steven, Mavin Macauley, Carl Peters and Ahmad Al-Mrabeh.

## Legend to Figure

The Twin Cycle Hypothesis. During chronic, excess calorie intake, particularly in the presence of muscle insulin resistance, the raised plasma insulin levels will expedite chronic excess calorie storage from carbohydrate via *de novo* lipogenesis. This tends to promote storage of fat in the liver very gradually over years. This will cause the liver to become increasingly resistant to insulin and a small increase in plasma glucose will tend to occur. In turn insulin secretion will increase to control plasma glucose down. The further increased insulin levels will bring about a self-reinforcing vicious cycle. Excess liver fat will inevitably lead to an increased rate of export of VLDL triglyceride from the liver. Amongst other tissue, islets will be exposed to higher rates of fatty acid supply and pancreas fat levels as a whole will increase. The increased exposure to fatty acid metabolites will bring about endoreticulum stress, and eventually beta cell de-differentiation, with relative inhibition of meal insulin secretion. The vicious cycles will interact over at least 10 years. At a personal threshold, the pancreas fat becomes too great a load, and plasma glucose levels will then rise relatively rapidly. The Figure is redrawn from reference [6] and previously published in this form [74].

