

Updates to DABAM: an open-source DataBase of Metrology for x-ray optical simulations.

R. Celestre^a, M. Sanchez del Rio^b and L. Rebuffi^c

^aSynchrotron SOLEIL, L'Orme des Merisiers, Départementale 128, 91190 Saint-Aubin, France.

^bESRF - the European Synchrotron, 71 Avenue des Martyrs, 38000 Grenoble, France.

^cArgonne National Laboratory, 9700 S. Cass Avenue, Lemont, 60439, IL, USA.

celestre@synchrotron-soleil.fr



Since its release in 2016, the database for metrology **DABAM** has been an **important tool** for accurate **beamline design** and tolerancing providing **metrology profiles** of mirrors for **optical simulations** [1]. Its **success** is due to the **collaborative efforts** of various authors and laboratories using an **open-source platform**. We propose to create the **DABAM2D** database by **extending DABAM** to include **2D maps** of **mirrors**, **refractive lenses** or any **free-form optics**. We expect to populate the database with profiles coming from both *ex-situ* and *at-wavelength* techniques. We invite x-ray optics **community** to **contribute with profiles** to this database.

1. In line with modern **open-source practices**, the **new database** is **not tied** to a specific **web repository**, allowing for the use of multiple **local** or **private repositories**. The **height profiles** are stored in a **HDF5 file** and a supplementary metadata **json file**:

```
<url-prefix>/data/dabam2d-XXXX.h5  
<url-prefix>/data/dabam2d-XXXX.txt
```

The data is in SI units. Each surface and its corresponding info file are sequentially numbered:

- 0xxx – single x-ray lenses;
- 1xxx – stacked lenses (CRL);
- 2xxx – mirror profiles;

The structure of the h5 file as well as the contents of the metadata file are described in detail in the GitHub repository (see QR code).




Fig. 1 DABAM2D maps can be visualised using the PaNOSC online H5Wasm viewer in a conventional web browser. Top row: examples of **single lens metrology** (profiles 0001, 0073 and 0084). Bottom row: examples of **CRL metrology** (profiles 1001, 1011 and 1010).

2. DABAM2D is fully **integrated** in the **OASYS** ecosystem [2], which interfaces ray-tracing codes **SHADOW 3** and **4** and wave-propagation packages like **SRW** and **WOFRY**.




Fig. 2 DABAM 2D mirror example. The top right window shows the OASYS preprocessor widget Surface/FE reader importing directly a map of a 2D mirror. The OASYS workspace show how a 2D profile file can be created, imported and used together with SHADOW 3, 4 and SRW workflows. Bottom left: simulation result using SHADOW 4 for an ideal beamline and using DABAM2D errors. This example is available in the GitHub repository (QR code).

3. Simulations using real **measured error profiles** are able to **reproduce** with high fidelity **what is observed** at the **beamlines**:




Fig. 4 (left) simulations of an ideal x-ray lens stack focusing at 8keV and (right) aberrated lens stack focusing using the metrology profiles 0001-0010. Reproduced from Celestre et al, *J. Synch. Rad.* 27(2), 629-643 (2020).




Fig. 5 Example of how the **simulations** are able to **capture the features** of an **aberrated** focused beam from an **x-ray lens stack**. Experimental data was collected at ID06/ESRF and the partially-coherent simulations were done using the CRL metrology profiles 1001, 1011 and 1010 in SRW. Reproduced from Celestre et al, *J. Synch. Rad.* 29(2), 629-643 (2022).




Fig. 6 Example of how the **simulations** are able to **capture the features** of an **aberrated** focused beam from an **x-ray lens stack**. Ptychographic reconstruction based on experimental data collected at ID01/ESRF (courtesy of E. Bellec and S. Leak). The coherent simulations were done using the lens metrology profiles 0022-0025 in SRW.

4. **Contributing to DABAM2D** is simple: prospective collaborators are invited to contact us and **submit** several profiles of **single lens profiles**, **lens stacks**, or **2D mirror maps**, ideally in the DABAM2D data structure, though this is not mandatory. **Contributors** will participate and **co-author** an **upcoming DABAM2D publication**.

5. We aim to collect **profiles** until late **October 2024** and submit a **manuscript** for **publication** with an extended author list before the end of **November 2024**.

6. The DABAM2D database is currently hosted at:

[GitHub.com/oasys-esrf-kit/dabam2d](https://github.com/oasys-esrf-kit/dabam2d)

Acknowledgemet

We acknowledge the ESRF for the beamtimes provided. T. Roth and R. Barrett helping and allowing the data collection and curation. RC recieved funding from the European Union's Horizon 2020 research and innovation programme (grant agreement No. 101007417) within the framework of the NFFA-Europe Pilot Joint Activities.

References

- [1] Sanchez del Rio et al, *J. Synchrotron Radiat.* 23 665–678 (2016).
- [2] Rebuffi and Sanchez del Rio, *Proc SPIE* 10388. 05 (2017).

