

fuzR: an R package providing biological trait data and utilities for marine benthic ecology

Olivier Beauchard and Karline Soetaert

Abstract

fuzR is an R-package that provides a global database of biological trait data and utilities for marine benthic ecology. The database includes 41 traits documented for 1893 macrozoobenthic species, and describing fitness components and ecosystem functions. The traits are organised in a fuzzy format that requires specific R functions to be easily handled. In this vignette, we show different applications of the package, particularly beneficial to species community ecology.

S1 The fuzR package

The document provides guidelines to handle fuzzy trait data by mean of the package fuzR. Such data are organised at two levels: numerical scores are attributed to modalities, themselves nested within traits. The handling of those complex data is facilitated by specifically designed functions in fuzR. Also, the package hosts the MacroTraits database intended to marine benthic ecologists. The extraction and use of the database are exemplified through common practices in functional ecology. Besides, the package remains useful for any dataset composed of fuzzy traits.

You can install the development version of fuzR like so:

```
#remotes::install_github("obeauchard/fuzR")
library(fuzR)
```

S2 Data extraction and manipulation

Trait data are typically used in taxa x trait modalities matrices, either for specific purposes of community ecology or for other practical matters. In MacroTraits, the biological traits are organised in a long-format table that requires special attention before conversion into long format since modalities are nested within traits. Firstly, let's visualise the trait long-format data frame:

```
data(MacroTraits)
print(head(MacroTraits$traits, 15))
```

##		Species	Trait.ID	Modality.ID	Score	Reference.ID
## 1	Abarenicola	pacifica	1	1	0	3291
## 2	Abarenicola	pacifica	1	2	2	3291
## 3	Abarenicola	pacifica	1	3	0	3291
## 4	Abarenicola	pacifica	1	4	0	3291
## 5	Abarenicola	pacifica	1	5	0	3291
## 6	Abarenicola	pacifica	2	1	0	3291

## 7	Abarenicola pacifica	2	2	0	3291
## 8	Abarenicola pacifica	2	3	1	3291
## 9	Abarenicola pacifica	2	4	1	3291
## 10	Abarenicola pacifica	2	5	0	3291
## 11	Abarenicola pacifica	3	1	0	3291
## 12	Abarenicola pacifica	3	2	0	3291
## 13	Abarenicola pacifica	3	3	1	3291
## 14	Abarenicola pacifica	3	4	0	3291
## 15	Abarenicola pacifica	3	5	0	3291

The label data frame:

```
print(head(MacroTraits$labels, 20))
```

##	Trait.ID	Modality.ID	Trait	Modality
## 1	1	1	Biome	Polar
## 2	1	2	Biome	Temperate
## 3	1	3	Biome	Tropical
## 4	1	4	Biome	Deep
## 5	1	5	Biome	HV-CS
## 6	2	1	Salinity	Oligohaline
## 7	2	2	Salinity	Mesohaline
## 8	2	3	Salinity	Polyhaline
## 9	2	4	Salinity	Euhaline
## 10	2	5	Salinity	Hyperhaline
## 11	3	1	Water depth	Extralittoral
## 12	3	2	Water depth	Supralittoral
## 13	3	3	Water depth	Mediolittoral
## 14	3	4	Water depth	Shore
## 15	3	5	Water depth	Shelf
## 16	3	6	Water depth	Slope
## 17	3	7	Water depth	Abyss
## 18	4	1	Sea floor affinity	Mainly hyperbenthic
## 19	4	2	Sea floor affinity	Occasionally hyperbenthic
## 20	4	3	Sea floor affinity	Epibenthic

S2.1 Endo-substratum engineering

As a first example of data manipulation and ordination, we choose four traits related to endo-substratum engineering: endo-bioconstruction type, endo-bioconstruction depth, ventilation/pumping and sediment biomixing type, indexed 33, 34, 36 and 37, respectively. The command `fuz.tab` takes the data frames `MacroTraits$traits` and `MacroTraits$labels` and returns a list of two objects: the species x trait modalities data frame (`$tab`) and the label data frame (`$lab`) with rows of the corresponding to columns of `tab`.

```
print(unique(MacroTraits$labels[c(1, 3)]))
```

##	Trait.ID	Trait
## 1	1	Biome
## 6	2	Salinity
## 11	3	Water depth

```

## 18      4      Sea floor affinity
## 23      5      Substratum
## 30      6      Substratum depth occupancy
## 36      7      Epi-bioconstruction occupancy
## 46      8      Life span
## 54      9      Age at maturity
## 60     10     Voltinism
## 64     11     Sexuality
## 68     12     Reproductive frequency
## 72     13     Fertilisation
## 75     14     Annual fecundity
## 81     15     Offspring type
## 84     16     Offspring size
## 89     17     Offspring protection
## 94     18     Offspring development
## 99     19     Offspring benthic stage duration
## 105    20     Offspring pelagic stage duration
## 111    21     Offspring settlement size
## 117    22     Body resistance
## 122    23     Body mass
## 128    24     Body length
## 134    25     Motility
## 139    26     Mobility
## 144    27     Rafting
## 147    28     Degree of attachment
## 150    29     Sociality
## 153    30     Epi-bioconstruction type
## 161    31     Epi-bioconstruction extension
## 167    32     Epi-bioconstruction size
## 174    33     Endo-bioconstruction type
## 182    34     Endo-bioconstruction depth
## 187    35     Endo-bioconstruction width
## 191    36     Ventilation/Pumping
## 195    37     Sediment biomixing type
## 200    38     Bioerosion
## 204    39     Biostabilisation
## 207    40     Feeding type
## 213    41     Foraging depth

```

```

w <- fuz.tab(data = MacroTraits$traits, lab = MacroTraits$labels,
  which.traits = c(37, 33, 34, 36))
indic <- w$lab$Trait.ID

```

The species x trait modalities data frame (w\$tab) is linked to the label data frame by its column names: “T” for “trait” and “M” for “modality”. The vector indic enables an easier selection of traits as blocks of modality columns afterward. Below, the two first traits.

```

print(head(w$tab)[indic %in% 1:2])

```

##	T1.M1	T1.M2	T1.M3	T1.M4	T1.M5	T2.M1	T2.M2	T2.M3	T2.M4
## Abarenicola pacifica	0	0	1	0	0	0	0	0	0
## Abatus cordatus	0	1	0	0	0	1	0	0	0
## Abdopus aculeatus	1	0	0	0	0	1	0	0	0

```
## Abludomelita obtusata      0      1      0      0      0      1      0      0      0
## Abra alba                  0      1      0      0      0      1      0      0      0
## Abra nitida                0      1      0      0      0      1      0      0      0
##                               T2.M5 T2.M6 T2.M7 T2.M8
## Abarenicola pacifica      0      1      0      0
## Abatus cordatus           0      0      0      0
## Abdopus aculeatus         0      0      0      0
## Abludomelita obtusata     0      0      0      0
## Abra alba                  0      0      0      0
## Abra nitida                0      0      0      0
```

```
print(w$lab[indic %in% 1:2,])
```

```
##      Column.name Trait.ID Modality.ID      Trait
## 1      T1.M1          1          1 Sediment biomixing type
## 2      T1.M2          1          2 Sediment biomixing type
## 3      T1.M3          1          3 Sediment biomixing type
## 4      T1.M4          1          4 Sediment biomixing type
## 5      T1.M5          1          5 Sediment biomixing type
## 6      T2.M1          2          1 Endo-bioconstruction type
## 7      T2.M2          2          2 Endo-bioconstruction type
## 8      T2.M3          2          3 Endo-bioconstruction type
## 9      T2.M4          2          4 Endo-bioconstruction type
## 10     T2.M5          2          5 Endo-bioconstruction type
## 11     T2.M6          2          6 Endo-bioconstruction type
## 12     T2.M7          2          7 Endo-bioconstruction type
## 13     T2.M8          2          8 Endo-bioconstruction type
##      Modality
## 1      None
## 2      Diffusion
## 3      Upward conveying
## 4      Downward conveying
## 5      Regeneration
## 6      None/Surficial
## 7      Rugosities/Pit
## 8      Chimney/Funnel
## 9      Tube
## 10     IJ-shaped burrow
## 11     UY-shaped burrow
## 12     Branched burrow
## 13     Anastomosed burrow
```

Fuzzy coding followed by Fuzzy Correspondence Analysis (FCA) is a typical operation on such a data table (Chevenet et al. 1994). This enables the identification of trait correlations along multivariate axes as biological gradients. This can be done with the ade4 package (Chessel et al. 2004).

```
library(ade4)
fuz <- prep.fuzzy.var(df = w$tab, col.blocks = table(indic))
fca <- dudi.fca(df = fuz, scannf = FALSE)
par(mfrow = c(2, 2), mar = rep(0.1, 4))
for(i in 1:max(w$lab$Trait.ID)){
  s.label(fca$li, cpoint = 0, clab = 0, cgrid = 1)
  points(fca$li, pch = 1, col = "grey60", cex = 1.2)
```

```

sub <- paste(letters[i], " ", unique(w$lab$Trait)[i], sep = " ")
s.distrib(fca$li, fuz[indic == i], clab = 0.8, cstar = 0, cpoint = 0,
  lab = w$lab$Modality[indic == i], sub = sub,
  possub = "topleft", csub = 1.2, add.p = TRUE)
}

```




Figure S1. Endo-substratum engineering pattern. Grey dots represent the 1893 species; due to identical trait profiles, many species have the same position. From (a) to (d), each panel represents the same species pattern, but grouped per modality of the specified trait. Trait modalities are positioned at the gravity centre of their respective species. The first axis (horizontal) opposes absence of engineering ability (a, “None”, right side of the axis) to burrowing species (left). The second axis (vertical) expresses the burrowing depth, associated to non-local transfer of sediment (a).

In Figure S1, the type of bioconstruction seems slightly specific to depth (Fig. S1b) and might be useful for some specific purposes. However, regarding bioturbation, fluxes of solutes through the sediment matrix (e.g., oxygen) depend on the type of burrow opening system toward the sediment-water interface whereby two or more openings enhance fluxes with less pressure against the burrow wall unlike a single one (“IJ-shaped”) induces greater effect on the surrounding sediment (Aller 1984, Meysman et al. 2005). Hence, a simplification is required here. The command `modif.trait` modifies a trait by merging its modalities according to a vector of new categories; new modality labels can also be created. The trait endo-bioconstruction type is modified according the criterion of burrow openness, respecting the previous order of the modalities. Note that the modality “Tube” is optional and does not characterise the burrow itself; tubes can have one or two openings towards the water column, so it is considered “None” in the new categorisation.

```
print(w$lab[-1][indic == 2,])
```

```
##      Trait.ID Modality.ID      Trait      Modality
## 6          2          1 Endo-bioconstruction type  None/Surficial
## 7          2          2 Endo-bioconstruction type  Rugosities/Pit
## 8          2          3 Endo-bioconstruction type  Chimney/Funnel
## 9          2          4 Endo-bioconstruction type      Tube
## 10         2          5 Endo-bioconstruction type  IJ-shaped burrow
## 11         2          6 Endo-bioconstruction type  UY-shaped burrow
## 12         2          7 Endo-bioconstruction type  Branched burrow
## 13         2          8 Endo-bioconstruction type  Anastomosed burrow
```

```
w <- fuz.modif(w$tab, w$lab, which.trait = 2,
              merge = c(1, 1, 3, 1, 3, 2, 3, 2),
              mod = c("None", "Open-ended", "Blind-ended"))
```

As previously, `w` is a list containing similar objects `w$tab` and `w$lab`, but accordingly modified. The new table can be processed by FCA again, within the same script for the graphical representation.

```
indic <- w$lab$Trait.ID
fuz <- prep.fuzzy.var(df = w$tab, col.blocks = table(indic))
fca <- dudi.fca(fuz, scan = FALSE)
par(mfrow = c(2, 2), mar = rep(0.1, 4))
for(i in 1:max(w$lab$Trait.ID)){
  s.label(fca$li, cpoint = 0, clab = 0, cgrid = 1)
  points(fca$li, pch = 1, col = "grey60", cex = 1.2)
  sub <- paste(letters[i], ") ", unique(w$lab$Trait)[i], sep = "")
  s.distri(fca$li, fuz[indic == i], clab = 0.8, cstar = 0, cpoint = 0,
           lab = w$lab$Modality[indic == i], sub = sub,
           possub = "topleft", csub = 1.2, add.p = TRUE)
}
```




Figure S2. Endo-substratum engineering pattern after modification of the trait “Endo-bioconstruction type”. See Figure S1 for details.

S2.2 Visualisation of biological gradients

The use of multiple biological traits enables the identification of synthetic biological gradients. Sometimes, the graphical representation of a single gradient is justified either when only a single one is significant or some aspects deserve to be specifically visualised. This can be the case with trade-offs that involve co-evolved traits. Typical trade-offs are found in reproductive strategies whereby energy allocation in offspring is balanced between quantity and size. For a given trait, the function `fuz.plot` distributes the modalities according to their positions along a predefined gradient as shown below.

```
#unique(MacroTraits$labels[c("Trait.ID", "Trait")])
w <- fuz.tab(which.traits = c(10, 14:18))
tab <- w$tab
lab <- w$lab
indic <- lab$Trait.ID
fuz <- prep.fuzzy.var(df = tab, col.blocks = table(indic))
fca <- dudi.fca(fuz, scanf = FALSE)
par(mfrow = c(3, 2), mar = c(4, 8, 3, 1))
for(i in 1:max(indic)){
```

```

# Order trait modalities according to their respective first axis score
x <- tab[indic == i][order(fca$co[,1][indic == i])]
# Similarly for modality labels
mod <- lab$Modality[indic == i][order(fca$co[,1][indic == i])]
# Plotting execution
fuz.plot(x = x, y = fca$li[,1], ylab = "", mod = mod)
# Add trait label
mtext(unique(lab$Trait)[i], cex = 1, adj = 0, line = 1)
}

```




Figure S3. Reproductive and developmental trait modality distributions along the first axis of Fuzzy Correspondence Analysis.

S3 Building indices by combining several fuzzy traits

S3.1 Reproductive allocation

Energy allocation into reproduction or self-maintenance represents an important trade-off in life-history strategies (Stearns 1992). Reproductive allocation can be calculated as the ratio of reproductive output (product of fecundity times offspring size) over body mass (Clarke 1987). The function `fuz.comb` enables an

easy combination of different fuzzy traits into a same formula. The argument `list.sco` is a list containing vectors of modality performance scores. Here, fecundity and body mass follow a log10 scale, so performance scores are attributed integer values. As offspring size is coded according to more arbitrary intervals, median values are attributed. For a species profile, a single value is obtained by summing the products of modality scores times their respective performance scores. By default, traits are equally weighted by dividing performance scores by their maximum. By this way, the operation results in a weighted mean. Before being combined into the formula, each trait is rescaled between 0 and 1. By default, the formula is the cumulative product of the selected traits, but any change is possible by rewriting the formula with the desired operators between traits that are noted "x1", "x2", ..., "xn" for n traits according to their order in the data frame `tab`.

```
#print(unique(MacroTraits$labels[c("Trait.ID", "Trait")]))
w <- fuz.tab(which.traits = c(14, 16, 23))
tab <- w$tab
lab <- w$lab
taxo <- MacroTraits$taxonomy
#identical(taxo$Species, rownames(tab))
print(lab[-1])
```

##	Trait.ID	Modality.ID	Trait	Modality
## 1	1	1	Annual fecundity	<10e2
## 2	1	2	Annual fecundity	10e2-10e3
## 3	1	3	Annual fecundity	10e3-10e4
## 4	1	4	Annual fecundity	10e4-10e5
## 5	1	5	Annual fecundity	10e5-10e6
## 6	1	6	Annual fecundity	>10e6
## 7	2	1	Offspring size	<0.1
## 8	2	2	Offspring size	0.1-0.5
## 9	2	3	Offspring size	0.5-1.5
## 10	2	4	Offspring size	1.5-5.0
## 11	2	5	Offspring size	>5.0
## 12	3	1	Body mass	<0.001
## 13	3	2	Body mass	0.001-0.010
## 14	3	3	Body mass	0.010-0.100
## 15	3	4	Body mass	0.100-1.000
## 16	3	5	Body mass	1.000-10.000
## 17	3	6	Body mass	>10.000

```
sco.list <- list(1:6,
                c(0.05, 0.3, 1, 3.25, 6),
                1:6)
y <- fuz.comb(tab, lab, list.sco = sco.list,
              formula = "x1 * x2 / x3")
```

The resulting synthetic and continuous trait can be represented per clade. With the taxonomy in `MacroTraits$taxonomy`, we choose orders containing a minimum of 30 species.

```
w <- taxo["Order"]
x <- model.matrix(data = w, ~ 0 + Order)
x <- data.frame(x)
colnames(x) <- unique(sort(w[,1]))
colnames(x) <- gsub("[unassigned] ", "", colnames(x), fixed = TRUE)
rownames(x) <- taxo$Species
```

```

x <- x[apply(x, 2, sum) >= 30]
x <- x[apply(x, 1, sum) > 0,]
y <- y[names(y) %in% rownames(x)]
taxo <- taxo[taxo$Species %in% rownames(x),]
x <- x[order(tapply(y, taxo$Order, mean))]]
#identical(rownames(x), names(y))
par(mar = c(5, 8, 1, 1))
ylab <- expression(paste("Reproductive allocation (",
                          bar(x) %+-% CI, ")", sep = ""))
fuz.plot(x, y, mean, "ci", ylim = c(0, 0.5),
         pch.p = "", ylab = ylab, cex.mod = 0.8)

```




Figure S4. Reproductive allocation as the ratio of reproductive output over body mass within various orders. Each order comprises a minimum of 30 species.

In Figure S4, reproductive allocation, as a dimensionless ratio, is a coarse approximation of the reality. However, the trend is quite consistent with general considerations (Bekeuma 1989, Giangrande et al. 1994, Marshall and Keough 2007, Kooijman and Augustine 2022). The gradient opposes high fecundity in sessile egg broadcasters (bivalves) to low fecundity in highly mobile and internal or external incubators (Malacostraca and Cephalopoda). The formers are characterised by less efficient fertilisation and experience fewer reproductive successes that need to be compensated by longer reproductive life span while the latters, mostly semelparous, experience a high reproductive success immediately after maturing and that does not require multiple reproductive attempts. Interestingly, body size remains independent as underlined by the proximity between tiny crustaceans (e.g., amphipods, mysids) and comparably gigantic octopuses.

S3.2 Intrinsic rate of natural increase

Intrinsic rate of natural increase (r) is another important descriptor of adaptation to disturbance frequency (Stearns 1992). Its empirical assessment is unrealistic for many species as it requires the follow up of several cohorts, which can take several decades or centuries. Savage et al. (2004) provided evidence that inversed age at maturity (α) is proportional to r . Recently, Beauchard and Soetaert (2026), using measurement on short-lived species estimated $r = 2.26 * 1/\alpha$. Similar to reproductive output, we can use the `fuz.comb` function to calculate r .

```
#print(unique(MacroTraits$labels[c("Trait.ID", "Trait")]))
w <- fuz.tab(which.traits = 9)
tab <- w$tab
lab <- w$lab
taxo <- MacroTraits$taxonomy
#identical(taxo$Species, rownames(tab))
print(lab)
```

	Column.name	Trait.ID	Modality.ID	Trait	Modality
## 1	T1.M1	1	1	Age at maturity	<6m
## 2	T1.M2	1	2	Age at maturity	6m-1y
## 3	T1.M3	1	3	Age at maturity	1-3y
## 4	T1.M4	1	4	Age at maturity	3-5y
## 5	T1.M5	1	5	Age at maturity	5-7y
## 6	T1.M6	1	6	Age at maturity	>7y

```
y <- fuz.comb(tab, lab,
              list.sco = list(c(0.25, 0.75, 1.5, 4, 6, 8)),
              formula = "2.26 / x1", sco.scale = FALSE, trait.scale = FALSE)
```

The variable can be represented with the same taxonomic partition as previously used (Fig. S5).

```
w <- taxo["Order"]
x <- model.matrix(data = w, ~ 0 + Order)
x <- data.frame(x)
colnames(x) <- unique(sort(w[,1]))
colnames(x) <- gsub("[unassigned] ", "", colnames(x), fixed = TRUE)
rownames(x) <- taxo$Species
x <- x[apply(x, 2, sum) >= 30]
x <- x[apply(x, 1, sum) > 0,]
y <- y[names(y) %in% rownames(x)]
taxo <- taxo[taxo$Species %in% rownames(x),]
x <- x[order(tapply(y, taxo$Order, mean))]
```

```
#identical(rownames(x), names(y))
par(mar = c(5, 8, 1, 1))
ylab <- expression(paste("Intrinsic rate of natural increase (",
                          bar(x) ~ year^-1 %+-% CI, ")", sep = ""))
fuz.plot(x, y, mean, "ci", ylim = c(0, 8),
         pch.p = "", ylab = ylab, cex.mod = 0.8)
```




Figure S5. Intrinsic rate of natural increase in various orders. Each order comprises a minimum of 30 species.

S3.3 Conversion of a fuzzy trait into a continuous one

The function `fuz.comb`, in its simplest use, is the conversion of a fuzzy trait into a continuous one. Let's consider the relationship between age at maturity and life span as an example. As these traits are continuous, we fix `sco.scale = FALSE` and `trait.scale = FALSE` in order to keep the magnitude measured in years.

```
#Age at maturity
w <- fuz.tab(which.traits = 9)
tab <- w$tab
lab <- w$lab
#print(lab)
alpha <- fuz.comb(tab, lab,
                  list.sco = list(c(0.25, 0.75, 1.5, 4, 6, 8)),
                  sco.scale = FALSE, trait.scale = FALSE)
w <- fuz.tab(which.traits = 8)
tab <- w$tab
lab <- w$lab
#print(lab)
#Life span
Tmax <- fuz.comb(tab, lab,
                 list.sco = list(c(0.25, 0.75, 1.5, 6.5, 15, 35, 75, 125)),
                 sco.scale = FALSE, trait.scale = FALSE)
```

In general, these two traits are correlated (Charnov 1993); see Figure S6.

```
w <- lm(log(alpha) ~ log(Tmax) - 1)
summary(w)
```

```
##
## Call:
## lm(formula = log(alpha) ~ log(Tmax) - 1)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.09654 -0.59246 -0.30478  0.03724  1.23567
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## log(Tmax)  0.379444   0.007651   49.59  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.6139 on 1892 degrees of freedom
## Multiple R-squared:  0.5652, Adjusted R-squared:  0.565
## F-statistic: 2460 on 1 and 1892 DF,  p-value: < 2.2e-16
```

```
w <- as.data.frame(table(Tmax, alpha))
w <- w[w$Freq > 0,]
x <- as.numeric(as.character(w$Tmax))
y <- as.numeric(as.character(w$alpha))
par(mar = c(4, 5, 1, 1), cex.axis = 0.8)
plot(log(x), log(y), bty = "n", pch = 20, cex = log(w$Freq),
      xlim = log(c(0, 200)+1), ylim = log(c(0, 12)+1),
      xaxt = "n", yaxt = "n", xlab = "", ylab = "")
clip(0, 5, 0, 2.5)
abline(lm(log(alpha) ~ log(Tmax) - 1))
axis(side = 1, at = log(c(1, 3, 10, 40, 150)), lab = c(0, 3, 10, 40, 150))
axis(side = 2, at = log(c(1, 2, 4, 7, 10)), lab = c(0, 2, 4, 7, 10))
mtext(side = 1, "Life span (years)", line = 2.3, cex = 1)
mtext(side = 2, "Age at maturity (years)", line = 2.2, cex = 1)
```




Figure S6. Relationship between life span and age at maturity after conversion from fuzzy to continuous format. Dot size proportional to number of species.

The reverse operation, the conversion of a continuous trait into a fuzzy one is also possible in `fuzR`. The function `fuz.conv` generates a series of binary variables from a continuous trait. The user is free to define interval boundaries. It is useful for technical purposes such as homogenising a data set, coping with outliers or circumventing problems of multivariate non-linearity. See `?fuz.conv`.

S3.3 Indices of ecosystem function

In the marine benthos, ecosystem functions represent also an important domain of trait index development. The mechanisms through which benthic species engineer the sea floor can be quantified by combining appropriate biological traits. Among engineering functions, bioturbation, the effects of animals that directly or indirectly impact biogeochemical transports through the sediment matrix, includes biomixing as solid particle displacement or reworking, and bioirrigation as water transport (Kristensen et al. 2012). Biodeposition is the transfert of materials from the water column to the sea floor (pelagic-benthic coupling) (Gili and Coma 1998).

S3.3.1 Biomixing

Sediment biomixing has been subject to quantification with particular interest (Swift 1993, Solan et al. 2004, Queirós et al. 2013). The trait “Biomixing type” includes the four types of sediment particule displacements

induced by the fauna (François et al. 1997): biodiffusion as random sediment particle displacements; upward and downward conveying as non-local and vertical transfers of particles (advection); regeneration as sudden upward displacement and later followed by sudden downward burrow collapse.

Solan et al. (2004) proposed an index called “bioturbation potential” (BP) that multiplies faunal abundance, mobility and a score of sediment reworking type (R) that combines biomixing type, motility and burying depth: 1 = epibenthos; 2 = surficial diffusor (1-2 cm); 3 = advector; 4 = deep diffusor; 5 = regenerator. R is considered to express the magnitude of sediment biomixing (also called “reworking”). Then, BP equals to the multiplication of observed faunal biomass, mobility and R.

Although this framework represents important advances and focus in current research, it exhibits some limitations. (1) While R is purely categorical, an organism can differently rework the sediment; for instance, a large crab can induce substantial bioffusion when walking on the sediment and regeneration when burying; callianassid shrimps, considered as specifically biodiffusors (Queirós et al. 2013), are also and particularly upward conveyors as shown by important “volcano-shaped” mounds around the burrow opening (Rowden et al. 1998). (2) So far, the quantification of R has been limited to European waters by expert judgement (Queirós et al. 2013), without possible generalisation in faunas from other parts of the world through a universal and easily applicable formula. (3) Some inconsistencies in category attribution may occur among the taxa currently documented.

The MacroTraits database provides an up-to-date fuzzy version of traits required to an equivalent of R, with detailed sediment depth distribution and mobility. In absence of field biomass, adult individual body mass at the species level is available; note that its log10 scale expressed through `sco.scale = 1:6` results in a similar effect than the square root of total community biomass in the formula from Solan et al. (2004). Here, we illustrate index computation for taxa typical of soft sediments (417 species).

```
#print(unique(MacroTraits$labels[c("Trait.ID", "Trait")]))
w <- fuz.tab(which.traits = c(5:6, 23, 26, 33, 36:37))
tab <- w$tab
lab <- w$lab
#Select soft-sediment species
print(lab[-1])
```

##	Trait.ID	Modality.ID	Trait	Modality
## 1	1	1	Substratum	Mud-Sandy mud
## 2	1	2	Substratum	Muddy sand
## 3	1	3	Substratum	Sand
## 4	1	4	Substratum	Mixed gravel
## 5	1	5	Substratum	Gravel
## 6	1	6	Substratum	Mixed soft-hard
## 7	1	7	Substratum	Hard
## 8	2	1	Substratum depth occupancy	0
## 9	2	2	Substratum depth occupancy	0-5
## 10	2	3	Substratum depth occupancy	5-15
## 11	2	4	Substratum depth occupancy	15-30
## 12	2	5	Substratum depth occupancy	>30
## 13	2	6	Substratum depth occupancy	Crevice/Den
## 14	3	1	Body mass	<0.001
## 15	3	2	Body mass	0.001-0.010
## 16	3	3	Body mass	0.010-0.100
## 17	3	4	Body mass	0.100-1.000
## 18	3	5	Body mass	1.000-10.000
## 19	3	6	Body mass	>10.000
## 20	4	1	Mobility	Immobile
## 21	4	2	Mobility	Limited

## 22	4	3	Mobility	Slow
## 23	4	4	Mobility	Fast
## 24	4	5	Mobility	Very fast
## 25	5	1	Endo-bioconstruction type	None/Surficial
## 26	5	2	Endo-bioconstruction type	Rugosities/Pit
## 27	5	3	Endo-bioconstruction type	Chimney/Funnel
## 28	5	4	Endo-bioconstruction type	Tube
## 29	5	5	Endo-bioconstruction type	IJ-shaped burrow
## 30	5	6	Endo-bioconstruction type	UY-shaped burrow
## 31	5	7	Endo-bioconstruction type	Branched burrow
## 32	5	8	Endo-bioconstruction type	Anastomosed burrow
## 33	6	1	Ventilation/Pumping	Null
## 34	6	2	Ventilation/Pumping	Potential/Low
## 35	6	3	Ventilation/Pumping	High
## 36	6	4	Ventilation/Pumping	Very high
## 37	7	1	Sediment biomixing type	None
## 38	7	2	Sediment biomixing type	Diffusion
## 39	7	3	Sediment biomixing type	Upward conveying
## 40	7	4	Sediment biomixing type	Downward conveying
## 41	7	5	Sediment biomixing type	Regeneration

```

hard <- tab[lab$Modality %in% c("Mixed soft-hard", "Hard")]
hard <- apply(hard, 1, sum)
tab <- tab[hard == 0,]
taxo <- MacroTraits$taxonomy
taxo <- taxo[taxo$Species %in% rownames(tab),]
taxo <- unique(taxo[c("Phylum", "Family", "Species")])
#print(unique(taxo[1:2][order(taxo[,1], taxo[,2]),]))
taxo <- taxo[taxo$Family %in% c(#Annelids
                              "Arenicolidae", "Capitellidae", "Cirratulidae",
                              "Glyceridae", "Magelonidae", "Nephtyidae",
                              "Onuphidae", "Opheliidae", "Oweniidae",
                              "Spionidae", "Terebellidae",
                              #Arthropods
                              "Ampeliscidae", "Axiidae", "Bathyporeiidae",
                              "Bodotriidae", "Callianassidae", "Callichiridae",
                              "Cirolanidae", "Corophiidae", "Crangonidae",
                              "Diastylidae", "Diogenidae", "Gammaridae",
                              "Geryonidae", "Haustoriidae", "Hippidae",
                              "Idoteidae", "Lithodidae", "Mysidae",
                              "Nephropidae", "Ocyropodidae", "Oregoniidae",
                              "Polybiidae", "Portunidae", "Sesarmidae",
                              "Upogebiidae", "Urothoidae", "Varunidae",
                              #Echinoderms
                              "Amphiuridae", "Asteroiidae", "Brissidae",
                              "Holothuriidae", "Loveniidae", "Ophiuridae",
                              #Bivalves
                              "Cardiidae", "Donacidae", "Macridae", "Nuculidae",
                              "Mesodesmatidae", "Pectinidae", "Pharidae",
                              "Pinnidae", "Semelidae", "Solenidae",
                              "Tellinidae", "Veneridae", "Yoldiidae"),]
#Excluding, Paphies ventricosa, bivalve with disproportionate biomass
tab <- tab[rownames(tab) != "Paphies ventricosa",]
taxo <- taxo[taxo$Species != "Paphies ventricosa",]

```



```
taxo$tax <- paste(taxo$Phylum, taxo$Family, sep = " - ")
tab <- tab[rownames(tab) %in% taxo$Species,]
```

Let's calculate bioturbation potential for each reworking mode by multiplying body mass, substratum depth occupancy and mobility. As the four reworking modes are calculated and scaled in a same way, their ranges can be compared (Figure S7).

```
#Data frame to store resulting indices
res <- data.frame(matrix(0, nrow(tab), 4))
colnames(res) <- lab$Modality[lab$Trait == "Sediment bioturbation type"][-1]
rownames(res) <- rownames(tab)
for(i in 1:4){
  #Consider successively each bioturbation type
  sco.bioturb <- matrix(c(0, 1, 0, 0, 0,
                        0, 0, 1, 0, 0,
                        0, 0, 0, 1, 0,
                        0, 0, 0, 0, 1),
                      nr = 4, byrow = TRUE)
  #Performance scores of each trait
  list.sco <- list(c(1, 2.5, 10, 22.5, 30, 0),
                  c(1:6),
                  c(0:4),
                  sco.bioturb[i,])
  #Combine the selected traits
  res[,i] <- fuz.comb(tab, lab, which.traits = c(2:4, 7),
                     list.sco = list.sco)
}
#Represent the non-null values of each index
par(mfrow = c(4, 1), mar = c(3, 3, 2, 1))
for(i in 1:4){
  w <- res[,i][res[,i] > 0]
  r <- range(pretty(c(min(res[res>0]), max(res))))
  hist(w, xlim = r, main = colnames(res)[i], breaks = seq(r[1], r[2], len = 20),
       freq = FALSE, col = 1, border = "white", ylim = c(0, 30))
}
```




Figure S7. Distributions of the four sediment biomixing modes. The same 417 species are considered in each panel.

If the reworking magnitude of the burrowing fauna must be considered based on those three traits, then biomixing type is not discriminant of the magnitude, except, to a lesser extent, regeneration (Fig. S7). Hence, in case of biogeochemical or species community compositional justification, each biomixing type can be separately investigated. Otherwise, a single biomixing index can result from the product of the four scores.

```
biomix <- fuz.comb(tab, lab, which.traits = c(2:4, 7),
  list.sco = list(c(1, 2.5, 10, 22.5, 30, 0),
    c(1:6),
    c(0:4),
    c(0, 1, 1, 1, 1)))

x <- taxo["Family"]
x <- model.matrix(data = x, ~ 0 + Family)
x <- data.frame(x)
```

```
w <- taxo[order(taxo$Family),]  
colnames(x) <- unique(w$tax)  
w <- unique(w[c("Phylum", "Family")])  
x <- x[order(w[,1], tapply(biomix, taxo$Family, mean))]  
par(mar = c(4, 9, 1, 1), cex.axis = 0.8)  
fuz.plot(x, biomix, cex.mod = 0.7, pos.mod = 0.8, cex.mct = 1.5, ylab = "Biomixing score")
```




Figure S8. Total biomixing score per family as the product of the four individual biomixing types (biodiffusion, upward conveying, downward conveying and regeneration). Grey dots, species; black dot, mean; bars extend from 25th to 75th percentile.

In Figure S8, consistent variations from immobile or small taxa such as large pen shells (Pinnidae) or tiny

crustacean (Idoteidae) to highly mobile and deep burrowing ones such as gerionid, fidler and magrove crabs (Ocypodidae, Varunidae), and mud shrimps (Callianassidae, Callichiridae, Upogebiidae). In between lies various combinations of trait performances. Rag worms (Nereidae), lug worms (Arenicolidae), brittle stars (Amphiuridae), urchins (Brissidae and Loveniidae) and razor clams (Pharidae, Solenidae) are slowly mobile, but deep and sometimes massive burrowers. Small amphipods (Ampeliscidae) are very active and some of them deep burrowers (Corophiidae, Haustoridae, Urotheidae).

S3.3.2 Bioirrigation

Bioirrigation is a function of ventilation or pumping ability in buried organisms, burrow type and water injection depth (Kristensen et al. 2012). Here, the distinction between “burying” and “burrowing” is crucial since the former state does not imply an endo-bioconstruction such as a burrow that can more easily transport water through the sediment matrix. Then, the type of burrow is assumed to be determinant depending on its connectedness with the sediment surface: water should be more flushed through a system that starts and ends at the sediment-water interface (e.g., “U-shaped”) than a system that ends within the sediment matrix (single or multiple ends). However, the absence of permanent or semi-permanent burrow does not exclude significant sediment irrigation through local pore water transport induced by body movements (Aller 1982, Kristensen 1988, Berg et al. 2001, Volkenborn et al. 2012, Volkenborn et al. 2016). Such an ability is expressed by the modality “Potential/Low” from the trait “Ventilation/Pumping”. The modality “None” is especially attributed to strictly epibenthic species, and “High” and “Very high” when documented or assumed, in general in deep gallery builders (Lindroth 1941, Kristensen 1988). “Very high” is specifically attributed to buried suspension-feeders that likely ventilate more due to feeding requirements as evidenced in nereid worms and mud shrimps (Kristensen 1981, Christensen et al. 2000, Kristensen and Kostka 2005, Kristensen et al. 2012).

Wrede et al. (2018) provided a first empirical quantification and assessment of a bioirrigation index. Faunal biomass is multiplied by injection depth and feeding type whereby suspension feeding, carnivory, deposit feeding and sub-surface suspension feeding are attributed the scores 1, 2, 3 and 4, respectively. Macro-Traits data enable the calculation of this index, with possible improvements. (1) As for biomixing, the fuzzy version of the data can provide more precise estimates. (2) The use of scored feeding types by Wrede et al. (2018) might be too arbitrary. Consistently, buried suspension-feeders are attributed the highest scores. However, there is no justification to attribute different scores between depositivore, suspensivore and carnivore. Therefore, we propose a simpler index that includes burrow type, injection depth and ventilation ability, the most complex part of the mechanism being expressed by the trait “Ventilation/Pumping”.

Firstly, the trait “Endo-bioconstruction type”, that includes burrow type, is modified in order to only express the three modalities of interest. The modality “Tube” becomes “None”; by this way, a species that builds a IJ-shaped burrow with a tube is attributed a 1/0/1 profile (i.e., for None/Open-ended/Blind-ended). Then, the weighted-mean performance score will account for it as tubes reduce water exchanges between the burrow and the sediment matrix (up to 10-40 % of flux in free solution; Aller 1983).

```
#print(lab)
w <- fuz.modif(tab, lab, which.trait = 5,
               merge = c(1, 1, 3, 1, 3, 2, 3, 2),
               mod = c("None", "Open-ended", "Blind-ended"))
tab <- w$tab
lab <- w$lab
print(lab[lab$Trait.ID == 5,])
```

##	Column.name	Trait.ID	Modality.ID	Trait	Modality
## 110	T5.M1	5	1	Endo-bioconstruction type	None
## 25	T5.M2	5	2	Endo-bioconstruction type	Open-ended
## 31	T5.M3	5	3	Endo-bioconstruction type	Blind-ended

Injection depth was expressed by substratum depth occupancy. Mobile species that build a burrow may not always occur at the maximum depth of the burrow, whereas they can inject water deeper. So, we fix `fun = c("max", rep("sum", 3))` whereby, for the first trait (depth occupancy), the score for maximum depth is selected while score for the three other traits remain a weighted mean as the sum of fuzzy and performance score products. Burrow type scoring is subject to debate. While Wrede et al. (2018) attributed highest score to blind-ended burrow, Renz et al. (2018) argued that open- or blind-ended type is not determinant as other factors may create context dependency. Here, we attribute highest score to blind-ended burrow since the induced increased pressure enhances radial advective exchanges with the surrounding sediment (Aller 1984, Meysman et al. 2005), effects being exacerbated over depth (Ziebis et al. 1996, Satoh and Okabe 2013).

```
#print(lab[-1])
bioirr <- fuz.comb(tab, lab, which.traits = c(2:3, 5, 6),
                  list.sco = list(c(0, 2.5, 10, 22.5, 30, 0),
                                c(1:6),
                                c(1:3),
                                c(1:4)),
                  fun = c("max", rep("sum", 3)))
x <- taxo["Family"]
x <- model.matrix(data = x, ~ 0 + Family)
x <- data.frame(x)
w <- taxo[order(taxo$Family),]
colnames(x) <- unique(w$tax)
w <- unique(w[c("Phylum", "Family")])
x <- x[order(w[,1], tapply(bioirr, taxo$Family, mean))]
par(mar = c(4, 9, 1, 1), cex.axis = 0.8)
fuz.plot(x, bioirr, cex.mod = 0.7, pos.mod = 0.8, cex.mct = 1.5, ylab = "Bioirrigation score")
```




Figure S9. Total bioirrigation score per family. Grey dots, species; black dot, mean; bars extend from 25th to 75th percentile.

The figure returns a consistent image of current considerations, especially the fact that only a few species among many produce disproportionate effects on the sediment (Solan et al. 2004). Two species, *Chaetopterus variopedatus* and *Urechis caupo*, with disproportionate scores (> 0.3 and > 0.5 , respectively),

were discarded for a better visualisation. Further improvements could include the mobility as several arthropods do not remain sedentary in their burrow, but continuously build new burrow systems (e.g., Callianassidae) while relic burrows can still ensure a minimum of passive irrigation (Aller 1984).

S3.3.3 Further considerations on ecosystem functions

It is important to keep in mind that a synthetic index as exemplified above remains a trait as it is calculated at the species level. In related applications from the literature, values of individual taxa are usually aggregated to obtain community scores (i.e., community-weighted mean, “CWM”). However, Peres-Neto et al. (2017) highlighted critical issues with CWM when correlating CWM scores with other variables measured at the sample level. Technically, CWM ignores within-community trait variations; experimentally, attributing random trait scores can yield substantially high correlation coefficients, hence leading to excessively inflated type I error. Instead, the Fourth-corner method or its combination with RLQ ordination must be considered for sound statistical validation (Dray et al. 2014, ter Braak et al. 2017). Regarding indices of benthic ecosystem functions, any trait combination should be validated by a significant relationship to empirical measurements (e.g., field or incubated core measurements of biomixing or bioirrigation). To our knowledge, there is no such validation in existing indices of benthic ecosystem functions (Queirós et al. 2015, Wrede et al. 2018). Therefore, the development of bioturbation indices is still an open research field. Further tests should proceed through an RLQ design: empirical measurements as table R (samples x measurement variable), faunal biomass as table L (samples x taxa) and trait index as table Q (taxa x index). Note that biomass used in our examples should not be part of tested indices as it should be replaced by experimentally measured biomasses in table L.

S4 Extraction of bibliographical sources

The source documentation related to specified species and trait combinations can be obtained through the function `get.ref`. The function simply combines the selection of trait IDs and the vector of desired species after having specified the data, label and reference data frames. The result is a long-format data frame.

```
sp <- c("Corophium volutator", "Lanice conchilega", "Macoma balthica")
w <- get.ref(sp = sp, which.traits = c(8, 9, 25, 40))
kable(w, caption = "Table S1. Example of output from the function get.ref().") %>%
  kable_styling(latex_options = "hold_position", font_size = 6) %>%
  column_spec(1, width = "3cm") %>%
  column_spec(2, width = "2cm") %>%
  column_spec(3, width = "8cm")
```

Table S1. Example of output from the function `get.ref()`.

Species	Trait	Reference
Corophium volutator	Life span	Fish JD, Mills A (1979) The reproductive biology of <i>Corophium volutator</i> and <i>C. arenarium</i> (Crustacea: Amphipoda). Journal of the Marine Biological Association of the United Kingdom 59:355–368
Corophium volutator	Age at maturity	Smit MGD, Kater BJ, Jak RG, van den Heuvel-Greve MJ (2006) Translating bioassay results to field population responses using a Leslie-matrix model for the marine amphipod <i>Corophium volutator</i> . Ecological Modelling 196:515–526
Corophium volutator	Motility	Meadows PS, Reid A (1966) The behaviour of <i>Corophium volutator</i> (Crustacea: Amphipoda). Journal of Zoology 150:1966:387–399
Corophium volutator	Feeding type	Jensen KT, Kristensen LD (1990) A field experiment on competition between <i>Corophium volutator</i> (Pallas) and <i>Corophium arenarium</i> Crawford (Crustacea: Amphipoda): effects on survival, reproduction and recruitment. Journal of Experimental Marine Biology and Ecology 137:1–24
Corophium volutator	Feeding type	Meadows PS, Reid A (1966) The behaviour of <i>Corophium volutator</i> (Crustacea: Amphipoda). Journal of Zoology 150:1966:387–399

Lanice conchilega	Life span	Rabaut M (2009) Lanice conchilega, fisheries and marine conservation: towards an ecosystem approach to marine management. PhD thesis, Ghent University, Ghent, 354 p
Lanice conchilega	Age at maturity	Rabaut M (2009) Lanice conchilega, fisheries and marine conservation: towards an ecosystem approach to marine management. PhD thesis, Ghent University, Ghent, 354 p
Lanice conchilega	Motility	Rabaut M (2009) Lanice conchilega, fisheries and marine conservation: towards an ecosystem approach to marine management. PhD thesis, Ghent University, Ghent, 354 p
Lanice conchilega	Feeding type	Rabaut M (2009) Lanice conchilega, fisheries and marine conservation: towards an ecosystem approach to marine management. PhD thesis, Ghent University, Ghent, 354 p
Macoma balthica	Life span	Gilbert MA (1973) Growth rate, longevity and maximum size of Macoma balthica (L.). Biological Bulletin 145:119–126
Macoma balthica	Age at maturity	Lammens JJ (1967) Growth and reproduction in a tidal flat population of Macoma balthica (L.). Netherlands Journal of Sea Research 3:315–382
Macoma balthica	Motility	Brafield AE, Newell GE (1961) The behaviour of Macoma balthica (L.). Journal of the Marine Biological Association of the United Kingdom 41:81–87
Macoma balthica	Feeding type	Olafsson EB (1986) Density dependence in suspension-feeding and deposit-feeding populations of the bivalve Macoma balthica: a field experiment. The Journal of Animal Ecology 55:517–526

S5 Case study on life-history strategies in the North Sea

Most functions in fuzR are not specific of the MacroTraits database. As an application example, we reproduce the analyses of Beauchard et al. (2022) on benthic data from the Dutch sector of the North Sea. The area is composed of two main habitats: deep and low hydrodynamics (“LD”), and shallow and high hydrodynamics (“HD”). Given those contrasting environmental conditions, the authors investigated the specificity of response trait combinations. A particular feature of these habitats is a similar number of taxa (152 and 167 for LD and HD, respectively) and a high number in common (129) that suggests the implication of the phenomenon of “spatial contingency” (Peres-Neto et al. 2012): two similar species pools can find different solutions in habitat occupancy under variable landscape heterogeneity. Indeed, variations in environment-trait relationships over the whole area and within each habitat were found to strongly differ. The higher environmental heterogeneity within HD induced more perceptible relationships and distinctions among life-history strategies. This simultaneous detection of recognized strategies was likely possible under regular spatial wave lengths of environmental heterogeneity on species composition, while at larger scale, or adding the low dynamics area with a dissimilar spatial heterogeneity pattern, the signal was altered.

In a more analytical context, this case study illustrates the complexity of applications combining the Fourth-corner and RLQ analyses (Dray et al., 2014). In its traditional use, RLQ relates environmental variables (table R) with biological traits (table Q) through species distributions (table L). Importantly, a significant R-Q pattern implies that R variables form gradients that match species distribution gradients, themselves corresponding to trait modality distributions along matching species gradients (Dolédec et al. 1996). In this application example, not all combinations of R and Q variables may lead to a significant RLQ pattern. Therefore, the user is encouraged to run the procedure with different combinations of R and Q variables to apprehend the need for deep data exploration before considering definitive RLQ outcomes. Besides, double constrained Correspondence Analysis, more recently developed (dc-CA; ter Braak et al. 2018, ter Braak and van Rossum 2025), may offer a better technical alternative if the dataset meets certain conditions. In cases of more sampling stations than environmental variables or more species than trait modalities, the modelling nature of dc-CA enables R and Q variable selection according to standard procedures in regression analysis (ter Braak and van Rossum 2025). However, the use of fuzzy traits implies a high number of Q variables as correlated predictors of table L in dc-CA, in which case RLQ may better perform (Dray et al. 2003, ter Braak et al. 2018).

S5.1 Data preparation

We use the MacroTraits data as an update of the original trait data (purely qualitative) to demonstrate the validity of their fuzzy version. MWT2022 contains the original habitat descriptors (`env`) and faunal distributions

(tabL.ind.complete) as individual organism densities.

```
library(raster)
```

```
## Loading required package: sp
```

```
data(MWTL2022)
fau <- MWTL2022$tabL.ind.complete
env <- MWTL2022$tabR
hab <- MWTL2022$hab
geo <- MWTL2022$geo[c("lon", "lat")]
bat <- MWTL2022$bat
```

We select most of the traits considered in the original publication and add voltinism and sexuality; motility is replaced by mobility, functionally more expressive. Although many taxa are identified at the species level, several are identified only at the genus level. The function `fuz.avg` enables the averaging of trait modality scores at the desired taxonomic level based on an existing taxa x trait data frame and a corresponding taxa x taxonomic level data frame. Within the investigated taxocenosis, the genera *Gammarus* and *Glycera* are represented in MacroTraits by some European species, but also by some others that occur outside European waters. Therefore, the argument `restrict.clade` limits the averaging to specified species biogeographically relevant. From phylum to species, the argument `max.lev` is attributed a value between 1 (phylum) and 6 (species). We limit the averaging at the genus level (5).

```
txn <- MacroTraits$taxonomy[8:13]
#print(unique(MacroTraits$labels[c(1, 3)]))
w <- fuz.tab(which.traits = c(8:21, 23, 26, 6, 40))
rc <- list(c("Gammarus locusta"),
           c("Glycera alba", "Glycera tridactyla"))
w <- fuz.avg(txc = colnames(fau), tab = w$tab, lab = w$lab,
             txn = txn, max.lev = 5, restrict.clade = rc,
             round.sco = 0.3)
```

Some trait modalities are marginally represented and some of them can be discarded.

```
w$tab <- w$tab[!w$lab$Modality %in%
               c("Crevice/Den", "Asexual seasonal", "Asexual continuous", "Symbiont-associated")]
w$lab <- w$lab[!w$lab$Modality %in%
               c("Crevice/Den", "Asexual seasonal", "Asexual continuous", "Symbiont-associated"),]
nb.tax <- apply(w$tab, 2, sum)
print(cbind(w$lab[-c(1, 3)], nb.tax))
```

##	Trait.ID	Trait	Modality	nb.tax
## 1	1	Life span	<6m	34
## 2	1	Life span	6m-1y	97
## 3	1	Life span	1-3y	94
## 4	1	Life span	3-10y	74
## 5	1	Life span	10-20y	24
## 6	1	Life span	20-50y	7
## 7	1	Life span	50-100y	2
## 8	1	Life span	>100y	2
## 9	2	Age at maturity	<6m	57
## 10	2	Age at maturity	6m-1y	185

## 11	2	Age at maturity	1-3y	116
## 12	2	Age at maturity	3-5y	32
## 13	2	Age at maturity	5-7y	3
## 14	2	Age at maturity	>7y	3
## 15	3	Voltinism	Semivoltine	160
## 16	3	Voltinism	Univoltine	88
## 17	3	Voltinism	Bivoltine	30
## 18	3	Voltinism	Multivoltine	2
## 19	4	Sexuality	Gonochorism	237
## 20	4	Sexuality	Homogamy	13
## 21	4	Sexuality	Protandry	8
## 22	5	Reproductive frequency	Sexual seasonal	210
## 23	5	Reproductive frequency	Sexual continuous	72
## 26	6	Fertilisation	Broadcasting	110
## 27	6	Fertilisation	Spermcasting	40
## 28	6	Fertilisation	Pairing	108
## 29	7	Annual fecundity	<10e2	78
## 30	7	Annual fecundity	10e2-10e3	63
## 31	7	Annual fecundity	10e3-10e4	54
## 32	7	Annual fecundity	10e4-10e5	54
## 33	7	Annual fecundity	10e5-10e6	20
## 34	7	Annual fecundity	>10e6	18
## 35	8	Offspring type	Egg	136
## 36	8	Offspring type	Larva	41
## 37	8	Offspring type	Juvenile	81
## 38	9	Offspring size	<0.1	60
## 39	9	Offspring size	0.1-0.5	104
## 40	9	Offspring size	0.5-1.5	43
## 41	9	Offspring size	1.5-5.0	59
## 42	10	Offspring protection	None	106
## 43	10	Offspring protection	Gel	22
## 44	10	Offspring protection	Capsule	15
## 45	10	Offspring protection	Bearing/Brooding	124
## 46	11	Offspring development	Planktotrophic	89
## 47	11	Offspring development	Lecithotrophic	56
## 48	11	Offspring development	Mixed planktotrophic	41
## 49	11	Offspring development	Mixed lecithotrophic	10
## 50	11	Offspring development	Internal	77
## 51	12	Offspring benthic stage duration	Null	106
## 52	12	Offspring benthic stage duration	<15	49
## 53	12	Offspring benthic stage duration	15-30	57
## 54	12	Offspring benthic stage duration	30-60	35
## 55	12	Offspring benthic stage duration	60-180	29
## 56	12	Offspring benthic stage duration	>180	4
## 57	13	Offspring pelagic stage duration	Null	113
## 58	13	Offspring pelagic stage duration	<15	34
## 59	13	Offspring pelagic stage duration	15-30	68
## 60	13	Offspring pelagic stage duration	30-60	63
## 61	13	Offspring pelagic stage duration	60-180	15
## 62	13	Offspring pelagic stage duration	>180	2
## 63	14	Offspring settlement size	<0.5	88
## 64	14	Offspring settlement size	0.5-1.5	102
## 65	14	Offspring settlement size	1.5-5	81
## 66	14	Offspring settlement size	5-10	3

## 67	14	Offspring settlement size	10-20	1
## 68	15	Body mass	<0.001	113
## 69	15	Body mass	0.001-0.010	66
## 70	15	Body mass	0.010-0.100	43
## 71	15	Body mass	0.100-1.000	34
## 72	15	Body mass	1.000-10.000	25
## 73	15	Body mass	>10.000	2
## 74	16	Mobility	Immobile	14
## 75	16	Mobility	Limited	95
## 76	16	Mobility	Slow	100
## 77	16	Mobility	Fast	117
## 78	16	Mobility	Very fast	12
## 79	17	Substratum depth occupancy	0	131
## 80	17	Substratum depth occupancy	0-5	183
## 81	17	Substratum depth occupancy	5-15	83
## 82	17	Substratum depth occupancy	15-30	27
## 83	17	Substratum depth occupancy	>30	10
## 85	18	Feeding type	Deposit feeding	137
## 86	18	Feeding type	Suspension feeding	103
## 87	18	Feeding type	Herbivory/Grazing	15
## 88	18	Feeding type	Carnivory/Scavenging	82

Some traits deserve to be simplified with a lower number of modalities. The function `fuz.modif` is used to this aim.

```
w <- fuz.modif(w$tab, w$lab, which.trait = 1, merge = c(1:5, 5, 5, 5),
  mod = c(w$lab$Modality[w$lab$Trait.ID == 1][1:4], ">10y"))
w <- fuz.modif(w$tab, w$lab, which.trait = 2, merge = c(1:4, 4, 4),
  mod = c(w$lab$Modality[w$lab$Trait.ID == 2][1:3], ">3y"))
w <- fuz.modif(w$tab, w$lab, which.trait = 3, merge = c(1:3, 3),
  mod = c(w$lab$Modality[w$lab$Trait.ID == 3][1:2], "Bi/Multivoltine"))
w <- fuz.modif(w$tab, w$lab, which.trait = 12, merge = c(1:3, 3, 3, 3),
  mod = c(w$lab$Modality[w$lab$Trait.ID == 12][1:2], ">15"))
w <- fuz.modif(w$tab, w$lab, which.trait = 13, merge = c(1:3, 3, 3, 3),
  mod = c(w$lab$Modality[w$lab$Trait.ID == 13][1:2], ">15"))
w <- fuz.modif(w$tab, w$lab, which.trait = 14, merge = c(1:3, 3, 3),
  mod = c(w$lab$Modality[w$lab$Trait.ID == 14][1:2], ">1.5"))
w <- fuz.modif(w$tab, w$lab, which.trait = 15, merge = c(1:5, 5),
  mod = c(w$lab$Modality[w$lab$Trait.ID == 15][1:4], ">1.000"))
w$lab$Modality[w$lab$Trait.ID == 9 & w$lab$Modality.ID == 4] <- ">1.5"
tabQ <- w$tab
labQ <- w$lab
indic <- labQ$Trait.ID
```

Some species with a wide geographic distribution have variable scores for life span and age at maturity. Those traits must be manually adjusted to the local populations.

```
tabQ[rownames(tabQ) == "Echinocardium cordatum",][indic == 2] <- c(0, 0, 0, 1)
tabQ[rownames(tabQ) == "Macoma balthica",][indic == 1] <- c(0, 0, 1, 1, 0)
```

S5.2 Whole area

The analyses are carried out with the *ade4* package. As shown in Beauchard et al. (2022), presence-absence yields the best analytical output in this analysis. The environment-trait relationships are explored through the combination of RLQ ordination with the Fourth-corner procedure (Dray et al. 2014).

```
#Presence-absence faunal data (table L)
tabL <- data.frame(ifelse(fau == 0, 0, 1), check.names = FALSE)
tabL <- tabL[colnames(tabL) %in% rownames(tabQ)]
#identical(colnames(tabL), rownames(tabQ))
#Faunal ordination
dudiL <- dudi.coa(tabL, scan = FALSE)
#Environmental ordination (table R)
dudiR <- dudi.hillsmith(env, row.w = dudiL$lw, scan = FALSE)
#Trait ordination (table Q)
fuz <- prep.fuzzy.var(df = tabQ, col.blocks = table(indic), row.w = dudiL$cw)
dudiQ <- dudi.fca(df = fuz, scannf = FALSE)
#RLQ ordination
rlq1 <- rlq(dudiR, dudiL, dudiQ, scannf = FALSE)
#RLQ pattern significance testing
print(randtest(rlq1, 999))
```

```
## class: krandtest lightkrandtest
## Monte-Carlo tests
## Call: randtest.rlq(xtest = rlq1, nrepet = 999)
##
## Number of tests: 2
##
## Adjustment method for multiple comparisons: none
## Permutation number: 999
##      Test      Obs   Std.Obs   Alter Pvalue
## 1 Model 2 0.03805548 38.640368 greater 0.001
## 2 Model 4 0.03805548 2.742712 greater 0.021
```

The test indicates that both R-L and L-Q relationships are significant ($p < 0.05$). The eigenvalue diagram displays a prominent first axis (Fig. S10). Therefore, the RLQ pattern can be considered unidimensional and only the first axis is interpreted.

```
layout(matrix(1:3, nc = 3), width = c(2, 1.3, 2))
par(mar = c(3, 5, 1, 1), cex.lab = 2, cex.axis = 1.2)
barplot(100*rlq1$eig/sum(rlq1$eig), col = 1, ylab = "% multidimensional variance")
fuz.plot(x = data.frame(factor(hab)), y = rlq1$lR[,1], mod = c("HD", "LD"), cex.mod = 1.5,
         amount = 0.1, horiz = FALSE, ylab = "First RLQ axis",
         pch.pt = 21, col.pt = 1, bg.pt = c("white", "grey70"), cex.pt = 2.5,
         pch.mct = 21, col.mct = "white", bg.mct = 1, cex.mct = 2.5,
         bar.lwd = 2)
par(mar = c(3, 1, 1, 1))
plot(geo, type = "n", xaxt = "n", yaxt = "n", bty = "n", xlab = "", ylab = "")
contour(bat, levels = seq(0, 80, by = 10), labcex = 1, add = TRUE)
s.value(geo, rlq1$lR[,1], cleg = 0, add.p = TRUE)
```




Figure S10. RLQ analysis of the whole area. Left panel, eigenvalue diagram showing a prominent first axis. Middle, sampling station first axis scores showing the distinction between high and low dynamics habitats (HD and LD, respectively; black dot, median score; bars extend from 25th to 75th percentiles). Right, sampling axis scores mapped over the area; black, positive scores; white, negative scores; square size proportional to deviation from 0. Contour lines, depth in meters.

The gradient is clearly spatialised, with low RLQ axis scores for the HD habitat, and high scores for LD, consistently with the habitat typology in Beauchard et al. (2022). This suggests that the two habitats are characterised by different trait compositions.

The `ade4` function `fourthcorner2` of the `ade4` package enables significance testing of coinertia between a block of variables and an RLQ axis, either a qualitative R variable (as a block of dummy variables) or a fuzzy trait (block of modality variables). We use this function to test habitat descriptors, either factors (testing coinertia) or single variables (testing Pearson's r squared). Then, p-values are corrected for multiple testing (`method = fdr`, false discovery rate).

```
p <- as.numeric()
indicR <- c(rep(1:4, rep(5, 4)), rep(5, 4), 6:8)
for(i in 1:ncol(env)){
  p[i] <- fourthcorner2(dudiR$tab[indicR == i], tabL,
                        rlq1$lQ[1], nrepet = 999)$trRLQ$pvalue
}
names(p) <- colnames(env)
p <- p.adjust(p, method = "fdr")
print(round(p, 3))
```

```
## depth current wave stratif sedim POM POC PP
## 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001
```

Then, we test the Pearson's r significance between each trait modality and the RLQ sampling station axis scores. This is done automatically by the function `fourthcorner.rlq` of the `ade4` package.

```
fc <- fourthcorner.rlq(rlq1, modeltype = 6, typetest = "Q.axes",
                        nrepet = 999, p.adjust.method.D = "fdr")
par(cex = 0.7)
plot(fc, stat = "D", col = c("white", "black", "dodgerblue3"))
```




Figure S11. Significance tests of correlations between trait modalities and RLQ axes (right, first axis; left, second axis). White, not significant ($p \geq 0.05$); blue, significant and positive correlation; black, significant and negative correlation.

As described in Beauchard et al. (2022), this first RLQ axis loosely opposes short-lived (Fig. S11, first blue cell) and early maturing species (second and third blue cells) to lately mature ones (first black cell, i.e., negatively correlated), suggesting also longer-lived ones, although long-lived modalities are not significant. This could simplistically be interpreted as an r-K gradient, but this analysis only reveals a fragment of a more complete biological pattern due to scale-dependent considerations as follows.

S5.3 Low dynamics

The RLQ pattern in the low dynamics habitat was not found significant (p-value around 0.3) due to the relative homogeneity of the deep and muddy area, which may not be selectively strong enough to generate variations in trait composition.

```

tabLL <- data.frame(iffelse(fau == 0, 0, 1)[hab == "LD",], check.names = FALSE)
tabLL <- tabLL[apply(tabLL, 2, sum) > 0]
tabQL <- tabQ[rownames(tabQ) %in% colnames(tabLL),]
tabLL <- tabLL[colnames(tabLL) %in% rownames(tabQL)]
#identical(colnames(tabLL), rownames(tabQL))
tabQL <- tabQL[apply(tabQ, 2, sum) > 0]
labQL <- labQ[labQ$Column.name %in% colnames(tabQL),]
indicL <- labQL$Trait.ID
tabRL <- env[hab == "LD",]
dudiLL <- dudi.coa(tabLL, scan = FALSE)
dudiRL <- dudi.hillsmith(tabRL, row.w = dudiLL$lw, scan = FALSE)
fuzL <- prep.fuzzy.var(df = tabQL, col.blocks = table(indicL), row.w = dudiLL$cw)
dudiQL <- dudi.fca(df = fuzL, scannf = FALSE)
rlqL <- rlq(dudiRL, dudiLL, dudiQL, scannf = FALSE)
print(randtest(rlqL, 999))

```

```

## class: krandtest lightkrandtest
## Monte-Carlo tests
## Call: randtest.rlq(xtest = rlqL, nrepet = 999)
##
## Number of tests: 2
##
## Adjustment method for multiple comparisons: none
## Permutation number: 999
##      Test      Obs   Std.Obs   Alter Pvalue
## 1 Model 2 0.008524049 6.4349392 greater 0.001
## 2 Model 4 0.008524049 0.5043754 greater 0.275

```

S5.4 High dynamics

Although this habitat is characterised by higher hydrodynamics and coarser sediments, its highly heterogeneous geomorphology creates alternations in abiotic conditions that induce local changes in community trait combinations. The test of the RLQ pattern is highly significant.

```

tabLH <- log(fau+1)[hab == "HD",]
tabLH <- tabLH[apply(tabLH, 2, sum) > 0]
tabQH <- tabQ[rownames(tabQ) %in% colnames(tabLH),]
tabLH <- tabLH[colnames(tabLH) %in% rownames(tabQH)]
#identical(colnames(tabLH), rownames(tabQH))
tabQH <- tabQH[apply(tabQ, 2, sum) > 0]
labQH <- labQ[labQ$Column.name %in% colnames(tabQH),]
indicH <- labQH$Trait.ID
tabRH <- env[hab == "HD",]
dudiLH <- dudi.coa(tabLH, scan = FALSE)
dudiRH <- dudi.hillsmith(tabRH, row.w = dudiLH$lw, scan = FALSE)
fuzH <- prep.fuzzy.var(df = tabQH, col.blocks = table(indicH), row.w = dudiLH$cw)
dudiQH <- dudi.fca(df = fuzH, scannf = FALSE)
rlqH <- rlq(dudiRH, dudiLH, dudiQH, scannf = FALSE)
print(randtest(rlqH, 999))

```

```

## class: krandtest lightkrandtest

```



```
## Monte-Carlo tests
## Call: randtest.rlq(xtest = rlqH, nrepet = 999)
##
## Number of tests: 2
##
## Adjustment method for multiple comparisons: none
## Permutation number: 999
##      Test      Obs   Std.Obs   Alter Pvalue
## 1 Model 2 0.03513899 16.355278 greater 0.001
## 2 Model 4 0.03513899 2.772274 greater 0.017
```

The pattern is expressed along two axes.

```
par(mfrow = c(1, 3), mar = c(1, 5, 1, 1), cex.lab = 2)
barplot(100*rlqH$eig/sum(rlqH$eig), col = 1, ylab = "% multidimensional variance")
par(mar = c(1, 1, 1, 1), cex.lab = 2)
for(i in 1:2){
  plot(geo, type = "n", xaxt = "n", yaxt = "n", bty = "n", xlab = "", ylab = "")
  contour(bat, levels = seq(0, 80, by = 10), labcex = 1, add = TRUE)
  s.value(geo[hab == "HD",], rlqH$lR[,i], cleg = 0,
    sub = paste("RLQ axis", i, sep = " "), csub = 2.5,
    possub = "bottomright", add.p = TRUE)
}
```




Figure S12. RLQ analysis of the high dynamics habitat. Left panel, eigenvalue diagram showing two axes that deserve interpretation. Middle and right, mapping of sampling station first and second axis scores, respectively. Black, positive scores; white, negative scores; square size proportional to deviation from 0. Contour lines, depth in meters.

Compared to the first analysis, the pattern reveals shorter spatial wave lengths of variations in functional composition (Fig. S11). As perfectly continuous abiotic gradients at the shelf scale are rarely encountered,

this suggests that trait pattern significance in marine benthic ecology may be spatially limited (see Fig. S11 middle and right panels; main variations of 50-100 km). Next to environmental variables (not shown), many more trait modalities are significantly correlated to both axes.

```
fc <- fourthcorner.rlq(rlqH, modeltype = 6, typetest = "Q.axes",
                       nrepet = 999, p.adjust.method.D = "fdr")
par(cex = 0.7)
plot(fc, stat = "D", col = c("white", "black", "dodgerblue3"))
```




Figure S13. Significance tests of correlations between trait modalities and RLQ axes (right, first axis; left, second axis). White, not significant ($p \geq 0.05$); blue, significant and positive correlation; black, significant and negative correlation.

The abiotic heterogeneity of the HD habitat enables more easily the detection of the three existing life-history strategies (Fig. S14). (1) A-strategists, adversity-selected species, right side of the figure. Very small, stress resistant through high mobility on mobile sand with short life and early maturity, important reproductive allocation (low fecundity, release of large juveniles as adult miniatures after internal incubation). (2) r-selected species, lower side of the pattern. Also short-lived with early maturity, but mixed development; disturbance resilient. (3) K-strategists, top-left corner; longer-lived species, some of them reaching maturity much later, infrequent reproductive successes, absence of parental cares, small and numerous offspring; many bivalves, with typical suspension feeding and planktotrophic development, and sometimes with large body mass; in general, vulnerable to disturbance.

```

par(mfrow = c(5, 4), mar = rep(0.1, 4))
for(i in 1:max(indicH)){
  s.label(rlqH$lQ, cpoint = 0, clab = 0)
  points(rlqH$lQ, pch = 20, cex = 2,
         col = adjustcolor("dodgerblue4", 0.4))
  s.distri(rlqH$lQ, fuzH[indicH == i], cstar = 0, cpoint = 0,
          lab = labQH$Modality[indicH == i], clab = 1.5,
          sub = unique(labQH$Trait)[i], possub = "topleft",
          csub = 2, add.p = TRUE)
}

```




Figure S14. RLQ analysis of the high dynamics habitat, trait representation along axis 1 (horizontal) and 2 (vertical). Each panel represents the same species pattern (blue dots) group by their respective modalities of the considered trait.

Conclusively, this illustrative case study validates the analytical relevance of updated traits in the MacroTraits database. Besides, the provided R code can be useful in other case studies from a comparative perspective, especially regarding the concept of spatial contingency that remains insufficiently investigated in the marine

benthos.

References

- Aller RC (1982) The effects of macrobenthos on chemical properties of marine sediment and overlying water. In McCall PL, Tevesz MJS (eds). *Animal–Sediment Relations*. Plenum Press, New York, pp 53–102
- Aller RC (1983) The importance of the diffusive permeability of animal burrow linings in determining marine sediment chemistry. *Journal of Marine Research* 41:299–322
- Aller RC (1984) The importance of relict burrow structures and burrow irrigation in controlling sedimentary solute distributions. *Geochimica et Cosmochimica Acta* 48:1929–1934
- Beauchard O, Mestdagh S, Koop L, Ysebaert T, Herman PMJ (2022) Benthic synecology in a soft sediment shelf: habitat contrasts and assembly rules of life strategies. *Marine Ecology Progress Series* 682:31–50
- Beauchard O, Soetaert S (2026) Bottom fishing assessment tool: an R package to model the effects of bottom trawling on the marine benthos. *Ecological Applications* <https://doi.org/10.1002/eap.70282>
- Berg P, Rysgaard S, Funch P, Sejr MK (2001) Effects of bioturbation on solutes and solids in marine sediments. *Aquatic Microbial Ecology* 26:81–94
- Beukema JJ (1989) Molluscan life spans and long-term cycles in benthic communities. *Oecologia* 80:570–570
- Charnov EL (1993) *Life history invariants. Some explorations of symmetry in evolutionary ecology*. Oxford University Press, Oxford, 167 p
- Chessel D, Dufour AB, Thioulouse J (2004) The ade4 package-I - One-table methods. *R News* 4:5–10
- Chevenet F, Dolédec S, Chessel D (1994) A fuzzy coding approach for the analysis of long-term ecological data *Freshwater Biology* 31:295–309
- Christensen B, Vedel A, Kristensen E (2000) Carbon and nitrogen fluxes in sediment inhabited by suspension-feeding (*Nereis diversicolor*) and non-suspension-feeding (*N. virens*) polychaetes. *Marine Ecology Progress Series* 192:203–217
- Clarke A (1987) Temperature, latitude and reproductive effort. *Marine Ecology Progress Series* 38:89–99
- Dray S, Chessel D, Thioulouse J (2003) Co-inertia analysis and the linking of ecological data tables. *Ecology* 84:3078–3089
- Dray S, Choler P, Dolédec S, Peres-Neto PR, Thuiller W, Pavoine S, ter Braak CJF (2014) Combining the fourth-corner and the RLQ methods for assessing trait responses to environmental variation. *Ecology* 95:14–21
- François F, Poggiale JC, Durbec JP, Stora G (1997) A new approach for the modelling of sediment reworking induced by a macrobenthic community. *Acta Biotheoretica* 45:295–319
- Giangrande A, Geraci S, Belmonte G (1994) Life-cycle and life-history diversity in marine invertebrates and the implications in community dynamics. *Oceanography and Marine Biology: An Annual Review* 32:305–333
- Gili JM, Coma R (1998) Benthic suspension feeders: their paramount role in littoral marine food webs. *Trends in Ecology and Evolution* 13:316–321
- Kooijman SALM, Augustine S (2022) The comparative energetics of the cephalopods: they neither grow nor reproduce fast. *Journal of Sea Research* 184:102205
- Kristensen E (1981) Direct measurement of ventilation and oxygen uptake in three species of tubiculous polychaetes (*Nereis* spp.). *Comparative Physiology* 145:45–50

- Kristensen E (1988) Benthic fauna and biogeochemical processes in marine sediments: microbial activities and fluxes. In Blackburn TH, Sørensen J (eds). Nitrogen cycling in coastal marine environments. John Wiley & Sons Ltd, Chichester, pp 275–299
- Kristensen E, Kostka JE (2005) Macrofaunal burrows and irrigation in marine sediment: microbiological and biogeochemical interactions. *Coastal and Estuarine Studies* 60:125–157
- Kristensen E, Penha-Lopes G, Delefosse M, Valdemarsen T, Quintana CO, Banta GT (2012) What is bioturbation? The need for a precise definition for fauna in aquatic sciences. *Marine Ecology Progress Series* 446:285–302
- Lindroth A (1941) Atmungsventilation der Polychäten. *Zeitschrift für Vergleichende Physiologie* 28:485–532
- Marshall DJ, Keough MJ (2007) The evolutionary ecology of offspring size in marine invertebrates. *Advances in Marine Biology* 53:1–60
- Meysman FJR, Galaktionov OS, Middelburg JJ (2005) Irrigation patterns in permeable sediments induced by burrow ventilation: a case study of *Arenicola marina*. *Marine Ecology Progress Series* 303:195–212
- Peres-Neto PR, Dray S, ter Braak CJF (2017) Linking trait variation to the environment: critical issues with community-weighted mean correlation resolved by the fourth-corner approach. *Ecography* 40:806–816
- Peres-Neto PR, Leibold MA, Dray S (2012) Assessing the effects of spatial contingency and environmental filtering on metacommunity phylogenetics. *Ecology* 93:14–30
- Queirós AM, Birchenough SNR, Bremner J, Godbold JA, Parker RE, Romero-Ramirez A, Reiss H, Solan M, Somerfield PJ, Van Colen C, Van Hoey G, Widdicombe S (2013) A bioturbation classification of European marine infaunal invertebrates. *Ecology and Evolution* 3:3958–3985
- Queirós AM, Stephens N, Cook R, Ravaglioli C, Nunes J, Dashfield S, Harris C, Tilstone GH, Fishwick J, Braeckman U, Somerfield PJ, Widdicombe S (2015) Can benthic community structure be used to predict the process of bioturbation in real ecosystems? *Progress in Oceanography* 137:559–569
- Renz JR, Powilleit M, Gogina M, Zettler ML, Morys C, Forster S (2018) Community bioirrigation potential (BIPc), an index to quantify the potential for solute exchange at the sediment-water interface. *Marine Environmental Research* 141:214–224
- Rowden AA, Jones MB, Morris AW (1998) The role of *Callinassa subterranea* (Montagu) (THALASSINIDEA) in sediment resuspension in the North Sea. *Continental Shelf Research* 18:1365–1380
- Satoh H, Okabe S (2013) Spatial and temporal oxygen dynamics in macrofaunal burrows in sediments: a review of analytical tools and observational evidence. *Microbes and Environments* 28:166–179
- Savage VM, Gillooly JF, Brown JH, West GB, Charnov EL (2004) Effects of body size and temperature on population growth. *The American Naturalist* 163:429–441
- Solan M, Cardinale BJ, Downing AL, Engelhardt KAM, Ruesink JL, Srivastava DS (2004) Extinction and ecosystem function in the marine benthos *Science* 306:1177–1180
- Stearns SC (1992) The evolution of life histories. Oxford University Press, Oxford, 249 p
- Swift DJ (1993) The macrobenthic infauna off Sellafield (north-eastern Irish Sea) with special reference to bioturbation. *Journal of the Marine Biological Association of the United Kingdom* 73:143–162
- ter Braak CJF, Peres-Neto P, Dray S (2017) A critical issue in model-based inference for studying trait-based community assembly and a solution. *PeerJ* 5:e2885
- ter Braak CJF, Šmilauer P, Dray S (2018) Algorithms and biplots for double constrained correspondence analysis. *Environmental and Ecological Statistics* 25:171–197
- ter Braak CJF, van Rossum BJ (2025) Linking multivariate trait variation to the environment: the advantages of double constrained correspondence analysis with the R package *douconca*. *Ecological Informatics* 88:103143

- Volkenborn N, Meile C, Polerecky L, Pilditch CA, Norkko A, Norkko J, Hewitt JE, Thrush SF, Wetthey DS, Woodin SA (2012) Intermittent bioirrigation and oxygen dynamics in permeable sediments: an experimental and modeling study of three tellinid bivalves. *Journal of Marine Research* 70:794–823
- Volkenborn N, Woodin SA, Wetthey DS, Polerecky L (2016) Bioirrigation in marine sediments. In Elias SA (ed). *Reference module in Earth systems and environmental sciences*. Elsevier, pp 1–10
- Wrede A, Beermann J, Dannheim J, Gutow L, Brey T (2018) Organism functional traits and ecosystem supporting services – A novel approach to predict bioirrigation. *Ecological Indicators* 91:737–743
- Ziebis W, Forster S, Huettel M, Jørgensen BB (1996) Complex burrows of the mud shrimp *Callinassa truncata* and their geochemical impact in the sea bed. *Nature* 382:619–622