

Cauchy functional equations

January 12, 2017



[Home Page](#)

[Title Page](#)

[Contents](#)



[Page 1 of 38](#)

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Cauchy functional equations

Ron Sahoo

Department of Mathematics

University of Louisville, Louisville, Kentucky 40292 USA

January, 2017



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 2 of 38

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Cauchy functional equations

The functional equations

$$f(x + y) = f(x) + f(y)$$

$$f(x + y) = f(x) f(y)$$

$$f(xy) = f(x) + f(y)$$

$$f(xy) = f(x) f(y)$$

for all $x, y \in \mathbb{R}$ are called Cauchy functional equations. Here

$f : \mathbb{R} \rightarrow \mathbb{R}$ is an unknown function.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 3 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

The first functional equation, namely

$$f(x + y) = f(x) + f(y) \quad (1)$$

for all $x, y \in \mathbb{R}$ is known as the additive Cauchy functional equation.

The functional equation (1) was first studied by A.M. Legendre (1791) and C.F. Gauss (1809) but A.L. Cauchy (1821) first found its general continuous solution.

[Home Page](#)[Title Page](#)[Contents](#)[Page 4 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

What are functional equations?

- Functional equations are equations in which the unknowns are functions.
- Functional equations is a field of mathematics which is over 270 years old.
- Functional equations appeared in the literature around the same time as the modern theory of functions.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 5 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- The field of functional equations includes differential equations, difference equations, integral equations, and iterations.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 6 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- To solve a functional equation means to find the unknown function (or functions).
- In order to find solutions, the unknown functions must often be restricted to a specific nature (such as **bounded, continuous, differentiable**).

[Home Page](#)[Title Page](#)[Contents](#)[Page 7 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Definition 1. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be an *additive function* if it satisfies the additive Cauchy functional equation $f(x + y) = f(x) + f(y)$ for all $x, y \in \mathbb{R}$.

Definition 2. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is called a *linear function* if and only if it is of the form $f(x) = cx$ where c is any real constant.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 8 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- The graph of a linear function $f(x) = cx$ is a non-vertical line that passes through the origin.
- The linear functions satisfy the additive Cauchy equations.
- Are there any other functions that satisfy the additive Cauchy equation?
- We begin by showing that the only continuous solutions of the additive Cauchy equation are those which are linear.

[Home Page](#)[Title Page](#)[Contents](#)[Page 9 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



This theorem was first proved by A. L. Cauchy in 1821.

Theorem 1. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying the Cauchy equation*

$$f(x + y) = f(x) + f(y), \text{ for all } x, y \in \mathbb{R}.$$

Then f is linear, that is $f(x) = mx$, where m is a real constant.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 10 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)




Proof. The Fundamental Theorem of Calculus says that if f is continuous on $[a, b]$, then the function g defined by

$$g(x) = \int_a^x f(t) dt \quad a \leq x \leq b$$

is continuous and differentiable on (a, b) , and $g'(x) = f(x)$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 11 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



$$g(x) = \int_0^x f(t) dt$$

$$g'(x) = f(x)$$

Illustration of Fundamental Theorem of Calculus



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 12 of 38

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

First, let us fix x and then integrate both side of the Cauchy equation with respect to y from 0 to 1 to get

$$\begin{aligned} f(x) &= \int_0^1 f(x) \, dy \\ &= \int_0^1 [f(x+y) - f(y)] \, dy \\ &= \int_x^{1+x} f(u) \, du - \int_0^1 f(y) \, dy \\ &= \int_d^{1+x} f(u) \, du - \int_d^x f(u) \, du - \int_0^1 f(y) \, dy. \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 13 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Hence

$$f(x) = \int_d^{1+x} f(u) du - \int_d^x f(u) du - \int_0^1 f(y) dy.$$

Since f is continuous, by the FTC, we get

$$\begin{aligned} f'(x) &= f(1+x) - f(x) \\ &= f(1) + f(x) - f(x) \\ &= f(1) \\ &= m \text{ (say).} \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 14 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Therefore solving the differential equation,

$$\frac{d}{dx}f(x) = m$$

we get

$$f(x) = m x + b,$$

where d is a constant of integration. It can be easily shown that $b = 0$ and thus $f(x) = mx$. This completes the proof of the theorem.

[Home Page](#)[Title Page](#)[Contents](#)

Page 15 of 38

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- Notice that in Theorem 1, we use the continuity of f to conclude that f is also integrable.
- The integrability of f forced the solution f of the additive Cauchy equation to be linear.
- Thus every integrable solution of the additive Cauchy equation is also linear.

[Home Page](#)[Title Page](#)[Contents](#)[Page 16 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Definition 3. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be **locally integrable** if and only if it is integrable over every finite interval.

Finite interval means interval of finite length. It is known that every locally integrable solution of the additive Cauchy equation is also linear.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 17 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



We give as a short proof of this using an argument provided by Shapiro (1973).

Assume f is a locally integrable solution of the additive Cauchy equation. Hence $f(x + y) = f(x) + f(y)$ holds for all x and y in \mathbb{R} .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 18 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Since f is additive and locally integrable, we get

$$\begin{aligned} y f(x) &= \int_0^y f(x) dz \\ &= \int_0^y [f(x+z) - f(z)] dz \\ &= \int_x^{x+y} f(u) du - \int_0^y f(z) dz \\ &= \int_0^{x+y} f(u) du - \int_0^x f(u) du - \int_0^y f(u) du. \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 19 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

The right side of the above equality is invariant under the interchange of x and y . Hence it follows that

$$y f(x) = x f(y)$$

for all $x, y \in \mathbb{R}$. Therefore, for $x \neq 0$, we obtain

$$\frac{f(x)}{x} = c,$$

where c is an arbitrary constant.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)

Page 20 of 38

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The last equality implies that $f(x) = cx$ for all $x \in \mathbb{R} \setminus \{0\}$.

Letting $x = 0$ and $y = 0$ in (1), we get $f(0) = 0$.

Together with this and the above, we conclude that f is a linear function in \mathbb{R} .

This completes the proof.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 21 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Although the proof of Theorem 1 is brief and involves only calculus, this proof is not very instructive.

We will present now a different proof which will help us to understand the behavior of the solution of the additive Cauchy equation a bit more.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 22 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



We begin with the following definition.

Definition 4. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is said to be *rationally homogeneous* if and only if

$$f(rx) = rf(x) \tag{2}$$

for all $x \in \mathbb{R}$ and all rational numbers r .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 23 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The following theorem shows that any solution of the additive Cauchy equation is rationally homogeneous.

Theorem2. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a solution of the additive Cauchy equation. Then f is rationally homogeneous. Moreover, f is linear on the set of rational numbers \mathbb{Q} .*

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 24 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Proof: Letting $x = 0 = y$ in (1) see that $f(0) = f(0) + f(0)$ and hence

$$f(0) = 0. \quad (3)$$

Substituting $y = -x$ in (1) and then using (3), we see that f is an odd function in \mathbb{R} , that is

$$f(-x) = -f(x) \quad (4)$$

for all $x \in \mathbb{R}$. Thus, so far, we have shown that a solution of the additive Cauchy equation is zero at the origin and it is an odd function.

[Home Page](#)[Title Page](#)[Contents](#)[«](#) [»](#)[◀](#) [▶](#)[Page 25 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Next, we will show that a solution of the additive Cauchy equation is rationally homogeneous. For any x ,

$$f(2x) = f(x + x) = f(x) + f(x) = 2f(x).$$

Hence

$$f(3x) = f(2x + x) = f(2x) + f(x) = 2f(x) + f(x) = 3f(x);$$

so in general (using induction)

$$f(nx) = n f(x) \tag{5}$$

for all positive integers n .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 26 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

If n is a negative integer, then $-n$ is a positive integer and by (5) and (4), we get

$$\begin{aligned} f(nx) &= f(-(-n)x) \\ &= -f(-nx) \\ &= -(-n) f(x) \\ &= n f(x). \end{aligned}$$

Thus, we have shown $f(nx) = n f(x)$ for all integers n and all $x \in \mathbb{R}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 27 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Next, let r be an arbitrary rational number. Hence, we have

$$r = \frac{k}{\ell}$$

where k is an integer and ℓ is a natural number. Further, $kx = \ell(rx)$. Using the integer homogeneity of f , we obtain

$$k f(x) = f(kx) = f(\ell(rx)) = \ell f(rx);$$

that is,

$$f(rx) = \frac{k}{\ell} f(x) = r f(x).$$

Thus, f is rationally homogeneous.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 28 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Further, letting $x = 1$ in the above equation and defining $c = f(1)$, we see that

$$f(r) = c r$$

for all rational numbers $r \in \mathbb{Q}$.

Hence, f is linear on the set of rational numbers and the proof is now complete.

[Home Page](#)[Title Page](#)[Contents](#)

Page 29 of 38

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Now we present the **second proof** of Theorem 1.

Proof: Let f be a continuous solution of the additive Cauchy equation. For any real number x there exists a sequence $\{r_n\}$ of rational numbers with $r_n \rightarrow x$.

Since f satisfies the additive Cauchy equation, by Theorem 2, f is linear on the set of rational numbers. That is,

$$f(r_n) = c r_n$$

for all n .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 30 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Now using the continuity of f , we get

$$\begin{aligned} f(x) &= f\left(\lim_{n \rightarrow \infty} r_n\right) \\ &= \lim_{n \rightarrow \infty} f(r_n) \\ &= \lim_{n \rightarrow \infty} c r_n \\ &= c x \end{aligned}$$

and the proof is now complete.

[Home Page](#)[Title Page](#)[Contents](#)

Page 31 of 38

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The following theorem is due to Darboux (1875).

Theorem 3. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a solution of the additive Cauchy equation (1). If f is continuous at a point, then it is continuous everywhere.*

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 32 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous at a point t in \mathbb{R} and let x be any arbitrary point. Hence, we have

$$\lim_{y \rightarrow t} f(y) = f(t).$$

Next, we show that f is continuous at x .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 33 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Consider

$$\begin{aligned}\lim_{y \rightarrow x} f(y) &= \lim_{y \rightarrow x} f(y - x + x - t + t) \\&= \lim_{y \rightarrow x} [f(y - x + t) + f(x - t)] \\&= \lim_{y \rightarrow x} f(y - x + t) + \lim_{y \rightarrow x} f(x - t) \\&= f(t) + f(x - t) \\&= f(t) + f(x) - f(t) \\&= f(x).\end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)

Page 34 of 38

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Thus we proved that f is continuous at the arbitrary point x and the arbitrariness of x implies that f is continuous everywhere in \mathbb{R} .

Now the proof of the theorem is complete.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 35 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The next theorem follows from Theorem 1 and Theorem 3.

Theorem 4. *Let f be a solution of the additive Cauchy functional equation (1). If f is continuous at a point, then f is linear; that is, $f(x) = cx$ for all $x \in \mathbb{R}$.*

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 36 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

References

- [1] B. Ebanks, P.K. Sahoo and W. Sander, Characterizations of Information Measures, World Scientific, Singapore, 1998.
- [2] P. K. Sahoo and Pl. Kannappan, Introduction to Functional Equations, CRC Press, Boca Raton, 2011.
- [3] P. K. Sahoo and T. Riedel, Mean Value Theorems and Functional Equations, World Scientific, Singapore, 1998.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 37 of 38](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Thank You



Home Page

Title Page

Contents



Page 38 of 38

Go Back

Full Screen

Close

Quit