

Introduction to functional equations

March 14, 2016



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 1 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Introduction to functional equations

Ron Sahoo

Department of Mathematics

University of Louisville, Louisville, Kentucky 40292 USA

March, 2016



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 2 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Introduction to functional equations

Abstract: In this introductory talk, we present an overview of the field of functional equations and then determine the solutions two simple functional equations, namely Sincov functional equation and Cauchy functional equation. This talk will be accessible to undergraduate and graduate students.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 3 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



What are functional equations?

- Functional equations are equations in which the unknowns are functions.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 4 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Some examples of functional equations are:

$$f(x + y) = f(x) + f(y)$$

$$f(x + y) = f(x)f(y)$$

$$f(xy) = f(x)f(y)$$

$$f(xy) = f(x) + f(y)$$

[Home Page](#)[Title Page](#)[Contents](#)

Page 5 of 66

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

$$f(x + y) = f(x)g(y) + g(x)f(y)$$

$$f(x + y) = f(x) + f(y) + f(x)f(y)$$

$$f(x + y) = 2f(x)f(y) - f(x - y)$$

$$f(x + y) = g(xy) + h(x - y)$$

[Home Page](#)[Title Page](#)[Contents](#)[Page 6 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



$$f(x + y) + f(x - y) = 2f(x) + 2f(y)$$

$$f(pr, qs) + f(ps, qr) = 2f(p, q) + 2f(r, s)$$

$$f(x, y) + f(x + y, z) = f(y, z) + f(x, y + z)$$

$$f(x, y) + f(y, z) = f(x, z)$$

[Home Page](#)[Title Page](#)[Contents](#)[Page 7 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- The field of functional equations includes differential equations, difference equations, integral equations, and iterations.

[Home Page](#)[Title Page](#)[Contents](#)[Page 8 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- To solve a functional equation means to find the unknown function (or functions).
- In order to find solutions, the unknown functions must often be restricted to a specific nature (such as **bounded, continuous, differentiable**).

[Home Page](#)[Title Page](#)[Contents](#)[Page 9 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- Functional equations is a field of mathematics which is over 270 years old.
- More than 15,500 papers have been published in this area (according to MathSciNet).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 10 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)


History of functional equations

- Functional equations appeared in the literature around the same time as the modern theory of functions.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 11 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- In 1747 and 1750, d'Alembert published three papers and these three papers were the first on functional equations.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 12 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Jean d'Alembert (1717-1783)



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 13 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Functional equations were studied by

- D'Alembert (1747)
- Euler (1768)
- Poisson (1804)
- Cauchy (1821)
- Abel (1823)
- Darboux (1875)

and many others.



[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 14 of 66

[Go Back](#)


[Full Screen](#)

[Close](#)


[Quit](#)



Jean d'Alembert
(1717-1783)



Leonard Euler
(1707-1783)



Simeon Poisson
(1781-1840)



[Home Page](#)

[Title Page](#)

[Contents](#)



[Page 15 of 66](#)

[Go Back](#)

[Full Screen](#)

[Close](#)


[Quit](#)



Augustin Cauchy
(1789-1857)



Niels Abel
(1802-1829)



Gaston Darboux
(1842-1917)



[Home Page](#)

[Title Page](#)

[Contents](#)



[Page 16 of 66](#)

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

- David Hilbert (1900) pointed out that while the theory of differential equations provides elegant and powerful techniques for solving functional equations, the differentiability assumptions are not inherently required.

[Home Page](#)[Title Page](#)[Contents](#)[Page 17 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



David Hilbert (1862-1943)



[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 18 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

- Motivated by Hilbert's suggestions many researchers in functional equations have treated various functional equations without any (or mild) regularity assumptions.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 19 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- This effort has given rise to the modern theory of functional equations. The theory of functional equations forms a modern mathematical discipline.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 20 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- This discipline has developed very rapidly in the last six decades largely due to the effort of Janos Aczél.



Professor Janos Aczél



[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 21 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Applications

Functional Equations have found applications in many areas of mathematics, sciences and engineering.

[Home Page](#)[Title Page](#)[Contents](#)

Page 22 of 66

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



For example, they are used in

- Algebra
- Combinatorics
- Economics
- Psychology
- Statistics
- Computer Science
- Dynamical Systems
- Dynamical Programming
- Information Theory
- Probability Theory

[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 23 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)



International Conferences

Each year, there are several conferences and seminars on functional equations. Among these the **ISFE** (started in 1962) and **ICFEI** (started in 1984) are premier conferences in this field.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 24 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



44th ISFE in Louisville, Kentucky, USA, June 14-21, 2006



[Home Page](#)

[Title Page](#)

[Contents](#)



[Page 25 of 66](#)




[Go Back](#)




[Full Screen](#)

[Close](#)

[Quit](#)

Few books on functional equations

[Home Page](#)[Title Page](#)[Contents](#)[Page 26 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[Page 27 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



[Home Page](#)

[Title Page](#)

[Contents](#)






[Page 28 of 66](#)



[Go Back](#)




[Full Screen](#)




[Close](#)

[Quit](#)



[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 29 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 30 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[Page 31 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Sincov Functional Equation

We will examine a very simple functional equation which has many applications.

This functional equation is the **Sincov functional equation**.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 32 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



- In 1903, the Russian mathematician D.M. Sincov studied the following functional equation

$$f(x, y) + f(y, z) = f(x, z) \quad (1)$$

for all $x, y, z \in \mathbb{R}$ (the set of real numbers).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 33 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Dmitrii Sintsov (1867-1946)



[Home Page](#)

[Title Page](#)

[Contents](#)



[Page 34 of 66](#)

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)




- Others like [Moritz Cantor](#) (1896) and [Gottlob Frege](#) (1874) treated this functional equation


$$f(x, y) + f(y, z) = f(x, z)$$

before Sincov.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 35 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Moritz Cantor (1829-1920)



Gottlob Frege (1848-1925)



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 36 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)



The functional equation (1), that is

$$f(x, y) + f(y, z) = f(x, z)$$

is known as the **Sincov functional equation**.

- The most general solution of the Sincov functional equation is $f(x, y) = \phi(x) - \phi(y)$, where ϕ is an **arbitrary function**.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 37 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The multiplicative version of (1) is the following:

$$f(x, y) f(y, z) = f(x, z) \quad (2)$$

for all $x, y, z \in \mathbb{R}$.

- The most general solution of (2) are $f(x, y) = 0$ and $f(x, y) = \frac{\phi(x)}{\phi(y)}$, where ϕ is a nonzero arbitrary function.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 38 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- The additive Sincov functional equation

$$f(x, y) + f(y, z) = f(x, z)$$

reduces to the well-known Cauchy functional equation

$$g(x + y) = g(x) + g(y)$$

if we take $f(x, y) = g(y - x)$.

To see this just substitute $u = y - x$ and $v = z - y$ so that

$$u + v = z - x.$$

[Home Page](#)[Title Page](#)[Contents](#)[Page 39 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Applications of Sincov Equation

- The Sincov functional equation arises in the axiomatic derivation of the **maximum entropy principle** (see Shore and Johnson (1980)).

[Home Page](#)[Title Page](#)[Contents](#)[Page 40 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- The Sincov functional equation is used in the **characterizations of information measures** on open domains (see Ebanks, Sahoo and Sander (1998)).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 41 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- It also arises in the study of **subgroup consistent poverty indices** in economics (see Foster and Shorrocks (1991)).
- Further, the Sincov functional equation has found applications in **actuarial mathematics** (see Shiu (1988)).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 42 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

In their book, Dodson and Poston (1977) define an affine space as follows:

- An *affine space* is a triple (S, V, f) where S is a set, V a vector space and $f : S^2 \rightarrow V$ such that

$$f(x, y) + f(y, z) = f(x, z)$$

for $x, y, z \in S$, and for all $x \in S$, the map $f_x(y) = f(x, y)$ is a bijection.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)

Page 43 of 66

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- In the definition of affine space, we again encounter the Sincov functional equation.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 44 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Cauchy Functional Equation

We will examine another very simple functional equation which has many applications.

This functional equation is the **Cauchy equation** and it is of the form

$$f(x + y) = f(x) + f(y), \quad \text{for all } x, y \in \mathbb{R}.$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 45 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



This theorem was first proved by A. L. Cauchy in 1821.

Theorem 1 *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying the Cauchy equation*

$$f(x + y) = f(x) + f(y), \text{ for all } x, y \in \mathbb{R}.$$

Then f is linear, that is $f(x) = mx$, where m is a real constant.

[Home Page](#)[Title Page](#)[Contents](#)

Page 46 of 66

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof. The Fundamental Theorem of Calculus says that if f is continuous on $[a, b]$, then the function g defined by


$$g(x) = \int_a^x f(t) dt \quad a \leq x \leq b$$

is continuous and differentiable on (a, b) , and $g'(x) = f(x)$.

[Home Page](#)[Title Page](#)[Contents](#)

Page 47 of 66

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



$$g(x) = \int_0^x f(t) dt$$

$$g'(x) = f(x)$$

Illustration of Fundamental Theorem of Calculus



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 48 of 66

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

First, let us fix x and then integrate both side of the Cauchy equation with respect to y from 0 to 1 to get

$$\begin{aligned} f(x) &= \int_0^1 f(x) dy \\ &= \int_0^1 [f(x+y) - f(y)] dy \\ &= \int_x^{1+x} f(u) du - \int_0^1 f(y) dy \\ &= \int_d^{1+x} f(u) du - \int_d^x f(u) du - \int_0^1 f(y) dy. \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 49 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Hence

$$f(x) = \int_d^{1+x} f(u) du - \int_d^x f(u) du - \int_0^1 f(y) dy.$$

Since f is continuous, by the FTC, we get

$$\begin{aligned} f'(x) &= f(1+x) - f(x) \\ &= f(1) + f(x) - f(x) \\ &= f(1) \\ &= m \text{ (say).} \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 50 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Therefore solving the differential equation,

$$\frac{d}{dx}f(x) = m$$

we get

$$f(x) = m x + b,$$

where d is a constant of integration. It can be easily shown that $b = 0$ and thus $f(x) = mx$.

[Home Page](#)[Title Page](#)[Contents](#)

Page 51 of 66

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

References

- [1] M. Cantor, Funktionalgleichungen mit drei von einander unabhängigen veränderlichen, *Z. Math. Phys.* 41 (1896), 161-163.
- [2] B. Ebanks, P.K. Sahoo and W. Sander, Characterizations of Information Measures, World Scientific, Singapore, 1998.
- [3] G. Frege, *Rechnungsmethoden, die sich auf eine Erweiterung des Grössebegriffes gründen*, Verlag Friedrich Frommann, Jena, 1874.
- [4] D. Gronau, A remark on Sincov's functional equation, *Notices of the South African Mathematical Society*, 31 (2000), 1-8.
- [5] P. K. Sahoo and Pl. Kannappan, Introduction to Functional Equations, CRC Press, Boca Raton, 2011.
- [6] P. K. Sahoo and T. Riedel, Mean Value Theorems and Functional Equations, World Scientific, Singapore, 1998.
- [7] D. M. Sincov, Notes sur la calcul fonctionnel (Russian), *Bull. Soc. Phys.-Math. Kazan*, 13 (1903), 48-72.
- [8] D. M. Sincov, Über eine funktionalgleichung, *Arch. Math. Phys.* 6 (1903), 216-217.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 52 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Thank You



Home Page

Title Page

Contents



Page 53 of 66

Go Back

Full Screen

Close

Quit

Appendix



Home Page

Title Page

Contents




Page 54 of 66

Go Back

Full Screen


Close

Quit

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 55 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Jean d'Alembert

[d'Alembert](#) was a French mathematician who was a pioneer in the study of differential equations and their use of in physics. He studied the equilibrium and motion of fluids.



Simeon Poisson

Poisson's most important works were a series of papers on definite integrals and his advances in Fourier series. This work was the foundation of later work in this area by Dirichlet and Riemann.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 56 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Home Page

Title Page

Contents



Page 57 of 66

Go Back

Full Screen


Close

Quit



Niels Abel


In 1824 [Abel](#) proved the impossibility of solving algebraically the general equation of the fifth degree.



Augustin Cauchy

Cauchy pioneered the study of analysis, both real and complex, and the theory of permutation groups. He also researched in convergence and divergence of infinite series, differential equations, determinants, probability and mathematical physics.


[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 58 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Leonard Euler

[Leonhard Euler](#) was a Swiss mathematician who made enormous contributions to a wide range of mathematics and physics including analytic geometry, trigonometry, geometry, calculus and number theory.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 59 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 60 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Gaston Darboux

[Darboux](#) made important contributions to differential geometry and analysis and the Darboux integral is named after him.



David Hilbert


[Hilbert](#)'s work in geometry had the greatest influence in that area after Euclid. A systematic study of the axioms of Euclidean geometry led Hilbert to propose 21 axioms and he analysed their significance. He made contributions to mathematics and physics.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 61 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 62 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Janós Aczél


[Janós Aczél](#) will be best remembered for his pioneering work on the theory of functional equations, with applications in many fields, such as information measures, index numbers, group decision making, aggregation, production functions, laws of science, theory of measurement and utility theory.



Dmitrii Sintsov


[Sintsov](#)'s main areas of interest were in the theory of conics and applications of this geometrical theory to the solution of differential equations and to the theory of nonholonomic differential geometry.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 63 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[Page 64 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Gottlob Frege

[Gottlob Frege](#) was one of the founders of modern symbolic logic putting forward the view that mathematics is reducible to logic. He is widely regarded as a logician on a par with Aristotle, Gödel, and Taraski.



Moritz Cantor

[Moritz Cantor](#) is best remembered for the four volume work *Vorlesungen ber Geschichte der Mathematik* which traces the history of mathematics up to 1799. The first volume was published in 1880 and the last volume appeared in 1908.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 65 of 66](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Thank You



Home Page

Title Page

Contents



Page 66 of 66

Go Back

Full Screen

Close

Quit