

A functional equation associated with elementary school arithmetic

February 15, 2017



[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 1 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

A functional equation associated with elementary school arithmetic

Ron Sahoo

Department of Mathematics

University of Louisville, Louisville, Kentucky 40292 USA

March, 2016



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 2 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Abstract: In this talk, I will present an interesting functional equation known as the cocycle equation. This equation has a long history. I will derive this equation using the traditional method of manual addition of two multi-digit numbers that students learn in elementary school arithmetic. The solutions of this equation will be presented on free abelian groups. This talk will assume only a basic knowledge of abstract algebra and therefore will be accessible to undergraduate and graduate students.

[Home Page](#)[Title Page](#)[Contents](#)[Page 3 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



- This talk is about a functional equation that arises from the elementary school arithmetic.
- We will see how this equation can be solved on a free abelian group.

[Home Page](#)[Title Page](#)[Contents](#)[Page 4 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Suppose we ask an elementary school student to add the number 25 to the number 18. The student will add the ones and will get 13. So he will write 3 and carry the digit 1. Then he will add the tens and the carry digit to get a 4. Therefore the sum he will have is 43.

[Home Page](#)[Title Page](#)[Contents](#)[Page 5 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

$$\begin{array}{r} 1 \\ 1 \quad 8 \\ + \quad 2 \quad 5 \\ \hline 4 \quad 3 \end{array}$$

Illustration of right to left addition of two-digit numbers



[Home Page](#)

[Title Page](#)

[Contents](#)



Page 6 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)



Let us create a formal group theoretic framework for the study of arithmetic.

Let us consider addition in the finite abelian group \mathbb{Z}_{100} .

This will serve as a model for addition of two-digit numbers.

[Home Page](#)[Title Page](#)[Contents](#)

Page 7 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

	1	
	1	8
+	2	5
<hr/>		
	4	3

Illustration of right to left addition of two-digit numbers

\mathbb{Z}_{10} is a subgroup of \mathbb{Z}_{100} consisting of the multiples of 10.

Let us call this subgroup H . It represents subgroup of “tens”.



[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#)

[▶▶](#)

[◀](#)

[▶](#)

Page 8 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

$$\begin{array}{r}
 \\
 \textcolor{red}{1} \\
 \textcolor{red}{2} \\
 \hline
 \textcolor{red}{4}
 \end{array}$$

Illustration of right to left addition of two-digit numbers

Also, \mathbb{Z}_{10} is a quotient group of \mathbb{Z}_{100} , namely, $\mathbb{Z}_{100}/\textcolor{red}{H}$. We call this group $\textcolor{blue}{G}$. This group $\textcolor{blue}{G}$ is a group of “ones”.



Home Page

Title Page

Contents

◀ ▶

◀ ▶

Page 9 of 100

Go Back

Full Screen

Close

Quit



For any $x \in \mathbb{Z}_{100}$, let us write

$$x = [a] [b],$$

where $a \in H$ and $b \in G$. This means we are thinking of a as the tens digit of x and b as the ones digit of x .

For example, we write the number 48 as $[4][8]$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 10 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



For any two elements $[a_1][b_1]$ and $[a_2][b_2]$ of \mathbb{Z}_{100} , their sum

$$[a_1][b_1] + [a_2][b_2]$$

in \mathbb{Z}_{100} can be written uniquely in the form $[a][b]$, with a and b are functions of a_1, a_2, b_1, b_2 .

[Home Page](#)[Title Page](#)[Contents](#)

Page 11 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The element b of G depends only on the elements b_1 and b_2 of G .

The ones digit of a sum depends only on the ones digits of the summands. This is the way one works from right to left when adding multi-digit numbers by hand.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 12 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The element a of H depends on the elements a_1, a_2, b_1, b_2 .

The tens digit of a sum depends on all four values a_1, a_2, b_1, b_2 because adding two multi-digit numbers is not merely adding columns.

[Home Page](#)[Title Page](#)[Contents](#)

Page 13 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The formula for a is

$$a = a_1 + a_2 + z(b_1, b_2) \quad (1)$$

where $z : G \times G \rightarrow H$ is a function expressing how the ones digits of the summands affects the tens digit of the sum.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)

Page 14 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The formula for b is

$$b = b_1 + b_2. \quad (2)$$

Here the addition symbol “+” is the group operation of G .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 15 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[Page 16 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#) $z(b_1, b_2)$ $[a_1]$ $[b_1]$ $+$ $[a_2]$ $[b_2]$

 $[a_1 + a_2 + z(b_1, b_2)] \quad [b_1 + b_2]$

Illustration of right to left addition of two-digit numbers

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 17 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

$$z(b_1 + b_2, b_3)$$

$$[a_1 + a_2 + z(b_1, b_2)]$$

$$[b_1 + b_2]$$

$$+$$

$$[a_3]$$

$$[b_3]$$

$$[a_1 + a_2 + a_3 + z(b_1, b_2) + z(b_1 + b_2, b_3)] \quad [b_1 + b_2 + b_3]$$

Illustration of right to left addition of two-digit numbers



Since the group operation of the group $(\mathbb{Z}_{100}, +)$ is associative, thus we have

$$([a_1][b_1] + [a_2][b_2]) + [a_3][b_3] = [a_1][b_1] + ([a_2][b_2] + [a_3][b_3])$$

for any $a_1, a_2, a_3 \in H$ and $b_1, b_2, b_3 \in G$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 18 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Using (1) and (2), we compute

$$\begin{aligned} & ([a_1][b_1] + [a_2][b_2]) + [a_3][b_3] \\ &= [a_1 + a_2 + a_3 + z(b_1, b_2) + z(b_1 + b_2, b_3)][b_1 + b_2 + b_3]. \end{aligned}$$

Similarly, we also have

$$\begin{aligned} & [a_1][b_1] + ([a_2][b_2] + [a_3][b_3]) \\ &= [a_1 + a_2 + a_3 + z(b_2, b_3) + z(b_1, b_2 + b_3)][b_1 + b_2 + b_3]. \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 19 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Hence the consideration of **the tens digit of the sum** yields the functional equation

$$z(b_1, b_2) + z(b_1 + b_2, b_3) = z(b_2, b_3) + z(b_1, b_2 + b_3)$$

for all $b_1, b_2, b_3 \in G$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 20 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The last functional equation can be rewritten as

$$F(a, b) + F(a + b, c) = F(b, c) + F(a, b + c) \quad (\text{CE})$$

for all $a, b, c \in G$ and $F : G \times G \rightarrow H$ is an unknown function to be determined.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 21 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

The functional equation (**CE**) is known as the **2-cocycle functional equation**. This equation has a long history, with connection to

- Factor Systems and Group Extension
- Cohomology Theory
- Information Theory
- Formal Groups
- Algebra of Polyhedra

[Home Page](#)[Title Page](#)[Contents](#)[Page 22 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- In 1956, S. Kurepa determined the differentiable solution of (CE) when $G = H = \mathbb{R}$ and proved that F is of the form

$$F(a, b) = f(a + b) - f(a) - f(b) \quad \forall a, b \in \mathbb{R} \quad (3)$$

where $f : G \rightarrow H$ is an arbitrary function.

- In 1959, J. Erdős proved that every symmetric solution $F : G^2 \rightarrow H$ of (CE) is of the form (3), where G is an abelian group and H is a divisible abelian group.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 23 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- In 1968, Jessen, Karpf and Thorup proved J. Erdős' result in a different way.
- Using some ideas from Jessen, Karpf and Thorup (1968), we will present the solution of the functional equation (CE) when G is a free abelian group.
- If H is a divisible abelian group and G is an abelian, then we also get the same solution.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 24 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Free Abelian Groups

First, we present the definition and some examples of free abelian group.

A free abelian group is an abelian group that has a basis in the sense that every element of the group can be written in one and only one way as a **finite linear combination** of elements of the basis with integer coefficients.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 25 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Not all abelian groups have basis, the ones they do are called free abelian groups.

Free abelian groups are the free groups that are abelian. In fact the only free groups that are abelian are the trivial group (free group of rank 0) and the cyclic group (free group of rank 1).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)

Page 26 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Example 1. Let $G = \mathbb{Z} \oplus \mathbb{Z} = \{ (a, b) \mid a, b \in \mathbb{Z} \}$. Then $(G, +)$ is a free abelian group with basis $\{(1, 0), (0, 1)\}$.

Example 2. $(\mathbb{Q}, +)$ is not a free abelian group.

Example 3. $(\mathbb{Z}_n, +)$ is not a free abelian group.

If G is a free abelian group with a basis X and $\text{card}(X) = n$, then G is isomorphic to $\mathbb{Z} \oplus \mathbb{Z} \oplus \cdots \oplus \mathbb{Z}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 27 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- For every nonempty set B , there exists a free abelian group with basis B .
- Subgroup of a free abelian group is a free abelian group.
- All free groups having B as basis are isomorphic.
- If G is a free abelian group with basis B , then for every arbitrary $f : B \rightarrow H$ (some abelian group) there exists unique homomorphism $F : G \rightarrow H$ which extends f .

[Home Page](#)[Title Page](#)[Contents](#)[Page 28 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Theorem 1 (J-K-T, 1968) *Let H be an abelian group and G be a free abelian group. Then every symmetric solution $F : G^2 \rightarrow H$ of*

$$F(a, b) + F(a + b, c) = F(b, c) + F(a, b + c) \quad (\text{CE})$$

for all $a, b, c \in G$ is of the form

$$F(a, b) = f(a + b) - f(a) - f(b), \quad (\text{S})$$

where $f : G \rightarrow H$ is an arbitrary function.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 29 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: It is easy to see that F given by (S) is symmetric and satisfies the 2-cocycle functional equation (CE).

Now we prove that F given by (S) is the only solution of the 2-cocycle functional equation (CE).

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 30 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Letting $b = 0$ in (CE), we get

$$F(a, 0) = F(0, c).$$

Letting $a = 0 = b$ in (CE), we get

$$F(0, c) = F(0, 0).$$

Hence

$$F(a, 0) = F(0, c) = F(0, 0). \quad (4)$$

for all $a, c \in G$.

[Home Page](#)[Title Page](#)[Contents](#)

Page 31 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Define a set W by

$$W = G \times H.$$

Let us define a binary operation “+” in W as

$$(a, x) + (b, y) = (a + b, x + y + F(a, b)) \quad (5)$$

for all $a, b \in G$ and all $x, y \in H$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 32 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



We show that $(W, +)$ is an abelian group. That is

- (1) $+$ is commutative;
- (2) W has an identity element;
- (3) each element in W has an inverse;
- (4) $+$ is associative.

[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#)

[▶▶](#)

[◀](#)

[▶](#)

Page 33 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Step 1. For each $a, b \in G$ and $x, y \in H$,

$$\begin{aligned}(a, x) + (b, y) &= (a + b, x + y + F(a, b)) \quad (\text{by definition}) \\ &= (b + a, y + x + F(a, b)) \quad (G \text{ and } H \text{ abelian}) \\ &= (b + a, y + x + F(b, a)) \quad (F \text{ symmetric}) \\ &= (b, y) + (a, x) \quad (\text{by definition})\end{aligned}$$

Hence the binary operation “+” is commutative in W .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 34 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Step 2. For each $a \in G$ and $x \in H$,

$$\begin{aligned}(a, x) + (0, -F(0, 0)) \\&= (a + 0, x - F(0, 0) + F(a, 0)) \\&= (a, x - F(0, 0) + F(0, 0)) \quad (\text{ by (CE) }) \\&= (a, x).\end{aligned}$$

Hence $(0, -F(0, 0))$ is the identity element in W . We denote this identity element by 0_W .

[Home Page](#)[Title Page](#)[Contents](#)[Page 35 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Step 3. For each $a \in G$ and $x \in H$,

$$\begin{aligned}(a, x) + (-a, -x - F(a, -a) - F(0, 0)) \\&= (a - a, x - x - F(a, -a) - F(0, 0) + F(a, -a)) \\&= (0, -F(0, 0)) \\&= 0_W.\end{aligned}$$

Therefore, every element (a, x) in W has an inverse $(-a, -x - F(a, -a) - F(0, 0))$ in W .

[Home Page](#)[Title Page](#)[Contents](#)[Page 36 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Step 4. For any $(a, x), (b, y), (c, z) \in W$, we have

$$\begin{aligned} & ((a, x) + (b, y)) + (c, z) \\ &= (a + b, x + y + F(a, b)) + (c, z) \\ &= (a + b + c, x + y + z + F(a, b) + F(a + b, c)) \\ &= (a + b + c, x + y + z + F(b, c) + F(a, b + c)) \\ &= (a, x) + (b + c, y + z + F(b, c)) \\ &= (a, x) + ((b, y) + (c, z)). \end{aligned}$$

Hence “+” is associative.

[Home Page](#)[Title Page](#)[Contents](#)[Page 37 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Consider the projection $\phi : W \rightarrow G$ defined by $\phi(a, x) = a$.

Then

$$\begin{aligned}\phi((a, x) + (b, y)) &= \phi(a + b, x + y + F(a, b)) \\ &= a + b \\ &= \phi(a, x) + \phi(b, y).\end{aligned}$$

Hence ϕ is a homomorphism from W onto G .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 38 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Next, we show a function $f : G \rightarrow H$ satisfies

$$F(a, b) = f(a + b) - f(a) - f(b) \quad (6)$$

for all $a, b \in G$ if and only if

$$K := \{(a, f(a)) \mid a \in G\} \quad (7)$$

is a subgroup of the abelian group W .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 39 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Suppose $F(a, b) = f(a + b) - f(a) - f(b)$ for all $a, b \in G$.

We want to show K is a subgroup of W . From (6), we have

$$F(a, -b) = f(a - b) - f(a) - f(-b)$$

$$F(b, -b) = f(0) - f(b) - f(-b)$$

$$F(0, 0) = f(0) - f(0) - f(0).$$

Hence

$$\begin{aligned} F(a, -b) - F(0, 0) - F(b, -b) \\ = f(a - b) - f(a) + f(b). \end{aligned} \quad (8)$$

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 40 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Let $(a, f(a))$ and $(b, f(b))$ be any two elements in K . We want to show that

$$(a - b, f(a - b)) \in K.$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 41 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Using (8), we compute

$$\begin{aligned} & (a, f(a)) - (b, f(b)) \\ &= (a, f(a)) + (-b, -f(b) - F(b, -b) - F(0, 0)) \\ &= (a - b, f(a) - f(b) - F(b, -b) - F(0, 0) + F(a, -b)) \\ &= (a - b, f(a) - f(b) + f(a - b) - f(a) + f(b)) \\ &= (a - b, f(a - b)) \in K. \end{aligned}$$

Hence K is a subgroup of W .

[Home Page](#)[Title Page](#)[Contents](#)

Page 42 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Next, we assume that K is a subgroup of W . We want to show that (6) holds.

Since $K < W$, therefore, whenever $(a, f(a))$ and $(b, f(b))$ are elements of K , we have

$$\begin{aligned} (a, f(a)) + (b, f(b)) &\in K \\ \Rightarrow (a + b, f(a) + f(b) + F(a, b)) &\in K. \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 43 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Hence from the definition of K , we have

$$f(a + b) = f(a) + f(b) + F(a, b)$$

which is

$$F(a, b) = f(a + b) - f(a) - f(b). \quad (6)$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 44 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Thus, the existence of a function $f : G \rightarrow H$ satisfying (6) is equivalent to the existence of a subgroup S of W which by ϕ is mapped bijectively onto G .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 45 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Since G is a free abelian group, G has a basis $\{a_i \mid i \in I\}$ in the sense that every element of G can be written in one and only one way as a **finite linear combination** of elements of the basis with integer coefficients.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 46 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Let x_i for $i \in I$ be arbitrary elements of the group H . Let

$$S = \langle \{(a_i, x_i) \mid i \in I\} \rangle$$

be the subgroup generated by the set $\{(a_i, x_i) \mid i \in I\}$. Then S is a subgroup of W . It is easy to see that the projection map ϕ maps S bijectively onto G .

[Home Page](#)[Title Page](#)[Contents](#)[Page 47 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Hence f is determined by its values

$$f(a_i) = x_i, \quad i \in I,$$

and these value can be arbitrarily chosen. This completes the proof of the theorem.

[Home Page](#)[Title Page](#)[Contents](#)

Page 48 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Theorem 2 (Hosszú, 1965) *Let H and G be abelian groups. Suppose H is uniquely 2-divisible. The most general solution $F : G^2 \rightarrow H$ of*

$$F(a, b) + F(a + b, c) = F(b, c) + F(a, b + c) \quad (\text{CE})$$

is of the form

$$F(a, b) = S(a, b) + A(a, b)$$

where S is a symmetric solution of (CE) and A is an anti-symmetric one, additive in both variables.

[Home Page](#)[Title Page](#)[Contents](#)

Page 49 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: Define

$$S(a, b) = \frac{1}{2} [F(a, b) + F(b, a)]$$

and

$$A(a, b) = \frac{1}{2} [F(a, b) - F(b, a)] .$$

Then $S(a, b) = S(b, a)$ and $A(a, b) = -A(b, a)$.

[Home Page](#)

[Title Page](#)

[Contents](#)



Page 50 of 100

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Also $S(a, b)$ and $A(a, b)$ are solutions of (CE). To see $A(a, b)$ is a solution of (CE), consider

$$\begin{aligned} & 2[A(a, b) + A(a + b, c)] \\ & \stackrel{\text{def}}{=} F(a, b) - F(b, a) + F(a + b, c) - F(c, a + b) \\ & = F(a, b) + F(a + b, c) - F(b, a) - F(c, a + b) \\ & \stackrel{(\text{CE})}{=} F(b, c) + F(a, b + c) - F(c, b) - F(c + b, a) \\ & = F(b, c) - F(c, b) + F(a, b + c) - F(b + c, a) \\ & = 2[A(b, c) + A(a, b + c)]. \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 51 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Similarly, it can be shown that $S(a, b)$ is a solution of (CE).

- Since A is antisymmetric, it is enough to show that A is additive in the first variable.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 52 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

$$2 A(a + b, c)$$

$$\stackrel{def}{=} F(a + b, c) - F(c, a + b)$$

$$\stackrel{\text{(CE)}}{=} F(a, b + c) + F(b, c) - F(a, b) \\ - [F(c + a, b) + F(c, a) - F(a, b)]$$

$$\stackrel{\text{(CE)}}{=} F(a, b + c) + F(b, c) \\ - [F(a, b + c) + F(c, b) - F(a, c) + F(c, a)]$$

$$\stackrel{def}{=} 2 A(b, c) + 2 A(a, c).$$

This completes the proof of the theorem.


[Home Page](#)
[Title Page](#)
[Contents](#)
[◀◀](#)
[▶▶](#)
[◀](#)
[▶](#)

Page 53 of 100

[Go Back](#)
[Full Screen](#)
[Close](#)
[Quit](#)



In 1995, Szekelyhidi found the bounded solution of the functional equation (CE) on a large class of groups known as amenable group.

To present his result, I will first remind you the definition of amenable group.

[Home Page](#)[Title Page](#)[Contents](#)[Page 54 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Amenable Groups

Let G be a finite group with $G = \{x_1, x_2, x_3, \dots, x_n\}$. Let $f : G \rightarrow \mathbb{C}$. Define

$$M(f) = \frac{1}{n} \sum_{k=1}^n f(x_k).$$

Then M satisfies

$$M(\alpha f + \beta g) = \alpha M(f) + \beta M(g).$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 55 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Hence M is a linear functional on the space of all complex valued functions f defined on G . When f is real-valued, it is easy to see that

$$\min\{f(x_1), \dots, f(x_n)\} \leq M(f) \leq \max\{f(x_1), \dots, f(x_n)\}$$

Moreover, it is also easy to check that

$$M(\overline{f}) = \overline{M(f)}$$

for all $f : G \rightarrow \mathbb{C}$.

[Home Page](#)[Title Page](#)[Contents](#)

Page 56 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Define

$$f_j(x_i) = f(x_i x_j) \quad \text{and} \quad {}_j f(x_i) = f(x_j x_i).$$

Then

$$\begin{aligned} M(f_j) &= \frac{1}{n} \sum_{i=1}^n f(x_i x_j) \\ &= \frac{1}{n} \sum_{i=1}^n f(x_i) = M(f). \end{aligned}$$

- Hence $M(f_j) = M(f)$ and M is right invariant.
- Similarly, $M({}_j f) = M(f)$ and M is left invariant.

[Home Page](#)[Title Page](#)[Contents](#)

Page 57 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Therefore $M(f)$ is an invariant mean for the finite group G .

This leads to the following definition of invariant mean for an arbitrary group.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 58 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Definition 1 *Let G be a group (or a semigroup). Let $B(G, \mathbb{C})$ be a space of all bounded complex valued functions defined on G with norm*

$$||f|| = \sup\{f(x) \mid x \in G\}.$$

A linear functional M in $B(G, \mathbb{C})$ is said to be a right invariant mean if the following conditions hold:

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 59 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



$$M(\overline{f}) = \overline{M(f)} \quad \forall f \in B(G, \mathbb{C}); \quad (9)$$

$$\inf\{f(x) \mid x \in G\} \leq M(f) \leq \sup\{f(x) \mid x \in G\} \quad (10)$$

for all real valued $f \in B(G, \mathbb{C})$;

$$M(f_x) = M(f) \quad (11)$$

for all $x \in G$ and $f \in B(G, \mathbb{C})$ where $f_x(t) = f(tx)$.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 60 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



A mean which is both **left** and **right** invariant is said to be **two-sided** invariant or just invariant.

A semigroup S that admits a left (right) invariant mean on $B(S, \mathbb{C})$ is called a left (right) amenable. If the semigroup S admits a two-sided invariant mean on $B(S, \mathbb{C})$, then S is said to be two-sided amenable or simply amenable.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 61 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

- The group of integers is amenable.
- Every finite group (semigroup) is amenable.
- Every commutative group (semigroup) is amenable.
- Every subgroup of an amenable group is amenable.
- If a group is left (right) amenable, then it is amenable.
- The free group on two generators is non-amenable.

[Home Page](#)[Title Page](#)[Contents](#)[Page 62 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Theorem 3 (Szekelyhidi, 1995) *Let G be a right amenable semigroup and let $F : G \times G \rightarrow \mathbb{C}$ be a bounded function satisfying*

$$F(x, y) + F(xy, z) = F(x, yz) + F(y, z)$$

for all $x, y, z \in G$. Then there exists a unique bounded function $f : G \rightarrow \mathbb{C}$ with

$$F(x, y) = f(x) + f(y) - f(xy) \quad \forall x, y \in G.$$

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 63 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: Let M denote any right invariant mean defined on the set of all bounded complex valued functions on G .

We write M_x to indicate that M is applied on the argument as a function of x .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)

Page 64 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



We apply M on both sides of

$$F(x, y) + F(xy, z) = F(x, yz) + F(y, z)$$

as functions of x , for any fixed $y, z \in G$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 65 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Thus

$$M_x[F(x, y)] + M_x[F(xy, z)] = M_x[F(x, yz)] + M_x[F(y, z)].$$

By right invariance of M and the fact that $M(1) = 1$, we obtain

$$M_x[F(x, y)] + M_x[F(x, z)] = M_x[F(x, yz)] + F(y, z).$$

[Home Page](#)[Title Page](#)[Contents](#)

Page 66 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Therefore

$$F(y, z) = M_x[F(x, y)] + M_x[F(x, z)] - M_x[F(x, yz)].$$

So defining

$$f(y) := M_x[F(x, y)] \quad \forall y \in G, \quad (12)$$

we have

$$F(y, z) = f(y) + f(z) - f(yz). \quad (13)$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 67 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



As F is bounded, it follows from (12) that f is bounded. Next we prove that f is unique. Suppose not. Then there exists g such that

$$F(y, z) = g(y) + g(z) - g(yz). \quad (14)$$

From (13) and (14), we have

$$(f - g)(yz) = (f - g)(y) + (f - g)(z) \quad \forall y, z \in G.$$

[Home Page](#)[Title Page](#)[Contents](#)[«](#) [»](#)[◀](#) [▶](#)[Page 68 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Since $f - g$ is a complex homomorphism of G , therefore it cannot be bounded unless it is zero. To see this, let $\phi = f - g$.

Since ϕ is a complex homomorphism of group G , therefore $\phi(x^n) = n\phi(x)$ for all $x \in G$. Hence

$$\phi(x) = \lim_{n \rightarrow \infty} \phi(x) = \lim_{n \rightarrow \infty} \frac{\phi(x^n)}{n} = 0$$

for each $x \in G$. Thus $\phi = 0$. Hence $f = g$. So f is unique.

This completes the proof of the theorem.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)

Page 69 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Generalized Cocycle Equation: In 1994, motivated by Szekelyhidi's result, Páles considered the following functional equation

$$F(x, y) + \frac{1}{n} \sum_{i=1}^n F(x\phi_i(y), z) = \frac{1}{n} \sum_{i=1}^n F(x, y\phi_i(z)) + F(y, z)$$

where $\phi_1, \phi_2, \dots, \phi_n : S \rightarrow S$ are pairwise distinct homomorphisms of the semigroup S and the set $\{\phi_1, \phi_2, \dots, \phi_n\}$ is a group under function composition.

[Home Page](#)[Title Page](#)[Contents](#)[«](#) [»](#)[◀](#) [▶](#)[Page 70 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

If $n = 1$ and $\phi_1(x) = x$, then the last functional equation reduces to 2-cocycle functional equation.

If $n = 2$, $\phi_1(x) = x$ and $\phi_2(x) = x^{-1}$, then the equation reduces to another interesting functional equation

$$\begin{aligned} F(x, y) + \frac{1}{2} [F(xy, z) + F(xy^{-1}, z)] \\ = \frac{1}{2} [F(xy, z) + F(x, yz^{-1})] + F(y, z). \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 71 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Theorem 4 (Páles, 1994) Suppose S is a semigroup and $f : S \rightarrow \mathbb{C}$ an arbitrary function. Then the function $F : S \times S \rightarrow \mathbb{C}$ defined by

$$F(x, y) = f(x) + f(y) - \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y)) \quad (15)$$

is a solution of the functional equation

$$F(x, y) + \frac{1}{n} \sum_{i=1}^n F(x\phi_i(y), z) = \frac{1}{n} \sum_{i=1}^n F(x, y\phi_i(z)) + F(y, z).$$

[Home Page](#)[Title Page](#)[Contents](#)[«](#) [»](#)[◀](#) [▶](#)[Page 72 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Proof. Computing the left side of the functional equation,
we get

$$\begin{aligned} & F(x,y) + \frac{1}{n} \sum_{i=1}^n F(x\phi_i(y), z) \\ & \stackrel{(15)}{=} f(x) + f(y) - \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y)) \\ & \quad + \frac{1}{n} \sum_{i=1}^n [f(x\phi_i(y)) + f(z) - \frac{1}{n} \sum_{j=1}^n f(x\phi_i(y)\phi_j(z))] \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 73 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

$$\begin{aligned}
&= f(x) + f(y) - \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y)) + \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y)) \\
&\quad + f(z) - \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n f(x\phi_i(y)\phi_j(z)) \\
&= f(x) + f(y) + f(z) - \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n f(x\phi_i(y)\phi_j(z)).
\end{aligned}$$


[Home Page](#)
[Title Page](#)
[Contents](#)
[◀◀](#)
[▶▶](#)
[◀](#)
[▶](#)

Page 74 of 100

[Go Back](#)
[Full Screen](#)
[Close](#)
[Quit](#)



Similarly, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n F(x, y\phi_i(z)) + F(y, z) \\ &= f(x) + f(y) + f(z) - \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n f(x\phi_j(y\phi_i(z))). \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[Page 75 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Hence

$$\begin{aligned} & F(x, y) + \frac{1}{n} \sum_{i=1}^n F(x\phi_i(y), z) - \frac{1}{n} \sum_{i=1}^n F(x, y\phi_i(z)) - F(y, z). \\ &= f(x) + f(y) + f(z) - \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n f(x\phi_i(y)\phi_j(z)) \\ &\quad - f(x) - f(y) - f(z) + \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n f(x\phi_j(y\phi_i(z))) \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 76 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 77 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

$$\begin{aligned} &= \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n [f(x\phi_j(y\phi_i(z))) - f(x\phi_i(y)\phi_j(z))] \\ &= \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n [f(x\phi_j(y)\phi_j(\phi_i(z)))) - f(x\phi_i(y)\phi_j(z))] \\ &= \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n [f(x\phi_j(y)\phi_i(z))) - f(x\phi_i(y)\phi_j(z))] = 0 \end{aligned}$$

where we have used the fact that ϕ_i are homomorphisms and $\{\phi_1, \phi_2, \dots, \phi_n\}$ is a group under composition. This proof is now complete.



In the next theorem, we present the bounded solutions of the generalized cocycle equation on right-amenable semigroups. The proof of this theorem is due to Páles but it is based on Szekelyhidi's proof.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 78 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Theorem 5 (Páles, 1994) *Let S be a right-amenable semi-group and $F : S \times S \rightarrow \mathbb{C}$ be a bounded function. Then F satisfies*

$$F(x, y) + \frac{1}{n} \sum_{i=1}^n F(x\phi_i(y), z) = \frac{1}{n} \sum_{i=1}^n F(x, y\phi_i(z)) + F(y, z)$$

for all $x, y, z \in S$ if and only if there exists a bounded function $f : S \rightarrow \mathbb{C}$ such that

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 79 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



$$F(x, y) = f(x) + f(y) - \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y))$$

for all $x, y \in S$.

Moreover, the representing function $f : S \rightarrow \mathbb{C}$ of F is unique.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 80 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Proof. Suppose F can be represented by (15), that is

$$F(x, y) = f(x) + f(y) - \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y))$$

with some bounded f . Then from (15), we see that F is bounded. By Theorem 4, $F(x, y)$ given by

$$F(x, y) = f(x) + f(y) - \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y))$$

is a solution of generalized cocycle equation.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 81 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

To prove the converse, let M be a right-invariant mean on the space of bounded complex-valued functions on S . We write M_x to indicate that M is applied on the argument as function of x .

Assume now that a bounded function F satisfies the generalized cocycle equation. Fix $y, z \in S$ and apply M_x on both sides of the generalized cocycle equation.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 82 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Then using the right-invariance and $M(1) = 1$, we obtain

$$\begin{aligned} M_x \left[F(x, y) + \frac{1}{n} \sum_{i=1}^n F(x\phi_i(y), z) \right] \\ = M_x \left[\frac{1}{n} \sum_{i=1}^n F(x, y\phi_i(z)) + F(y, z) \right] \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 83 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Hence

$$\begin{aligned} M_x[F(x, y)] + \frac{1}{n} \sum_{i=1}^n M_x[F(x\phi_i(y), z)] \\ = \frac{1}{n} \sum_{i=1}^n M_x[F(x, y\phi_i(z))] + F(y, z) \end{aligned}$$

Since $M_x[F(x\phi_i(y), z)] = M_x[F(x, z)]$, we have

$$\begin{aligned} M_x[F(x, y)] + M_x[F(x, z)] \\ = \frac{1}{n} \sum_{i=1}^n M_x[F(x, y\phi_i(z))] + F(y, z) \end{aligned}$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 84 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Define a function $f : S \rightarrow \mathbb{C}$ by

$$f(u) = M_x[F(x, u)]. \quad (16)$$

Then from the last equality, we obtain

$$F(x, y) = f(x) + f(y) - \frac{1}{n} \sum_{i=1}^n f(x\phi_i(y)).$$

which is (15).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 85 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



It remains to prove that f is uniquely determined. Suppose not. Then there exist $f_1, f_2 : S \rightarrow \mathbb{C}$ such that

$$F(x, y) = f_1(x) + f_1(y) - \frac{1}{n} \sum_{i=1}^n f_1(x\phi_i(y)) \quad (17)$$

and

$$F(x, y) = f_2(x) + f_2(y) - \frac{1}{n} \sum_{i=1}^n f_2(x\phi_i(y)). \quad (18)$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 86 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Let $g := f_1 - f_2$. Then (17) and (18) imply

$$g(x) + g(y) = \frac{1}{n} \sum_{i=1}^n g(x\phi_i(y)). \quad (19)$$

Applying M_x to both sides of (19), we get

$$M_x[g(x)] + g(y) = \frac{1}{n} \sum_{i=1}^n M_x[g(x\phi_i(y))].$$

which is

$$M_x[g(x)] + g(y) = M_x[g(x)].$$

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 87 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Therefore $g(y) = 0$ for all $y \in S$. Hence

$$f_1(y) = f_2(y) \quad \forall y \in S.$$

This shows that f is unique and the proof of the theorem is now complete.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 88 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



- Based on the functional equation (CE), we make several remarks.

Remark 1 *The function F satisfying (CE) and the conditions $F(x, e) = 0 = F(e, x)$ is called a cocycle.*

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 89 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Remark 2 *The set of all cocycles from $G \times G$ to H is denoted by $\mathbb{Z}^2(G, H)$. Hence*

$$\mathbb{Z}^2(G, H) = \left\{ F : G \times G \rightarrow H \mid \right. \\ \left. F \in S(G, H) \text{ \& } F(a, e) = 0 = F(e, b) \right\},$$

where $S(G, H)$ is the set of solution of 2-cocycle functional equation (CE).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 90 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Recall that the real-valued solution of 2-cocycle equation on free abelian groups is of the form

$$F(a, b) = f(ab) - f(a) - f(b) \quad \forall a, b \in G.$$

Remark 3 $\mathbb{Z}^2(G, H)$ is an abelian group under point-wise addition of cocycles.

[Home Page](#)[Title Page](#)[Contents](#)[Page 91 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Remark 4 Given a function $f : G \rightarrow H$ the map

$$(a, b) \mapsto f(a + b) - f(a) - f(b)$$

is called the coboundary of f if $f(0) = 0$.

- Note that this map measures how much f deviates being a homomorphism of G into H .

[Home Page](#)[Title Page](#)[Contents](#)[Page 92 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Remark5 *The set of all coboundaries from $G \times G$ to H is denoted by $\mathbb{B}^2(G, H)$.*

Remark6 $\mathbb{B}^2(G, H)$ *is an abelian group under point-wise addition of coboundaries.*

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 93 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Remark 7 $\mathbb{B}^2(G, H)$ is a normal subgroup of $\mathbb{Z}^2(G, H)$.

Remark 8 The quotient group

$$\mathbb{H}^2(G, H) = \mathbb{Z}^2(G, H) / \mathbb{B}^2(G, H)$$

is called the second cohomology group of G .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 94 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Remark9 *Szekelyhidi's theorem can be rephrased as: For any discrete amenable group G , the second bounded cohomology group $\mathbb{H}_b^2(G, \mathbb{C}) = 0$.*

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 95 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Remark 10 *Ramsey (2005) proved the following result: For any discrete amenable group G , the bounded cohomology group $\mathbb{H}_b^n(G, \mathbb{R}) = 0$ for any $n \in \mathbb{N}$.*

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 96 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Remark 11 *We have studied (see Faiziev and Sahoo (2005)) the second bounded cohomology of amalgamated product of groups.*

[Home Page](#)[Title Page](#)[Contents](#)

Page 97 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

References

- [1] B. Ebanks and H. Stetkaer, Continuous cocycles on locally compact groups, *Aequat. Math.*, 78 (2009), 123-145.
- [2] B. Ebanks and H. Stetkaer, Continuous cocycles on locally compact groups: II, *Aequat. Math.*, 80 (2010), 57-83.
- [3] J. Erdos, A remark on the paper "On some functional equations" by S. Kurepa, *Glasnik Mat.*, 14 (1959), 3-5.
- [4] V.A. Faiziev and P.K. Sahoo, Pseudocharacters and the problem of expressibility for some groups, *Journal of Algebra*, 250 (2002), 603-637.
- [5] V.A. Faiziev and P.K. Sahoo, Remark on the second bounded cohomology of amalgamated product of groups, *Sarajevo J. Math.* 1 (13) (2005), 27-48.
- [6] D.C. Isaksen, A cohomological viewpoint of elementary school arithmetic, *Monthly* 109 (2002), 796-805.
- [7] M. Hosszu, On a functional equation treated by S. Kurepa, *Glasnik Mat.*, 18 (1963), 59-60.
- [8] M. Hosszu, On a class of functional equations, *Publ. De L'Institute Mat.*, 3 (1963), 53-55.
- [9] D.C. Isaksen, A cohomological viewpoint on elementary school arithmetic, *Monthly*, 109 (2002), 796-805.
- [10] B. Jessen, J. Karpf and A. Thorup, Some functional equations in groups and rings, *Math. Scand.* 22 (1968), 257-265.
- [11] S. Kurepa, On some functional equations, *Glasnik Mat.*, 11 (1956), 3-5.
- [12] Z. Pales, Bounded solutions and stability of functional equations for two variables, *Results Math.* 26 (1994), 360-365.
- [13] L. Szekelyhidi, Stability properties of functional equations in several variables, *Publ. Math. Debrecen*, 47 (1995), 95-100.

[Home Page](#)[Title Page](#)[Contents](#)[Page 98 of 100](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Thank You



Home Page

Title Page

Contents



Page 99 of 100

Go Back

Full Screen

Close

Quit



A group G is said to be divisible if for every $x \in G$ and for every positive integer n there exists a $y \in G$ such that $ny = x$.

Example 1. $(\mathbb{Z}, +)$ is not a divisible group.

Example 2. $(\mathbb{Q}, +)$ is a divisible group.

Example 3. $(\mathbb{R}, +)$ is a divisible group.

[Home Page](#)[Title Page](#)[Contents](#)

Page 100 of 100

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)