

Remaining Cauchy functional equations

January 23, 2017



[Home Page](#)

[Title Page](#)

[Contents](#)



[Page 1 of 56](#)

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Remaining Cauchy functional equations

Ron Sahoo

Department of Mathematics

University of Louisville, Louisville, Kentucky 40292 USA

January, 2017



[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#)

[▶▶](#)

[◀](#)

[▶](#)

[Page 2 of 56](#)

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Introduction

In Chapter 5 of his book *Cours d'Analyse*, A. L. Cauchy (1821) also studied three other functional equations, namely,

$$f(x + y) = f(x)f(y), \quad (1)$$

$$f(xy) = f(x) + f(y) \quad (2)$$

and

$$f(xy) = f(x)f(y) \quad (3)$$

besides the additive Cauchy functional equation

$$f(x + y) = f(x) + f(y) \quad (4)$$

for all $x, y \in \mathbb{R}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 3 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

In this lecture, we solve the remaining Cauchy functional equations.

The general solution of each of these functional equations is determined in terms of the additive function.

Finally, using the general solution, the continuous solution is provided for each of these functional equations.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 4 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Solution of Exponential Cauchy Equation

First, we determine the general solution of the exponential Cauchy functional equation (1) without assuming any regularity condition such as continuity, boundedness or differentiability on the unknown function f .

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 5 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Theorem 1 . *If the functional equation (1), that is,*

$$f(x + y) = f(x)f(y),$$

holds for all real numbers x and y , then the general solutions of (1) are given by

$$f(x) = e^{A(x)} \quad \text{and} \quad f(x) = 0 \quad \forall x \in \mathbb{R}, \quad (5)$$

where $A : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function and e is the Napierian base of logarithm.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 6 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: It is easy to see that $f(x) = 0$ for all $x \in \mathbb{R}$ is a solution of (1), that is

$$f(x + y) = f(x) f(y) \quad \text{for all } x, y \in \mathbb{R}.$$

Hence from now on we suppose that $f(x)$ is not identically zero.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 7 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

To complete the rest of the proof of the theorem, we need to show

- (a) f is nowhere zero,
- (b) f is strictly positive,
- (c) $A := \ln f$ is additive.

[Home Page](#)[Title Page](#)[Contents](#)

Page 8 of 56

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



We claim that $f(x)$ is nowhere zero. Suppose not. Then there exists a y_o such that $f(y_o) = 0$. From (1), we get

$$f(y) = f((y - y_o) + y_o) = f(y - y_o) f(y_o) = 0$$

for all $y \in \mathbb{R}$.

This is a contradiction to our assumption that $f(x)$ is not identically zero. Hence $f(x)$ is nowhere zero.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 9 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Letting $x = \frac{t}{2} = y$ in $f(x + y) = f(x) f(y)$, we see that

$$f(t) = f\left(\frac{t}{2}\right)^2$$

for all $t \in \mathbb{R}$. Hence $f(x)$ is a strictly positive. Now taking natural logarithm of both sides of $f(x + y) = f(x) f(y)$, we obtain

$$\ln f(x + y) = \ln f(x) + \ln f(y).$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 10 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Defining $A : \mathbb{R} \rightarrow \mathbb{R}$ by $A(x) = \ln f(x)$, we have

$$A(x + y) = A(x) + A(y). \quad (6)$$

Hence we have the asserted solution $f(x) = e^{A(x)}$ and the proof is now complete.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 11 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The following corollary is obvious from the above theorem.

Corollary 1 *If the equation (1), that is, $f(x+y) = f(x)f(y)$, holds for all real numbers x and y , then the general continuous solutions of (1) are given by*

$$f(x) = e^{cx} \quad \text{and} \quad f(x) = 0 \quad \forall x \in \mathbb{R}, \quad (7)$$

where c is an arbitrary real constant.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 12 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Next we give the definition of exponential functions.

Definition 1 A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is called a (real-valued) real exponential function if it satisfies $f(x + y) = f(x) f(y)$ for all $x, y \in \mathbb{R}$.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 13 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Let n be a positive integer. Suppose the functional equation

$$f(x + y + nxy) = f(x) f(y) \quad (8)$$

holds for all reals $x > -\frac{1}{n}$ and all $y > -\frac{1}{n}$.

- When $n \rightarrow 0$, the functional equation (8) reduces to the exponential Cauchy functional equation.
- This equation was studied by Thielman (1949).

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 14 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Theorem2 . *Every solution f of the functional equation*

$$f(x + y + nxy) = f(x) f(y)$$

for all reals $x > -\frac{1}{n}$ and all $y > -\frac{1}{n}$ is of the form

$$f(x) = 0 \quad \text{or} \quad f(x) = e^{A(\ln(1+nx))}, \quad (9)$$

where $A : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 15 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Proof: We write the functional equation (8) as

$$f\left(\frac{(1+nx)(1+ny)-1}{n}\right) = f(x)f(y). \quad (10)$$

Next we define $0 < 1+nx = e^u$ and $0 < 1+ny = e^v$ so that $u = \ln(1+nx)$ and $v = \ln(1+ny)$.

Now rewriting (10), we obtain

$$f\left(\frac{e^{u+v}-1}{n}\right) = f\left(\frac{e^u-1}{n}\right) f\left(\frac{e^v-1}{n}\right) \quad (11)$$

for all $u, v \in \mathbb{R}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 16 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Letting

$$\phi(u) = f \left(\frac{e^u - 1}{n} \right) \quad (12)$$

in (11), we have

$$\phi(u + v) = \phi(u) \phi(v) \quad (13)$$

for all $u, v \in \mathbb{R}$.

[Home Page](#)[Title Page](#)[Contents](#)

Page 17 of 56

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Hence by Theorem 1, we have

$$\phi(x) = e^{A(x)} \quad \text{or} \quad \phi(x) = 0 \quad \forall x \in \mathbb{R}, \quad (14)$$

where $A : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function and e is the Napierian base of logarithm.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 18 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Therefore from

$$\phi(u) = f\left(\frac{e^u - 1}{n}\right)$$

and $\phi(x) = e^{A(x)}$ or $\phi(x) = 0$, we obtain

$$f(x) = 0 \quad \text{or} \quad f(x) = e^{A(\ln(1+nx))},$$

where $A : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function. The proof of the theorem is now complete.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)

Page 19 of 56

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The following corollary is obvious.

Corollary 2 *Every continuous solution f of the functional equation (8) holding for all reals $x > -\frac{1}{n}$ and all $y > -\frac{1}{n}$ is of the form*

$$f(x) = 0 \quad \text{or} \quad f(x) = (1 + nx)^k, \quad (15)$$

where k is an arbitrary constant.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 20 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Solution of Logarithmic Cauchy Equation

Now we consider the second Cauchy functional equation (2), that is

$$f(xy) = f(x) + f(y) \quad \text{for all } x, y \in \mathbb{R} \setminus \{0\}.$$

This functional equation is known as the logarithmic Cauchy equation.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 21 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Theorem 3 . *If the functional equation (2), that is,*

$$f(xy) = f(x) + f(y)$$

holds for all $x, y \in \mathbb{R} \setminus \{0\}$, then the general solution of (2) is given by

$$f(x) = A(\ln |x|) \quad \forall x \in \mathbb{R} \setminus \{0\}, \quad (16)$$

where A is an additive function.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 22 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: To prove this theorem, we have to show

(a) f is even on $\mathbb{R} \setminus \{0\}$, and

(b) $A(s) := \ln(e^s)$ is additive on \mathbb{R} .

[Home Page](#)

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 23 of 56

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

Substitute $x = t$ and $y = t$ in $f(xy) = f(x) + f(y)$ to get

$$f(t^2) = 2f(t).$$

Letting $x = -t$ and $y = -t$ in $f(xy) = f(x) + f(y)$, we have

$$f(t^2) = 2f(-t).$$

Hence we see that

$$f(t) = f(-t) \quad \forall t \in \mathbb{R} \setminus \{0\}. \quad (17)$$

[Home Page](#)[Title Page](#)[Contents](#)[Page 24 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Next, suppose the functional equation $f(xy) = f(x) + f(y)$ holds for all $x > 0$ and $y > 0$. Let

$$x = e^s \quad \text{and} \quad y = e^t \quad (18)$$

so that

$$s = \ln x \quad \text{and} \quad t = \ln y. \quad (19)$$

Note that $s, t \in \mathbb{R}$ since $x, y \in \mathbb{R}_+$ where

$$\mathbb{R}_+ = \{x \in \mathbb{R} \mid x > 0\}.$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 25 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Substituting $x = e^s$ and $y = e^t$ in $f(xy) = f(x) + f(y)$, we obtain

$$f(e^{s+t}) = f(e^s) + f(e^t).$$

Defining

$$A(s) = f(e^s) \tag{20}$$

and using the last equation we have

$$A(s + t) = A(s) + A(t)$$

for all $s, t \in \mathbb{R}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 26 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Hence from the definition $A(s) = f(e^s)$, we have

$$f(x) = A(\ln x) \quad \forall x \in \mathbb{R}_+. \quad (21)$$

Since $f(t) = f(-t)$, we see that the general solution of $f(xy) = f(x) + f(y)$ is

$$f(x) = A(\ln |x|) \quad \forall x \in \mathbb{R} \setminus \{0\}$$

and the proof is now complete.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 27 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The following corollary is a consequence of the last theorem.

Corollary 3 *The general solution of the functional equation*

$f(xy) = f(x) + f(y)$ for all $x, y \in \mathbb{R}_+$ is given by

$$f(x) = A(\ln x), \quad (22)$$

where $A : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function.

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 28 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



The following result is also trivial.

Corollary 4 *The general solution of the functional equation*
 $f(xy) = f(x) + f(y)$ *for all* $x, y \in \mathbb{R}$ *is given by*

$$f(x) = 0 \quad \forall x \in \mathbb{R}. \quad (23)$$

Proof: Substitute $y = 0$ in (2) to get $f(0) = f(x) + f(0)$ and hence we have the asserted solution. **QED**

[Home Page](#)[Title Page](#)[Contents](#)[<<](#)[>>](#)[<](#)[>](#)[Page 29 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Corollary 5 *The general continuous solution of the functional equation $f(xy) = f(x) + f(y)$ for all x, y in $\mathbb{R} \setminus \{0\}$ is given by*

$$f(x) = c \ln |x| \quad \forall x \in \mathbb{R} \setminus \{0\}, \quad (24)$$

where c is an arbitrary real constant.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 30 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Definition 2 A function $f : \mathbb{R}_+ \rightarrow \mathbb{R}$ is called a logarithmic function if it satisfies $f(xy) = f(x) + f(y)$ for all $x, y \in \mathbb{R}_+$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 31 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Solution of Multiplicative Cauchy Equation

Now we treat the last Cauchy equation (3), that is

$$f(xy) = f(x)f(y) \quad \text{for all } x, y \in \mathbb{R}.$$

This equation is the most complicated of the three equations considered in this chapter.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 32 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



In the following theorem we need the notion of the signum function. The signum function is denoted by $\operatorname{sgn}(x)$ and defined as

$$\operatorname{sgn}(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0. \end{cases} \quad (25)$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 33 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

It is easy to see that

$$|\operatorname{sgn}(x)| = \begin{cases} 1 & \text{if } x \in \mathbb{R} \setminus \{0\} \\ 0 & \text{if } x = 0. \end{cases}$$

Hence the following function

$$f(x) = \begin{cases} e^{A(\ln |x|)} & \text{if } x \in \mathbb{R} \setminus \{0\} \\ 0 & \text{if } x = 0. \end{cases}$$

can be compactly written as $f(x) = e^{A(\ln |x|)} |\operatorname{sgn}(x)|$.

[Home Page](#)[Title Page](#)[Contents](#)[Page 34 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Theorem4. *The general solutions of $f(xy) = f(x)f(y)$ holding for all $x, y \in \mathbb{R}$ are given by*

$$f(x) = 0, \quad (26)$$

$$f(x) = 1, \quad (27)$$

$$f(x) = e^{A(\ln |x|)} |sgn(x)|, \quad (28)$$

and

$$f(x) = e^{A(\ln |x|)} sgn(x), \quad (29)$$

where $A : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function and e is the Napierian base of logarithm.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 35 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: The steps to prove this theorems are the following:

Step 1. Show $f(0) = 0$, $f(0) = 1$, $f(1) = 0$ and $f(1) = 1$

Step 2. $f(x) \geq 0$ for all $x \in \mathbb{R}_+$

Step 3. If $f(x_o) = 0$, the $f(x) = 0$ for all $x \in \mathbb{R}$

Step 4. If $f(x) \neq 0$ for all $x \in \mathbb{R} \setminus \{0\}$, then $f(x) = 1$

[Home Page](#)[Title Page](#)[Contents](#)

Page 36 of 56

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Step 5. If $f(0) = 0$, then $f(x)$ is nowhere zero on $\mathbb{R} \setminus \{0\}$

Step 6. If $f(x)$ is nowhere zero on $\mathbb{R} \setminus \{0\}$, then the map $A(s) := \ln f(e^s)$ is additive on \mathbb{R}

Step 7. Show $f(1) = 1$ since $f(1) = 0$ yields a contradiction

Step 8. Show $f(1) = 1$ implies $f(-1) = 1$ or $f(-1) = -1$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 37 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Step 9. If $f(1) = 1$, then

$$f(x) = e^A(\ln(|x|)) |sgn(x)|$$

for all $x \in \mathbb{R} \setminus \{0\}$

Step 10. If $f(-1) = -1$, then show

$$f(x) = e^A(\ln(|x|)) sgn(x)$$

for all $x \in \mathbb{R} \setminus \{0\}$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 38 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Letting $x = 0 = y$ in (3), we obtain $f(0)[1 - f(0)] = 0$ and hence either

$$f(0) = 0 \quad \text{or} \quad f(0) = 1. \quad (30)$$

Substituting $x = 1 = y$ in (3), we have $f(1)[1 - f(1)] = 0$ and hence either

$$f(1) = 0 \quad \text{or} \quad f(1) = 1. \quad (31)$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 39 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Let x be a positive real number, that is $x > 0$. Then (3), that is, $f(xy) = f(x)f(y)$ implies

$$f(x) = f(\sqrt{x})^2 \geq 0. \quad (32)$$

Suppose there exists an $x_0 \in \mathbb{R}$, $x_0 \neq 0$ such that $f(x_0) = 0$.

Let $x \in \mathbb{R}$ be an arbitrary real number. Then from (3) we have

$$f(x) = f\left(x_0 \frac{x}{x_0}\right) = f(x_0) f\left(\frac{x}{x_0}\right) = 0$$

for all $x \in \mathbb{R}$ and we obtain the solution (26).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 40 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



From now on we suppose that $f(x) \neq 0$ for all $x \in \mathbb{R} \setminus \{0\}$.

From (30) we have either $f(0) = 0$ or $f(0) = 1$. If $f(0) = 1$, then letting $y = 0$ in (3), we obtain

$$f(0) = f(x)f(0)$$

and hence

$$f(x) = 1.$$

for all $x \in \mathbb{R}$. Thus we have the asserted solution (27).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 41 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Next we consider the case $f(0) = 0$. In this case we claim that f is nowhere zero in $\mathbb{R} \setminus \{0\}$. Suppose not. Then there exists a y_o in $\mathbb{R} \setminus \{0\}$ such that $f(y_o) = 0$. Letting $y = y_o$ in (3), we have

$$f(xy_o) = f(x)f(y_o) = 0.$$

Hence

$$f(x) = 0 \quad \forall x \in \mathbb{R} \setminus \{0\}$$

which is a contradiction to our assumption that f is not identically zero. Thus f is nowhere zero in $\mathbb{R} \setminus \{0\}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 42 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

From the fact that f is nowhere zero in $\mathbb{R} \setminus \{0\}$ and (32), we have

$$f(x) > 0 \quad \text{for} \quad x > 0. \quad (33)$$

Let

$$x = e^s \quad \text{and} \quad y = e^t \quad (34)$$

so that

$$s = \ln x \quad \text{and} \quad t = \ln y. \quad (35)$$

Note that $s, t \in \mathbb{R}$ since $x, y \in \mathbb{R}_+$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 43 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Substituting (34) into (3), we obtain

$$f(e^{s+t}) = f(e^s)f(e^t).$$

Since $f(t) > 0$ for all $t > 0$, taking the natural logarithm of both sides of the last equation, we have

$$A(s+t) = A(s) + A(t),$$

where

$$A(s) = \ln f(e^s) \quad \forall s \in \mathbb{R}. \quad (36)$$

[Home Page](#)[Title Page](#)[Contents](#)[«](#) [»](#)[◀](#) [▶](#)[Page 44 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Thus A is an additive function. From (36) and (35), we obtain

$$f(x) = e^{A(\ln |x|)} \quad \forall x \in \mathbb{R}_+. \quad (37)$$

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 45 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

From (31) we see that either $f(1) = 0$ or $f(1) = 1$. If $f(1) = 0$, then letting $y = 1$ in (3), we obtain

$$f(x) = 0 \quad \forall x \in \mathbb{R} \setminus \{0\}$$

contrary to our assumption that f is not identically zero on $\mathbb{R} \setminus \{0\}$. Hence $f(1) = 1$. Now letting $x = -1 = y$ in (3), we get $f(1) = f(-1)^2$ and hence

$$f(-1) = 1 \quad \text{or} \quad f(-1) = -1. \quad (38)$$

[Home Page](#)[Title Page](#)[Contents](#)[«](#) [»](#)[◀](#) [▶](#)[Page 46 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

If $f(-1) = 1$, then letting $y = -1$ in (3), we have

$$f(-x) = f(x)f(-1) = f(x)$$

for all $x \in \mathbb{R} \setminus \{0\}$. Thus (37) yields $f(x) = e^{A(\ln |x|)}$ for all $x \in \mathbb{R} \setminus \{0\}$. Since $f(0) = 0$, we have

$$f(x) = \begin{cases} e^{A(\ln |x|)} & \text{if } x \in \mathbb{R} \setminus \{0\} \\ 0 & \text{if } x = 0 \end{cases}$$

which is the asserted solution (28).

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 47 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

If $f(-1) = -1$, then letting $y = -1$ in (3), we have

$$f(-x) = f(x)f(-1) = -f(x)$$

for all $x \in \mathbb{R} \setminus \{0\}$. Hence (37) yields

$$f(x) = \begin{cases} e^{A(\ln |x|)} & \text{if } x > 0 \\ -e^{A(\ln |x|)} & \text{if } x < 0 \end{cases}$$

for all $x \in \mathbb{R} \setminus \{0\}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 48 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Together with the fact that $f(0) = 0$, we have

$$f(x) = \begin{cases} e^{A(\ln |x|)} & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -e^{A(\ln |x|)} & \text{if } x < 0 \end{cases}$$

which is the asserted solution (29). Now the proof of the theorem is complete.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 49 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Corollary 6 *The general continuous solution of the equation*

$f(xy) = f(x)f(y)$ for all $x, y \in \mathbb{R}$ is given by

$$f(x) = 0, \quad (39)$$

$$f(x) = 1, \quad (40)$$

$$f(x) = |x|^\alpha, \quad (41)$$

and

$$f(x) = |x|^\alpha \operatorname{sgn}(x), \quad (42)$$

where α is an arbitrary positive real constant.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 50 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Proof: By Theorem 4 either $f = 0$, or $f = 1$, or f has the form (28) or (29), where $A : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function. Since f is continuous and $A(t) = \ln f(e^t)$, A is also continuous on \mathbb{R} . Therefore $A(t) = \alpha t$, where $\alpha \in \mathbb{R}$ is an arbitrary constant. Hence from (28) and (29), we get

$$f(x) = |x|^\alpha \quad \text{and} \quad f(x) = |x|^\alpha \operatorname{sgn}(x),$$

respectively.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 51 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

The only thing remaining to be shown is $\alpha > 0$. If we had $\alpha = 0$, then (41) will yield $f(x) = 1$ for $x \neq 0$, and by continuity of f we must have $f(0) = 1$. Hence we will have $f = 1$, already listed in (40). Formula (42) with $\alpha = 0$ yields

$$f(x) = 1 \quad \text{for} \quad x > 0$$

and

$$f(x) = -1 \quad \text{for} \quad x < 0$$

and thus f cannot be continuous.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 52 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Similarly if $\alpha < 0$, then f given by (41) and (42) satisfies

$$\lim_{x \rightarrow 0^+} f(x) = \infty$$

and hence cannot be continuous at 0. Now the proof of the corollary is complete.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#)[▶▶](#)[◀](#)[▶](#)[Page 53 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



Definition 3 A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is called a *multiplicative function* if it satisfies $f(xy) = f(x) f(y)$ for all $x, y \in \mathbb{R}$.

[Home Page](#)[Title Page](#)[Contents](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)[Page 54 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

References

- [1] P. K. Sahoo and Pl. Kannappan, Introduction to Functional Equations, CRC Press, Boca Raton, 2011.
- [2] P. K. Sahoo and T. Riedel, Mean Value Theorems and Functional Equations, World Scientific, Singapore, 1998.
- [3] B. Ebanks, P.K. Sahoo and W. Sander, Characterizations of Information Measures, World Scientific, Singapore, 1998.

[Home Page](#)[Title Page](#)[Contents](#)[Page 55 of 56](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)

Thank You



Home Page

Title Page

Contents



Page 56 of 56

Go Back

Full Screen

Close

Quit