Convergence And Completeness

Erik An

<2025-12-11 Thu>

Contents

1 1. Convergence of Sequences (Metric Spaces) 2
1.1 Metric Spaces (The Distance Interface) 2
1.1.1 [Definition 1.1.1: Metric] 2
1.2 Convergence of Sequences 3
1.2.1 [Definition 1.1.3: Convergence] 3
1.2.2 [Theorem 1.1.5: Uniqueness of Limit] 3

2 1.2 Completeness 4
2.1 Cauchy Sequences & Completeness 4
2.1.1 [Definition 1.2.1: Cauchy Sequence] 4
2.1.2 [Definition 1.2.5: Complete Metric Space] 5
2.2 Subsequences & Compactness 5
2.2.1 [Theorem 1.2.9: Bolzano-Weierstrass] 5

1 1. Convergence of Sequences (Metric Spaces)

1.1 Metric Spaces (The Distance Interface)

1.1.1 [Definition 1.1.1: Metric]

1. Layer 1: The Logic & Spec View (QA Engineer)
   - **Concept:** A "Metric" is an Interface specification for measuring "cost" or "distance" between two objects.
   - **Input:** Two objects x, y from a set Ω.
   - **Output:** A non-negative float representing distance.
   - **Invariants (Unit Tests):**
     (a) **Identity:** Distance is 0 if and only if objects are identical ($x = y$).
     (b) **Symmetry:** Direction doesn't matter ($dist(x, y) = dist(y, x)$).
     (c) **Triangle Inequality:** The direct path is always the optimal (or equal) cost compared to a detour through z.
   - **Edge Cases:** If Ω is empty, the definition is vacuously true.
2. Layer 2: The Feynman View (Intuition)
   - Think of a **Ruler** or a **GPS**.
   - **Triangle Inequality ($d(x, z) \leq d(x, y) + d(y, z)$):** This is the core physical intuition. If walking from Home to Work ($x \to z$) is 5km, walking Home to Coffee Shop to Work ($x \to y \to z$) cannot be **shorter** than 5km. It can only be longer or equal.
   - Without a valid metric, we have no concept of "near" or "far," which means we cannot define limits.
3. Layer 3: Formal Source of Truth Given a set $\Omega \neq \emptyset$, a mapping $d_\Omega : \Omega \times \Omega \to [0, \infty)$ is a metric if:
   (a) **Symmetry:** $\forall x, y \in \Omega, d_\Omega(x, y) = d_\Omega(y, x)$
   (b) **Triangular Inequality:** $\forall x, y, z \in \Omega$,
   $$d_\Omega(x, z) \leq d_\Omega(x, y) + d_\Omega(y, z)$$
   (c) **Definiteness:** $\forall x, y \in \Omega$,
   $$d_\Omega(x, y) = 0 \iff x = y$$

1.2 Convergence of Sequences

1.2.1 [Definition 1.1.3: Convergence]

1. Layer 1: The Logic & Spec View (QA Engineer)

   - **Goal:** To validate that a stream of data (a_n) stabilizes at a final value a.
   - **The Challenger (User):** Provides an arbitrary error tolerance $\epsilon > 0$ (e.g., "I need precision to 10^{-6}").
   - **The Responder (System):** Must find a timestamp N (index).
   - **The Guarantee:** For **all** times $n \geq N$, the error $|a_n - a|$ is strictly less than ϵ.
   - **Control Flow:**
     - Loop: 'For all' $\epsilon \ldots$
     - Action: 'Find' $N \ldots$
     - Check: 'Verify' sequence stays inside bounds after N.

2. Layer 2: The Feynman View (Intuition)

   - **The "Sniper" Analogy:**
     - The limit a is the bullseye.
     - The sequence terms are shots fired.
     - To be "convergent" means no matter how tiny the target ring (ϵ) is drawn, eventually (N) the shooter hits inside that ring **every single time** thereafter.
   - **Key distinction:** It doesn't matter if the first million shots are wild. It only matters what happens closer to infinity (the "tail").

3. Layer 3: Formal Source of Truth Let (Ω, d_Ω) be a metric space. A sequence $(a_n)_{n\in\mathbb{N}} \subseteq \Omega$ converges to $a \in \Omega$ if:

   $$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ such that } \forall n \geq N : d_\Omega(a_n, a) \leq \epsilon$$

   Notation: $\lim_{n\to\infty} a_n = a$ or $a_n \to a$.

1.2.2 [Theorem 1.1.5: Uniqueness of Limit]

1. Layer 1: The Logic & Spec View (QA Engineer)

   - **Goal:** Verify system determinism.

- **Assertion:** A sequence cannot settle in two different states simultaneously.
- **Proof Logic (Contradiction):**
  - Assume it converges to both a and a'.
  - Set tolerance ϵ to be less than half the distance between a and a'.
  - The sequence would have to be in two disjoint bubbles at the same time. Impossible.

2. Layer 2: The Feynman View (Intuition)
   - **"You can't be in two places at once."**
   - If you are "very close" to New York and "very close" to London at the same time, New York and London must be the same place.
3. Layer 3: Formal Source of Truth If $a_n \to a$ and $a_n \to a'$, then $a = a'$.
   Proof Argument:

$$d_\Omega(a, a') \leq d_\Omega(a, a_N) + d_\Omega(a_N, a') \leq 2\epsilon$$

   Since ϵ is arbitrary, $d_\Omega(a, a') = 0 \implies a = a'$.

2 1.2 Completeness

2.1 Cauchy Sequences & Completeness

2.1.1 [Definition 1.2.1: Cauchy Sequence]

1. Layer 1: The Logic & Spec View (QA Engineer)
   - **Goal:** Check for stability **without knowing the answer**.
   - **Concept:** "Internal Consistency" test.
   - **The Test:** Do the elements get closer to **each other** as time goes on?
   - **Difference from Convergence:** Convergence checks $distance(a_n, \text{Target})$. Cauchy checks $distance(a_n, a_m)$.
2. Layer 2: The Feynman View (Intuition)
   - **The "Huddle":** The sequence members are crowding together. They stop moving apart.

- **The Tragedy of the Rationals:** A sequence can huddle together perfectly (be Cauchy) but have nowhere to go.
  - **Example:** The sequence $1, 1.4, 1.41, 1.414\ldots$ (approximating $\sqrt{2}$).
  - In $\mathbb{Q}$ (fractions), this sequence is trying to land on a "hole". It is Cauchy, but does not converge **in** $\mathbb{Q}$.

3. Layer 3: Formal Source of Truth A sequence (a_n) is a Cauchy sequence if:

$$\forall \epsilon > 0, \exists N \in \mathbb{N} \text{ such that } \forall n, m \geq N : d_\Omega(a_n, a_m) \leq \epsilon$$

2.1.2 [Definition 1.2.5: Complete Metric Space]

1. Layer 1: The Logic & Spec View (QA Engineer)
   - **Goal:** System Integrity Certification.
   - **Definition:** A space is "Complete" if it has no "holes".
   - **Rule:** 'Strict Mode'. Every sequence that passes the "Internal Consistency Check" (Cauchy) must successfully return a valid limit object (Convergent).
2. Layer 2: The Feynman View (Intuition)
   - **Filling the Holes:** $\mathbb{R}$ (Real numbers) is just $\mathbb{Q}$ (Rationals) with all the holes filled in.
   - If you walk towards a cliff edge in a Complete world, the ground effectively generates underneath your feet so you land on something.
3. Layer 3: Formal Source of Truth A metric space Ω is called **complete** if every Cauchy sequence in Ω is convergent in Ω.
   - **Axiom:** $\mathbb{R}$ is complete.
   - **Lemma:** $\mathbb{Q}$ is NOT complete.

2.2 Subsequences & Compactness

2.2.1 [Theorem 1.2.9: Bolzano-Weierstrass]

1. Layer 1: The Logic & Spec View (QA Engineer)
   - **Goal:** Handle infinite data in finite memory.

- **Input:** A sequence that is **Bounded** (stuck in a box).
- **Output:** A subsequence that **Converges**.
- **Logic:** If you generate infinite points but limit the range of values (bounds), you are forced to revisit the same neighborhood infinitely often.

2. Layer 2: The Feynman View (Intuition)

   - **Infinite Pigeonhole Principle:**
     - Imagine an infinite number of pigeons flying into a finite grid of cages.
     - At least one specific area must get crowded with infinite pigeons.
     - If we just track those specific pigeons, they converge to a specific spot.

3. Layer 3: Formal Source of Truth If $(a_n)_{n\in\mathbb{N}} \subseteq \mathbb{R}$ is a **bounded** sequence, then it contains a **convergent subsequence** $(a_{n_j})_{j\in\mathbb{N}}$.