

Correlated Equilibria & Learning

Omar Boufous

Supervised by

Rachid El-Azouzi², Mikael Touati¹, Eitan Altman^{2,3}, Mustapha Bouhtou¹

¹Orange, Châtillon, France

²CERI/LIA, Université d'Avignon, Avignon, France

³INRIA Sophia Antipolis, France

December 20, 2024

PhD Thesis Defense

Presentation Outline

1. Background & Preliminaries

- 1.1 Introduction
- 1.2 Related Work
- 1.3 Notations & Model

2. Learning Correlated Equilibria

- 2.1 Online Learning with Regret
- 2.2 Algorithm Description
- 2.3 Stochastic Model
- 2.4 Simulation Results
- 2.5 Time-varying Game
- 2.6 Application to a Congestion Game

3. Constrained Correlated Equilibria

- 3.1 Definition & Example
- 3.2 Properties of Constrained Correlated Equilibrium Strategies
- 3.3 Canonical Representation and Existence of Constrained Correlated Equilibria
- 3.4 Computation of Constrained Correlated Equilibria
- 3.5 Simulation Results

4. Conclusions & Possible Directions of Research

Presentation Outline

1. Background & Preliminaries

- 1.1 Introduction
- 1.2 Related Work
- 1.3 Notations & Model

2. Learning Correlated Equilibria

- 2.1 Online Learning with Regret
- 2.2 Algorithm Description
- 2.3 Stochastic Model
- 2.4 Simulation Results
- 2.5 Time-varying Game
- 2.6 Application to a Congestion Game

3. Constrained Correlated Equilibria

- 3.1 Definition & Example
- 3.2 Properties of Constrained Correlated Equilibrium Strategies
- 3.3 Canonical Representation and Existence of Constrained Correlated Equilibria
- 3.4 Computation of Constrained Correlated Equilibria
- 3.5 Simulation Results

4. Conclusions & Possible Directions of Research

Presentation Outline

1. Background & Preliminaries

- 1.1 Introduction
- 1.2 Related Work
- 1.3 Notations & Model

2. Learning Correlated Equilibria

- 2.1 Online Learning with Regret
- 2.2 Algorithm Description
- 2.3 Stochastic Model
- 2.4 Simulation Results
- 2.5 Time-varying Game
- 2.6 Application to a Congestion Game

3. Constrained Correlated Equilibria

- 3.1 Definition & Example
- 3.2 Properties of Constrained Correlated Equilibrium Strategies
- 3.3 Canonical Representation and Existence of Constrained Correlated Equilibria
- 3.4 Computation of Constrained Correlated Equilibria
- 3.5 Simulation Results

4. Conclusions & Possible Directions of Research

Presentation Outline

1. Background & Preliminaries

- 1.1 Introduction
- 1.2 Related Work
- 1.3 Notations & Model

2. Learning Correlated Equilibria

- 2.1 Online Learning with Regret
- 2.2 Algorithm Description
- 2.3 Stochastic Model
- 2.4 Simulation Results
- 2.5 Time-varying Game
- 2.6 Application to a Congestion Game

3. Constrained Correlated Equilibria

- 3.1 Definition & Example
- 3.2 Properties of Constrained Correlated Equilibrium Strategies
- 3.3 Canonical Representation and Existence of Constrained Correlated Equilibria
- 3.4 Computation of Constrained Correlated Equilibria
- 3.5 Simulation Results

4. Conclusions & Possible Directions of Research

Presentation Outline

1. Background & Preliminaries

- 1.1 Introduction
- 1.2 Related Work
- 1.3 Notations & Model

2. Learning Correlated Equilibria

- 2.1 Online Learning with Regret
- 2.2 Algorithm Description
- 2.3 Stochastic Model
- 2.4 Simulation Results
- 2.5 Time-varying Game
- 2.6 Application to a Congestion Game

3. Constrained Correlated Equilibria

- 3.1 Definition & Example
- 3.2 Properties of Constrained Correlated Equilibrium Strategies
- 3.3 Canonical Representation and Existence of Constrained Correlated Equilibria
- 3.4 Computation of Constrained Correlated Equilibria
- 3.5 Simulation Results

4. Conclusions & Possible Directions of Research

Presentation Outline

1. Background & Preliminaries

- 1.1 Introduction
- 1.2 Related Work
- 1.3 Notations & Model

2. Learning Correlated Equilibria

- 2.1 Online Learning with Regret
- 2.2 Algorithm Description
- 2.3 Stochastic Model
- 2.4 Simulation Results
- 2.5 Time-varying Game
- 2.6 Application to a Congestion Game

3. Constrained Correlated Equilibria

- 3.1 Definition & Example
- 3.2 Properties of Constrained Correlated Equilibrium Strategies
- 3.3 Canonical Representation and Existence of Constrained Correlated Equilibria
- 3.4 Computation of Constrained Correlated Equilibria
- 3.5 Simulation Results

4. Conclusions & Possible Directions of Research

Presentation Outline

1. Background & Preliminaries

- 1.1 Introduction
- 1.2 Related Work
- 1.3 Notations & Model

2. Learning Correlated Equilibria

- 2.1 Online Learning with Regret
- 2.2 Algorithm Description
- 2.3 Stochastic Model
- 2.4 Simulation Results
- 2.5 Time-varying Game
- 2.6 Application to a Congestion Game

3. Constrained Correlated Equilibria

- 3.1 Definition & Example
- 3.2 Properties of Constrained Correlated Equilibrium Strategies
- 3.3 Canonical Representation and Existence of Constrained Correlated Equilibria
- 3.4 Computation of Constrained Correlated Equilibria
- 3.5 Simulation Results

4. Conclusions & Possible Directions of Research

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
- ▶ Simple **implementation** using a correlation device
- ▶ Many **applications** in engineering, economics, etc.

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
- ▶ Simple **implementation** using a correlation device
- ▶ Many **applications** in engineering, economics, etc.



Figure 1: Robert John Aumann

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
- ▶ Simple **implementation** using a correlation device
- ▶ Many **applications** in engineering, economics, etc.



Figure 1: Robert John Aumann

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
- ▶ Simple **implementation** using a correlation device
- ▶ Many **applications** in engineering, economics, etc.



Figure 1: Robert John Aumann

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
 - ▶ Simple **implementation** using a correlation device
 - ▶ Many **applications** in engineering, economics, etc.



Figure 1: Robert John Aumann

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
- ▶ Simple **implementation** using a correlation device
- ▶ Many **applications** in engineering, economics, etc.



Figure 1: Robert John Aumann

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
- ▶ Simple **implementation** using a correlation device
- ▶ Many **applications** in engineering, economics, etc.



Figure 1: Robert John Aumann

Introduction

- ▶ A **generalization** of Nash equilibria with strong connections to Bayesian rationality introduced by Aumann
- ▶ A solution concept with appealing **computational properties**
- ▶ Basic **learning dynamics** naturally lead to correlated equilibria
- ▶ **Canonical** form characterizing the set of correlated equilibrium distributions
- ▶ Simple **implementation** using a correlation device
- ▶ Many **applications** in engineering, economics, etc.



Figure 1: Robert John Aumann

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Related Work

Correlated Equilibria

- ▶ Defined in (Aumann, 1974) and (Aumann, 1987)
- ▶ A proof of existence and a generalization to infinite games in (Hart & Schmeidler, 1989)
- ▶ Extensive form correlated equilibrium defined in (Von Stengel & Forges, 2008)
- ▶ Other extensions in (Forges, 2020), (Brandenburger & Dekel, 1992) and (Grant & Stauber, 2022)

Learning Correlated Equilibria

- ▶ Regret-matching (Hart & Mas-Colell, 2000) implying the convergence to the set of correlated equilibria
- ▶ Correlated Q-learning for selection of specific correlated equilibrium points (Greenwald & Hall, 2003)
- ▶ Convergence to the social welfare maximizing correlated equilibrium (Borowski et al., 2015)
- ▶ Learning algorithms for specific applications e.g., resource allocation game (Cigler & Faltings, 2011)

Constraints & Correlated Equilibria

- ▶ (Debreu, 1952) defines the concept of generalized equilibrium
- ▶ (Arrow & Debreu, 1954) shows a proof of existence
- ▶ (Rosen, 1965) shows existence and uniqueness of equilibria in concave games with coupled constraints
- ▶ (Bernasconi et al., 23) introduces the concept of constrained Phi-equilibrium

Many Other Theoretical and Applied Contributions

- ▶ Applications in energy systems and smart grids (Kulkarni, 2017), resource allocation problems (Pang et al., 2008) (Nowak et al., 2018)

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's set of strategies is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the set of strategy profiles is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's **set of strategies** is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the **set of strategy profiles** is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's **set of strategies** is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the **set of strategy profiles** is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's **set of strategies** is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the **set of strategy profiles** is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Model & Definitions

Finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$

- ▶ Set of players \mathcal{N}
- ▶ Action set \mathcal{A}_i for each $i \in \mathcal{N}$
- ▶ Utility function $u_i : \prod_{i \in \mathcal{N}} \mathcal{A}_i = \mathcal{A} \rightarrow \mathbb{R}$

Correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$

- ▶ A sample space Ω
- ▶ Partition \mathcal{P}_i of Ω for each $i \in \mathcal{N}$
- ▶ Probability distribution q over Ω

Player i 's **set of strategies** is $\mathcal{S}_{i,d} = \{f_i : \Omega \rightarrow \mathcal{A}_i \text{ s.t. } f_i \text{ is } \mathcal{P}_i\text{-measurable}\}$ and the **set of strategy profiles** is $\mathcal{S}_d = \mathcal{S}_{1,d} \times \dots \times \mathcal{S}_{n,d}$.

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (1)$$

A correlated equilibrium distribution of G (Forges, 2020) is a probability distribution $p \in \Delta(\mathcal{A})$ such that,

$$\forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \quad \sum_{a \in \mathcal{A}} p(a) u_i(a_i, a_{-i}) \geq \sum_{a \in \mathcal{A}} p(a) u_i(\beta_i(a_i), a_{-i}) \quad (2)$$

Part 1 : Learning Correlated Equilibria

Online Learning with Regret

The game G is played repeatedly through time $t = 1, 2, 3, \dots$. Each step t , action profile $\mathbf{a}^t = (a_1^t, \dots, a_n^t)$ is played

Definition 3 – Regret (Hart & Mas-Colell, 2000)

The regret player i experiences at time t for any pair of actions $j, k \in \mathcal{A}_i$ is

$$R_i^t(j, k) = \max \left(0, \frac{1}{t} \sum_{\tau \leq t: a_i^\tau = j} [u_i(k, \mathbf{a}_{-i}^\tau) - u_i(\mathbf{a}^\tau)] \right) = \max \left(0, D_i^t(j, k) \right) \quad (3)$$

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i: k \neq j} p_i^{t+1}(k). \end{cases}$$

Proposition 1 (Hart & Mas-Colell, 2000)

Let $(\mathbf{a}^t)_{t=1,2,\dots}$ be a sequence of plays (i.e., $\mathbf{a}^t \in \mathcal{A}$ for all t) and let $\varepsilon \geq 0$. Then: $\limsup_{t \rightarrow \infty} R_i^t(j, k) \leq \varepsilon$ for

every $i \in \mathcal{N}$ and every $j, k \in \mathcal{A}_i$ with $j \neq k$, if and only if the sequence of empirical distributions q^t converges to the set of correlated ε -equilibria.

Online Learning with Regret

The game G is played repeatedly through time $t = 1, 2, 3, \dots$. Each step t , action profile $\mathbf{a}^t = (a_1^t, \dots, a_n^t)$ is played

Definition 3 – Regret (Hart & Mas-Colell, 2000)

The regret player i experiences at time t for any pair of actions $j, k \in \mathcal{A}_i$ is

$$R_i^t(j, k) = \max \left(0, \frac{1}{t} \sum_{\tau \leq t: a_i^\tau = j} [u_i(k, \mathbf{a}_{-i}^\tau) - u_i(\mathbf{a}^\tau)] \right) = \max \left(0, D_i^t(j, k) \right) \quad (3)$$

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i: k \neq j} p_i^{t+1}(k). \end{cases}$$

Proposition 1 (Hart & Mas-Colell, 2000)

Let $(\mathbf{a}^t)_{t=1,2,\dots}$ be a sequence of plays (i.e., $\mathbf{a}^t \in \mathcal{A}$ for all t) and let $\varepsilon \geq 0$. Then: $\limsup_{t \rightarrow \infty} R_i^t(j, k) \leq \varepsilon$ for every $i \in \mathcal{N}$ and every $j, k \in \mathcal{A}_i$ with $j \neq k$, if and only if the sequence of empirical distributions q^t converges to the set of correlated ε -equilibria.

Online Learning with Regret

The game G is played repeatedly through time $t = 1, 2, 3, \dots$. Each step t , action profile $\mathbf{a}^t = (a_1^t, \dots, a_n^t)$ is played

Definition 3 – Regret (Hart & Mas-Colell, 2000)

The regret player i experiences at time t for any pair of actions $j, k \in \mathcal{A}_i$ is

$$R_i^t(j, k) = \max \left(0, \frac{1}{t} \sum_{\tau \leq t: \mathbf{a}_i^\tau = j} [u_i(k, \mathbf{a}_{-i}^\tau) - u_i(\mathbf{a}^\tau)] \right) = \max \left(0, D_i^t(j, k) \right) \quad (3)$$

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i: k \neq j} p_i^{t+1}(k). \end{cases}$$

Proposition 1 (Hart & Mas-Colell, 2000)

Let $(\mathbf{a}^t)_{t=1,2,\dots}$ be a sequence of plays (i.e., $\mathbf{a}^t \in \mathcal{A}$ for all t) and let $\varepsilon \geq 0$. Then: $\limsup_{t \rightarrow \infty} R_i^t(j, k) \leq \varepsilon$ for every $i \in \mathcal{N}$ and every $j, k \in \mathcal{A}_i$ with $j \neq k$, if and only if the sequence of empirical distributions q^t converges to the set of correlated ε -equilibria.

Online Learning with Regret

The game G is played repeatedly through time $t = 1, 2, 3, \dots$. Each step t , action profile $\mathbf{a}^t = (a_1^t, \dots, a_n^t)$ is played

Definition 3 – Regret (Hart & Mas-Colell, 2000)

The regret player i experiences at time t for any pair of actions $j, k \in \mathcal{A}_i$ is

$$R_i^t(j, k) = \max \left(0, \frac{1}{t} \sum_{\tau \leq t: \mathbf{a}_i^\tau = j} [u_i(k, \mathbf{a}_{-i}^\tau) - u_i(\mathbf{a}^\tau)] \right) = \max \left(0, D_i^t(j, k) \right) \quad (3)$$

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i: k \neq j} p_i^{t+1}(k). \end{cases}$$

Proposition 1 (Hart & Mas-Colell, 2000)

Let $(\mathbf{a}^t)_{t=1,2,\dots}$ be a sequence of plays (i.e., $\mathbf{a}^t \in \mathcal{A}$ for all t) and let $\varepsilon \geq 0$. Then: $\limsup_{t \rightarrow \infty} R_i^t(j, k) \leq \varepsilon$ for every $i \in \mathcal{N}$ and every $j, k \in \mathcal{A}_i$ with $j \neq k$, if and only if the sequence of empirical distributions q^t converges to the set of correlated ε -equilibria.

Online Learning with Regret

The game G is played repeatedly through time $t = 1, 2, 3, \dots$. Each step t , action profile $\mathbf{a}^t = (a_1^t, \dots, a_n^t)$ is played

Definition 3 – Regret (Hart & Mas-Colell, 2000)

The regret player i experiences at time t for any pair of actions $j, k \in \mathcal{A}_i$ is

$$R_i^t(j, k) = \max \left(0, \frac{1}{t} \sum_{\tau \leq t: \mathbf{a}_i^\tau = j} [u_i(k, \mathbf{a}_{-i}^\tau) - u_i(\mathbf{a}^\tau)] \right) = \max \left(0, D_i^t(j, k) \right) \quad (3)$$

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i: k \neq j} p_i^{t+1}(k). \end{cases}$$

Proposition 1 (Hart & Mas-Colell, 2000)

Let $(\mathbf{a}^t)_{t=1,2,\dots}$ be a sequence of plays (i.e., $\mathbf{a}^t \in \mathcal{A}$ for all t) and let $\varepsilon \geq 0$. Then: $\limsup_{t \rightarrow \infty} R_i^t(j, k) \leq \varepsilon$ for every $i \in \mathcal{N}$ and every $j, k \in \mathcal{A}_i$ with $j \neq k$, if and only if the sequence of empirical distributions q^t converges to the set of correlated ε -equilibria.

Online Learning with Regret

The game G is played repeatedly through time $t = 1, 2, 3, \dots$. Each step t , action profile $\mathbf{a}^t = (a_1^t, \dots, a_n^t)$ is played

Definition 3 – Regret (Hart & Mas-Colell, 2000)

The regret player i experiences at time t for any pair of actions $j, k \in \mathcal{A}_i$ is

$$R_i^t(j, k) = \max \left(0, \frac{1}{t} \sum_{\tau \leq t: a_i^\tau = j} [u_i(k, \mathbf{a}_{-i}^\tau) - u_i(\mathbf{a}^\tau)] \right) = \max \left(0, D_i^t(j, k) \right) \quad (3)$$

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i: k \neq j} p_i^{t+1}(k). \end{cases}$$

Proposition 1 (Hart & Mas-Colell, 2000)

Let $(\mathbf{a}^t)_{t=1,2,\dots}$ be a sequence of plays (i.e., $\mathbf{a}^t \in \mathcal{A}$ for all t) and let $\varepsilon \geq 0$. Then: $\limsup_{t \rightarrow \infty} R_i^t(j, k) \leq \varepsilon$ for

every $i \in \mathcal{N}$ and every $j, k \in \mathcal{A}_i$ with $j \neq k$, if and only if the sequence of empirical distributions q^t converges to the set of correlated ε -equilibria.

Online Learning with Regret

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i: k \neq j} p_i^{t+1}(k). \end{cases}$$

Regret-Matching algorithm applied to a 3-by-2 game

	D	E
A	(2, 29)	(16, 7)
B	(4, 7)	(6, 13)
C	(4, 4)	(6, 6)

Table 1: Utility matrix.

- ▶ The empirical distribution does not seem to converge towards a correlated equilibrium point
- ▶ We propose a new dynamic called Correlated Perturbed Regret Minimization (CPRM)

Online Learning with Regret

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i : k \neq j} p_i^{t+1}(k). \end{cases}$$

Regret-Matching algorithm applied to a 3-by-2 game

	D	E
A	(2, 29)	(16, 7)
B	(4, 7)	(6, 13)
C	(4, 4)	(6, 6)

Table 1: Utility matrix.

- ▶ The empirical distribution does not seem to converge towards a correlated equilibrium point
- ▶ We propose a new dynamic called Correlated Perturbed Regret Minimization (CPRM)

Online Learning with Regret

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i : k \neq j} p_i^{t+1}(k). \end{cases}$$

Regret-Matching algorithm applied to a 3-by-2 game

	D	E
A	(2, 29)	(16, 7)
B	(4, 7)	(6, 13)
C	(4, 4)	(6, 6)

Table 1: Utility matrix.




Figure 2: Evolution of the empirical probability distribution.

- ▶ The empirical distribution does not seem to converge towards a correlated equilibrium point
- ▶ We propose a new dynamic called Correlated Perturbed Regret Minimization (CPRM)

Online Learning with Regret

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i : k \neq j} p_i^{t+1}(k). \end{cases}$$

Regret-Matching algorithm applied to a 3-by-2 game

	D	E
A	(2, 29)	(16, 7)
B	(4, 7)	(6, 13)
C	(4, 4)	(6, 6)

Table 1: Utility matrix.




Figure 2: Evolution of the empirical probability distribution.

- ▶ The empirical distribution does not seem to converge towards a correlated equilibrium point
- ▶ We propose a new dynamic called Correlated Perturbed Regret Minimization (CPRM)

Online Learning with Regret

Regret-Matching algorithm (Hart & Mas-Colell, 2000)

$$\begin{cases} p_i^{t+1}(k) = \frac{1}{\mu} R_i^t(j, k), \\ p_i^{t+1}(j) = 1 - \sum_{k \in \mathcal{A}_i : k \neq j} p_i^{t+1}(k). \end{cases}$$

Regret-Matching algorithm applied to a 3-by-2 game

	D	E
A	(2, 29)	(16, 7)
B	(4, 7)	(6, 13)
C	(4, 4)	(6, 6)

Table 1: Utility matrix.




Figure 2: Evolution of the empirical probability distribution.

- ▶ The empirical distribution does not seem to converge towards a correlated equilibrium point
- ▶ We propose a new dynamic called Correlated Perturbed Regret Minimization (CPRM)

CPRM Algorithm

Inspired from the learning algorithm introduced in (Young, H. P., 2009) for learning pure Nash equilibria.

At time t , each player i is characterized by a mood $m_i^t \in \{\text{syn}, \text{asyn}\}$



*asynchronous
player*



*synchronous
player*

CPRM Algorithm

Inspired from the learning algorithm introduced in (Young, H. P., 2009) for learning pure Nash equilibria.

At time t , each player i is characterized by a mood $m_i^t \in \{\text{syn}, \text{asyn}\}$



*asynchronous
player*



*synchronous
player*

CPRM Algorithm

Inspired from the learning algorithm introduced in (Young, H. P., 2009) for learning pure Nash equilibria.

At time t , each player i is characterized by a mood $m_i^t \in \{\text{syn}, \text{asyn}\}$



*asynchronous
player*



*synchronous
player*

CPRM Algorithm

Inspired from the learning algorithm introduced in (Young, H. P., 2009) for learning pure Nash equilibria.

At time t , each player i is characterized by a mood $m_i^t \in \{\text{syn}, \text{asyn}\}$



*asynchronous
player*




*synchronous
player*

CPRM Algorithm

$$\mathbf{a} = (a_1, a_2, a_3, \dots, a_n)$$

drawn from \mathbf{q}^t




Players implementing a
regret-minimizing strategy

Players play the actions drawn
from the empirical distribution

CPRM Algorithm

$$\mathbf{a} = (a_1, a_2, a_3, \dots, a_n)$$

drawn from \mathbf{q}^t




In the long run

↪ We expect players to play the action profiles drawn from the current empirical distributions to stabilize the play at an equilibrium point.

CPRM Algorithm

$$\mathbf{a} = (a_1, a_2, a_3, \dots, a_n)$$

drawn from \mathbf{q}^t



In the long run

↪ We expect players to play the action profiles drawn from the current empirical distributions to stabilize the play at an equilibrium point.

Algorithm Analysis

Mood dynamics




Characteristics of the stochastic process

- ▶ Transition probabilities depend on the regrets and a perturbation parameter $\varepsilon > 0$
- ▶ The stochastic process $X^t = (X_1^t, \dots, X_n^t)$ describing players' states is a perturbed non-homogeneous Markov chain
- ▶ The Markov chain is defined over a infinitely countable state spaces
- ▶ For a regularly perturbed homogenous Markov chain, the process spends almost all time in the stochastically stable states when $\varepsilon \rightarrow 0$

Algorithm Analysis

Mood dynamics



Characteristics of the stochastic process

- ▶ Transition probabilities depend on the regrets and a perturbation parameter $\varepsilon > 0$
- ▶ The stochastic process $X^t = (X_1^t, \dots, X_n^t)$ describing players' states is a perturbed non-homogeneous Markov chain
- ▶ The Markov chain is defined over a infinitely countable state spaces
- ▶ For a regularly perturbed homogenous Markov chain, the process spends almost all time in the stochastically stable states when $\varepsilon \rightarrow 0$

Algorithm Analysis

Mood dynamics



Characteristics of the stochastic process

- ▶ Transition probabilities depend on the regrets and a perturbation parameter $\varepsilon > 0$
- ▶ The stochastic process $X^t = (X_1^t, \dots, X_n^t)$ describing players' states is a perturbed non-homogeneous Markov chain
- ▶ The Markov chain is defined over a infinitely countable state spaces
- ▶ For a regularly perturbed homogenous Markov chain, the process spends almost all time in the stochastically stable states when $\varepsilon \rightarrow 0$

Algorithm Analysis

Mood dynamics



Characteristics of the stochastic process

- ▶ Transition probabilities depend on the regrets and a perturbation parameter $\varepsilon > 0$
- ▶ The stochastic process $\mathbf{X}^t = (X_1^t, \dots, X_n^t)$ describing players' states is a perturbed non-homogeneous Markov chain
- ▶ The Markov chain is defined over a infinitely countable state spaces
- ▶ For a regularly perturbed homogenous Markov chain, the process spends almost all time in the stochastically stable states when $\varepsilon \rightarrow 0$

Algorithm Analysis

Mood dynamics



Characteristics of the stochastic process

- ▶ Transition probabilities depend on the regrets and a perturbation parameter $\varepsilon > 0$
- ▶ The stochastic process $\mathbf{X}^t = (X_1^t, \dots, X_n^t)$ describing players' states is a perturbed non-homogeneous Markov chain
- ▶ The Markov chain is defined over a infinitely countable state spaces
- ▶ For a regularly perturbed homogenous Markov chain, the process spends almost all time in the stochastically stable states when $\varepsilon \rightarrow 0$

Algorithm Analysis

Mood dynamics



Characteristics of the stochastic process

- ▶ Transition probabilities depend on the regrets and a perturbation parameter $\varepsilon > 0$
- ▶ The stochastic process $\mathbf{X}^t = (X_1^t, \dots, X_n^t)$ describing players' states is a perturbed non-homogeneous Markov chain
- ▶ The Markov chain is defined over a infinitely countable state spaces
- ▶ For a regularly perturbed homogenous Markov chain, the process spends almost all time in the stochastically stable states when $\varepsilon \rightarrow 0$

Simulation Results

	D	E
A	(2, 29)	(16, 7)
B	(4, 7)	(6, 13)
C	(4, 4)	(6, 6)

Table 2: Utility matrix.

Simulation Results






Figure 3: Evolution of the empirical distribution over action profiles.


Simulation Results



(a) CPRM algorithm.



(b) Regret minimizing strategy.



(c) Regret-matching.

Figure 4: Evolution of players' regrets for the three algorithms.

Time-varying Game

		D	E			D	E
A	(2, 29)	(16, 7)		A	(9, 4, 0)	(4, 1, 4)	
B	(4, 7)	(6, 13)		B	(8, 0, 1)	(6, 7, 2)	
C	(4, 4)	(6, 6)		C	(11, 9, 3)	(2, 0, 4)	

↓

		D	E			D	E
A	(2, 29, 2)	(16, 7, 8)		A	(9, 4, 0)	(4, 1, 4)	
B	(4, 7, 2)	(6, 13, 0)		B	(8, 0, 1)	(6, 7, 2)	
C	(4, 4, 1)	(6, 6, 5)		C	(11, 9, 3)	(2, 0, 4)	

X Y

↓

		D	E			D	E
A	(2, 29)	(16, 7)		A	(9, 4, 0)	(4, 1, 4)	
B	(4, 7)	(6, 13)		B	(8, 0, 1)	(6, 7, 2)	
C	(4, 4)	(6, 6)		C	(11, 9, 3)	(2, 0, 4)	

↓

		D	E			D	E
A	(2, 29, 2)	(16, 7, 8)		A	(9, 4, 0)	(4, 1, 4)	
B	(4, 7, 2)	(6, 13, 0)		B	(8, 0, 1)	(6, 7, 2)	
C	(4, 4, 1)	(6, 6, 5)		C	(11, 9, 3)	(2, 0, 4)	

X Y

Figure 5: Sequence of stage games in the dynamic case.




Figure 6: Evolution of maximal regrets




Figure 7: Evolution of the empirical distribution

Application to a Congestion Game




Figure 8: Network graph

Player	Source node	Destination node
1	B	F
2	B	E
3	D	B
4	F	E

Figure 9: Source-destination nodes for each player




Figure 10: Evolution of the empirical distribution




Figure 11: Evolution of maximal regrets

Part 2 : Constrained Correlated Equilibria

Constrained Correlated Equilibrium Strategies

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (4)$$

	P	A
P	8, 8	3, 10
A	10, 3	0, 0

Table 3: Utility matrix of the game of Chicken.

- ▶ $\Omega = \{H, M, L\}$
- ▶ $\mathcal{P}_1 = \{\{H\}, \{M, L\}\}$ and $\mathcal{P}_2 = \{\{H, M\}, \{L\}\}$
- ▶ $q(H) = q(M) = q(L) = 1/3$

Constrained Correlated Equilibrium Strategies

Definition 1 – Correlated equilibrium (Aumann, 1974)

A correlated equilibrium of G is a pair (d, α^*) where $\alpha^* : \Omega \rightarrow \mathcal{A}$ is a correlated strategy profile such that

$$\forall i \in \mathcal{N}, \forall \alpha'_i \in \mathcal{S}_{i,d} \quad \sum_{\omega \in \Omega} q(\omega) u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) \geq \sum_{\omega \in \Omega} q(\omega) u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \quad (4)$$

	P	A
P	8, 8	3, 10
A	10, 3	0, 0

Table 3: Utility matrix of the game of Chicken.

- ▶ $\Omega = \{H, M, L\}$
- ▶ $\mathcal{P}_1 = \{\{H\}, \{M, L\}\}$ and $\mathcal{P}_2 = \{\{H, M\}, \{L\}\}$
- ▶ $q(H) = q(M) = q(L) = 1/3$

Constrained Correlated Equilibrium Strategies

	P	A
P	8, 8	3, 10
A	10, 3	0, 0

Table 4: Utility matrix of the game of Chicken.

- ▶ $\Omega = \{H, M, L\}$
- ▶ $\mathcal{P}_1 = \{\{H\}, \{M, L\}\}$ and $\mathcal{P}_2 = \{\{H, M\}, \{L\}\}$
- ▶ $q(H) = q(M) = q(L) = 1/3$

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Figure 12: Extension of the game of Chicken.

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Table 2: Constrained extension of the game of Chicken.

Constrained Correlated Equilibrium Strategies

	P	A
P	8, 8	3, 10
A	10, 3	0, 0

Table 4: Utility matrix of the game of Chicken.

- ▶ $\Omega = \{H, M, L\}$
- ▶ $\mathcal{P}_1 = \{\{H\}, \{M, L\}\}$ and $\mathcal{P}_2 = \{\{H, M\}, \{L\}\}$
- ▶ $q(H) = q(M) = q(L) = 1/3$

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Figure 12: Extension of the game of Chicken.

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Table 2: Constrained extension of the game of Chicken.

Constrained Correlated Equilibrium Strategies

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Figure 13: Extension of the game of Chicken.

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Figure 14: Constrained extension of the game of Chicken.

Definition 4 – Constrained Correlated equilibrium (Boufous et al., 2024)

A constrained correlated equilibrium of G is a triplet $(d, \mathcal{R}_d, \alpha^*)$ where d is a correlation device, \mathcal{R}_d is a coupled constraint set in G_d and α^* a correlated strategy profile such that, $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any $\alpha'_i \in \mathcal{S}_{i,d}$ such that $(\alpha'_i, \alpha^*_{-i}) \in \mathcal{R}_d$,

$$\sum_{\omega \in \Omega} q(\omega) [u_i(\alpha_i^*(\omega), \alpha^*_{-i}(\omega)) - u_i(\alpha'_i(\omega), \alpha^*_{-i}(\omega))] \geq 0 \quad (5)$$

Constrained Correlated Equilibrium Strategies

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Figure 13: Extension of the game of Chicken.

	$L \mapsto P$ $L^c \mapsto P$	$L \mapsto A$ $L^c \mapsto A$	$L \mapsto A$ $L^c \mapsto P$	$L \mapsto P$ $L^c \mapsto A$
$H \mapsto P$ $H^c \mapsto P$	8, 8	3, 10	6.33, 8.67	4.67, 9.33
$H \mapsto A$ $H^c \mapsto A$	10, 3	0, 0	6.67, 2	3.33, 1
$H \mapsto A$ $H^c \mapsto P$	8.67, 6.33	2, 6.67	7, 7	3.67, 6
$H \mapsto P$ $H^c \mapsto A$	9.33, 4.67	1, 3.33	6, 3.67	4.33, 4.33

Figure 14: Constrained extension of the game of Chicken.

Definition 4 – Constrained Correlated equilibrium (Boufous et al., 2024)

A constrained correlated equilibrium of G is a triplet $(d, \mathcal{R}_d, \alpha^*)$ where d is a correlation device, \mathcal{R}_d is a coupled constraint set in G_d and α^* a correlated strategy profile such that, $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any $\alpha'_i \in \mathcal{S}_{i,d}$ such that $(\alpha'_i, \alpha^*_{-i}) \in \mathcal{R}_d$,

$$\sum_{\omega \in \Omega} q(\omega) [u_i(\alpha^*_i(\omega), \alpha^*_{-i}(\omega)) - u_i(\alpha'_i(\omega), \alpha^*_{-i}(\omega))] \geq 0 \quad (5)$$

Properties of Constrained Correlated Equilibrium Strategies

Consider a finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$, a correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$ and a constraint set $\mathcal{R}_d \subset \mathcal{S}_d$.

Proposition 1 : If (d, α^*) is a correlated equilibrium and $\alpha^* \in \mathcal{R}_d$, then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 2 : If $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any α'_i s.t. $(\alpha'_i, \alpha_{-i}^*) \in \mathcal{R}_d$, for any $\omega \in \Omega$

$$\sum_{\omega' \in P_i(\omega)} q(\omega') [u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega'), \alpha_{-i}^*(\omega'))] \geq 0 \quad (6)$$

then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 3 : The triplet $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium if and only if $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any $\alpha'_i \in \mathcal{S}_{i,d}$,

$$\sum_{\omega \in \Omega} q(\omega) [u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega))] \geq 0 \text{ or } (\alpha'_i, \alpha_{-i}^*) \notin \mathcal{R}_d \quad (7)$$

Properties of Constrained Correlated Equilibrium Strategies

Consider a finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$, a correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$ and a constraint set $\mathcal{R}_d \subset \mathcal{S}_d$.

Proposition 1 : If (d, α^*) is a correlated equilibrium and $\alpha^* \in \mathcal{R}_d$, then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 2 : If $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any α'_i s.t. $(\alpha'_i, \alpha_{-i}^*) \in \mathcal{R}_d$, for any $\omega \in \Omega$

$$\sum_{\omega' \in P_i(\omega)} q(\omega') [u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega'), \alpha_{-i}^*(\omega'))] \geq 0 \quad (6)$$

then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 3 : The triplet $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium if and only if $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any $\alpha'_i \in \mathcal{S}_{i,d}$,

$$\sum_{\omega \in \Omega} q(\omega) [u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega))] \geq 0 \text{ or } (\alpha'_i, \alpha_{-i}^*) \notin \mathcal{R}_d \quad (7)$$

Properties of Constrained Correlated Equilibrium Strategies

Consider a finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$, a correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$ and a constraint set $\mathcal{R}_d \subset \mathcal{S}_d$.

Proposition 1 : If (d, α^*) is a correlated equilibrium and $\alpha^* \in \mathcal{R}_d$, then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 2 : If $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any α'_i s.t. $(\alpha'_i, \alpha_{-i}^*) \in \mathcal{R}_d$, for any $\omega \in \Omega$

$$\sum_{\omega' \in P_i(\omega)} q(\omega') [u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega'), \alpha_{-i}^*(\omega'))] \geq 0 \quad (6)$$

then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 3 : The triplet $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium if and only if $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any $\alpha'_i \in \mathcal{S}_{i,d}$,

$$\sum_{\omega \in \Omega} q(\omega) [u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega))] \geq 0 \text{ or } (\alpha'_i, \alpha_{-i}^*) \notin \mathcal{R}_d \quad (7)$$

Properties of Constrained Correlated Equilibrium Strategies

Consider a finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$, a correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$ and a constraint set $\mathcal{R}_d \subset \mathcal{S}_d$.

Proposition 1 : If (d, α^*) is a correlated equilibrium and $\alpha^* \in \mathcal{R}_d$, then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 2 : If $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any α'_i s.t. $(\alpha'_i, \alpha_{-i}^*) \in \mathcal{R}_d$, for any $\omega \in \Omega$

$$\sum_{\omega' \in P_i(\omega)} q(\omega') \left[u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega'), \alpha_{-i}^*(\omega')) \right] \geq 0 \quad (6)$$

then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 3 : The triplet $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium if and only if $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any $\alpha'_i \in \mathcal{S}_{i,d}$,

$$\sum_{\omega \in \Omega} q(\omega) \left[u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \right] \geq 0 \quad \text{or} \quad (\alpha'_i, \alpha_{-i}^*) \notin \mathcal{R}_d \quad (7)$$

Properties of Constrained Correlated Equilibrium Strategies

Consider a finite non-cooperative game $G = (\mathcal{N}, (\mathcal{A}_i)_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$, a correlation device $d = (\Omega, (\mathcal{P}_i)_{i \in \mathcal{N}}, q)$ and a constraint set $\mathcal{R}_d \subset \mathcal{S}_d$.

Proposition 1 : If (d, α^*) is a correlated equilibrium and $\alpha^* \in \mathcal{R}_d$, then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 2 : If $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any α'_i s.t. $(\alpha'_i, \alpha_{-i}^*) \in \mathcal{R}_d$, for any $\omega \in \Omega$

$$\sum_{\omega' \in P_i(\omega)} q(\omega') \left[u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega'), \alpha_{-i}^*(\omega')) \right] \geq 0 \quad (6)$$

then $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium.

Proposition 3 : The triplet $(d, \mathcal{R}_d, \alpha^*)$ is a constrained correlated equilibrium if and only if $\alpha^* \in \mathcal{R}_d$ and for any $i \in \mathcal{N}$, for any $\alpha'_i \in \mathcal{S}_{i,d}$,

$$\sum_{\omega \in \Omega} q(\omega) \left[u_i(\alpha_i^*(\omega), \alpha_{-i}^*(\omega)) - u_i(\alpha'_i(\omega), \alpha_{-i}^*(\omega)) \right] \geq 0 \quad \text{or} \quad (\alpha'_i, \alpha_{-i}^*) \notin \mathcal{R}_d \quad (7)$$

Constraints induced by a set of feasible probability distributions

- ▶ Let $\mathcal{C} \subseteq \Delta(\mathcal{A})$ be a set of probability distributions. For each correlation device d ,

$$\mathcal{R}_d = \{\alpha \in \mathcal{S}_d \mid p_\alpha \in \mathcal{C}\} \quad (8)$$

- ▶ Performance measures in many applications can be expressed in terms of probability distribution over action profiles. Examples include applications in smart grids, wireless networks, etc.

- ▶ Constraints on the social welfare :
$$\sum_{i \in \mathcal{N}} \sum_{a \in \mathcal{A}} p(a) u_i(a) \geq D_1$$
- ▶ Constraints on the Nash product :
$$\prod_{i \in \mathcal{N}} \left(\sum_{a \in \mathcal{A}} p(a) u_i(a) \right) \geq D_2$$

Theorem 1 – Characterization of the Set of Constrained Correlated Equilibrium Distributions

Let G be a finite non-cooperative game and \mathcal{C} a set of feasible probability distributions. The distribution $p \in \Delta(\mathcal{A})$ is a constrained correlated equilibrium distribution if and only if for any player $i \in \mathcal{N}$, for any strategy $\beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i$, if $z_{\beta_i, p} \in \mathcal{C}$, then

$$\sum_{a \in \mathcal{A}} p(a) [u_i(a_i, a_{-i}) - u_i(\beta_i(a_i), a_{-i})] \geq 0 \quad (9)$$

where $z_{\beta_i, p}(a) = \sum_{b_i \in \mathcal{A}_i} p(b_i, a_{-i}) \mathbb{1}_{\beta_i(b_i) = a_i}$ for any $a \in \mathcal{A}$ is the distribution resulting from player i 's unilateral deviation β_i .

Constraints induced by a set of feasible probability distributions

- ▶ Let $\mathcal{C} \subseteq \Delta(\mathcal{A})$ be a set of probability distributions. For each correlation device d ,

$$\mathcal{R}_d = \{\alpha \in \mathcal{S}_d \mid p_\alpha \in \mathcal{C}\} \quad (8)$$

- ▶ Performance measures in many applications can be expressed in terms of probability distribution over action profiles. Examples include applications in smart grids, wireless networks, etc.

- ▶ Constraints on the social welfare : $\sum_{i \in \mathcal{N}} \sum_{a \in \mathcal{A}} p(a) u_i(a) \geq D_1$
- ▶ Constraints on the Nash product : $\prod_{i \in \mathcal{N}} \left(\sum_{a \in \mathcal{A}} p(a) u_i(a) \right) \geq D_2$

Theorem 1 – Characterization of the Set of Constrained Correlated Equilibrium Distributions

Let G be a finite non-cooperative game and \mathcal{C} a set of feasible probability distributions. The distribution $p \in \Delta(\mathcal{A})$ is a constrained correlated equilibrium distribution if and only if for any player $i \in \mathcal{N}$, for any strategy $\beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i$, if $z_{\beta_i, p} \in \mathcal{C}$, then

$$\sum_{a \in \mathcal{A}} p(a) [u_i(a_i, a_{-i}) - u_i(\beta_i(a_i), a_{-i})] \geq 0 \quad (9)$$

where $z_{\beta_i, p}(a) = \sum_{b_i \in \mathcal{A}_i} p(b_i, a_{-i}) \mathbb{1}_{\beta_i(b_i) = a_i}$ for any $a \in \mathcal{A}$ is the distribution resulting from player i 's unilateral deviation β_i .

Constraints induced by a set of feasible probability distributions

- ▶ Let $\mathcal{C} \subseteq \Delta(\mathcal{A})$ be a set of probability distributions. For each correlation device d ,

$$\mathcal{R}_d = \{\alpha \in \mathcal{S}_d \mid p_\alpha \in \mathcal{C}\} \quad (8)$$

- ▶ Performance measures in many applications can be expressed in terms of probability distribution over action profiles. Examples include applications in smart grids, wireless networks, etc.

- ▶ Constraints on the social welfare : $\sum_{i \in \mathcal{N}} \sum_{a \in \mathcal{A}} p(a) u_i(a) \geq D_1$
- ▶ Constraints on the Nash product : $\prod_{i \in \mathcal{N}} \left(\sum_{a \in \mathcal{A}} p(a) u_i(a) \right) \geq D_2$

Theorem 1 – Characterization of the Set of Constrained Correlated Equilibrium Distributions

Let G be a finite non-cooperative game and \mathcal{C} a set of feasible probability distributions. The distribution $p \in \Delta(\mathcal{A})$ is a constrained correlated equilibrium distribution if and only if for any player $i \in \mathcal{N}$, for any strategy $\beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i$, if $z_{\beta_i, p} \in \mathcal{C}$, then

$$\sum_{a \in \mathcal{A}} p(a) [u_i(a_i, a_{-i}) - u_i(\beta_i(a_i), a_{-i})] \geq 0 \quad (9)$$

where $z_{\beta_i, p}(a) = \sum_{b_i \in \mathcal{A}_i} p(b_i, a_{-i}) \mathbb{1}_{\beta_i(b_i) = a_i}$ for any $a \in \mathcal{A}$ is the distribution resulting from player i 's unilateral deviation β_i .

Constraints induced by a set of feasible probability distributions

- ▶ Let $\mathcal{C} \subseteq \Delta(\mathcal{A})$ be a set of probability distributions. For each correlation device d ,

$$\mathcal{R}_d = \{\alpha \in \mathcal{S}_d \mid p_\alpha \in \mathcal{C}\} \quad (8)$$

- ▶ Performance measures in many applications can be expressed in terms of probability distribution over action profiles. Examples include applications in smart grids, wireless networks, etc.

- ▶ Constraints on the social welfare :
$$\sum_{i \in \mathcal{N}} \sum_{a \in \mathcal{A}} p(a) u_i(a) \geq D_1$$
- ▶ Constraints on the Nash product :
$$\prod_{i \in \mathcal{N}} \left(\sum_{a \in \mathcal{A}} p(a) u_i(a) \right) \geq D_2$$

Theorem 1 – Characterization of the Set of Constrained Correlated Equilibrium Distributions

Let G be a finite non-cooperative game and \mathcal{C} a set of feasible probability distributions. The distribution $p \in \Delta(\mathcal{A})$ is a constrained correlated equilibrium distribution if and only if for any player $i \in \mathcal{N}$, for any strategy $\beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i$, if $z_{\beta_i, p} \in \mathcal{C}$, then

$$\sum_{a \in \mathcal{A}} p(a) [u_i(a_i, a_{-i}) - u_i(\beta_i(a_i), a_{-i})] \geq 0 \quad (9)$$

where $z_{\beta_i, p}(a) = \sum_{b_i \in \mathcal{A}_i} p(b_i, a_{-i}) \mathbb{1}_{\beta_i(b_i) = a_i}$ for any $a \in \mathcal{A}$ is the distribution resulting from player i 's unilateral deviation β_i .

Constraints induced by a set of feasible probability distributions

- ▶ Let $\mathcal{C} \subseteq \Delta(\mathcal{A})$ be a set of probability distributions. For each correlation device d ,

$$\mathcal{R}_d = \{\alpha \in \mathcal{S}_d \mid p_\alpha \in \mathcal{C}\} \quad (8)$$

- ▶ Performance measures in many applications can be expressed in terms of probability distribution over action profiles. Examples include applications in smart grids, wireless networks, etc.

- ▶ Constraints on the social welfare :
$$\sum_{i \in \mathcal{N}} \sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) u_i(\mathbf{a}) \geq D_1$$
- ▶ Constraints on the Nash product :
$$\prod_{i \in \mathcal{N}} \left(\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) u_i(\mathbf{a}) \right) \geq D_2$$

Theorem 1 – Characterization of the Set of Constrained Correlated Equilibrium Distributions

Let G be a finite non-cooperative game and \mathcal{C} a set of feasible probability distributions. The distribution $p \in \Delta(\mathcal{A})$ is a constrained correlated equilibrium distribution if and only if for any player $i \in \mathcal{N}$, for any strategy $\beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i$, if $z_{\beta_i, p} \in \mathcal{C}$, then

$$\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) [u_i(a_i, \mathbf{a}_{-i}) - u_i(\beta_i(a_i), \mathbf{a}_{-i})] \geq 0 \quad (9)$$

where $z_{\beta_i, p}(\mathbf{a}) = \sum_{b_i \in \mathcal{A}_i} p(b_i, \mathbf{a}_{-i}) \mathbb{1}_{\beta_i(b_i) = a_i}$ for any $\mathbf{a} \in \mathcal{A}$ is the distribution resulting from player i 's unilateral deviation β_i .

Constraints induced by a set of feasible probability distributions

- ▶ Let $\mathcal{C} \subseteq \Delta(\mathcal{A})$ be a set of probability distributions. For each correlation device d ,

$$\mathcal{R}_d = \{\alpha \in \mathcal{S}_d \mid p_\alpha \in \mathcal{C}\} \quad (8)$$

- ▶ Performance measures in many applications can be expressed in terms of probability distribution over action profiles. Examples include applications in smart grids, wireless networks, etc.

- ▶ Constraints on the social welfare :
$$\sum_{i \in \mathcal{N}} \sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) u_i(\mathbf{a}) \geq D_1$$
- ▶ Constraints on the Nash product :
$$\prod_{i \in \mathcal{N}} \left(\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) u_i(\mathbf{a}) \right) \geq D_2$$

Theorem 1 – Characterization of the Set of Constrained Correlated Equilibrium Distributions

Let G be a finite non-cooperative game and \mathcal{C} a set of feasible probability distributions. The distribution $p \in \Delta(\mathcal{A})$ is a constrained correlated equilibrium distribution if and only if for any player $i \in \mathcal{N}$, for any strategy $\beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i$, if $z_{\beta_i, p} \in \mathcal{C}$, then

$$\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) [u_i(a_i, \mathbf{a}_{-i}) - u_i(\beta_i(a_i), \mathbf{a}_{-i})] \geq 0 \quad (9)$$

where $z_{\beta_i, p}(\mathbf{a}) = \sum_{b_i \in \mathcal{A}_i} p(b_i, \mathbf{a}_{-i}) \mathbb{1}_{\beta_i(b_i) = a_i}$ for any $\mathbf{a} \in \mathcal{A}$ is the distribution resulting from player i 's unilateral deviation β_i .

Constraints induced by a set of feasible probability distributions

- ▶ Let $\mathcal{C} \subseteq \Delta(\mathcal{A})$ be a set of probability distributions. For each correlation device d ,

$$\mathcal{R}_d = \{\alpha \in \mathcal{S}_d \mid p_\alpha \in \mathcal{C}\} \quad (8)$$

- ▶ Performance measures in many applications can be expressed in terms of probability distribution over action profiles. Examples include applications in smart grids, wireless networks, etc.

- ▶ Constraints on the social welfare :
$$\sum_{i \in \mathcal{N}} \sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) u_i(\mathbf{a}) \geq D_1$$
- ▶ Constraints on the Nash product :
$$\prod_{i \in \mathcal{N}} \left(\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) u_i(\mathbf{a}) \right) \geq D_2$$

Theorem 1 – Characterization of the Set of Constrained Correlated Equilibrium Distributions

Let G be a finite non-cooperative game and \mathcal{C} a set of feasible probability distributions. The distribution $p \in \Delta(\mathcal{A})$ is a constrained correlated equilibrium distribution if and only if for any player $i \in \mathcal{N}$, for any strategy $\beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i$, if $z_{\beta_i, p} \in \mathcal{C}$, then

$$\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) [u_i(a_i, \mathbf{a}_{-i}) - u_i(\beta_i(a_i), \mathbf{a}_{-i})] \geq 0 \quad (9)$$

where $z_{\beta_i, p}(\mathbf{a}) = \sum_{b_i \in \mathcal{A}_i} p(b_i, \mathbf{a}_{-i}) \mathbb{1}_{\beta_i(b_i) = a_i}$ for any $\mathbf{a} \in \mathcal{A}$ is the distribution resulting from player i 's unilateral deviation β_i .

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Existence of Constrained Correlated Equilibria

Consider the following two-player game:

	L	R
U	$(2, 2)$	$(1, 1)$
D	$(3, 0)$	$(0, 5)$

Figure 13: Two-player game in matrix form.

Let $\mathcal{C} \subset \Delta(\mathcal{A})$ be a feasible set of probability distributions such that,

$$\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid p(U, L) = 1 \text{ or } p(U, R) = 1 \text{ or } p(D, L) = 1 \text{ or } p(D, R) = 1\} \quad (10)$$

- ▶ The set of feasible strategies is the set of pure action profiles
- ▶ The players must play a correlated strategy profile inducing a pure action profile in G
- ▶ There does not exist a constrained correlated equilibrium for this game

Theorem 2 - Existence of Constrained Correlated Equilibria

Let G be a finite non-cooperative game and \mathcal{C} a feasible set of probability distributions. If the intersection of \mathcal{C} with the set of correlated equilibrium distributions of the game G is non-empty, then a constrained correlated equilibrium of G exists.

Constrained Correlated Equilibria of the Mixed Extension

Define the mixed extension of G by the game $\Delta G = (\mathcal{N}, (\Delta(\mathcal{A}_i))_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$ where $\Delta(\mathcal{A}_i) = \{p \in \mathbb{R}_+^{|\mathcal{A}_i|} \mid \sum_{a_i \in \mathcal{A}_i} p(a_i) = 1\}$ is the set of probability distributions on \mathcal{A}_i .

Let d be a correlation device and ΔG_d be the extension of ΔG by d . The set of strategies of player i in ΔG_d is

$$\tilde{\mathcal{S}}_{i,d} = \{\gamma_i : \Omega \rightarrow \Delta(\mathcal{A}_i) \mid \gamma_i \text{ is } \mathcal{P}_i\text{-measurable}\} \quad (11)$$

For any correlated strategy profile $\gamma \in \tilde{\mathcal{S}}_d = \times_{i \in \mathcal{N}} \tilde{\mathcal{S}}_{i,d}$, the utility function of player i is $\tilde{u}_i : \tilde{\mathcal{S}}_d \rightarrow \mathbb{R}$ such that,

$$\tilde{u}_i(\gamma_i, \gamma_{-i}) = \sum_{\omega \in \Omega} q(\omega) u_i(\gamma_i(\omega), \alpha_{-i}(\omega)) \quad (12)$$

Theorem 2 : Let G be a finite non-cooperative game, \mathcal{C} a feasible set of probability distributions, d a correlation device and $\gamma^* \in \tilde{\mathcal{S}}_d$ a correlated strategy profile. If $(d, \tilde{\mathcal{R}}_d, \gamma^*)$ is a constrained correlated equilibrium of ΔG , then it exists a constrained correlated equilibrium $(d', \mathcal{R}_{d'}, \alpha^*)$ of G such that $p_{\gamma^*} = p_{\alpha^*}$.

→ The set of constrained correlated equilibrium distributions of the game ΔG is included in the set of constrained correlated equilibrium distributions of G .

Constrained Correlated Equilibria of the Mixed Extension

Define the mixed extension of G by the game $\Delta G = (\mathcal{N}, (\Delta(\mathcal{A}_i))_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$ where $\Delta(\mathcal{A}_i) = \{p \in \mathbb{R}_+^{|\mathcal{A}_i|} \mid \sum_{a_i \in \mathcal{A}_i} p(a_i) = 1\}$ is the set of probability distributions on \mathcal{A}_i .

Let d be a correlation device and ΔG_d be the extension of ΔG by d . The set of strategies of player i in ΔG_d is

$$\tilde{\mathcal{S}}_{i,d} = \{\gamma_i : \Omega \rightarrow \Delta(\mathcal{A}_i) \mid \gamma_i \text{ is } \mathcal{P}_i\text{-measurable}\} \quad (11)$$

For any correlated strategy profile $\gamma \in \tilde{\mathcal{S}}_d = \times_{i \in \mathcal{N}} \tilde{\mathcal{S}}_{i,d}$, the utility function of player i is $\tilde{u}_i : \tilde{\mathcal{S}}_d \rightarrow \mathbb{R}$ such that,

$$\tilde{u}_i(\gamma_i, \gamma_{-i}) = \sum_{\omega \in \Omega} q(\omega) u_i(\gamma_i(\omega), \alpha_{-i}(\omega)) \quad (12)$$

Theorem 2 : Let G be a finite non-cooperative game, \mathcal{C} a feasible set of probability distributions, d a correlation device and $\gamma^* \in \tilde{\mathcal{S}}_d$ a correlated strategy profile. If $(d, \tilde{\mathcal{R}}_d, \gamma^*)$ is a constrained correlated equilibrium of ΔG , then it exists a constrained correlated equilibrium $(d', \mathcal{R}_{d'}, \alpha^*)$ of G such that $p_{\gamma^*} = p_{\alpha^*}$.

→ The set of constrained correlated equilibrium distributions of the game ΔG is included in the set of constrained correlated equilibrium distributions of G .

Constrained Correlated Equilibria of the Mixed Extension

Define the mixed extension of G by the game $\Delta G = (\mathcal{N}, (\Delta(\mathcal{A}_i))_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$ where $\Delta(\mathcal{A}_i) = \{p \in \mathbb{R}_+^{|\mathcal{A}_i|} \mid \sum_{a_i \in \mathcal{A}_i} p(a_i) = 1\}$ is the set of probability distributions on \mathcal{A}_i .

Let d be a correlation device and ΔG_d be the extension of ΔG by d . The set of strategies of player i in ΔG_d is

$$\tilde{\mathcal{S}}_{i,d} = \{\gamma_i : \Omega \rightarrow \Delta(\mathcal{A}_i) \mid \gamma_i \text{ is } \mathcal{P}_i\text{-measurable}\} \quad (11)$$

For any correlated strategy profile $\gamma \in \tilde{\mathcal{S}}_d = \times_{i \in \mathcal{N}} \tilde{\mathcal{S}}_{i,d}$, the utility function of player i is $\tilde{u}_i : \tilde{\mathcal{S}}_d \rightarrow \mathbb{R}$ such that,

$$\tilde{u}_i(\gamma_i, \gamma_{-i}) = \sum_{\omega \in \Omega} q(\omega) u_i(\gamma_i(\omega), \alpha_{-i}(\omega)) \quad (12)$$

Theorem 2 : Let G be a finite non-cooperative game, \mathcal{C} a feasible set of probability distributions, d a correlation device and $\gamma^* \in \tilde{\mathcal{S}}_d$ a correlated strategy profile. If $(d, \tilde{\mathcal{R}}_d, \gamma^*)$ is a constrained correlated equilibrium of ΔG , then it exists a constrained correlated equilibrium $(d', \mathcal{R}_{d'}, \alpha^*)$ of G such that $p_{\gamma^*} = p_{\alpha^*}$.

→ The set of constrained correlated equilibrium distributions of the game ΔG is included in the set of constrained correlated equilibrium distributions of G .

Constrained Correlated Equilibria of the Mixed Extension

Define the mixed extension of G by the game $\Delta G = (\mathcal{N}, (\Delta(\mathcal{A}_i))_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$ where $\Delta(\mathcal{A}_i) = \{p \in \mathbb{R}_+^{|\mathcal{A}_i|} \mid \sum_{a_i \in \mathcal{A}_i} p(a_i) = 1\}$ is the set of probability distributions on \mathcal{A}_i .

Let d be a correlation device and ΔG_d be the extension of ΔG by d . The set of strategies of player i in ΔG_d is

$$\tilde{\mathcal{S}}_{i,d} = \{\gamma_i : \Omega \rightarrow \Delta(\mathcal{A}_i) \mid \gamma_i \text{ is } \mathcal{P}_i\text{-measurable}\} \quad (11)$$

For any correlated strategy profile $\gamma \in \tilde{\mathcal{S}}_d = \times_{i \in \mathcal{N}} \tilde{\mathcal{S}}_{i,d}$, the utility function of player i is $\tilde{u}_i : \tilde{\mathcal{S}}_d \rightarrow \mathbb{R}$ such that,

$$\tilde{u}_i(\gamma_i, \gamma_{-i}) = \sum_{\omega \in \Omega} q(\omega) u_i(\gamma_i(\omega), \alpha_{-i}(\omega)) \quad (12)$$

Theorem 2 : Let G be a finite non-cooperative game, \mathcal{C} a feasible set of probability distributions, d a correlation device and $\gamma^* \in \tilde{\mathcal{S}}_d$ a correlated strategy profile. If $(d, \tilde{\mathcal{R}}_d, \gamma^*)$ is a constrained correlated equilibrium of ΔG , then it exists a constrained correlated equilibrium $(d', \mathcal{R}_{d'}, \alpha^*)$ of G such that $p_{\gamma^*} = p_{\alpha^*}$.

→ The set of constrained correlated equilibrium distributions of the game ΔG is included in the set of constrained correlated equilibrium distributions of G .

Constrained Correlated Equilibria of the Mixed Extension

Define the mixed extension of G by the game $\Delta G = (\mathcal{N}, (\Delta(\mathcal{A}_i))_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$ where $\Delta(\mathcal{A}_i) = \{p \in \mathbb{R}_+^{|\mathcal{A}_i|} \mid \sum_{a_i \in \mathcal{A}_i} p(a_i) = 1\}$ is the set of probability distributions on \mathcal{A}_i .

Let d be a correlation device and ΔG_d be the extension of ΔG by d . The set of strategies of player i in ΔG_d is

$$\tilde{\mathcal{S}}_{i,d} = \{\gamma_i : \Omega \rightarrow \Delta(\mathcal{A}_i) \mid \gamma_i \text{ is } \mathcal{P}_i\text{-measurable}\} \quad (11)$$

For any correlated strategy profile $\gamma \in \tilde{\mathcal{S}}_d = \times_{i \in \mathcal{N}} \tilde{\mathcal{S}}_{i,d}$, the utility function of player i is $\tilde{u}_i : \tilde{\mathcal{S}}_d \rightarrow \mathbb{R}$ such that,

$$\tilde{u}_i(\gamma_i, \gamma_{-i}) = \sum_{\omega \in \Omega} q(\omega) u_i(\gamma_i(\omega), \alpha_{-i}(\omega)) \quad (12)$$

Theorem 2 : Let G be a finite non-cooperative game, \mathcal{C} a feasible set of probability distributions, d a correlation device and $\gamma^* \in \tilde{\mathcal{S}}_d$ a correlated strategy profile. If $(d, \tilde{\mathcal{R}}_d, \gamma^*)$ is a constrained correlated equilibrium of ΔG , then it exists a constrained correlated equilibrium $(d', \mathcal{R}_{d'}, \alpha^*)$ of G such that $p_{\gamma^*} = p_{\alpha^*}$.

→ The set of constrained correlated equilibrium distributions of the game ΔG is included in the set of constrained correlated equilibrium distributions of G .

Constrained Correlated Equilibria of the Mixed Extension

Define the mixed extension of G by the game $\Delta G = (\mathcal{N}, (\Delta(\mathcal{A}_i))_{i \in \mathcal{N}}, (u_i)_{i \in \mathcal{N}})$ where

$\Delta(\mathcal{A}_i) = \{p \in \mathbb{R}_+^{|\mathcal{A}_i|} \mid \sum_{a_i \in \mathcal{A}_i} p(a_i) = 1\}$ is the set of probability distributions on \mathcal{A}_i .

Let d be a correlation device and ΔG_d be the extension of ΔG by d . The set of strategies of player i in ΔG_d is

$$\tilde{\mathcal{S}}_{i,d} = \{\gamma_i : \Omega \rightarrow \Delta(\mathcal{A}_i) \mid \gamma_i \text{ is } \mathcal{P}_i\text{-measurable}\} \quad (11)$$

For any correlated strategy profile $\gamma \in \tilde{\mathcal{S}}_d = \times_{i \in \mathcal{N}} \tilde{\mathcal{S}}_{i,d}$, the utility function of player i is $\tilde{u}_i : \tilde{\mathcal{S}}_d \rightarrow \mathbb{R}$ such that,

$$\tilde{u}_i(\gamma_i, \gamma_{-i}) = \sum_{\omega \in \Omega} q(\omega) u_i(\gamma_i(\omega), \alpha_{-i}(\omega)) \quad (12)$$

Theorem 2 : Let G be a finite non-cooperative game, \mathcal{C} a feasible set of probability distributions, d a correlation device and $\gamma^* \in \tilde{\mathcal{S}}_d$ a correlated strategy profile. If $(d, \tilde{\mathcal{R}}_d, \gamma^*)$ is a constrained correlated equilibrium of ΔG , then it exists a constrained correlated equilibrium $(d', \mathcal{R}_{d'}, \alpha^*)$ of G such that $p_{\gamma^*} = p_{\alpha^*}$.

→ The set of constrained correlated equilibrium distributions of the game ΔG is included in the set of constrained correlated equilibrium distributions of G .

Computation of Constrained Correlated Equilibria

$$\begin{aligned} & \text{maximize } 0 \\ \text{s.t. } & p(a) \geq 0 \quad \forall a \in \mathcal{A}, \quad \sum_{a \in \mathcal{A}} p(a) = 1 \quad (13) \\ & \forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \end{aligned}$$

$$\sum_{a \in \mathcal{A}} p(a) [u_i(a) - u_i(\beta_i(a_i), a_{-i})] \geq 0. \quad (14)$$

$$\begin{aligned} & \text{maximize } 0 \\ \text{s.t. } & p(a) \geq 0 \quad \forall a \in \mathcal{A}, \quad \sum_{a \in \mathcal{A}} p(a) = 1 \quad (15) \\ & \forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \text{ s.t. } z_{\beta_i, p} \in \mathcal{C} \end{aligned}$$

$$\sum_{a \in \mathcal{A}} p(a) [u_i(a) - u_i(\beta_i(a_i), a_{-i})] \geq 0, \quad (16)$$

$$p \in \mathcal{C} \quad (17)$$

Assuming linear constraints e.g., $\mathcal{C} = \{p \in \Delta(\mathcal{A}) \mid Fp \leq 0\}$, we show that the problem of computing a constrained correlated equilibrium can be formulated as a Mixed-Integer Linear Program.

Computation of Constrained Correlated Equilibria

$$\begin{aligned} & \text{maximize } 0 \\ \text{s.t. } & p(\mathbf{a}) \geq 0 \quad \forall \mathbf{a} \in \mathcal{A}, \quad \sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) = 1 \quad (13) \\ & \forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \end{aligned}$$

$$\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) [u_i(\mathbf{a}) - u_i(\beta_i(a_i), \mathbf{a}_{-i})] \geq 0. \quad (14)$$

$$\begin{aligned} & \text{maximize } 0 \\ \text{s.t. } & p(\mathbf{a}) \geq 0 \quad \forall \mathbf{a} \in \mathcal{A}, \quad \sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) = 1 \quad (15) \\ & \forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \text{ s.t. } \mathbf{z}_{\beta_i, \mathbf{p}} \in \mathcal{C} \end{aligned}$$

$$\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) [u_i(\mathbf{a}) - u_i(\beta_i(a_i), \mathbf{a}_{-i})] \geq 0, \quad (16)$$

$$\mathbf{p} \in \mathcal{C} \quad (17)$$

Assuming linear constraints e.g., $\mathcal{C} = \{\mathbf{p} \in \Delta(\mathcal{A}) \mid F\mathbf{p} \leq 0\}$, we show that the problem of computing a constrained correlated equilibrium can be formulated as a Mixed-Integer Linear Program.

Computation of Constrained Correlated Equilibria

$$\begin{aligned} & \text{maximize } 0 \\ \text{s.t. } & p(\mathbf{a}) \geq 0 \quad \forall \mathbf{a} \in \mathcal{A}, \quad \sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) = 1 \quad (13) \\ & \forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \end{aligned}$$

$$\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) [u_i(\mathbf{a}) - u_i(\beta_i(a_i), \mathbf{a}_{-i})] \geq 0. \quad (14)$$

$$\begin{aligned} & \text{maximize } 0 \\ \text{s.t. } & p(\mathbf{a}) \geq 0 \quad \forall \mathbf{a} \in \mathcal{A}, \quad \sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) = 1 \quad (15) \\ & \forall i \in \mathcal{N}, \forall \beta_i : \mathcal{A}_i \rightarrow \mathcal{A}_i \text{ s.t. } \mathbf{z}_{\beta_i, \mathbf{p}} \in \mathcal{C} \end{aligned}$$

$$\sum_{\mathbf{a} \in \mathcal{A}} p(\mathbf{a}) [u_i(\mathbf{a}) - u_i(\beta_i(a_i), \mathbf{a}_{-i})] \geq 0, \quad (16)$$

$$\mathbf{p} \in \mathcal{C} \quad (17)$$

Assuming linear constraints e.g., $\mathcal{C} = \{\mathbf{p} \in \Delta(\mathcal{A}) \mid F\mathbf{p} \leq 0\}$, we show that the problem of computing a constrained correlated equilibrium distribution **can be formulated as a Mixed-Integer Linear Program**.

Simulation Example

	P	A
P	8, 8	3, 10
A	10, 3	0, 0

Table 3: Game of Chicken

$$SW(q) = \sum_{i \in \mathcal{N}} u_i(q) \quad (18)$$

where $u_i(q) = \sum_{a \in \mathcal{A}} q(a) u_i(a)$

- ▶ Feasible utilities in green
- ▶ CE utilities in yellow
- ▶ CCE utilities in red




Figure 14: Sets of Utilities of the two players with $SW \geq 12$

Simulation Results




Figure 15: Sets of Utilities of the two players with $SW \geq 13$




Figure 16: Sets of Utilities of the two players with $SW \geq 14$




Figure 17: Sets of Utilities of the two players with $SW \geq 15$

- ▶ The set may not be convex
- ▶ There are constrained correlated equilibria outside the set of correlated equilibrium distributions

Simulation Results




Figure 15: Sets of Utilities of the two players with $SW \geq 13$




Figure 16: Sets of Utilities of the two players with $SW \geq 14$




Figure 17: Sets of Utilities of the two players with $SW \geq 15$

- ▶ The set may not be convex
- ▶ There are constrained correlated equilibria outside the set of correlated equilibrium distributions

Simulation Results




Figure 15: Sets of Utilities of the two players with $SW \geq 13$




Figure 16: Sets of Utilities of the two players with $SW \geq 14$




Figure 17: Sets of Utilities of the two players with $SW \geq 15$

- ▶ The set may not be convex
- ▶ There are constrained correlated equilibria outside the set of correlated equilibrium distributions

Conclusion & Highlight of Contributions

▶ A learning rule to converge towards a correlated equilibrium distribution

- ▶ Empirical evidence of convergence of the distribution to a constrained correlated equilibrium
- ▶ A Markov chain models the learning algorithm
- ▶ The algorithm is adaptive to changes in games e.g., payoff functions or number of players

▶ Arbitrary constraints and arbitrary correlation device

▶ Characterizations of Equilibrium Strategies

- ▶ Definition through generalized Nash equilibrium and correlated equilibrium with alternative characterizations

▶ Properties and relationship to (unconstrained) Correlated Equilibria

- ▶ A feasible correlated equilibrium is a constrained correlated equilibrium
- ▶ The set of constrained correlated equilibrium distributions may not be convex
- ▶ There exist constrained correlated equilibrium distributions outside the set of correlated equilibrium distributions

▶ Constraints on probability distributions

▶ Characterization of constrained correlated equilibrium distributions

- ▶ Closed-form expression of the set of constrained correlated equilibrium distribution

▶ Conditions of Existence

- ▶ There does not always exist a constrained correlated equilibrium in finite games

▶ Constrained correlated equilibrium distributions of the mixed extension of a game

- ▶ Every constrained correlated equilibrium distribution of ΔG is a constrained correlated equilibrium distribution of G

▶ Computation of linearly constrained correlated equilibria

Conclusion & Highlight of Contributions

▶ A learning rule to converge towards a correlated equilibrium distribution

- ▶ Empirical evidence of convergence of the distribution to a constrained correlated equilibrium
- ▶ A Markov chain models the learning algorithm
- ▶ The algorithm is adaptive to changes in games e.g., payoff functions or number of players

▶ Arbitrary constraints and arbitrary correlation device

▶ **Characterizations of Equilibrium Strategies**

- ▶ Definition through generalized Nash equilibrium and correlated equilibrium with alternative characterizations

▶ **Properties and relationship to (unconstrained) Correlated Equilibria**

- ▶ A feasible correlated equilibrium is a constrained correlated equilibrium
- ▶ The set of constrained correlated equilibrium distributions may not be convex
- ▶ There exist constrained correlated equilibrium distributions outside the set of correlated equilibrium distributions

▶ Constraints on probability distributions

▶ **Characterization of constrained correlated equilibrium distributions**

- ▶ Closed-form expression of the set of constrained correlated equilibrium distribution

▶ **Conditions of Existence**

- ▶ There does not always exist a constrained correlated equilibrium in finite games

▶ **Constrained correlated equilibrium distributions of the mixed extension of a game**

- ▶ Every constrained correlated equilibrium distribution of ΔG is a constrained correlated equilibrium distribution of G

▶ **Computation of linearly constrained correlated equilibria**

Conclusion & Highlight of Contributions

- ▶ **A learning rule to converge towards a correlated equilibrium distribution**
 - ▶ Empirical evidence of convergence of the distribution to a constrained correlated equilibrium
 - ▶ A Markov chain models the learning algorithm
 - ▶ The algorithm is adaptive to changes in games e.g., payoff functions or number of players
- ▶ **Arbitrary constraints and arbitrary correlation device**
 - ▶ **Characterizations of Equilibrium Strategies**
 - ▶ Definition through generalized Nash equilibrium and correlated equilibrium with alternative characterizations
 - ▶ **Properties and relationship to (unconstrained) Correlated Equilibria**
 - ▶ A feasible correlated equilibrium is a constrained correlated equilibrium
 - ▶ The set of constrained correlated equilibrium distributions may not be convex
 - ▶ There exist constrained correlated equilibrium distributions outside the set of correlated equilibrium distributions
- ▶ **Constraints on probability distributions**
 - ▶ **Characterization of constrained correlated equilibrium distributions**
 - ▶ Closed-form expression of the set of constrained correlated equilibrium distribution
 - ▶ **Conditions of Existence**
 - ▶ There does not always exist a constrained correlated equilibrium in finite games
 - ▶ **Constrained correlated equilibrium distributions of the mixed extension of a game**
 - ▶ Every constrained correlated equilibrium distribution of ΔG is a constrained correlated equilibrium distribution of G
 - ▶ **Computation of linearly constrained correlated equilibria**

Open Perspectives

▶ Research Questions

- ▶ Are there any theoretical guarantees of convergence to the a correlated equilibrium distribution?
- ▶ Connection to Bayesian rationality still holds in the presence of constraints ?
- ▶ What happens in the case of infinite games ?
- ▶ Existence problem should be studied for weaker or alternative assumptions
- ▶ Study of the problem of computing constrained correlated equilibria (complexity, feasibility)

▶ Some Unexplored Topics

- ▶ Learning rule leading to a specific correlated equilibrium (e.g., social welfare maximizing)
- ▶ Learning with constraints
- ▶ Subjective correlated equilibrium

▶ Potential applications of constrained correlated equilibria and learning correlated equilibria in various fields

- ▶ Mobile Edge Computing Systems
- ▶ Wireless Networks with constraints
- ▶ Engineering and beyond

Open Perspectives

► Research Questions

- Are there any theoretical guarantees of convergence to the a correlated equilibrium distribution?
- Connection to Bayesian rationality still holds in the presence of constraints ?
- What happens in the case of infinite games ?
- Existence problem should be studied for weaker or alternative assumptions
- Study of the problem of computing constrained correlated equilibria (complexity, feasibility)

► Some Unexplored Topics

- Learning rule leading to a specific correlated equilibrium (e.g., social welfare maximizing)
- Learning with constraints
- Subjective correlated equilibrium

► Potential applications of constrained correlated equilibria and learning correlated equilibria in various fields

- Mobile Edge Computing Systems
- Wireless Networks with constraints
- Engineering and beyond

Open Perspectives

▶ Research Questions

- ▶ Are there any theoretical guarantees of convergence to the a correlated equilibrium distribution?
- ▶ Connection to Bayesian rationality still holds in the presence of constraints ?
- ▶ What happens in the case of infinite games ?
- ▶ Existence problem should be studied for weaker or alternative assumptions
- ▶ Study of the problem of computing constrained correlated equilibria (complexity, feasibility)

▶ Some Unexplored Topics

- ▶ Learning rule leading to a specific correlated equilibrium (e.g., social welfare maximizing)
- ▶ Learning with constraints
- ▶ Subjective correlated equilibrium

▶ Potential applications of constrained correlated equilibria and learning correlated equilibria in various fields

- ▶ Mobile Edge Computing Systems
- ▶ Wireless Networks with constraints
- ▶ Engineering and beyond

Open Perspectives

▶ Research Questions

- ▶ Are there any theoretical guarantees of convergence to the a correlated equilibrium distribution?
- ▶ Connection to Bayesian rationality still holds in the presence of constraints ?
- ▶ What happens in the case of infinite games ?
- ▶ Existence problem should be studied for weaker or alternative assumptions
- ▶ Study of the problem of computing constrained correlated equilibria (complexity, feasibility)

▶ Some Unexplored Topics

- ▶ Learning rule leading to a specific correlated equilibrium (e.g., social welfare maximizing)
- ▶ Learning with constraints
- ▶ Subjective correlated equilibrium

▶ Potential applications of constrained correlated equilibria and learning correlated equilibria in various fields

- ▶ Mobile Edge Computing Systems
- ▶ Wireless Networks with constraints
- ▶ Engineering and beyond

Thank you!



Questions?

For more information:

omar.t.boufous@gmail.com

Bibliography

- R. J. Aumann. *Subjectivity and correlation in randomized strategies*. Journal of Mathematical Economics, 1974.
- R. J. Aumann. *Correlated equilibrium as an expression of bayesian rationality*. Econometrica, 1987.
- S. Hart & D. Schmeidler. *Existence of correlated equilibria*. Mathematics of Operations Research, 1989.
- B. von Stengel & F. Forges. *Extensive-form correlated equilibrium: Definition and computational complexity*. Mathematics of Operations Research, 2008.
- F. Forges. *Correlated equilibria and communication in games*. Complex Social and Behavioral Systems: Game Theory and Agent-Based Models, 2020.
- A. Brandenburger, E. Dekel, and John Geanakoplos. *Correlated equilibrium with generalized information structures*. Games and Economic Behavior, 1992.
- S. Grant and R. Stauber. *Delegation and ambiguity in correlated equilibrium*. Games and Economic Behavior, 2022.
- G. Debreu. *A social equilibrium existence theorem*. Proceedings of the National Academy of Sciences, 1952.
- K. J. Arrow and G Debreu. *Existence of an equilibrium for a competitive economy*. Econometrica: Journal of the Econometric Society, 1954.
- J. B. Rosen. *Existence and Uniqueness of Equilibrium Points for Concave N-Person Games*. Econometrica, 1965.
- Young, H. P. (2009). *Learning by trial and error*. Games and economic behavior.

Bibliography

Ankur A Kulkarni. *Games and teams with shared constraints*. Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences, 375(2100):20160302, 2017.

Jong-Shi Pang, Gesualdo Scutari, Francisco Facchinei, and Chaoxiong Wang. *Distributed power allocation with rate constraints in gaussian parallel interference channels*. IEEE Transactions on Information Theory, 54(8):3471–3489, 2008.

Daniel Nowak, Tobias Mahn, Hussein Al-Shatri, Alexandra Schwartz, and Anja Klein. *A generalized nash game for mobile edge computation offloading*. In 2018 6th IEEE International Conference on Mobile Cloud Computing, Services, and Engineering (MobileCloud), pages 95–102. IEEE, 2018.

Computation of Constrained Correlated Equilibria

Mixed-Integer Linear Program

$$\mathbf{p} \geq 0, \Sigma_{\mathbf{a} \in \mathcal{A}} \mathbf{p}(\mathbf{a}) = 1, F\mathbf{p} \leq 0 \quad (19)$$

$$\left[U_i - U_i B_{\beta_i} \right] \mathbf{p} \leq M_i \times \mathbf{y}_{\beta_i} \quad \forall i \in \mathcal{N}, \quad \forall \beta_i \quad (20)$$

$$FB_{\beta_i} \mathbf{p} \geq -K \left(1 - \mathbf{y}_{\beta_i} \right) + \delta \quad \forall i \in \mathcal{N}, \quad \forall \beta_i \quad (21)$$

$$FB_{\beta_i} \mathbf{p} \leq K \mathbf{y}_{\beta_i} \quad \forall i \in \mathcal{N}, \quad \forall \beta_i \quad (22)$$

$$\mathbf{y}_{\beta_i} \in \{0, 1\} \quad \forall i \in \mathcal{N}, \quad \forall \beta_i \quad (23)$$