

ImPRESS Statistical Analysis Mock-up Report

Fully planned efficacy and safety results of the Newly Diagnosed Glioblastoma Cohort

ImPRESS Statistical Team

25 June 2026

Contents

Introduction	4
Randomisation Status	4
Cohort Scope	4
Baseline Characteristics	4
Primary Efficacy Analyses (MRI)	11
Candidate dose-response models	12
Mean relative cerebral blood flow (PTA-ROI)	12
Mean relative cerebral blood flow (ROI2)	17
Mean relative solid stress (PTA-ROI)	22
Mean relative solid stress (ROI2)	27
Other MRI Endpoints up to Day 43	32
Mean DSC K2 leakage	32
Mean relative DSC extravasation	35
Mean relative apparent diffusion coefficient	37
Mean relative cerebral blood	39
Mean relative direction value	41
Mean relative fractional anisotropy	43
Mean relative restricted spectrum imaging	45
Mean relative transit time	47
Mean relative vessel caliber	49
Mean relative vessel size index	51
Mean relative vortex area	53
Mean tissue elasticity	55
Mean tissue relative viscosity	57
Mean tissue stiffness	59
Mean tissue viscosity	61
Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3	63
Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3	65
MRI Endpoints at Cycle 11 (Day 141)	67
Mean DSC K2 leakage	67
Mean relative DSC extravasation	69
Mean relative apparent diffusion coefficient	70
Mean relative cerebral blood	71

Mean relative cerebral blood flow (PTA-ROI)	72
Mean relative cerebral blood flow (ROI2)	73
Mean relative direction value	74
Mean relative fractional anisotropy	75
Mean relative restricted spectrum imaging	76
Mean relative solid stress (PTA-ROI)	77
Mean relative solid stress (ROI2)	78
Mean relative transit time	79
Mean relative vessel caliber	80
Mean relative vessel size index	81
Mean relative vortex area	82
Mean tissue elasticity	83
Mean tissue relative viscosity	84
Mean tissue stiffness	85
Mean tissue viscosity	86
Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3	87
Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3	88
Neurologic Performance	89
Karnofsky Performance Scale	89
ECOG Score	92
Total NANO score	95
Quality of Life Outcomes	98
EORTC QLQ-C30 score	98
EORTC QLQ-BN20 score	100
Steroid Use	102
Steroid use	102
RANO Responses	105
State of disease according to the RANO criteria	105
Best overall radiographic response	106
Biomarker Endpoints (Serum Cytokines / Angiogenic Factors)	107
Fractalkine	107
Granzyme B	109
IFN-g	111
IL-1b	113
IL-6	115
IL-8	117
MCP-1	119
MMP-10	121
PDGF-DD	123
PECAM-1	125
TGF-alpha	127
Thrombospondin-2	129
VEGF-A	131
VEGF-C	133
Angiopoietin-1	135
Angiopoietin-2	137
ANGPTL4	139
CA9	141
CD163	143
Collagen IV alpha 1	145

EGF	147
ErbB2	149
FGF Basic	151
IL-12 p70	153
IL-2	155
IP-10 (CXCL10)	157
I-TAC (CXCL11)	159
MIG (CXCL9)	161
PIGF	163
Tenascin C	165
TNFRSF18 (GITR)	167
TNFSF6 (FasL)	169
VEGFR2	171
Biomarker Endpoints — Long-term (Cycle 15+)	173
Fractalkine	173
Granzyme B	175
IFN-g	177
IL-1b	179
IL-6	181
IL-8	183
MCP-1	185
MMP-10	187
PDGF-DD	189
PECAM-1	191
TGF-alpha	193
Thrombospondin-2	195
VEGF-A	197
VEGF-C	199
Angiopoietin-1	201
Angiopoietin-2	203
ANGPTL4	205
CA9	207
CD163	209
Collagen IV alpha 1	211
EGF	213
ErbB2	215
FGF Basic	217
IL-12 p70	219
IL-2	221
IP-10 (CXCL10)	223
I-TAC (CXCL11)	225
MIG (CXCL9)	227
PIGF	229
Tenascin C	231
TNFRSF18 (GITR)	233
TNFSF6 (FasL)	235
VEGFR2	237
Time-to-Event Endpoints	239
Progression-free survival	240
6-month progression-free survival	242
Overall survival	244
Overall survival at 12 months	246

Overall survival at 24 months	248
Safety	250
Extent of Exposure	250
Adverse Events	252
Vital Signs	256
Conventional Laboratory Parameters	265
Deaths	267

Introduction

The ImPRESS study is a prospective investigation of losartan as an adjunctive therapy for patients with glioblastoma. According to Protocol Version 8.0 (14-Nov-2025), the trial collects comprehensive demographic, clinical, imaging, and quality-of-life information prior to treatment initiation. The statistical objectives for the interim analyses focus on characterising baseline status, neurological function, and tumour burden before randomisation to evaluate comparability across planned treatment arms.

Randomisation Status

The present dataset is built on sham randomisation generated from the most recent export (see `_targets.R` workflow). No true treatment assignments have been unlocked. All findings in this document should therefore be interpreted as preliminary and descriptive only. Final efficacy comparisons will not be conducted until the database is locked and genuine randomisation codes are released.

Cohort Scope

The results herein are restricted to participants in cohort code 2 as configured in `config/cfg.yml`. Subjects from other cohorts are excluded from all summaries, figures, and tables in this draft. A combined-cohort analysis can be delivered once recruitment and cleaning are complete for all strata.

Baseline Characteristics

This section summarises baseline characteristics for the newly diagnosed glioblastoma cohort using the `adb1` ADaM dataset.

Demographics and Baseline characteristics

	all obs
Age (years)	
n	54
Mean (SD)	52.6 (10.2)
Median	55.0
Min - Max	26.0 - 69.0
Sex	
n	54
Female	29 (53.7%)
Male	25 (46.3%)

Demographics and Baseline characteristics

	all obs
Cohort	
n	54
Newly diagnosed glioblastoma	54 (100%)
Country	
n	54
NOR	54 (100%)
Childbearing potential?	
n	29
No	16 (55.2%)
Yes	13 (44.8%)
Diagnosis	
n	54
Newly diagnosed glioblastoma	54 (100%)
Time from surgery to randomisation (days)	
n	54
Mean (SD)	27.4 (7.3)
Median	25.5
Min - Max	13.0 - 48.0
IDH1/2 mutation	
n	54
Negative	47 (87%)
Positive	7 (13%)
MGMT status	
n	54
Negative	30 (55.6%)
Positive	24 (44.4%)
Pregnancy test result	
n	13
Negative	13 (100%)
ECOG Score	
n	54
Mean (SD)	0.3 (0.5)
Median	0.0
Min - Max	0.0 - 2.0

Demographics and Baseline characteristics

	all obs
ECOG Score (categorical)	
n	54
ECOG 0	40 (74.1%)
ECOG 1	13 (24.1%)
ECOG 2	1 (1.9%)
Karnofsky Performance Scale	
n	54
Mean (SD)	95.0 (8.0)
Median	100.0
Min - Max	60.0 - 100.0
Karnofsky Performance Scale (categorical)	
n	54
KPS 100	34 (63%)
KPS 60	1 (1.9%)
KPS 80	4 (7.4%)
KPS 90	15 (27.8%)
Total NANO score	
n	54
Mean (SD)	1.0 (1.3)
Median	0.0
Min - Max	0.0 - 4.0
EORTC QLQ-C30 score	
n	53
Mean (SD)	56.3 (8.6)
Median	54.0
Min - Max	41.0 - 77.0
EORTC QLQ-BN20 score	
n	53
Mean (SD)	30.8 (6.6)
Median	30.0
Min - Max	20.0 - 49.0
Systolic blood pressure	
n	54
Mean (SD)	130.0 (15.4)

Demographics and Baseline characteristics

	all obs
Median	128.5
Min - Max	106.0 - 180.0
Diastolic blood pressure	
n	54
Mean (SD)	81.9 (7.1)
Median	82.0
Min - Max	67.0 - 97.0
Pulse Rate	
n	54
Mean (SD)	80.6 (15.6)
Median	78.0
Min - Max	54.0 - 114.0
Weight (kg)	
n	54
Mean (SD)	78.8 (13.2)
Median	78.3
Min - Max	48.2 - 116.0
BMI	
n	51
Mean (SD)	25.0 (4.2)
Median	23.9
Min - Max	18.6 - 39.9
Height (cm)	
n	51
Mean (SD)	176.5 (8.8)
Median	175.0
Min - Max	158.0 - 194.0
Has the patient received chemotherapy	
n	54
No	54 (100%)
Has the patient received immunotherapy	
n	54
No	54 (100%)
Has the patient received radiation therapy to the brain?	

Demographics and Baseline characteristics

	all obs
n	54
No	54 (100%)
CNS surgery	
n	54
No	3 (5.6%)
Yes	51 (94.4%)
Steroids at time of MRI	
n	54
No	42 (77.8%)
Yes	12 (22.2%)
Mean relative cerebral blood flow (PTA-ROI)	
n	54
Mean (SD)	1.5 (0.4)
Median	1.4
Min - Max	0.7 - 2.4
Mean relative cerebral blood flow (ROI2)	
n	47
Mean (SD)	1.1 (0.5)
Median	1.0
Min - Max	0.2 - 2.2
Mean relative solid stress (PTA-ROI)	
n	41
Mean (SD)	0.8 (0.1)
Median	0.8
Min - Max	0.5 - 1.1
Mean relative solid stress (ROI2)	
n	32
Mean (SD)	0.9 (0.2)
Median	0.9
Min - Max	0.5 - 1.4
Mean relative apparent diffusion coefficient	
n	52
Mean (SD)	1.4 (0.2)
Median	1.4

Demographics and Baseline characteristics

	all obs
Min - Max	1.0 - 2.0
Mean relative fractional anisotropy	
n	52
Mean (SD)	0.5 (0.1)
Median	0.5
Min - Max	0.3 - 1.0
Mean relative restricted spectrum imaging	
n	50
Mean (SD)	0.5 (0.2)
Median	0.5
Min - Max	0.2 - 1.1
Mean DSC K2 leakage	
n	54
Mean (SD)	0.2 (1.6)
Median	0.4
Min - Max	-3.9 - 4.5
Mean relative cerebral blood	
n	54
Mean (SD)	1.7 (0.5)
Median	1.7
Min - Max	1.0 - 2.8
Mean relative transit time	
n	54
Mean (SD)	1.4 (0.3)
Median	1.4
Min - Max	0.9 - 2.3
Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3	
n	54
Mean (SD)	6.3 (6.5)
Median	4.3
Min - Max	0.0 - 26.3
Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3	
n	54
Mean (SD)	13.5 (14.3)

Demographics and Baseline characteristics

	all obs
Median	8.5
Min - Max	0.1 - 57.4
Mean tissue elasticity	
n	41
Mean (SD)	0.8 (0.1)
Median	0.8
Min - Max	0.5 - 1.2
Mean tissue relative viscosity	
n	41
Mean (SD)	0.9 (0.2)
Median	0.9
Min - Max	0.0 - 1.1
Mean tissue stiffness	
n	41
Mean (SD)	0.8 (0.1)
Median	0.8
Min - Max	0.5 - 1.1
Mean tissue viscosity	
n	41
Mean (SD)	0.7 (0.2)
Median	0.7
Min - Max	-0.0 - 1.1
Mean relative DSC extravasation	
n	28
Mean (SD)	1.2 (0.8)
Median	1.0
Min - Max	0.7 - 5.1
Mean relative direction value	
n	28
Mean (SD)	0.4 (1.4)
Median	0.3
Min - Max	-2.7 - 5.4
Mean relative vessel caliber	
n	28

Demographics and Baseline characteristics


	all obs
Mean (SD)	1.0 (0.3)
Median	1.0
Min - Max	0.5 - 1.9
Mean relative vessel size index	
n	28
Mean (SD)	1.4 (0.6)
Median	1.2
Min - Max	0.6 - 2.6
Mean relative vortex area	
n	28
Mean (SD)	1.0 (0.1)
Median	1.0
Min - Max	0.8 - 1.1

Primary Efficacy Analyses (MRI)

The primary variable is analysed with the MCP-Mod methodology as described in the protocol and SAP. We pre-specify four candidate dose-response models (linear, piecewise linear, Emax, sigmoid Emax) and derive optimal contrasts using the estimated variance-covariance from the mixed model. The MCP step tests for a dose-response signal across candidate models with multiplicity control; the Mod step presents the model fit by AIC and reports ED values. Results below are grouped by MRI endpoint (primary and sensitivity (ROI2)). All analyses are based on data up to and including Day 43 (Week 6).

Candidate dose-response models

Figure 1: Pre-specified candidate dose-response models



Mean relative cerebral blood flow (PTA-ROI)


The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot

Mean Response over Time for Mean relative cerebral blood flow (PTA-ROI)



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.663	-0.817	-0.634	-0.865
25 mg	-0.158	0.045	-0.315	0.268
50 mg	0.097	0.259	0.317	0.275
100 mg	0.725	0.514	0.631	0.322

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
sigEmax	-0.10	0.69
linear	-0.60	0.85
emax	-0.79	0.89
linInt	-1.05	0.93

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	6.92	50.00	70.00	90.00



Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linInt	8.00	12.50	17.50	22.50
emax	8.17	0.10	0.23	0.89
sigEmax	10.17	0.62	0.71	0.88


Note Fitted models are presented with AIC and ED for reference


Estimated means by dose

Treatment	Estimated Change from Baseline (95% CI)
0 mg	0.06 (-0.04 to 0.16)
25 mg	-0.05 (-0.17 to 0.07)
50 mg	0.07 (-0.05 to 0.18)
100 mg	-0.01 (-0.12 to 0.11)

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.06 (-0.17 to 0.05)

Mean relative cerebral blood flow (ROI2)


The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot

Mean Response over Time for Mean relative cerebral blood flow (ROI2)



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.719	-0.826	-0.707	-0.863
25 mg	-0.081	0.071	-0.203	0.231
50 mg	0.120	0.257	0.305	0.279
100 mg	0.680	0.497	0.605	0.353

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
sigEmax	1.53	0.11
linear	1.44	0.12
emax	1.08	0.21
linInt	0.64	0.37

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	5.00	50.00	70.00	90.00



Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
emax	7.28	37.50	58.33	84.37
linInt	8.00	57.64	74.58	91.53
sigEmax	8.45	50.04	54.46	62.29


Note Fitted models are presented with AIC and ED for reference


Estimated means by dose

Treatment	Estimated Change from Baseline (95% CI)
0 mg	0.04 (-0.11 to 0.19)
25 mg	-0.03 (-0.22 to 0.15)
50 mg	0.08 (-0.09 to 0.26)
100 mg	0.15 (-0.01 to 0.32)

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.04 (-0.10 to 0.18)


Mean relative solid stress (PTA-ROI)

The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.707	-0.835	-0.635	-0.865
25 mg	-0.157	0.065	-0.342	0.290
50 mg	0.205	0.351	0.450	0.328
100 mg	0.658	0.419	0.527	0.247

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
sigEmax	0.48	0.45
linear	-0.06	0.67
emax	-0.67	0.86
linInt	-1.29	0.96

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linInt	8.00	12.50	17.50	22.50



Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	9.39	50.00	70.00	90.00
emax	9.75	0.10	0.23	0.89
sigEmax	11.74	0.62	0.71	0.88


Note Fitted models are presented with AIC and ED for reference


Estimated means by dose

Treatment	Estimated Change from Baseline (95% CI)
0 mg	0.03 (-0.01 to 0.07)
25 mg	-0.04 (-0.08 to 0.01)
50 mg	0.02 (-0.03 to 0.06)
100 mg	0.01 (-0.04 to 0.06)

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.03 (-0.07 to 0.02)

Mean relative solid stress (ROI2)


The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot

Mean Response over Time for Mean relative solid stress (ROI2)



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.669	-0.819	-0.596	-0.862
25 mg	-0.224	0.008	-0.392	0.260
50 mg	0.220	0.382	0.461	0.363
100 mg	0.673	0.428	0.528	0.239

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
sigEmax	-0.23	0.74
linear	-0.58	0.85
emax	-0.68	0.87
linInt	-0.81	0.9

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	5.12	50.00	70.00	90.00



Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
emax	6.80	0.10	0.23	0.89
linInt	8.00	12.50	17.50	22.50
sigEmax	8.80	0.62	0.71	0.88


Note Fitted models are presented with AIC and ED for reference


Estimated means by dose

Treatment	Estimated Change from Baseline (95% CI)
0 mg	0.03 (-0.08 to 0.13)
25 mg	-0.05 (-0.17 to 0.06)
50 mg	0.01 (-0.10 to 0.12)
100 mg	-0.04 (-0.17 to 0.09)

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.05 (-0.17 to 0.07)


Other MRI Endpoints up to Day 43

For remaining MRI endpoints, we present estimated means by dose, contrasts vs 0 mg, and a plot with confidence intervals.

Mean DSC K2 leakage

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.14 (-0.57 to 0.28)
25 mg	-0.15 (-0.67 to 0.36)
50 mg	-0.39 (-0.89 to 0.12)
100 mg	-0.64 (-1.13 to -0.15)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.01 (-0.77 to 0.75)
50 mg vs 0 mg	-0.24 (-0.99 to 0.51)
100 mg vs 0 mg	-0.49 (-1.22 to 0.23)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.27 (-0.75 to 0.21)

Mean relative DSC extravasation

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.08 (-0.37 to 0.21)
25 mg	0.11 (-0.17 to 0.39)
50 mg	-0.01 (-0.30 to 0.29)
100 mg	0.01 (-0.31 to 0.33)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.19 (-0.33 to 0.71)
50 mg vs 0 mg	0.07 (-0.45 to 0.59)
100 mg vs 0 mg	0.09 (-0.46 to 0.64)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.12 (-0.24 to 0.48)

Mean relative apparent diffusion coefficient

A total of 51 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.16 (0.10 to 0.22)
25 mg	0.17 (0.10 to 0.24)
50 mg	0.15 (0.08 to 0.22)
100 mg	0.15 (0.08 to 0.22)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.01 (-0.09 to 0.11)
50 mg vs 0 mg	-0.01 (-0.11 to 0.09)
100 mg vs 0 mg	-0.01 (-0.10 to 0.09)

Plot



Supplementary analysis: any dose vs 0 mg


Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.01 (-0.07 to 0.06)

Mean relative cerebral blood

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start

Mean Response over Time for Mean relative cerebral blood




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.01 (-0.10 to 0.12)
25 mg	-0.10 (-0.23 to 0.02)
50 mg	0.04 (-0.09 to 0.17)
100 mg	-0.09 (-0.21 to 0.03)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.11 (-0.31 to 0.08)
50 mg vs 0 mg	0.03 (-0.16 to 0.22)
100 mg vs 0 mg	-0.10 (-0.29 to 0.09)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.06 (-0.19 to 0.06)

Mean relative direction value

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-44.52 (-3952.76 to 3863.72)
25 mg	2613.06 (-1286.41 to 6512.53)
50 mg	370.06 (-3538.82 to 4278.94)
100 mg	1263.10 (-2973.21 to 5499.42)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	2657.57 (-4493.68 to 9808.83)
50 mg vs 0 mg	414.58 (-6507.53 to 7336.68)
100 mg vs 0 mg	1307.62 (-5835.93 to 8451.17)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	1380.11 (-3384.17 to 6144.38)

Mean relative fractional anisotropy

A total of 51 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.05 (-0.08 to -0.02)
25 mg	-0.03 (-0.06 to 0.01)
50 mg	-0.04 (-0.08 to -0.01)
100 mg	-0.07 (-0.11 to -0.04)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.02 (-0.03 to 0.07)
50 mg vs 0 mg	0.01 (-0.04 to 0.05)
100 mg vs 0 mg	-0.03 (-0.07 to 0.02)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.00 (-0.03 to 0.03)

Mean relative restricted spectrum imaging

A total of 49 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.11 (-0.15 to -0.06)
25 mg	-0.11 (-0.17 to -0.06)
50 mg	-0.07 (-0.12 to -0.02)
100 mg	-0.10 (-0.15 to -0.05)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.01 (-0.08 to 0.07)
50 mg vs 0 mg	0.04 (-0.04 to 0.11)
100 mg vs 0 mg	0.01 (-0.06 to 0.07)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.01 (-0.03 to 0.05)

Mean relative transit time

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.06 (-0.13 to 0.02)
25 mg	-0.08 (-0.17 to 0.00)
50 mg	-0.06 (-0.14 to 0.02)
100 mg	-0.07 (-0.15 to 0.01)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.03 (-0.16 to 0.10)
50 mg vs 0 mg	0.00 (-0.13 to 0.12)
100 mg vs 0 mg	-0.02 (-0.14 to 0.10)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.02 (-0.10 to 0.06)

Mean relative vessel caliber

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.16 (-0.03 to 0.34)
25 mg	-0.09 (-0.28 to 0.11)
50 mg	0.15 (-0.05 to 0.35)
100 mg	0.11 (-0.11 to 0.34)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.24 (-0.55 to 0.06)
50 mg vs 0 mg	-0.01 (-0.32 to 0.31)
100 mg vs 0 mg	-0.05 (-0.38 to 0.29)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.11 (-0.31 to 0.10)

Mean relative vessel size index

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.35 (-0.04 to 0.74)
25 mg	-0.12 (-0.53 to 0.29)
50 mg	0.36 (-0.07 to 0.79)
100 mg	0.28 (-0.20 to 0.75)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.47 (-1.13 to 0.19)
50 mg vs 0 mg	0.01 (-0.67 to 0.68)
100 mg vs 0 mg	-0.08 (-0.80 to 0.64)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.19 (-0.64 to 0.25)

Mean relative vortex area

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.01 (-0.04 to 0.06)
25 mg	-0.02 (-0.07 to 0.04)
50 mg	0.02 (-0.04 to 0.08)
100 mg	0.02 (-0.04 to 0.08)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.03 (-0.11 to 0.05)
50 mg vs 0 mg	0.01 (-0.08 to 0.10)
100 mg vs 0 mg	0.01 (-0.08 to 0.10)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.01 (-0.06 to 0.05)

Mean tissue elasticity

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.02 (-0.02 to 0.06)
25 mg	-0.02 (-0.06 to 0.03)
50 mg	0.03 (-0.02 to 0.07)
100 mg	0.03 (-0.02 to 0.08)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.04 (-0.11 to 0.03)
50 mg vs 0 mg	0.01 (-0.06 to 0.07)
100 mg vs 0 mg	0.01 (-0.07 to 0.09)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.01 (-0.05 to 0.04)

Mean tissue relative viscosity

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.03 (-0.02 to 0.07)
25 mg	-0.08 (-0.13 to -0.03)
50 mg	0.02 (-0.03 to 0.06)
100 mg	-0.04 (-0.10 to 0.01)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.11 (-0.20 to -0.03)
50 mg vs 0 mg	-0.01 (-0.09 to 0.07)
100 mg vs 0 mg	-0.07 (-0.16 to 0.02)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.07 (-0.13 to -0.01)

Mean tissue stiffness

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.03 (-0.01 to 0.07)
25 mg	-0.04 (-0.08 to 0.01)
50 mg	0.02 (-0.03 to 0.06)
100 mg	0.01 (-0.04 to 0.06)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.06 (-0.13 to 0.01)
50 mg vs 0 mg	-0.01 (-0.07 to 0.06)
100 mg vs 0 mg	-0.01 (-0.09 to 0.06)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.03 (-0.07 to 0.02)

Mean tissue viscosity

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.07 (-0.01 to 0.14)
25 mg	-0.07 (-0.16 to 0.01)
50 mg	0.05 (-0.03 to 0.13)
100 mg	-0.03 (-0.12 to 0.07)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.14 (-0.26 to -0.02)
50 mg vs 0 mg	-0.02 (-0.14 to 0.11)
100 mg vs 0 mg	-0.09 (-0.23 to 0.04)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.08 (-0.16 to 0.00)

Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.00 (-1.50 to 1.49)
25 mg	0.57 (-1.21 to 2.35)
50 mg	-0.99 (-2.77 to 0.78)
100 mg	-0.30 (-2.00 to 1.40)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.57 (-1.63 to 2.78)
50 mg vs 0 mg	-0.99 (-3.22 to 1.24)
100 mg vs 0 mg	-0.29 (-2.35 to 1.76)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.25 (-1.55 to 1.06)

Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	7.89 (2.97 to 12.82)
25 mg	8.07 (2.22 to 13.93)
50 mg	8.07 (2.22 to 13.92)
100 mg	1.87 (-3.76 to 7.50)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.18 (-7.47 to 7.83)
50 mg vs 0 mg	0.18 (-7.51 to 7.87)
100 mg vs 0 mg	-6.02 (-13.22 to 1.17)

Plot



Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-2.24 (-6.84 to 2.37)

MRI Endpoints at Cycle 11 (Day 141)

For all MRI endpoints, we report estimated change from baseline at Cycle 11 (day 141), grouped by the extended treatment variable. The summaries include mean estimates, contrasts versus the reference group, and a plot with confidence intervals.

Mean DSC K2 leakage

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.85 (-2.14 to 0.44)
losartan 25 mg	-0.56 (-2.45 to 1.32)
losartan 50 mg	-2.60 (-4.39 to -0.80)
losartan 100 mg	-1.14 (-2.37 to 0.09)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.29 (-2.48 to 3.06)
losartan 50 mg - losartan 0 mg	-1.75 (-4.41 to 0.92)
losartan 100 mg - losartan 0 mg	-0.29 (-2.47 to 1.89)

Plot



Mean relative DSC extravasation

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.38 (-0.91 to 0.16)
losartan 25 mg	0.18 (-0.48 to 0.84)
losartan 50 mg	0.35 (-0.15 to 0.85)
losartan 100 mg	0.03 (-0.40 to 0.46)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.55 (-0.52 to 1.63)
losartan 50 mg - losartan 0 mg	0.73 (-0.18 to 1.63)
losartan 100 mg - losartan 0 mg	0.41 (-0.47 to 1.28)

Plot



Mean relative apparent diffusion coefficient

A total of 43 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.31 (0.20 to 0.42)
losartan 25 mg	0.34 (0.18 to 0.51)
losartan 50 mg	0.10 (-0.04 to 0.24)
losartan 100 mg	0.23 (0.13 to 0.33)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.04 (-0.20 to 0.28)
losartan 50 mg - losartan 0 mg	-0.21 (-0.42 to 0.01)
losartan 100 mg - losartan 0 mg	-0.07 (-0.26 to 0.11)

Plot



Mean relative cerebral blood

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.24 (-0.43 to -0.04)
losartan 25 mg	-0.09 (-0.38 to 0.19)
losartan 50 mg	-0.05 (-0.32 to 0.21)
losartan 100 mg	-0.32 (-0.51 to -0.14)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.15 (-0.27 to 0.56)
losartan 50 mg - losartan 0 mg	0.19 (-0.21 to 0.59)
losartan 100 mg - losartan 0 mg	-0.09 (-0.41 to 0.24)

Plot



Mean relative cerebral blood flow (PTA-ROI)

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.17 (-0.32 to -0.02)
losartan 25 mg	-0.09 (-0.30 to 0.13)
losartan 50 mg	-0.08 (-0.28 to 0.12)
losartan 100 mg	-0.21 (-0.35 to -0.07)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.08 (-0.24 to 0.40)
losartan 50 mg - losartan 0 mg	0.09 (-0.21 to 0.40)
losartan 100 mg - losartan 0 mg	-0.04 (-0.29 to 0.21)

Plot



Mean relative cerebral blood flow (ROI2)

A total of 40 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.26 (-0.41 to -0.11)
losartan 25 mg	-0.35 (-0.59 to -0.11)
losartan 50 mg	-0.24 (-0.44 to -0.03)
losartan 100 mg	-0.34 (-0.48 to -0.20)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.09 (-0.43 to 0.26)
losartan 50 mg - losartan 0 mg	0.02 (-0.28 to 0.33)
losartan 100 mg - losartan 0 mg	-0.08 (-0.33 to 0.17)

Plot



Mean relative direction value

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	59.78 (-7.67 to 127.22)
losartan 25 mg	-1.55 (-81.99 to 78.89)
losartan 50 mg	1.14 (-62.00 to 64.27)
losartan 100 mg	0.06 (-52.02 to 52.15)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-61.33 (-195.07 to 72.40)
losartan 50 mg - losartan 0 mg	-58.64 (-177.84 to 60.55)
losartan 100 mg - losartan 0 mg	-59.72 (-166.37 to 46.93)

Plot



Mean relative fractional anisotropy

A total of 40 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.05 (-0.13 to 0.03)
losartan 25 mg	-0.11 (-0.23 to 0.00)
losartan 50 mg	0.00 (-0.10 to 0.10)
losartan 100 mg	-0.06 (-0.14 to 0.02)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.07 (-0.24 to 0.11)
losartan 50 mg - losartan 0 mg	0.05 (-0.11 to 0.20)
losartan 100 mg - losartan 0 mg	-0.01 (-0.14 to 0.13)

Plot



Mean relative restricted spectrum imaging

A total of 40 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.24 (-0.31 to -0.16)
losartan 25 mg	-0.17 (-0.27 to -0.06)
losartan 50 mg	-0.10 (-0.20 to -0.01)
losartan 100 mg	-0.17 (-0.24 to -0.10)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.07 (-0.08 to 0.23)
losartan 50 mg - losartan 0 mg	0.13 (-0.01 to 0.28)
losartan 100 mg - losartan 0 mg	0.07 (-0.05 to 0.19)

Plot



Mean relative solid stress (PTA-ROI)

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.05 (-0.14 to 0.05)
losartan 25 mg	-0.10 (-0.26 to 0.06)
losartan 50 mg	0.10 (-0.03 to 0.23)
losartan 100 mg	0.06 (-0.04 to 0.16)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.05 (-0.27 to 0.16)
losartan 50 mg - losartan 0 mg	0.15 (-0.05 to 0.36)
losartan 100 mg - losartan 0 mg	0.10 (-0.07 to 0.28)

Plot



Mean relative solid stress (ROI2)

A total of 23 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.04 (-0.14 to 0.05)
losartan 25 mg	0.04 (-0.13 to 0.21)
losartan 50 mg	-0.08 (-0.21 to 0.05)
losartan 100 mg	-0.01 (-0.11 to 0.09)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.08 (-0.16 to 0.33)
losartan 50 mg - losartan 0 mg	-0.04 (-0.23 to 0.16)
losartan 100 mg - losartan 0 mg	0.04 (-0.13 to 0.21)

Plot



Mean relative transit time

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.11 (-0.21 to -0.01)
losartan 25 mg	-0.07 (-0.21 to 0.08)
losartan 50 mg	-0.07 (-0.21 to 0.06)
losartan 100 mg	-0.06 (-0.15 to 0.03)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.04 (-0.17 to 0.25)
losartan 50 mg - losartan 0 mg	0.04 (-0.17 to 0.24)
losartan 100 mg - losartan 0 mg	0.05 (-0.12 to 0.21)

Plot



Mean relative vessel caliber

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.05 (-0.39 to 0.29)
losartan 25 mg	0.20 (-0.24 to 0.65)
losartan 50 mg	0.01 (-0.33 to 0.35)
losartan 100 mg	0.20 (-0.09 to 0.49)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.25 (-0.44 to 0.94)
losartan 50 mg - losartan 0 mg	0.06 (-0.53 to 0.65)
losartan 100 mg - losartan 0 mg	0.25 (-0.30 to 0.79)

Plot



Mean relative vessel size index

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.01 (-0.61 to 0.63)
losartan 25 mg	0.61 (-0.20 to 1.42)
losartan 50 mg	0.06 (-0.56 to 0.68)
losartan 100 mg	0.21 (-0.33 to 0.76)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.60 (-0.64 to 1.85)
losartan 50 mg - losartan 0 mg	0.05 (-1.03 to 1.12)
losartan 100 mg - losartan 0 mg	0.21 (-0.83 to 1.24)

Plot



Mean relative vortex area

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.03 (-0.05 to 0.11)
losartan 25 mg	0.03 (-0.08 to 0.14)
losartan 50 mg	-0.01 (-0.09 to 0.07)
losartan 100 mg	0.05 (-0.02 to 0.12)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.00 (-0.16 to 0.17)
losartan 50 mg - losartan 0 mg	-0.04 (-0.18 to 0.10)
losartan 100 mg - losartan 0 mg	0.02 (-0.11 to 0.15)

Plot



Mean tissue elasticity

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.02 (-0.11 to 0.07)
losartan 25 mg	-0.07 (-0.21 to 0.07)
losartan 50 mg	0.12 (0.00 to 0.24)
losartan 100 mg	0.05 (-0.04 to 0.15)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.05 (-0.24 to 0.15)
losartan 50 mg - losartan 0 mg	0.14 (-0.05 to 0.33)
losartan 100 mg - losartan 0 mg	0.07 (-0.09 to 0.24)

Plot



Mean tissue relative viscosity

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.05 (-0.12 to 0.03)
losartan 25 mg	-0.06 (-0.17 to 0.06)
losartan 50 mg	-0.01 (-0.10 to 0.09)
losartan 100 mg	0.08 (0.00 to 0.16)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.01 (-0.18 to 0.17)
losartan 50 mg - losartan 0 mg	0.04 (-0.11 to 0.19)
losartan 100 mg - losartan 0 mg	0.13 (-0.01 to 0.26)

Plot



Mean tissue stiffness

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.05 (-0.14 to 0.05)
losartan 25 mg	-0.10 (-0.26 to 0.06)
losartan 50 mg	0.10 (-0.03 to 0.23)
losartan 100 mg	0.06 (-0.04 to 0.16)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.05 (-0.27 to 0.16)
losartan 50 mg - losartan 0 mg	0.15 (-0.05 to 0.36)
losartan 100 mg - losartan 0 mg	0.10 (-0.07 to 0.28)

Plot



Mean tissue viscosity

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.02 (-0.16 to 0.12)
losartan 25 mg	-0.09 (-0.31 to 0.14)
losartan 50 mg	0.07 (-0.11 to 0.25)
losartan 100 mg	0.08 (-0.06 to 0.23)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.07 (-0.40 to 0.26)
losartan 50 mg - losartan 0 mg	0.09 (-0.19 to 0.37)
losartan 100 mg - losartan 0 mg	0.10 (-0.15 to 0.35)

Plot



Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	3.68 (0.10 to 7.27)
losartan 25 mg	2.63 (-2.66 to 7.92)
losartan 50 mg	-2.03 (-7.10 to 3.04)
losartan 100 mg	1.55 (-1.82 to 4.91)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-1.05 (-8.78 to 6.68)
losartan 50 mg - losartan 0 mg	-5.71 (-13.30 to 1.87)
losartan 100 mg - losartan 0 mg	-2.14 (-8.10 to 3.83)

Plot



Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	25.97 (10.30 to 41.63)
losartan 25 mg	18.46 (-4.59 to 41.52)
losartan 50 mg	21.61 (0.02 to 43.20)
losartan 100 mg	11.05 (-3.87 to 25.97)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-7.50 (-41.30 to 26.30)
losartan 50 mg - losartan 0 mg	-4.36 (-36.70 to 27.99)
losartan 100 mg - losartan 0 mg	-14.91 (-41.23 to 11.41)

Plot




Neurologic Performance

Change from baseline in KPS, ECOG, and NANO scores at Days 15/43 and model-based estimates at Days 155/239 are reported below for the newly diagnosed glioblastoma cohort. Note that since the day 29 is not measured for any of the measures, the dose-response analyses are based on Days 15 and 43 only for the early timepoints. There might therefore be a limited ability to detect dose-response relationships at these early visits because we are lacking one contrast.

Karnofsky Performance Scale

A total of 54 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Days 15/43 (change from baseline)

Dose (mg)	Estimated change from baseline (95% CI)
0	-0.31 (-2.10 to 1.47)
25	-1.13 (-3.06 to 0.80)
50	0.23 (-1.75 to 2.20)
100	1.00 (-0.93 to 2.92)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	-0.82 (-3.99 to 2.36)
losartan 50 mg vs losartan 0 mg	0.54 (-2.69 to 3.77)
losartan 100 mg vs losartan 0 mg	1.31 (-1.86 to 4.48)

Plot




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-1.86 (-4.93 to 1.21)
155	losartan 25 mg	-6.97 (-11.07 to -2.88)
155	losartan 50 mg	0.21 (-4.14 to 4.57)
155	losartan 100 mg	3.09 (0.20 to 5.98)
239	losartan 0 mg	0.09 (-3.26 to 3.43)
239	losartan 25 mg	-0.86 (-6.14 to 4.42)
239	losartan 50 mg	0.56 (-3.80 to 4.91)
239	losartan 100 mg	3.21 (0.25 to 6.16)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	-5.12 (-11.28 to 1.05)
155	losartan 50 mg - losartan 0 mg	2.07 (-4.44 to 8.59)
155	losartan 100 mg - losartan 0 mg	4.95 (-0.17 to 10.07)
239	losartan 25 mg - losartan 0 mg	-0.95 (-8.48 to 6.58)
239	losartan 50 mg - losartan 0 mg	0.47 (-6.24 to 7.18)
239	losartan 100 mg - losartan 0 mg	3.12 (-2.29 to 8.53)


Plot



ECOG Score

A total of 54 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Days 15/43 (change from baseline)

Dose (mg)	Estimated change from baseline (95% CI)
0	-0.09 (-0.29 to 0.10)
25	0.04 (-0.16 to 0.23)
50	0.11 (-0.09 to 0.31)
100	0.12 (-0.08 to 0.31)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	0.13 (-0.22 to 0.48)
losartan 50 mg vs losartan 0 mg	0.21 (-0.15 to 0.57)
losartan 100 mg vs losartan 0 mg	0.21 (-0.14 to 0.57)

Plot




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-0.11 (-0.36 to 0.14)
155	losartan 25 mg	0.50 (0.16 to 0.84)
155	losartan 50 mg	-0.16 (-0.53 to 0.20)
155	losartan 100 mg	0.05 (-0.18 to 0.28)
239	losartan 0 mg	-0.09 (-0.36 to 0.19)
239	losartan 25 mg	-0.01 (-0.46 to 0.44)
239	losartan 50 mg	-0.02 (-0.38 to 0.35)
239	losartan 100 mg	-0.11 (-0.36 to 0.13)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	0.62 (0.11 to 1.12)
155	losartan 50 mg - losartan 0 mg	-0.05 (-0.59 to 0.49)
155	losartan 100 mg - losartan 0 mg	0.17 (-0.24 to 0.58)
239	losartan 25 mg - losartan 0 mg	0.08 (-0.56 to 0.72)
239	losartan 50 mg - losartan 0 mg	0.07 (-0.49 to 0.63)
239	losartan 100 mg - losartan 0 mg	-0.03 (-0.47 to 0.42)


Plot



Total NANO score

A total of 54 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Days 15/43 (change from baseline)

Dose (mg)	Estimated change from baseline (95% CI)
0	-0.17 (-0.48 to 0.13)
25	0.08 (-0.25 to 0.41)
50	-0.21 (-0.55 to 0.12)
100	-0.19 (-0.52 to 0.15)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	0.26 (-0.28 to 0.80)
losartan 50 mg vs losartan 0 mg	-0.04 (-0.59 to 0.51)
losartan 100 mg vs losartan 0 mg	-0.01 (-0.56 to 0.53)

Plot




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-0.03 (-0.49 to 0.43)
155	losartan 25 mg	0.40 (-0.23 to 1.02)
155	losartan 50 mg	0.00 (-0.63 to 0.63)
155	losartan 100 mg	-0.33 (-0.77 to 0.11)
239	losartan 0 mg	0.03 (-0.48 to 0.53)
239	losartan 25 mg	0.34 (-0.54 to 1.23)
239	losartan 50 mg	-0.37 (-1.03 to 0.28)
239	losartan 100 mg	-0.61 (-1.06 to -0.17)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	0.43 (-0.51 to 1.37)
155	losartan 50 mg - losartan 0 mg	0.03 (-0.91 to 0.98)
155	losartan 100 mg - losartan 0 mg	-0.29 (-1.07 to 0.48)
239	losartan 25 mg - losartan 0 mg	0.31 (-0.92 to 1.54)
239	losartan 50 mg - losartan 0 mg	-0.40 (-1.40 to 0.60)
239	losartan 100 mg - losartan 0 mg	-0.64 (-1.46 to 0.17)

Plot



Quality of Life Outcomes

Model based estimates of change from baseline in QLQ-C30 and QLQ-BN20 scores at Days 155/239 are reported below for the newly diagnosed glioblastoma cohort. Note that even if the QoL outcomes were collected at Day 43 it is not possible to make a meaningful analysis of the effect of losartan as all participants received the treatment at that time point.

EORTC QLQ-C30 score

A total of 19 subjects were included in the analysis for this endpoint.


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-3.84 (-9.82 to 2.13)
155	losartan 25 mg	4.90 (-2.01 to 11.81)
155	losartan 50 mg	1.35 (-5.52 to 8.23)
155	losartan 100 mg	-4.54 (-9.34 to 0.27)
239	losartan 0 mg	4.16 (-1.82 to 10.13)
239	losartan 25 mg	8.78 (0.24 to 17.32)
239	losartan 50 mg	-2.31 (-9.19 to 4.56)
239	losartan 100 mg	-7.15 (-12.48 to -1.82)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	8.74 (-2.45 to 19.94)
155	losartan 50 mg - losartan 0 mg	5.20 (-5.77 to 16.16)
155	losartan 100 mg - losartan 0 mg	-0.69 (-10.04 to 8.65)
239	losartan 25 mg - losartan 0 mg	4.62 (-8.09 to 17.33)
239	losartan 50 mg - losartan 0 mg	-6.47 (-17.44 to 4.50)
239	losartan 100 mg - losartan 0 mg	-11.31 (-21.09 to -1.53)

Plot



EORTC QLQ-BN20 score

A total of 18 subjects were included in the analysis for this endpoint.


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-1.15 (-5.86 to 3.57)
155	losartan 25 mg	1.83 (-3.61 to 7.28)
155	losartan 50 mg	-2.92 (-8.53 to 2.69)
155	losartan 100 mg	-0.16 (-4.19 to 3.86)
239	losartan 0 mg	-1.15 (-5.86 to 3.57)
239	losartan 25 mg	5.03 (-1.25 to 11.30)
239	losartan 50 mg	-4.92 (-10.53 to 0.69)
239	losartan 100 mg	-2.94 (-7.18 to 1.31)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	2.98 (-5.72 to 11.69)
155	losartan 50 mg - losartan 0 mg	-1.77 (-10.76 to 7.22)
155	losartan 100 mg - losartan 0 mg	0.98 (-6.67 to 8.64)
239	losartan 25 mg - losartan 0 mg	6.17 (-3.28 to 15.63)
239	losartan 50 mg - losartan 0 mg	-3.77 (-12.76 to 5.22)
239	losartan 100 mg - losartan 0 mg	-1.79 (-9.62 to 6.04)

Plot



Steroid Use

Binary steroid use is analysed with longitudinal logistic regression and population-averaged marginal risk differences for early (Days 15/29/43) and late (Days 155/239) visits in the newly diagnosed glioblastoma cohort.

Steroid use

A total of 54 subjects were included in the analysis.


Days 15/29/43 (by dose)

Treatment	Risk (95% CI)
losartan 0 mg	0.223 (0.211 to 0.235)
losartan 25 mg	0.226 (0.221 to 0.230)
losartan 50 mg	0.227 (0.218 to 0.237)
losartan 100 mg	0.226 (0.222 to 0.230)

Risk differences vs 0 mg

Comparison	Risk difference (95% CI)
100 - 0	0.003 (-0.010 to 0.016)
25 - 0	0.003 (-0.010 to 0.016)
50 - 0	0.005 (-0.014 to 0.023)

Plot




Days 155/239 (population-averaged risk)

Visit	Treatment	Risk (95% CI)
155	losartan 0 mg	0.172 (-0.026 to 0.371)
239	losartan 0 mg	0.118 (-0.055 to 0.291)
155	losartan 25 mg	0.495 (0.129 to 0.862)
239	losartan 25 mg	0.000 (0.000 to 0.000)
155	losartan 50 mg	0.220 (-0.075 to 0.514)
239	losartan 50 mg	0.244 (-0.083 to 0.570)
155	losartan 100 mg	0.243 (0.030 to 0.456)
239	losartan 100 mg	0.284 (0.041 to 0.528)

Risk differences vs reference

Visit	Comparison	Risk difference (95% CI)
155	losartan 100 mg - losartan 0 mg	0.067 (-0.209 to 0.342)
155	losartan 25 mg - losartan 0 mg	0.294 (-0.088 to 0.675)
155	losartan 50 mg - losartan 0 mg	0.045 (-0.287 to 0.377)
239	losartan 100 mg - losartan 0 mg	0.162 (-0.126 to 0.450)
239	losartan 25 mg - losartan 0 mg	-0.142 (-0.319 to 0.036)
239	losartan 50 mg - losartan 0 mg	0.124 (-0.226 to 0.475)

Plot



RANO Responses

Ordinal RANO outcomes are analysed using proportional odds models within the configured cohort. For State of disease according to the RANO criteria, visits at Days 141 and 225 are included; for Best overall radiographic response, only Day 225 is included. Proportional odds ratios and confidence intervals are reported below.

State of disease according to the RANO criteria

Visit 141

Comparison	Propoprtional Odds Ratio (95% CI)
losartan 25 vs 0 mg	0.867 (0.135 to 5.052)
losartan 50 vs 0 mg	0.475 (0.056 to 2.952)
losartan 100 vs 0 mg	2.166 (0.560 to 8.883)

Visit 225

Comparison	Propoprtional Odds Ratio (95% CI)
losartan 25 vs 0 mg	4.000 (0.425 to 91.460)
losartan 50 vs 0 mg	0.750 (0.106 to 4.936)
losartan 100 vs 0 mg	6.500 (1.125 to 54.420)

Best overall radiographic response

Visit 225

Comparison	Propoprtional Odds Ratio (95% CI)
losartan 25 vs 0 mg	3.557 (0.329 to 53.896)
losartan 50 vs 0 mg	0.702 (0.090 to 5.863)
losartan 100 vs 0 mg	2.423 (0.416 to 16.369)

Biomarker Endpoints (Serum Cytokines / Angiogenic Factors)

Serum cytokine and angiogenic-factor concentrations from Bio-Plex 19-plex and 14-plex panels are analysed as exploratory biomarker endpoints. For each analyte, a linear mixed model with random subject intercept is fitted with change from Screening baseline as the response. Where the log-scale model fits better (lower AIC), the analysis is performed on the log-concentration scale and results are labelled accordingly.

Biomarker samples are collected at screening (baseline), Day 15 (Cycle 2), and Day 43 (Cycle 4) for the early analysis, and again at Cycle 17 (Day 225) and later for the long-term analysis. The early analysis uses the stepped-wedge treatment assignment; the long-term analysis uses the assigned long-term dose (`trt02p`). The early analyses results are presented first, then the long term.

A formal model selection and residual diagnostic assessment is available in a separate document (`impress_lb_model_fit.pdf`).

Fractalkine

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.021 (-0.275 to 0.232)
25 mg	-0.125 (-0.390 to 0.140)
50 mg	0.110 (-0.151 to 0.371)
100 mg	0.276 (0.018 to 0.534)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.10 (-0.58 to 0.37)
losartan 50 mg vs losartan 0 mg0	0.13 (-0.34 to 0.60)
losartan 100 mg vs losartan 0 mg0	0.30 (-0.17 to 0.76)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1464.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	218.6	TRUE

Plot



Granzyme B

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.127 (-0.315 to 0.061)
25 mg	-0.001 (-0.208 to 0.207)
50 mg	-0.364 (-0.578 to -0.151)
100 mg	-0.047 (-0.256 to 0.162)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.13 (-0.20 to 0.46)
losartan 50 mg vs losartan 0 mg0	-0.24 (-0.58 to 0.11)
losartan 100 mg vs losartan 0 mg0	0.08 (-0.26 to 0.42)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	823.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	156.1	TRUE

Plot



IFN-g

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.218 (-0.551 to 0.115)
25 mg	-0.121 (-0.485 to 0.243)
50 mg	-0.506 (-0.880 to -0.131)
100 mg	0.088 (-0.281 to 0.458)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.10 (-0.49 to 0.68)
losartan 50 mg vs losartan 0 mg0	-0.29 (-0.90 to 0.32)
losartan 100 mg vs losartan 0 mg0	0.31 (-0.31 to 0.92)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	625.1	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	267.2	TRUE

Plot



IL-1b

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.134 (-0.503 to 0.235)
25 mg	-0.035 (-0.449 to 0.379)
50 mg	-0.079 (-0.515 to 0.358)
100 mg	-0.014 (-0.440 to 0.412)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.10 (-0.51 to 0.71)
losartan 50 mg vs losartan 0 mg0	0.06 (-0.61 to 0.72)
losartan 100 mg vs losartan 0 mg0	0.12 (-0.54 to 0.78)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	514.8	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	288.5	TRUE

Plot



IL-6

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.016 (-0.530 to 0.561)
25 mg	-0.343 (-0.924 to 0.237)
50 mg	-0.304 (-0.893 to 0.286)
100 mg	-0.099 (-0.683 to 0.485)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.36 (-1.36 to 0.64)
losartan 50 mg vs losartan 0 mg0	-0.32 (-1.32 to 0.69)
losartan 100 mg vs losartan 0 mg0	-0.11 (-1.13 to 0.90)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	391.9	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	363.9	TRUE

Plot



IL-8

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.104 (-0.220 to 0.012)
25 mg	-0.005 (-0.135 to 0.125)
50 mg	-0.110 (-0.245 to 0.024)
100 mg	-0.067 (-0.199 to 0.064)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.10 (-0.10 to 0.30)
losartan 50 mg vs losartan 0 mg0	-0.01 (-0.22 to 0.21)
losartan 100 mg vs losartan 0 mg0	0.04 (-0.17 to 0.25)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	426	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	64	TRUE

Plot



MCP-1

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.068 (-0.302 to 0.166)
25 mg	-0.128 (-0.395 to 0.139)
50 mg	-0.263 (-0.535 to 0.009)
100 mg	0.007 (-0.263 to 0.277)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.06 (-0.46 to 0.34)
losartan 50 mg vs losartan 0 mg0	-0.19 (-0.62 to 0.23)
losartan 100 mg vs losartan 0 mg0	0.07 (-0.36 to 0.50)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1149.7	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	198.6	TRUE

Plot



MMP-10

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.025 (-0.069 to 0.119)
25 mg	0.151 (0.048 to 0.254)
50 mg	0.013 (-0.092 to 0.117)
100 mg	0.052 (-0.051 to 0.155)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.13 (-0.04 to 0.29)
losartan 50 mg vs losartan 0 mg0	-0.01 (-0.19 to 0.16)
losartan 100 mg vs losartan 0 mg0	0.03 (-0.15 to 0.20)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1253.0	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	21.9	TRUE

Plot



PDGF-DD

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.241 (-0.473 to -0.010)
25 mg	-0.271 (-0.530 to -0.012)
50 mg	-0.196 (-0.465 to 0.074)
100 mg	-0.400 (-0.664 to -0.137)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.03 (-0.43 to 0.37)
losartan 50 mg vs losartan 0 mg0	0.05 (-0.38 to 0.47)
losartan 100 mg vs losartan 0 mg0	-0.16 (-0.58 to 0.26)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1106.4	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	197.4	TRUE

Plot



PECAM-1

A total of 52 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.299 (-0.616 to 0.018)
25 mg	-0.261 (-0.594 to 0.071)
50 mg	-0.181 (-0.534 to 0.173)
100 mg	-0.184 (-0.510 to 0.142)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.04 (-0.55 to 0.62)
losartan 50 mg vs losartan 0 mg0	0.12 (-0.49 to 0.73)
losartan 100 mg vs losartan 0 mg0	0.11 (-0.46 to 0.69)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	2306.1	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	252.8	TRUE

Plot



TGF-alpha

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.181 (-0.138 to 0.500)
25 mg	0.079 (-0.271 to 0.428)
50 mg	-0.207 (-0.565 to 0.151)
100 mg	0.076 (-0.272 to 0.424)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.10 (-0.67 to 0.47)
losartan 50 mg vs losartan 0 mg0	-0.39 (-0.98 to 0.20)
losartan 100 mg vs losartan 0 mg0	-0.11 (-0.69 to 0.48)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	544.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	260.4	TRUE

Plot



Thrombospondin-2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.113 (-0.241 to 0.016)
25 mg	-0.012 (-0.156 to 0.133)
50 mg	-0.094 (-0.248 to 0.060)
100 mg	-0.122 (-0.272 to 0.027)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.10 (-0.11 to 0.31)
losartan 50 mg vs losartan 0 mg0	0.02 (-0.21 to 0.25)
losartan 100 mg vs losartan 0 mg0	-0.01 (-0.24 to 0.22)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1881.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	83.3	TRUE

Plot



VEGF-A

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.372 (-0.598 to -0.146)
25 mg	-0.351 (-0.603 to -0.099)
50 mg	-0.248 (-0.505 to 0.009)
100 mg	-0.436 (-0.689 to -0.184)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.02 (-0.38 to 0.42)
losartan 50 mg vs losartan 0 mg0	0.12 (-0.29 to 0.54)
losartan 100 mg vs losartan 0 mg0	-0.06 (-0.48 to 0.35)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1080.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	193.2	TRUE

Plot



VEGF-C

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.108 (-0.279 to 0.062)
25 mg	-0.080 (-0.278 to 0.119)
50 mg	-0.037 (-0.243 to 0.168)
100 mg	-0.248 (-0.448 to -0.048)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.03 (-0.26 to 0.31)
losartan 50 mg vs losartan 0 mg0	0.07 (-0.23 to 0.38)
losartan 100 mg vs losartan 0 mg0	-0.14 (-0.44 to 0.16)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1477.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	138.1	TRUE

Plot



Angiopoietin-1

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.144 (-0.377 to 0.088)
25 mg	-0.249 (-0.512 to 0.013)
50 mg	-0.081 (-0.341 to 0.179)
100 mg	-0.316 (-0.573 to -0.060)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.11 (-0.53 to 0.32)
losartan 50 mg vs losartan 0 mg0	0.06 (-0.37 to 0.49)
losartan 100 mg vs losartan 0 mg0	-0.17 (-0.60 to 0.26)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1920.0	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	198.4	TRUE

Plot



Angiopoietin-2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.014 (-0.095 to 0.067)
25 mg	-0.075 (-0.164 to 0.014)
50 mg	-0.055 (-0.146 to 0.037)
100 mg	-0.037 (-0.125 to 0.052)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.06 (-0.21 to 0.08)
losartan 50 mg vs losartan 0 mg0	-0.04 (-0.19 to 0.11)
losartan 100 mg vs losartan 0 mg0	-0.02 (-0.17 to 0.13)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1605.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	-6.1	TRUE

Plot



ANGPTL4

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.064 (-0.163 to 0.035)
25 mg	-0.047 (-0.156 to 0.062)
50 mg	-0.031 (-0.144 to 0.083)
100 mg	-0.028 (-0.138 to 0.083)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.02 (-0.16 to 0.19)
losartan 50 mg vs losartan 0 mg0	0.03 (-0.15 to 0.22)
losartan 100 mg vs losartan 0 mg0	0.04 (-0.14 to 0.22)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	2350.7	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	32.5	TRUE

Plot



CA9

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.062 (-0.217 to 0.341)
25 mg	0.043 (-0.266 to 0.353)
50 mg	0.016 (-0.307 to 0.340)
100 mg	0.128 (-0.188 to 0.444)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.02 (-0.50 to 0.46)
losartan 50 mg vs losartan 0 mg0	-0.05 (-0.55 to 0.46)
losartan 100 mg vs losartan 0 mg0	0.07 (-0.44 to 0.57)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1075	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	234	TRUE

Plot



CD163

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.074 (-0.019 to 0.167)
25 mg	0.046 (-0.057 to 0.149)
50 mg	0.165 (0.058 to 0.273)
100 mg	0.102 (-0.003 to 0.207)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.03 (-0.19 to 0.13)
losartan 50 mg vs losartan 0 mg0	0.09 (-0.08 to 0.26)
losartan 100 mg vs losartan 0 mg0	0.03 (-0.14 to 0.20)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	2658.8	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	21.1	TRUE

Plot



Collagen IV alpha 1

A total of 52 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.073 (-0.017 to 0.163)
25 mg	0.061 (-0.037 to 0.160)
50 mg	0.072 (-0.033 to 0.177)
100 mg	0.143 (0.042 to 0.243)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.01 (-0.17 to 0.14)
losartan 50 mg vs losartan 0 mg0	0.00 (-0.17 to 0.16)
losartan 100 mg vs losartan 0 mg0	0.07 (-0.09 to 0.23)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1325.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	11.9	TRUE

Plot



EGF

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.901 (-1.691 to -0.112)
25 mg	-0.598 (-1.464 to 0.267)
50 mg	-0.689 (-1.583 to 0.205)
100 mg	-0.694 (-1.568 to 0.181)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.30 (-1.09 to 1.70)
losartan 50 mg vs losartan 0 mg0	0.21 (-1.24 to 1.66)
losartan 100 mg vs losartan 0 mg0	0.21 (-1.23 to 1.64)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	872.2	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	434.3	TRUE

Plot



ErbB2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.052 (-0.028 to 0.132)
25 mg	-0.039 (-0.130 to 0.052)
50 mg	0.083 (-0.012 to 0.179)
100 mg	0.002 (-0.090 to 0.095)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.09 (-0.23 to 0.04)
losartan 50 mg vs losartan 0 mg0	0.03 (-0.11 to 0.18)
losartan 100 mg vs losartan 0 mg0	-0.05 (-0.19 to 0.09)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1701.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	-7.7	TRUE

Plot



FGF Basic

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.280 (-0.525 to -0.034)
25 mg	-0.281 (-0.554 to -0.008)
50 mg	-0.100 (-0.383 to 0.183)
100 mg	-0.251 (-0.532 to 0.030)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.00 (-0.43 to 0.43)
losartan 50 mg vs losartan 0 mg0	0.18 (-0.27 to 0.63)
losartan 100 mg vs losartan 0 mg0	0.03 (-0.42 to 0.48)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	648.4	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	208.9	TRUE

Plot



IL-12 p70

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.269 (-0.028 to 0.567)
25 mg	-0.140 (-0.474 to 0.194)
50 mg	0.255 (-0.103 to 0.613)
100 mg	0.149 (-0.204 to 0.501)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.41 (-0.84 to 0.02)
losartan 50 mg vs losartan 0 mg0	-0.01 (-0.50 to 0.47)
losartan 100 mg vs losartan 0 mg0	-0.12 (-0.61 to 0.37)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1273.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	243.4	TRUE

Plot



IL-2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.114 (-0.658 to 0.431)
25 mg	-0.340 (-0.926 to 0.246)
50 mg	0.053 (-0.542 to 0.647)
100 mg	-0.096 (-0.678 to 0.486)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.23 (-1.22 to 0.76)
losartan 50 mg vs losartan 0 mg0	0.17 (-0.84 to 1.17)
losartan 100 mg vs losartan 0 mg0	0.02 (-0.98 to 1.01)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	787.2	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	363.0	TRUE

Plot



IP-10 (CXCL10)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.182 (-0.025 to 0.390)
25 mg	0.235 (0.008 to 0.461)
50 mg	0.150 (-0.084 to 0.384)
100 mg	0.273 (0.044 to 0.502)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.05 (-0.31 to 0.42)
losartan 50 mg vs losartan 0 mg0	-0.03 (-0.41 to 0.35)
losartan 100 mg vs losartan 0 mg0	0.09 (-0.29 to 0.47)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1152.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	174.7	TRUE

Plot



I-TAC (CXCL11)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.028 (-0.129 to 0.072)
25 mg	-0.026 (-0.140 to 0.088)
50 mg	-0.024 (-0.145 to 0.097)
100 mg	0.003 (-0.115 to 0.121)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.00 (-0.15 to 0.16)
losartan 50 mg vs losartan 0 mg0	0.00 (-0.17 to 0.18)
losartan 100 mg vs losartan 0 mg0	0.03 (-0.14 to 0.20)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1174.0	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	34.1	TRUE

Plot



MIG (CXCL9)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.326 (-0.144 to 0.796)
25 mg	-0.298 (-0.821 to 0.224)
50 mg	-0.114 (-0.662 to 0.434)
100 mg	0.212 (-0.326 to 0.749)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.62 (-1.42 to 0.17)
losartan 50 mg vs losartan 0 mg0	-0.44 (-1.29 to 0.41)
losartan 100 mg vs losartan 0 mg0	-0.11 (-0.96 to 0.73)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1450.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	334.5	TRUE

Plot



PIGF

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.069 (-0.003 to 0.142)
25 mg	0.063 (-0.018 to 0.144)
50 mg	0.037 (-0.048 to 0.122)
100 mg	0.054 (-0.030 to 0.137)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.01 (-0.13 to 0.12)
losartan 50 mg vs losartan 0 mg0	-0.03 (-0.16 to 0.10)
losartan 100 mg vs losartan 0 mg0	-0.02 (-0.15 to 0.12)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	790.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	-26.0	TRUE

Plot



Tenascin C

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.022 (-0.025 to 0.069)
25 mg	0.068 (0.016 to 0.120)
50 mg	0.023 (-0.031 to 0.078)
100 mg	0.028 (-0.027 to 0.083)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.05 (-0.03 to 0.13)
losartan 50 mg vs losartan 0 mg0	0.00 (-0.08 to 0.09)
losartan 100 mg vs losartan 0 mg0	0.01 (-0.08 to 0.09)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1754.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	-113.2	TRUE

Plot



TNFRSF18 (GITR)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.027 (-0.289 to 0.235)
25 mg	-0.238 (-0.531 to 0.054)
50 mg	0.018 (-0.290 to 0.325)
100 mg	-0.101 (-0.401 to 0.200)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.21 (-0.65 to 0.23)
losartan 50 mg vs losartan 0 mg0	0.05 (-0.43 to 0.52)
losartan 100 mg vs losartan 0 mg0	-0.07 (-0.54 to 0.40)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1513.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	222.2	TRUE

Plot



TNFSF6 (FasL)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.024 (-0.143 to 0.095)
25 mg	-0.037 (-0.168 to 0.094)
50 mg	-0.022 (-0.160 to 0.116)
100 mg	-0.039 (-0.175 to 0.097)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.01 (-0.21 to 0.19)
losartan 50 mg vs losartan 0 mg0	0.00 (-0.21 to 0.22)
losartan 100 mg vs losartan 0 mg0	-0.01 (-0.23 to 0.20)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	878.8	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	66.4	TRUE

Plot



VEGFR2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.005 (-0.049 to 0.060)
25 mg	-0.013 (-0.074 to 0.049)
50 mg	0.059 (-0.007 to 0.126)
100 mg	0.060 (-0.004 to 0.124)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.02 (-0.10 to 0.07)
losartan 50 mg vs losartan 0 mg0	0.05 (-0.04 to 0.15)
losartan 100 mg vs losartan 0 mg0	0.05 (-0.04 to 0.15)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1749.2	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	-84.0	TRUE

Plot



Biomarker Endpoints — Long-term (Cycle 15+)

Model-based estimates of change from baseline at Cycle 15 and later are reported below for analytes with available long-term samples. The same scale (linear or log-transformed) as selected for the early analysis is used for each analyte.

Fractalkine

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.036 (-0.594 to 0.667)
losartan 25 mg	0.533 (-0.629 to 1.695)
losartan 50 mg	0.190 (-0.558 to 0.937)
losartan 100 mg	0.469 (-0.065 to 1.003)

Differences vs 0 mg by long-term dose


Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.50 (-1.09 to 2.09)
losartan 50 mg - losartan 0 mg	0.15 (-1.04 to 1.35)
losartan 100 mg - losartan 0 mg	0.43 (-0.59 to 1.45)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	519.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	100.6	TRUE

Plot

Fractalkine



Granzyme B

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-1.200 (-1.967 to -0.433)
losartan 25 mg	-0.755 (-2.161 to 0.651)
losartan 50 mg	-0.333 (-1.239 to 0.574)
losartan 100 mg	-0.216 (-0.857 to 0.424)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.45 (-1.53 to 2.42)
losartan 50 mg - losartan 0 mg	0.87 (-0.58 to 2.32)
losartan 100 mg - losartan 0 mg	0.98 (-0.24 to 2.20)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	238.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	113.9	TRUE

Plot



IFN-g

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.132 (-0.576 to 0.312)
losartan 25 mg	-1.425 (-2.220 to -0.630)
losartan 50 mg	0.004 (-0.516 to 0.525)
losartan 100 mg	-0.229 (-0.597 to 0.138)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-1.29 (-2.41 to -0.17)
losartan 50 mg - losartan 0 mg	0.14 (-0.71 to 0.98)
losartan 100 mg - losartan 0 mg	-0.10 (-0.80 to 0.61)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	107.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	75.9	TRUE

Plot



IL-1b

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.417 (0.010 to 0.825)
losartan 25 mg	-0.514 (-1.279 to 0.250)
losartan 50 mg	-0.046 (-0.541 to 0.449)
losartan 100 mg	-0.022 (-0.365 to 0.321)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.93 (-1.97 to 0.11)
losartan 50 mg - losartan 0 mg	-0.46 (-1.25 to 0.33)
losartan 100 mg - losartan 0 mg	-0.44 (-1.09 to 0.21)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	101.9	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	71.1	TRUE

Plot



IL-6

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.065 (-0.839 to 0.710)
losartan 25 mg	-1.253 (-2.665 to 0.160)
losartan 50 mg	-0.491 (-1.436 to 0.454)
losartan 100 mg	-0.283 (-0.948 to 0.383)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-1.19 (-3.16 to 0.78)
losartan 50 mg - losartan 0 mg	-0.43 (-1.90 to 1.05)
losartan 100 mg - losartan 0 mg	-0.22 (-1.47 to 1.04)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	104.9	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	115.0	TRUE

Plot



IL-8

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.217 (-0.785 to 0.350)
losartan 25 mg	0.672 (-0.376 to 1.719)
losartan 50 mg	0.060 (-0.611 to 0.732)
losartan 100 mg	0.041 (-0.434 to 0.516)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.89 (-0.58 to 2.36)
losartan 50 mg - losartan 0 mg	0.28 (-0.80 to 1.35)
losartan 100 mg - losartan 0 mg	0.26 (-0.64 to 1.16)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	157.0	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	93.5	TRUE

Plot



MCP-1

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.048 (-0.251 to 0.154)
losartan 25 mg	-0.177 (-0.565 to 0.211)
losartan 50 mg	0.123 (-0.118 to 0.364)
losartan 100 mg	-0.067 (-0.238 to 0.104)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.13 (-0.67 to 0.41)
losartan 50 mg - losartan 0 mg	0.17 (-0.21 to 0.55)
losartan 100 mg - losartan 0 mg	-0.02 (-0.34 to 0.30)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	366.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	23.9	TRUE

Plot



MMP-10

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.037 (-0.170 to 0.245)
losartan 25 mg	0.137 (-0.247 to 0.521)
losartan 50 mg	-0.021 (-0.266 to 0.223)
losartan 100 mg	-0.002 (-0.177 to 0.173)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.10 (-0.42 to 0.62)
losartan 50 mg - losartan 0 mg	-0.06 (-0.45 to 0.34)
losartan 100 mg - losartan 0 mg	-0.04 (-0.38 to 0.30)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	440.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	24.3	TRUE

Plot



PDGF-DD

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.040 (-0.323 to 0.403)
losartan 25 mg	-0.574 (-1.242 to 0.095)
losartan 50 mg	-0.245 (-0.679 to 0.190)
losartan 100 mg	-0.332 (-0.639 to -0.025)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.61 (-1.54 to 0.31)
losartan 50 mg - losartan 0 mg	-0.29 (-0.98 to 0.41)
losartan 100 mg - losartan 0 mg	-0.37 (-0.95 to 0.21)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	358.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	63.7	TRUE

Plot



PECAM-1

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.264 (-0.274 to 0.803)
losartan 25 mg	-0.319 (-1.296 to 0.659)
losartan 50 mg	-0.088 (-0.726 to 0.549)
losartan 100 mg	0.165 (-0.285 to 0.615)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.58 (-1.95 to 0.79)
losartan 50 mg - losartan 0 mg	-0.35 (-1.37 to 0.67)
losartan 100 mg - losartan 0 mg	-0.10 (-0.96 to 0.76)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	832.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	89.8	TRUE

Plot



TGF-alpha

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.334 (-0.659 to -0.009)
losartan 25 mg	-0.508 (-1.101 to 0.085)
losartan 50 mg	0.070 (-0.324 to 0.465)
losartan 100 mg	-0.040 (-0.316 to 0.236)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.17 (-1.00 to 0.65)
losartan 50 mg - losartan 0 mg	0.40 (-0.22 to 1.03)
losartan 100 mg - losartan 0 mg	0.29 (-0.23 to 0.81)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	139.8	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	56.3	TRUE

Plot



Thrombospondin-2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.154 (-0.351 to 0.043)
losartan 25 mg	-0.078 (-0.447 to 0.290)
losartan 50 mg	-0.128 (-0.370 to 0.113)
losartan 100 mg	0.033 (-0.132 to 0.199)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.08 (-0.43 to 0.58)
losartan 50 mg - losartan 0 mg	0.03 (-0.36 to 0.41)
losartan 100 mg - losartan 0 mg	0.19 (-0.13 to 0.50)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	659.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	21.7	TRUE

Plot



VEGF-A

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.036 (-0.437 to 0.365)
losartan 25 mg	-0.562 (-1.290 to 0.167)
losartan 50 mg	-0.060 (-0.535 to 0.415)
losartan 100 mg	-0.359 (-0.698 to -0.020)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.53 (-1.53 to 0.48)
losartan 50 mg - losartan 0 mg	-0.02 (-0.78 to 0.74)
losartan 100 mg - losartan 0 mg	-0.32 (-0.97 to 0.32)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	381.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	69.9	TRUE

Plot



VEGF-C

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.020 (-0.343 to 0.304)
losartan 25 mg	-0.324 (-0.929 to 0.282)
losartan 50 mg	-0.157 (-0.545 to 0.232)
losartan 100 mg	-0.267 (-0.540 to 0.006)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.30 (-1.14 to 0.53)
losartan 50 mg - losartan 0 mg	-0.14 (-0.75 to 0.48)
losartan 100 mg - losartan 0 mg	-0.25 (-0.76 to 0.27)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	497.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	55.9	TRUE

Plot



Angiopoietin-1

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.204 (-0.486 to 0.077)
losartan 25 mg	-0.715 (-1.242 to -0.189)
losartan 50 mg	-0.308 (-0.645 to 0.028)
losartan 100 mg	-0.361 (-0.600 to -0.123)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.51 (-1.24 to 0.22)
losartan 50 mg - losartan 0 mg	-0.10 (-0.64 to 0.43)
losartan 100 mg - losartan 0 mg	-0.16 (-0.61 to 0.29)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	627.7	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	46.6	TRUE

Plot



Angiopoietin-2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.035 (-0.090 to 0.159)
losartan 25 mg	-0.152 (-0.378 to 0.073)
losartan 50 mg	0.031 (-0.117 to 0.178)
losartan 100 mg	0.055 (-0.048 to 0.158)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.19 (-0.51 to 0.13)
losartan 50 mg - losartan 0 mg	0.00 (-0.24 to 0.24)
losartan 100 mg - losartan 0 mg	0.02 (-0.18 to 0.22)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	552.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-10.4	TRUE

Plot



ANGPTL4

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.023 (-0.175 to 0.128)
losartan 25 mg	0.020 (-0.246 to 0.286)
losartan 50 mg	-0.118 (-0.295 to 0.059)
losartan 100 mg	0.030 (-0.094 to 0.154)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.04 (-0.33 to 0.41)
losartan 50 mg - losartan 0 mg	-0.09 (-0.39 to 0.20)
losartan 100 mg - losartan 0 mg	0.05 (-0.19 to 0.30)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	822.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	1.7	TRUE

Plot



CA9

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.307 (-0.353 to 0.966)
losartan 25 mg	0.424 (-0.698 to 1.546)
losartan 50 mg	-0.537 (-1.329 to 0.255)
losartan 100 mg	0.131 (-0.389 to 0.651)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.12 (-1.45 to 1.69)
losartan 50 mg - losartan 0 mg	-0.84 (-2.18 to 0.50)
losartan 100 mg - losartan 0 mg	-0.18 (-1.21 to 0.86)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	354.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	99.5	TRUE

Plot



CD163

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.013 (-0.177 to 0.202)
losartan 25 mg	-0.078 (-0.420 to 0.263)
losartan 50 mg	0.240 (0.016 to 0.464)
losartan 100 mg	0.124 (-0.035 to 0.283)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.09 (-0.57 to 0.39)
losartan 50 mg - losartan 0 mg	0.23 (-0.13 to 0.58)
losartan 100 mg - losartan 0 mg	0.11 (-0.19 to 0.42)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	968.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	18.7	TRUE

Plot



Collagen IV alpha 1

A total of 33 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.181 (0.017 to 0.346)
losartan 25 mg	0.021 (-0.269 to 0.312)
losartan 50 mg	0.033 (-0.154 to 0.220)
losartan 100 mg	0.105 (-0.027 to 0.236)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.16 (-0.57 to 0.25)
losartan 50 mg - losartan 0 mg	-0.15 (-0.45 to 0.15)
losartan 100 mg - losartan 0 mg	-0.08 (-0.33 to 0.18)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	459.6	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	6.3	TRUE

Plot



EGF

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.144 (-0.408 to 0.696)
losartan 25 mg	-1.185 (-2.192 to -0.178)
losartan 50 mg	-0.429 (-1.088 to 0.231)
losartan 100 mg	-0.496 (-0.962 to -0.030)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-1.33 (-2.73 to 0.07)
losartan 50 mg - losartan 0 mg	-0.57 (-1.62 to 0.48)
losartan 100 mg - losartan 0 mg	-0.64 (-1.52 to 0.24)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	306.0	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	92.3	TRUE

Plot



ErbB2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.084 (-0.060 to 0.227)
losartan 25 mg	0.002 (-0.263 to 0.267)
losartan 50 mg	0.097 (-0.081 to 0.276)
losartan 100 mg	0.059 (-0.064 to 0.182)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.08 (-0.45 to 0.29)
losartan 50 mg - losartan 0 mg	0.01 (-0.26 to 0.29)
losartan 100 mg - losartan 0 mg	-0.02 (-0.26 to 0.21)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	606.0	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	0.8	TRUE

Plot



FGF Basic

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.095 (-0.353 to 0.542)
losartan 25 mg	0.042 (-0.800 to 0.884)
losartan 50 mg	0.146 (-0.389 to 0.680)
losartan 100 mg	-0.059 (-0.444 to 0.325)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.05 (-1.21 to 1.10)
losartan 50 mg - losartan 0 mg	0.05 (-0.80 to 0.90)
losartan 100 mg - losartan 0 mg	-0.15 (-0.88 to 0.57)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	274.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	77.9	TRUE

Plot



IL-12 p70

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.170 (-0.921 to 0.581)
losartan 25 mg	0.560 (-0.949 to 2.070)
losartan 50 mg	0.536 (-0.345 to 1.417)
losartan 100 mg	-0.004 (-0.628 to 0.620)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.73 (-1.41 to 2.87)
losartan 50 mg - losartan 0 mg	0.71 (-0.72 to 2.13)
losartan 100 mg - losartan 0 mg	0.17 (-1.01 to 1.34)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	442.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	111.5	TRUE

Plot



IL-2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.111 (-1.002 to 0.780)
losartan 25 mg	-1.271 (-2.907 to 0.365)
losartan 50 mg	-0.164 (-1.244 to 0.916)
losartan 100 mg	0.171 (-0.577 to 0.920)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-1.16 (-3.41 to 1.09)
losartan 50 mg - losartan 0 mg	-0.05 (-1.78 to 1.67)
losartan 100 mg - losartan 0 mg	0.28 (-1.14 to 1.70)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	350.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	124.4	TRUE

Plot



IP-10 (CXCL10)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.021 (-0.224 to 0.267)
losartan 25 mg	-0.005 (-0.468 to 0.459)
losartan 50 mg	0.080 (-0.213 to 0.373)
losartan 100 mg	0.150 (-0.058 to 0.358)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.03 (-0.67 to 0.62)
losartan 50 mg - losartan 0 mg	0.06 (-0.41 to 0.53)
losartan 100 mg - losartan 0 mg	0.13 (-0.26 to 0.52)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	338.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	37.0	TRUE

Plot



I-TAC (CXCL11)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.020 (-0.234 to 0.195)
losartan 25 mg	-0.330 (-0.723 to 0.062)
losartan 50 mg	-0.087 (-0.346 to 0.172)
losartan 100 mg	-0.129 (-0.311 to 0.053)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.31 (-0.86 to 0.23)
losartan 50 mg - losartan 0 mg	-0.07 (-0.48 to 0.34)
losartan 100 mg - losartan 0 mg	-0.11 (-0.45 to 0.23)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	421.8	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	28.1	TRUE

Plot



MIG (CXCL9)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.311 (-0.848 to 0.225)
losartan 25 mg	-0.897 (-1.869 to 0.075)
losartan 50 mg	0.284 (-0.365 to 0.933)
losartan 100 mg	0.076 (-0.374 to 0.526)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.59 (-1.94 to 0.77)
losartan 50 mg - losartan 0 mg	0.60 (-0.44 to 1.64)
losartan 100 mg - losartan 0 mg	0.39 (-0.46 to 1.24)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	490.9	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	89.7	TRUE

Plot



PIGF

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.063 (-0.073 to 0.199)
losartan 25 mg	-0.148 (-0.395 to 0.100)
losartan 50 mg	0.002 (-0.161 to 0.165)
losartan 100 mg	0.149 (0.033 to 0.264)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.21 (-0.56 to 0.13)
losartan 50 mg - losartan 0 mg	-0.06 (-0.32 to 0.20)
losartan 100 mg - losartan 0 mg	0.09 (-0.13 to 0.30)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	279.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-3.2	TRUE

Plot



Tenascin C

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.034 (-0.129 to 0.061)
losartan 25 mg	-0.038 (-0.205 to 0.128)
losartan 50 mg	0.016 (-0.094 to 0.125)
losartan 100 mg	0.020 (-0.062 to 0.102)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.00 (-0.24 to 0.23)
losartan 50 mg - losartan 0 mg	0.05 (-0.12 to 0.22)
losartan 100 mg - losartan 0 mg	0.05 (-0.11 to 0.21)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	622.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-30.3	TRUE

Plot



TNFRSF18 (GITR)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-1.018 (-1.862 to -0.173)
losartan 25 mg	-1.984 (-3.516 to -0.452)
losartan 50 mg	-0.060 (-1.075 to 0.955)
losartan 100 mg	-0.080 (-0.790 to 0.629)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.97 (-3.10 to 1.17)
losartan 50 mg - losartan 0 mg	0.96 (-0.67 to 2.58)
losartan 100 mg - losartan 0 mg	0.94 (-0.41 to 2.28)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	500.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	120.8	TRUE

Plot



TNFSF6 (FasL)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.081 (-0.252 to 0.091)
losartan 25 mg	-0.182 (-0.493 to 0.128)
losartan 50 mg	0.081 (-0.122 to 0.284)
losartan 100 mg	-0.127 (-0.271 to 0.017)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.10 (-0.53 to 0.33)
losartan 50 mg - losartan 0 mg	0.16 (-0.16 to 0.49)
losartan 100 mg - losartan 0 mg	-0.05 (-0.32 to 0.23)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	281.9	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	12.1	TRUE

Plot



VEGFR2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.013 (-0.136 to 0.111)
losartan 25 mg	-0.074 (-0.298 to 0.150)
losartan 50 mg	0.052 (-0.095 to 0.198)
losartan 100 mg	0.041 (-0.064 to 0.146)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.06 (-0.37 to 0.25)
losartan 50 mg - losartan 0 mg	0.06 (-0.17 to 0.30)
losartan 100 mg - losartan 0 mg	0.05 (-0.15 to 0.25)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	646.0	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-10.2	TRUE

Plot




Time-to-Event Endpoints

Kaplan-Meier curves are shown with the at-risk table below. Overall log-rank p-values are reported for each endpoint, followed by Cox regression hazard ratios versus 0 mg.

The proportional-hazards assumption underlying the Cox hazard ratios is assessed with standard diagnostics (scaled Schoenfeld residual tests and plots, and complementary log-log survival plots) in a separate document ([impress_tte_ph_check.pdf](#)).

Progression-free survival



Number at risk

Dose	0	250	500	750	1000	1250
0 mg	18	10	5	2	1	1
25 mg	9	3	1	0	0	0
50 mg	9	4	1	0	0	0
100 mg	18	13	7	2	1	0

Days since randomisation


Log-rank p-value

Overall log-rank p-value: 0.213

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	1.64 (0.69 to 3.88)	0.260
50 mg vs 0 mg	1.28 (0.54 to 3.03)	0.575
100 mg vs 0 mg	0.68 (0.33 to 1.40)	0.296

6-month progression-free survival




Log-rank p-value

Overall log-rank p-value: 0.305

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	1.61 (0.36 to 7.20)	0.533
50 mg vs 0 mg	0.56 (0.06 to 5.01)	0.603
100 mg vs 0 mg	0.24 (0.03 to 2.16)	0.203

Overall survival



Number at risk

Dose	0	250	500	750	1000
0 mg	18	18	8	2	0
25 mg	9	6	2	0	0
50 mg	9	8	4	2	0
100 mg	18	17	11	4	1

Days since randomisation


Log-rank p-value

Overall log-rank p-value: 0.075

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	2.07 (0.84 to 5.10)	0.112
50 mg vs 0 mg	0.80 (0.28 to 2.26)	0.669
100 mg vs 0 mg	0.63 (0.26 to 1.49)	0.289

Overall survival at 12 months




Log-rank p-value

Overall log-rank p-value: 0.171

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	1.99 (0.53 to 7.41)	0.307
50 mg vs 0 mg	0.44 (0.05 to 3.80)	0.458
100 mg vs 0 mg	0.39 (0.08 to 2.00)	0.258

Overall survival at 24 months



Log-rank p-value

Overall log-rank p-value: 0.075

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	2.07 (0.84 to 5.10)	0.112
50 mg vs 0 mg	0.80 (0.28 to 2.26)	0.669
100 mg vs 0 mg	0.63 (0.26 to 1.49)	0.289

Safety

Safety analyses follow SAP §7 and are descriptive/exploratory only, by treatment dose group and over time, with no formal hypothesis testing. The Safety Set is nominally all participants who received at least one dose of losartan, analysed by treatment actually received. **In this sham mock-up the dosing export (`_EX.csv`) is empty, so the Safety Set is approximated by the Full Analysis Set and all safety outputs are grouped by planned dose.** Assessment by treatment actually received will be added at database lock and unblinding. Conventional laboratory parameters are limited to creatinine, the only safety-lab assay populated in the current export.

Extent of Exposure

N/% exposed per planned dose level, treatment duration overall and by long-term dose, short-term vs continued treatment, dose switches / early discontinuation, and duration categories, for the newly diagnosed glioblastoma cohort.

A total of 54 subjects are included.

Subjects exposed per planned dose level

Dose (mg)	Short-term (randomised)	Long-term (maintenance)
0	0 (0.0%)	18 (33.3%)
25	18 (33.3%)	9 (16.7%)
50	18 (33.3%)	9 (16.7%)
100	18 (33.3%)	18 (33.3%)

Treatment duration (overall)

Group	N	Median (range), days	Mean (SD), days
Overall	54	450 (15-1065)	471.2 (256.8)

Treatment duration by long-term dose

Long-term dose (mg)	N	Median (range), days	Mean (SD), days
0	18	439 (40-786)	441.4 (265.0)
25	9	361 (60-736)	326.0 (207.5)
50	9	483 (15-772)	484.4 (259.1)
100	18	620 (111-1065)	567.1 (248.1)

Treatment continuation and changes

Measure	N (%)
Continued to long-term treatment	52 (96.3%)
Short-term assignment only	2 (3.7%)
Switched dose (short- vs long-term)	18 (33.3%)
Discontinued treatment early	5 (9.3%)

Duration categories

Duration category	N (%)
≤ 14 days	0 (0.0%)
$> 14 - \leq 42$ days	2 (3.7%)
> 42 days	52 (96.3%)

Adverse Events

Treatment-emergent adverse event (TEAE) incidence (subjects with at least one event), coded with MedDRA, split by onset into an **early** phase (onset Day ≤ 44) and a **late** phase (onset Day ≥ 45). Early-phase events are grouped by the stepped-wedge dose in effect at onset (a 0 mg group captures events before a subject's randomised step started; columns are not mutually exclusive, and each denominator is the number of subjects ever at that dose during Days 1-44). Late-phase events are grouped by the long-term planned dose (denominator = subjects per dose in the analysis set).

Early phase (onset Day ≤ 44 , by stepped-wedge dose)

A total of 17 subjects had at least one TEAE in this phase.

Overall summary

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Subjects in analysis set	N=54	N=18	N=18	N=18	N=54
Subjects with ≥ 1 TEAE	7 (13.0%)	5 (27.8%)	2 (11.1%)	4 (22.2%)	17 (31.5%)
with ≥ 1 serious TEAE	5 (9.3%)	3 (16.7%)	1 (5.6%)	2 (11.1%)	11 (20.4%)
with ≥ 1 treatment-related TEAE	1 (1.9%)	1 (5.6%)	0 (0.0%)	1 (5.6%)	3 (5.6%)
with ≥ 1 TEAE leading to discontinuation	0 (0.0%)	1 (5.6%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
with ≥ 1 fatal TEAE	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)

Subjects by maximum severity

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Mild	1 (1.9%)	3 (16.7%)	1 (5.6%)	1 (5.6%)	6 (11.1%)
Moderate	3 (5.6%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	4 (7.4%)
Severe	3 (5.6%)	2 (11.1%)	1 (5.6%)	2 (11.1%)	8 (14.8%)
Life-threatening	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Death	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)

Subjects by relationship to treatment

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Unrelated	4 (7.4%)	4 (22.2%)	2 (11.1%)	2 (11.1%)	12 (22.2%)
Probable related	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (5.6%)	2 (3.7%)
Not applicable	2 (3.7%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (3.7%)
Possibly related	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Unlikely related	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)

Incidence by system organ class and preferred term

SOC / Preferred term	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Endocrine disorders	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Inappropriate antidiuretic hormone secretion	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Eye disorders	0 (0.0%)	1 (5.6%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Retinal detachment	0 (0.0%)	1 (5.6%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Gastrointestinal disorders	0 (0.0%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (1.9%)
Lower gastrointestinal haemorrhage	0 (0.0%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (1.9%)
Infections and infestations	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Wound infection	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Investigations	1 (1.9%)	3 (16.7%)	1 (5.6%)	1 (5.6%)	5 (9.3%)
Blood creatinine increased	0 (0.0%)	1 (5.6%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Neutrophil count decreased	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Platelet count decreased	1 (1.9%)	2 (11.1%)	1 (5.6%)	0 (0.0%)	4 (7.4%)
Nervous system disorders	4 (7.4%)	1 (5.6%)	0 (0.0%)	1 (5.6%)	6 (11.1%)
Brain oedema	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Dizziness	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Epilepsy	1 (1.9%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	2 (3.7%)
Headache	1 (1.9%)	1 (5.6%)	0 (0.0%)	0 (0.0%)	2 (3.7%)
Social circumstances	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Loss of personal independence in daily activities	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Vascular disorders	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Hypotension	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)

Late phase (onset Day $\geq$ 45, by long-term dose)

A total of 21 subjects had at least one TEAE in this phase.

Overall summary

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Subjects in analysis set	N=18	N=9	N=9	N=18	N=54
Subjects with $\geq$ 1 TEAE	5 (27.8%)	4 (44.4%)	5 (55.6%)	7 (38.9%)	21 (38.9%)
with $\geq$ 1 serious TEAE	5 (27.8%)	4 (44.4%)	3 (33.3%)	4 (22.2%)	16 (29.6%)
with $\geq$ 1 treatment-related TEAE	1 (5.6%)	0 (0.0%)	1 (11.1%)	3 (16.7%)	5 (9.3%)
with $\geq$ 1 TEAE leading to discontinuation	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
with $\geq$ 1 fatal TEAE	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)

Subjects by maximum severity

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Mild	2 (11.1%)	0 (0.0%)	1 (11.1%)	4 (22.2%)	7 (13.0%)
Moderate	1 (5.6%)	1 (11.1%)	0 (0.0%)	1 (5.6%)	3 (5.6%)
Severe	2 (11.1%)	2 (22.2%)	4 (44.4%)	2 (11.1%)	10 (18.5%)
Life-threatening	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Death	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)

Subjects by relationship to treatment

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Definite	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Unrelated	4 (22.2%)	4 (44.4%)	5 (55.6%)	5 (27.8%)	18 (33.3%)
Probable related	1 (5.6%)	0 (0.0%)	1 (11.1%)	1 (5.6%)	3 (5.6%)
Possibly related	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Unlikely related	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)

Incidence by system organ class and preferred term

SOC / Preferred term	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
General disorders and administration site conditions	2 (11.1%)	1 (11.1%)	0 (0.0%)	1 (5.6%)	4 (7.4%)
Gait disturbance	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Pain	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Pyrexia	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Sudden death	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Infections and infestations	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Wound infection	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)

SOC / Preferred term	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Investigations	0 (0.0%)	1 (11.1%)	3 (33.3%)	1 (5.6%)	5 (9.3%)
Alanine aminotransferase increased	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)
Gamma-glutamyltransferase increased	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Neutrophil count decreased	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)
Platelet count decreased	0 (0.0%)	0 (0.0%)	1 (11.1%)	1 (5.6%)	2 (3.7%)
Nervous system disorders	4 (22.2%)	3 (33.3%)	3 (33.3%)	5 (27.8%)	15 (27.8%)
Ataxia	1 (5.6%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	2 (3.7%)
Dizziness	0 (0.0%)	0 (0.0%)	1 (11.1%)	2 (11.1%)	3 (5.6%)
Epilepsy	1 (5.6%)	0 (0.0%)	1 (11.1%)	1 (5.6%)	3 (5.6%)
Generalised tonic-clonic seizure	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)
Haemorrhage intracranial	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Headache	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Partial seizures	3 (16.7%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	4 (7.4%)
Seizure	0 (0.0%)	2 (22.2%)	2 (22.2%)	0 (0.0%)	4 (7.4%)
Somnolence	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Syncope	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Psychiatric disorders	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Confusional state	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Renal and urinary disorders	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Renal failure	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Respiratory, thoracic and mediastinal disorders	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Pulmonary embolism	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)

Vital Signs

Systolic and diastolic blood pressure and pulse rate are summarised descriptively by dose group and visit (observed and change from baseline) and analysed with a repeated- measures linear mixed model analogous to the efficacy endpoints: change from baseline with baseline value, nominal visit and treatment group as fixed effects and a subject random intercept. Short-term (Days 15/29/43) and long-term (Days 155/239) windows are reported.

Systolic BP

A total of 54 subjects contributed to the model-based analyses.

Descriptive summary by dose and visit

Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	3	128.0 (19.7)	0.0 (0.0)
15	0 mg	36	123.1 (14.5)	-4.5 (13.4)
15	25 mg	6	124.8 (16.4)	-17.3 (10.5)
15	50 mg	6	122.5 (13.0)	-9.7 (7.3)
15	100 mg	6	121.2 (7.1)	-8.2 (5.7)
29	0 mg	18	114.6 (12.2)	-15.3 (10.2)
29	25 mg	12	117.8 (15.2)	-14.5 (12.6)
29	50 mg	11	117.2 (11.3)	-12.7 (14.6)
29	100 mg	12	115.8 (8.5)	-11.7 (14.8)
43	25 mg	18	115.7 (13.9)	-15.8 (18.0)
43	50 mg	16	112.6 (9.6)	-14.2 (12.6)
43	100 mg	18	120.2 (14.2)	-10.2 (15.7)
57	25 mg	1	116.0 (NA)	-19.0 (NA)
71	0 mg	17	116.9 (9.2)	-14.0 (13.4)
71	25 mg	8	122.9 (21.9)	-9.0 (16.1)
71	50 mg	8	124.1 (9.7)	-2.0 (8.6)
71	100 mg	18	123.3 (13.6)	-7.0 (16.3)
99	0 mg	17	118.6 (11.5)	-12.1 (13.5)
99	25 mg	8	122.6 (24.1)	-9.2 (11.0)
99	50 mg	8	120.2 (14.5)	-5.9 (14.9)
99	100 mg	18	119.6 (13.2)	-10.8 (14.3)
127	0 mg	14	115.6 (9.9)	-13.6 (15.0)
127	25 mg	6	124.5 (24.6)	-9.5 (13.2)
127	50 mg	8	120.9 (13.1)	-5.2 (21.1)
127	100 mg	14	128.1 (15.8)	-2.1 (18.0)
141	0 mg	1	124.0 (NA)	-19.0 (NA)
155	0 mg	14	119.9 (11.9)	-9.3 (11.6)
155	25 mg	8	122.8 (22.3)	-9.1 (9.4)
155	50 mg	8	123.1 (9.8)	-3.0 (14.0)
155	100 mg	18	123.7 (12.1)	-7.0 (15.4)
183	0 mg	14	123.3 (8.7)	-5.9 (16.6)
183	25 mg	6	123.5 (24.9)	-8.7 (5.6)
183	50 mg	8	124.9 (15.1)	-1.2 (16.6)
183	100 mg	16	123.0 (12.3)	-6.1 (20.7)
211	0 mg	13	113.5 (10.4)	-15.3 (15.8)
211	25 mg	5	119.8 (18.1)	-14.4 (9.7)
211	50 mg	7	124.9 (6.6)	-3.0 (12.2)
211	100 mg	13	121.8 (12.8)	-5.0 (19.3)
239	0 mg	11	122.4 (12.7)	-7.5 (11.7)


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
239	25 mg	4	122.8 (32.4)	-12.8 (11.7)
239	50 mg	7	125.0 (5.4)	-2.9 (14.3)
239	100 mg	15	123.7 (14.9)	-4.5 (23.3)
253	0 mg	46	125.5 (15.8)	-3.9 (15.4)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	-12.00 (-15.64 to -8.37)
25 mg	-13.17 (-17.47 to -8.86)
50 mg	-11.53 (-15.94 to -7.11)
100 mg	-8.44 (-12.73 to -4.16)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg0	-1.16 (-7.74 to 5.41)
losartan 50 mg vs losartan 0 mg0	0.47 (-6.17 to 7.12)
losartan 100 mg vs losartan 0 mg0	3.56 (-2.98 to 10.10)




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-9.49 (-16.17 to -2.80)
155	losartan 25 mg	-7.90 (-16.76 to 0.95)
155	losartan 50 mg	-4.85 (-13.72 to 4.02)
155	losartan 100 mg	-6.32 (-12.34 to -0.30)
239	losartan 0 mg	-7.25 (-14.59 to 0.09)
239	losartan 25 mg	-9.62 (-21.39 to 2.15)
239	losartan 50 mg	-3.53 (-12.85 to 5.79)
239	losartan 100 mg	-5.23 (-11.58 to 1.12)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	1.58 (-11.85 to 15.02)
155	losartan 50 mg - losartan 0 mg	4.63 (-8.81 to 18.08)
155	losartan 100 mg - losartan 0 mg	3.17 (-7.72 to 14.06)
239	losartan 25 mg - losartan 0 mg	-2.37 (-19.15 to 14.40)
239	losartan 50 mg - losartan 0 mg	3.72 (-10.64 to 18.08)
239	losartan 100 mg - losartan 0 mg	2.02 (-9.73 to 13.76)



Plot

Diastolic BP

A total of 54 subjects contributed to the model-based analyses.

Descriptive summary by dose and visit


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	3	72.7 (5.1)	0.0 (0.0)
15	0 mg	36	78.0 (7.4)	-2.4 (8.2)
15	25 mg	6	82.2 (5.0)	-4.2 (4.7)
15	50 mg	6	77.8 (7.7)	-6.0 (5.8)
15	100 mg	6	78.7 (8.9)	-5.7 (8.7)
29	0 mg	18	74.5 (8.0)	-5.9 (8.0)
29	25 mg	12	75.2 (7.6)	-6.3 (7.4)
29	50 mg	11	75.7 (6.1)	-7.5 (6.2)
29	100 mg	12	76.6 (6.0)	-5.4 (7.8)
43	25 mg	18	73.4 (7.8)	-8.7 (9.3)
43	50 mg	16	73.6 (6.5)	-8.6 (6.7)
43	100 mg	18	76.1 (6.5)	-4.2 (7.9)
57	25 mg	1	74.0 (NA)	-10.0 (NA)
71	0 mg	17	75.8 (5.4)	-7.8 (7.2)
71	25 mg	8	78.1 (8.2)	-3.1 (9.2)
71	50 mg	8	80.4 (5.3)	-1.2 (6.7)
71	100 mg	18	76.5 (9.6)	-3.7 (7.3)
99	0 mg	17	77.2 (8.0)	-6.5 (9.8)
99	25 mg	8	78.9 (11.5)	-2.4 (11.6)
99	50 mg	8	80.0 (10.7)	-1.6 (9.5)
99	100 mg	18	76.1 (8.7)	-4.1 (8.5)
127	0 mg	14	75.0 (7.7)	-8.1 (9.1)
127	25 mg	6	80.7 (11.1)	-2.0 (8.2)
127	50 mg	8	80.0 (8.9)	-1.6 (10.8)
127	100 mg	14	77.1 (7.0)	-3.1 (8.5)
141	0 mg	1	77.0 (NA)	-7.0 (NA)
155	0 mg	14	77.1 (7.7)	-6.1 (6.5)
155	25 mg	8	78.2 (10.6)	-3.0 (6.1)
155	50 mg	8	81.2 (5.4)	-0.4 (9.1)
155	100 mg	18	78.4 (7.9)	-1.4 (7.0)
183	0 mg	14	79.4 (5.8)	-3.8 (7.4)
183	25 mg	6	79.7 (9.2)	-0.7 (5.2)
183	50 mg	8	78.9 (7.7)	-2.8 (7.9)
183	100 mg	16	79.1 (8.3)	-1.4 (8.1)
211	0 mg	13	73.5 (5.5)	-8.8 (5.2)
211	25 mg	5	74.8 (6.9)	-8.2 (3.1)
211	50 mg	7	82.1 (6.1)	-0.4 (6.8)
211	100 mg	13	77.6 (7.0)	-3.1 (6.5)
239	0 mg	11	78.9 (7.3)	-5.2 (5.6)
239	25 mg	4	75.8 (6.5)	-7.2 (4.8)
239	50 mg	7	82.3 (5.0)	-0.3 (8.4)
239	100 mg	15	79.0 (8.4)	-1.0 (8.0)
253	0 mg	46	80.5 (9.0)	-1.7 (8.3)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	-5.77 (-7.97 to -3.56)
25 mg	-7.51 (-10.13 to -4.89)
50 mg	-6.03 (-8.73 to -3.33)
100 mg	-3.23 (-5.85 to -0.62)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	-1.74 (-5.62 to 2.14)
losartan 50 mg vs losartan 0 mg	-0.26 (-4.22 to 3.69)
losartan 100 mg vs losartan 0 mg	2.53 (-1.34 to 6.40)



Plot


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-5.45 (-8.94 to -1.95)
155	losartan 25 mg	-3.16 (-7.76 to 1.44)
155	losartan 50 mg	-0.38 (-4.98 to 4.22)
155	losartan 100 mg	-2.17 (-5.33 to 0.98)
239	losartan 0 mg	-3.97 (-7.79 to -0.15)
239	losartan 25 mg	-6.11 (-12.10 to -0.12)

Visit	Treatment	Estimated change from baseline (95% CI)
239	losartan 50 mg	0.39 (-4.43 to 5.21)
239	losartan 100 mg	-1.62 (-4.93 to 1.69)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	2.29 (-4.72 to 9.29)
155	losartan 50 mg - losartan 0 mg	5.07 (-1.93 to 12.07)
155	losartan 100 mg - losartan 0 mg	3.27 (-2.47 to 9.01)
239	losartan 25 mg - losartan 0 mg	-2.14 (-10.73 to 6.44)
239	losartan 50 mg - losartan 0 mg	4.36 (-3.07 to 11.79)
239	losartan 100 mg - losartan 0 mg	2.35 (-3.81 to 8.51)



Plot

Pulse rate

A total of 54 subjects contributed to the model-based analyses.

Descriptive summary by dose and visit


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	3	71.7 (4.5)	0.0 (0.0)
15	0 mg	36	74.4 (11.5)	-6.9 (14.2)
15	25 mg	6	80.8 (16.5)	-1.3 (9.2)
15	50 mg	6	66.5 (6.8)	-0.7 (5.6)
15	100 mg	6	79.7 (7.3)	-8.5 (14.1)
29	0 mg	18	78.1 (13.0)	-4.5 (14.3)
29	25 mg	12	77.0 (16.5)	-3.2 (12.3)
29	50 mg	11	77.2 (13.0)	1.2 (9.3)
29	100 mg	12	76.8 (9.1)	-4.2 (13.3)
43	25 mg	18	73.4 (14.7)	-7.8 (13.5)
43	50 mg	16	72.4 (11.0)	-7.3 (13.5)
43	100 mg	18	75.7 (12.2)	-3.6 (11.9)
57	25 mg	1	60.0 (NA)	6.0 (NA)
71	0 mg	17	73.0 (12.3)	-11.5 (11.4)
71	25 mg	8	74.6 (9.1)	-3.9 (12.6)
71	50 mg	8	70.8 (7.8)	-8.4 (12.9)
71	100 mg	18	76.2 (13.0)	-3.1 (15.4)
99	0 mg	17	76.4 (14.7)	-7.5 (15.0)
99	25 mg	8	76.8 (11.1)	-1.8 (14.9)
99	50 mg	8	75.8 (9.7)	-3.4 (16.9)
99	100 mg	18	75.7 (8.8)	-3.6 (14.0)
127	0 mg	14	73.8 (12.5)	-12.1 (14.6)
127	25 mg	6	71.8 (11.1)	-4.3 (9.6)
127	50 mg	8	71.8 (13.0)	-7.4 (20.4)
127	100 mg	14	74.1 (11.6)	-6.1 (14.8)
141	0 mg	1	65.0 (NA)	-4.0 (NA)
155	0 mg	14	74.4 (12.9)	-11.6 (13.6)
155	25 mg	8	75.4 (11.3)	-3.1 (8.0)
155	50 mg	8	72.2 (8.4)	-6.9 (12.5)
155	100 mg	18	75.7 (13.1)	-4.3 (14.6)
183	0 mg	14	73.0 (11.1)	-12.9 (14.9)
183	25 mg	6	73.0 (9.3)	-7.0 (12.5)
183	50 mg	8	73.9 (7.3)	-5.2 (16.5)
183	100 mg	16	74.8 (11.1)	-6.2 (12.5)
211	0 mg	13	73.8 (10.0)	-14.6 (14.2)
211	25 mg	5	72.8 (13.5)	-8.8 (14.5)
211	50 mg	7	73.9 (14.2)	-2.6 (21.4)
211	100 mg	13	74.3 (11.6)	-7.3 (13.2)
239	0 mg	11	72.9 (11.5)	-15.8 (10.5)
239	25 mg	4	87.5 (24.8)	3.8 (28.2)
239	50 mg	7	72.9 (11.7)	-3.6 (18.9)
239	100 mg	15	76.0 (12.1)	-4.5 (15.7)
253	0 mg	45	77.1 (12.6)	-5.2 (16.2)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	-3.46 (-6.78 to -0.13)
25 mg	-5.66 (-9.64 to -1.68)
50 mg	-6.98 (-11.08 to -2.88)
100 mg	-4.81 (-8.79 to -0.83)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	-2.20 (-7.86 to 3.46)
losartan 50 mg vs losartan 0 mg	-3.52 (-9.31 to 2.27)
losartan 100 mg vs losartan 0 mg	-1.36 (-7.02 to 4.30)



Plot


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-9.11 (-15.11 to -3.11)
155	losartan 25 mg	-5.14 (-13.04 to 2.76)
155	losartan 50 mg	-8.51 (-16.40 to -0.63)
155	losartan 100 mg	-5.11 (-10.46 to 0.25)
239	losartan 0 mg	-11.71 (-18.37 to -5.04)
239	losartan 25 mg	2.85 (-7.77 to 13.48)

Visit	Treatment	Estimated change from baseline (95% CI)
239	losartan 50 mg	-6.95 (-15.30 to 1.41)
239	losartan 100 mg	-5.75 (-11.43 to -0.07)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	3.97 (-8.09 to 16.04)
155	losartan 50 mg - losartan 0 mg	0.60 (-11.45 to 12.65)
155	losartan 100 mg - losartan 0 mg	4.01 (-5.76 to 13.78)
239	losartan 25 mg - losartan 0 mg	14.56 (-0.62 to 29.74)
239	losartan 50 mg - losartan 0 mg	4.76 (-8.26 to 17.79)
239	losartan 100 mg - losartan 0 mg	5.96 (-4.68 to 16.59)



Plot

Conventional Laboratory Parameters

Conventional clinical laboratory parameters (creatinine in this export) are summarised descriptively and with the same repeated-measures mixed model as vital signs, plus categorisation against reference ranges (counts/proportions outside the normal range).

Creatinine

Reference-range categorisation by dose

trt	LOW	NORMAL	HIGH
0 mg	3 (3.0%)	97 (97.0%)	0 (0.0%)
25 mg	1 (3.4%)	27 (93.1%)	1 (3.4%)
50 mg	0 (0.0%)	30 (100.0%)	0 (0.0%)
100 mg	1 (2.6%)	36 (94.7%)	1 (2.6%)

Descriptive summary by dose and visit


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	54	70.4 (12.9)	0.0 (0.0)
15	0 mg	35	74.1 (10.8)	2.9 (8.1)
15	25 mg	6	69.8 (17.6)	8.2 (15.6)
15	50 mg	6	71.8 (12.0)	2.5 (3.7)
15	100 mg	6	67.7 (8.2)	-0.6 (1.8)
43	25 mg	18	73.2 (21.0)	5.1 (7.4)
43	50 mg	17	75.1 (9.1)	6.3 (4.3)
43	100 mg	18	76.9 (14.9)	1.9 (5.9)
225	0 mg	11	73.7 (11.8)	7.1 (6.4)
225	25 mg	5	75.0 (16.4)	1.2 (13.8)
225	50 mg	7	76.9 (6.6)	6.3 (9.4)
225	100 mg	14	74.1 (13.5)	0.7 (7.4)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	3.62 (0.30 to 6.95)
25 mg	4.42 (0.97 to 7.86)
50 mg	4.62 (1.19 to 8.06)
100 mg	2.48 (-0.98 to 5.95)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg0	0.80 (-5.33 to 6.92)
losartan 50 mg vs losartan 0 mg0	1.00 (-5.05 to 7.05)
losartan 100 mg vs losartan 0 mg0	-1.14 (-7.21 to 4.93)



Deaths

Deaths are summarised by long-term planned dose group (in addition to the overall survival analysis in the time-to-event section) and listed with date and cause.

A total of 34 deaths were observed among 54 subjects.

Deaths by dose group

Long-term dose	N	Deaths n (%)
0 mg	18	12 (66.7%)
25 mg	9	8 (88.9%)
50 mg	9	5 (55.6%)
100 mg	18	9 (50.0%)

Listing of deaths

Subject	Dose	Death date	Days from start	Cause
ImPRESS-ImPR-2005	0 mg	2021-04-20	307	Progressive disease
ImPRESS-ImPR-2001	0 mg	2021-06-29	527	Progressive disease
ImPRESS-ImPR-2022	0 mg	2022-08-28	455	Progressive disease
ImPRESS-ImPR-2030	0 mg	2022-11-20	356	Progressive disease
ImPRESS-ImPR-2029	0 mg	2023-01-13	437	Progressive disease
ImPRESS-ImPR-2033	0 mg	2023-02-26	370	Progressive disease
ImPRESS-ImPR-2034	0 mg	2023-03-18	347	Progressive disease
ImPRESS-ImPR-2041	0 mg	2024-01-18	303	Progressive disease
ImPRESS-ImPR-2038	0 mg	2024-04-24	448	Progressive disease
ImPRESS-ImPR-2050	0 mg	2025-03-28	430	Progressive disease
ImPRESS-ImPR-2052	0 mg	2025-04-26	312	Progressive disease
ImPRESS-ImPR-2048	0 mg	2025-09-09	665	Progressive disease
ImPRESS-ImPR-2015	25 mg	2022-01-11	429	Progressive disease
ImPRESS-ImPR-2011	25 mg	2022-02-17	540	Progressive disease
ImPRESS-ImPR-2016	25 mg	2022-03-21	456	Progressive disease
ImPRESS-ImPR-2035	25 mg	2023-05-07	362	Progressive disease
ImPRESS-ImPR-2036	25 mg	2023-06-15	205	Progressive disease
ImPRESS-ImPR-2046	25 mg	2023-07-29	67	Progressive disease
ImPRESS-ImPR-2040	25 mg	2023-09-05	182	Other cause
ImPRESS-ImPR-2037	25 mg	2024-02-02	387	Progressive disease
ImPRESS-ImPR-2019	50 mg	2022-01-01	300	Progressive disease
ImPRESS-ImPR-2013	50 mg	2022-02-01	484	Progressive disease
ImPRESS-ImPR-2009	50 mg	2022-02-04	551	Progressive disease
ImPRESS-ImPR-2023	50 mg	2022-08-28	432	Progressive disease
ImPRESS-ImPR-2044	50 mg	2024-07-14	433	Progressive disease
ImPRESS-ImPR-2014	100 mg	2021-03-09	134	Progressive disease
ImPRESS-ImPR-2004	100 mg	2021-07-29	430	Progressive disease
ImPRESS-ImPR-2002	100 mg	2021-08-31	583	Progressive disease
ImPRESS-ImPR-2012	100 mg	2021-11-03	421	Progressive disease
ImPRESS-ImPR-2008	100 mg	2021-11-10	465	Progressive disease
ImPRESS-ImPR-2026	100 mg	2022-10-30	419	Progressive disease
ImPRESS-ImPR-2031	100 mg	2022-12-06	331	Progressive disease
ImPRESS-ImPR-2027	100 mg	2023-09-07	717	Progressive disease
ImPRESS-ImPR-2045	100 mg	2024-06-15	401	Progressive disease