

ImPRESS Statistical Analysis Mock-up Report

Fully planned efficacy and safety results of the Newly Diagnosed Glioblastoma Cohort

ImPRESS Statistical Team

20 June 2026

Contents

Introduction	4
Randomisation Status	4
Cohort Scope	4
Baseline Characteristics	4
Primary Efficacy Analyses (MRI)	11
Candidate dose-response models	12
Mean relative cerebral blood flow (PTA-ROI)	12
Mean relative cerebral blood flow (ROI2)	17
Mean relative solid stress (PTA-ROI)	22
Mean relative solid stress (ROI2)	27
Other MRI Endpoints up to Day 43	32
Mean DSC K2 leakage	32
Mean relative DSC extravasation	35
Mean relative apparent diffusion coefficient	37
Mean relative cerebral blood	39
Mean relative direction value	41
Mean relative fractional anisotropy	43
Mean relative restricted spectrum imaging	45
Mean relative transit time	47
Mean relative vessel caliber	49
Mean relative vessel size index	51
Mean relative vortex area	53
Mean tissue elasticity	55
Mean tissue relative viscosity	57
Mean tissue stiffness	59
Mean tissue viscosity	61
Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3	63
Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3	65
MRI Endpoints at Cycle 11 (Day 141)	67
Mean DSC K2 leakage	67
Mean relative DSC extravasation	69
Mean relative apparent diffusion coefficient	70
Mean relative cerebral blood	71

Mean relative cerebral blood flow (PTA-ROI)	72
Mean relative cerebral blood flow (ROI2)	73
Mean relative direction value	74
Mean relative fractional anisotropy	75
Mean relative restricted spectrum imaging	76
Mean relative solid stress (PTA-ROI)	77
Mean relative solid stress (ROI2)	78
Mean relative transit time	79
Mean relative vessel caliber	80
Mean relative vessel size index	81
Mean relative vortex area	82
Mean tissue elasticity	83
Mean tissue relative viscosity	84
Mean tissue stiffness	85
Mean tissue viscosity	86
Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3	87
Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3	88
Neurologic Performance	89
Karnofsky Performance Scale	89
ECOG Score	92
Total NANO score	95
Quality of Life Outcomes	98
EORTC QLQ-C30 score	98
EORTC QLQ-BN20 score	100
Steroid Use	102
Steroid use	102
RANO Responses	105
State of disease according to the RANO criteria	105
Best overall radiographic response	106
Biomarker Endpoints (Serum Cytokines / Angiogenic Factors)	107
Fractalkine	107
Granzyme B	109
IFN-g	111
IL-1b	113
IL-6	115
IL-8	117
MCP-1	119
MMP-10	121
PDGF-DD	123
PECAM-1	125
TGF-alpha	127
Thrombospondin-2	129
VEGF-A	131
VEGF-C	133
Angiopoietin-1	135
Angiopoietin-2	137
ANGPTL4	139
CA9	141
CD163	143
Collagen IV alpha 1	145

EGF	147
ErbB2	149
FGF Basic	151
IL-12 p70	153
IL-2	155
IP-10 (CXCL10)	157
I-TAC (CXCL11)	159
MIG (CXCL9)	161
PIGF	163
Tenascin C	165
TNFRSF18 (GITR)	167
TNFSF6 (FasL)	169
VEGFR2	171
Biomarker Endpoints — Long-term (Cycle 15+)	173
Fractalkine	173
Granzyme B	175
IFN-g	177
IL-1b	179
IL-6	181
IL-8	183
MCP-1	185
MMP-10	187
PDGF-DD	189
PECAM-1	191
TGF-alpha	193
Thrombospondin-2	195
VEGF-A	197
VEGF-C	199
Angiopoietin-1	201
Angiopoietin-2	203
ANGPTL4	205
CA9	207
CD163	209
Collagen IV alpha 1	211
EGF	213
ErbB2	215
FGF Basic	217
IL-12 p70	219
IL-2	221
IP-10 (CXCL10)	223
I-TAC (CXCL11)	225
MIG (CXCL9)	227
PIGF	229
Tenascin C	231
TNFRSF18 (GITR)	233
TNFSF6 (FasL)	235
VEGFR2	237
Time-to-Event Endpoints	239
Progression-free survival	240
6-month progression-free survival	242
Overall survival	244
Overall survival at 12 months	246

Overall survival at 24 months	248
Safety	250
Extent of Exposure	250
Adverse Events	252
Vital Signs	256
Conventional Laboratory Parameters	265
Deaths	267

Introduction

The ImPRESS study is a prospective investigation of losartan as an adjunctive therapy for patients with glioblastoma. According to Protocol Version 7.0 (17-Dec-2024), the trial collects comprehensive demographic, clinical, imaging, and quality-of-life information prior to treatment initiation. The statistical objectives for the interim analyses focus on characterising baseline status, neurological function, and tumour burden before randomisation to evaluate comparability across planned treatment arms.

Randomisation Status

The present dataset is built on sham randomisation generated from the most recent export (see `_targets.R` workflow). No true treatment assignments have been unlocked. All findings in this document should therefore be interpreted as preliminary and descriptive only. Final efficacy comparisons will not be conducted until the database is locked and genuine randomisation codes are released.

Cohort Scope

The results herein are restricted to participants in cohort code 2 as configured in `config/cfg.yml`. Subjects from other cohorts are excluded from all summaries, figures, and tables in this draft. A combined-cohort analysis can be delivered once recruitment and cleaning are complete for all strata.

Baseline Characteristics

This section summarises baseline characteristics for the newly diagnosed glioblastoma cohort using the `adb1` ADaM dataset.

Demographics and Baseline characteristics

	all obs
Age (years)	
n	54
Mean (SD)	52.6 (10.2)
Median	55.0
Min - Max	26.0 - 69.0
Sex	
n	54
Female	29 (53.7%)
Male	25 (46.3%)

Demographics and Baseline characteristics

	all obs
Cohort	
n	54
Newly diagnosed glioblastoma	54 (100%)
Country	
n	54
NOR	54 (100%)
Childbearing potential?	
n	29
No	16 (55.2%)
Yes	13 (44.8%)
Diagnosis	
n	54
Newly diagnosed glioblastoma	54 (100%)
Time from surgery to randomisation (days)	
n	54
Mean (SD)	27.4 (7.3)
Median	25.5
Min - Max	13.0 - 48.0
IDH1/2 mutation	
n	54
Negative	47 (87%)
Positive	7 (13%)
MGMT status	
n	54
Negative	30 (55.6%)
Positive	24 (44.4%)
Pregnancy test result	
n	13
Negative	13 (100%)
ECOG Score	
n	54
Mean (SD)	0.3 (0.5)
Median	0.0
Min - Max	0.0 - 2.0

Demographics and Baseline characteristics

	all obs
ECOG Score (categorical)	
n	54
ECOG 0	40 (74.1%)
ECOG 1	13 (24.1%)
ECOG 2	1 (1.9%)
Karnofsky Performance Scale	
n	54
Mean (SD)	95.0 (8.0)
Median	100.0
Min - Max	60.0 - 100.0
Karnofsky Performance Scale (categorical)	
n	54
KPS 100	34 (63%)
KPS 60	1 (1.9%)
KPS 80	4 (7.4%)
KPS 90	15 (27.8%)
Total NANO score	
n	54
Mean (SD)	1.0 (1.3)
Median	0.0
Min - Max	0.0 - 4.0
EORTC QLQ-C30 score	
n	53
Mean (SD)	56.3 (8.6)
Median	54.0
Min - Max	41.0 - 77.0
EORTC QLQ-BN20 score	
n	53
Mean (SD)	30.8 (6.6)
Median	30.0
Min - Max	20.0 - 49.0
Systolic blood pressure	
n	54
Mean (SD)	130.0 (15.4)

Demographics and Baseline characteristics

	all obs
Median	128.5
Min - Max	106.0 - 180.0
Diastolic blood pressure	
n	54
Mean (SD)	81.9 (7.1)
Median	82.0
Min - Max	67.0 - 97.0
Pulse Rate	
n	54
Mean (SD)	80.6 (15.6)
Median	78.0
Min - Max	54.0 - 114.0
Weight (kg)	
n	54
Mean (SD)	78.8 (13.2)
Median	78.3
Min - Max	48.2 - 116.0
BMI	
n	51
Mean (SD)	25.0 (4.2)
Median	23.9
Min - Max	18.6 - 39.9
Height (cm)	
n	51
Mean (SD)	176.5 (8.8)
Median	175.0
Min - Max	158.0 - 194.0
Has the patient received chemotherapy	
n	54
No	54 (100%)
Has the patient received immunotherapy	
n	54
No	54 (100%)
Has the patient received radiation therapy to the brain?	

Demographics and Baseline characteristics

	all obs
n	54
No	54 (100%)
CNS surgery	
n	54
No	3 (5.6%)
Yes	51 (94.4%)
Steroids at time of MRI	
n	54
No	42 (77.8%)
Yes	12 (22.2%)
Mean relative cerebral blood flow (PTA-ROI)	
n	54
Mean (SD)	1.5 (0.4)
Median	1.4
Min - Max	0.7 - 2.4
Mean relative cerebral blood flow (ROI2)	
n	47
Mean (SD)	1.1 (0.5)
Median	1.0
Min - Max	0.2 - 2.2
Mean relative solid stress (PTA-ROI)	
n	41
Mean (SD)	0.8 (0.1)
Median	0.8
Min - Max	0.5 - 1.1
Mean relative solid stress (ROI2)	
n	32
Mean (SD)	0.9 (0.2)
Median	0.9
Min - Max	0.5 - 1.4
Mean relative apparent diffusion coefficient	
n	52
Mean (SD)	1.4 (0.2)
Median	1.4

Demographics and Baseline characteristics

	all obs
Min - Max	1.0 - 2.0
Mean relative fractional anisotropy	
n	52
Mean (SD)	0.5 (0.1)
Median	0.5
Min - Max	0.3 - 1.0
Mean relative restricted spectrum imaging	
n	50
Mean (SD)	0.5 (0.2)
Median	0.5
Min - Max	0.2 - 1.1
Mean DSC K2 leakage	
n	54
Mean (SD)	0.2 (1.6)
Median	0.4
Min - Max	-3.9 - 4.5
Mean relative cerebral blood	
n	54
Mean (SD)	1.7 (0.5)
Median	1.7
Min - Max	1.0 - 2.8
Mean relative transit time	
n	54
Mean (SD)	1.4 (0.3)
Median	1.4
Min - Max	0.9 - 2.3
Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3	
n	54
Mean (SD)	6.3 (6.5)
Median	4.3
Min - Max	0.0 - 26.3
Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3	
n	54
Mean (SD)	13.5 (14.3)

Demographics and Baseline characteristics

	all obs
Median	8.5
Min - Max	0.1 - 57.4
Mean tissue elasticity	
n	41
Mean (SD)	0.8 (0.1)
Median	0.8
Min - Max	0.5 - 1.2
Mean tissue relative viscosity	
n	41
Mean (SD)	0.9 (0.2)
Median	0.9
Min - Max	0.0 - 1.1
Mean tissue stiffness	
n	41
Mean (SD)	0.8 (0.1)
Median	0.8
Min - Max	0.5 - 1.1
Mean tissue viscosity	
n	41
Mean (SD)	0.7 (0.2)
Median	0.7
Min - Max	-0.0 - 1.1
Mean relative DSC extravasation	
n	28
Mean (SD)	1.2 (0.8)
Median	1.0
Min - Max	0.7 - 5.1
Mean relative direction value	
n	28
Mean (SD)	0.4 (1.4)
Median	0.3
Min - Max	-2.7 - 5.4
Mean relative vessel caliber	
n	28

Demographics and Baseline characteristics


	all obs
Mean (SD)	1.0 (0.3)
Median	1.0
Min - Max	0.5 - 1.9
Mean relative vessel size index	
n	28
Mean (SD)	1.4 (0.6)
Median	1.2
Min - Max	0.6 - 2.6
Mean relative vortex area	
n	28
Mean (SD)	1.0 (0.1)
Median	1.0
Min - Max	0.8 - 1.1

Primary Efficacy Analyses (MRI)

The primary variable is analysed with the MCP-Mod methodology as described in the protocol and SAP. We pre-specify four candidate dose-response models (linear, piecewise linear, Emax, sigmoid Emax) and derive optimal contrasts using the estimated variance-covariance from the mixed model. The MCP step tests for a dose-response signal across candidate models with multiplicity control; the Mod step presents the model fit by AIC and reports ED values. Results below are grouped by MRI endpoint (primary and sensitivity (ROI2)). All analyses are based on data up to and including Day 43 (Week 6).

Candidate dose-response models

Figure 1: Pre-specified candidate dose-response models




Mean relative cerebral blood flow (PTA-ROI)

The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.642	-0.819	-0.588	-0.865
25 mg	-0.194	0.047	-0.381	0.311
50 mg	0.101	0.265	0.343	0.262
100 mg	0.735	0.507	0.626	0.292

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
linear	2.14	0.032
sigEmax	1.93	0.052
emax	1.85	0.062
linInt	1.39	0.14

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	4.11	50.00	70.00	90.00

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
emax	6.40	37.50	58.33	84.37
linInt	8.00	61.36	76.82	92.27
sigEmax	8.07	58.74	74.96	91.44

Estimated means by dose

Treatment

Estimated Change from Baseline (95% CI)

0 mg

-0.03 (-0.13 to 0.07)

25 mg

0.00 (-0.11 to 0.12)

50 mg

0.02 (-0.09 to 0.14)



100 mg


0.12 (0.01 to 0.24)


Note:

Fitted models are presented with AIC and ED for reference

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.08 (-0.04 to 0.20)


Mean relative cerebral blood flow (ROI2)

The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.740	-0.850	-0.659	-0.863
25 mg	-0.101	0.133	-0.307	0.344
50 mg	0.210	0.322	0.441	0.286
100 mg	0.631	0.395	0.526	0.234

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
linInt	-0.11	0.69
emax	-0.20	0.73
sigEmax	-0.28	0.75
linear	-0.29	0.76

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	4.03	50.00	70.00	90.00

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
emax	6.04	37.50	58.33	84.37
linInt	8.00	70.14	82.08	94.03
sigEmax	8.00	56.76	61.75	70.52

Estimated means by dose

Treatment

Estimated Change from Baseline (95% CI)

0 mg

0.07 (-0.08 to 0.22)

25 mg

0.07 (-0.09 to 0.24)

50 mg

0.06 (-0.10 to 0.23)



100 mg


0.04 (-0.16 to 0.23)


Note:

Fitted models are presented with AIC and ED for reference

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.01 (-0.15 to 0.13)


Mean relative solid stress (PTA-ROI)

The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.673	-0.840	-0.600	-0.860
25 mg	-0.154	0.124	-0.367	0.378
50 mg	0.112	0.251	0.344	0.231
100 mg	0.715	0.465	0.623	0.251

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
linear	-1.24	0.96
linInt	-1.70	0.99
sigEmax	-1.72	0.99
emax	-1.76	0.99

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linInt	8.00	33.16	39.90	46.63

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	8.98	50.00	70.00	90.00
emax	9.30	11.57	23.39	54.08
sigEmax	10.12	26.25	28.57	32.70

Estimated means by dose

Treatment

Estimated Change from Baseline (95% CI)

0 mg

0.03 (-0.01 to 0.08)

25 mg

0.02 (-0.03 to 0.06)

50 mg

-0.04 (-0.09 to 0.01)



100 mg


0.01 (-0.04 to 0.06)


Note:

Fitted models are presented with AIC and ED for reference

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.04 (-0.08 to 0.01)

Mean relative solid stress (ROI2)


The primary analysis follows the MCP-Mod methodology as detailed in the SAP, using pre-specified candidate dose-response models.

A total of 54 subjects were included in the analysis for this endpoint.

The four candidate models (linear, piecewise linear, Emax, sigmoid Emax) are shown in Figure 1.

Line plot

Mean Response over Time for Mean relative solid stress (ROI2)



Optimal contrasts

Treatment	linear	emax	sigEmax	linInt
0 mg	-0.593	-0.829	-0.468	-0.853
25 mg	-0.278	0.074	-0.519	0.428
50 mg	0.126	0.270	0.381	0.211
100 mg	0.745	0.485	0.605	0.214

MCP-Mod tests

Candidate model	t-statistic	Adjusted p-value
linInt	0.07	0.66
emax	-0.25	0.77
linear	-0.49	0.84
sigEmax	-0.64	0.88

Fitted model parameters (AIC/ED)

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
linear	4.40	50.0	70.00	90.00

Model	AIC-value	ED ₅₀	ED ₇₀	ED ₉₀
emax	6.45	37.5	58.33	84.37
linInt	8.00	12.5	17.50	22.50
sigEmax	8.17	43.6	47.45	54.30

Estimated means by dose

Treatment

Estimated Change from Baseline (95% CI)

0 mg

-0.01 (-0.12 to 0.10)

25 mg

0.02 (-0.08 to 0.12)

50 mg

-0.03 (-0.15 to 0.10)



100 mg


-0.04 (-0.16 to 0.09)


Note:

Fitted models are presented with AIC and ED for reference

Fitted dose-response curves







Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.00 (-0.12 to 0.13)


Other MRI Endpoints up to Day 43

For remaining MRI endpoints, we present estimated means by dose, contrasts vs 0 mg, and a plot with confidence intervals.

Mean DSC K2 leakage

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.19 (-0.61 to 0.24)
25 mg	-0.39 (-0.88 to 0.10)
50 mg	-0.61 (-1.11 to -0.12)
100 mg	-0.11 (-0.61 to 0.39)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.20 (-0.95 to 0.54)
50 mg vs 0 mg	-0.43 (-1.20 to 0.34)
100 mg vs 0 mg	0.08 (-0.67 to 0.83)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.18 (-0.67 to 0.31)

Mean relative DSC extravasation

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.01 (-0.32 to 0.31)
25 mg	0.14 (-0.12 to 0.39)
50 mg	-0.04 (-0.31 to 0.24)
100 mg	-0.18 (-0.56 to 0.20)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.14 (-0.36 to 0.65)
50 mg vs 0 mg	-0.03 (-0.57 to 0.51)
100 mg vs 0 mg	-0.17 (-0.79 to 0.45)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.03 (-0.35 to 0.42)

Mean relative apparent diffusion coefficient

A total of 51 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.15 (0.09 to 0.21)
25 mg	0.16 (0.09 to 0.23)
50 mg	0.15 (0.08 to 0.22)
100 mg	0.16 (0.09 to 0.23)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.01 (-0.09 to 0.11)
50 mg vs 0 mg	0.00 (-0.10 to 0.10)
100 mg vs 0 mg	0.01 (-0.09 to 0.11)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.01 (-0.05 to 0.07)

Mean relative cerebral blood

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.07 (-0.18 to 0.03)
25 mg	-0.09 (-0.21 to 0.04)
50 mg	-0.01 (-0.13 to 0.12)
100 mg	0.06 (-0.07 to 0.19)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.01 (-0.20 to 0.18)
50 mg vs 0 mg	0.07 (-0.13 to 0.27)
100 mg vs 0 mg	0.14 (-0.06 to 0.33)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.06 (-0.07 to 0.19)

Mean relative direction value

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	4599.49 (609.51 to 8589.47)
25 mg	-893.17 (-4232.87 to 2446.54)
50 mg	1359.80 (-2173.79 to 4893.38)
100 mg	-2139.66 (-7109.18 to 2829.86)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-5492.66 (-12051.33 to 1066.01)
50 mg vs 0 mg	-3239.70 (-10007.37 to 3527.98)
100 mg vs 0 mg	-6739.15 (-14799.22 to 1320.91)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-4835.72 (-9669.36 to -2.08)

Mean relative fractional anisotropy

A total of 51 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.05 (-0.08 to -0.02)
25 mg	-0.02 (-0.05 to 0.02)
50 mg	-0.07 (-0.10 to -0.03)
100 mg	-0.06 (-0.09 to -0.02)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.03 (-0.02 to 0.08)
50 mg vs 0 mg	-0.02 (-0.06 to 0.03)
100 mg vs 0 mg	-0.01 (-0.06 to 0.04)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.00 (-0.03 to 0.03)

Mean relative restricted spectrum imaging

A total of 49 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.08 (-0.12 to -0.03)
25 mg	-0.12 (-0.17 to -0.07)
50 mg	-0.12 (-0.17 to -0.07)
100 mg	-0.09 (-0.14 to -0.04)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.05 (-0.12 to 0.02)
50 mg vs 0 mg	-0.04 (-0.11 to 0.03)
100 mg vs 0 mg	-0.02 (-0.09 to 0.05)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.04 (-0.08 to 0.01)

Mean relative transit time

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.03 (-0.11 to 0.04)
25 mg	-0.14 (-0.22 to -0.06)
50 mg	-0.05 (-0.13 to 0.04)
100 mg	-0.06 (-0.15 to 0.02)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.10 (-0.22 to 0.02)
50 mg vs 0 mg	-0.01 (-0.14 to 0.11)
100 mg vs 0 mg	-0.03 (-0.15 to 0.09)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.05 (-0.13 to 0.03)

Mean relative vessel caliber

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.06 (-0.13 to 0.25)
25 mg	0.10 (-0.09 to 0.30)
50 mg	0.18 (-0.02 to 0.37)
100 mg	-0.11 (-0.39 to 0.17)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.05 (-0.26 to 0.35)
50 mg vs 0 mg	0.12 (-0.20 to 0.44)
100 mg vs 0 mg	-0.17 (-0.56 to 0.23)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.03 (-0.18 to 0.25)

Mean relative vessel size index

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.09 (-0.32 to 0.50)
25 mg	0.20 (-0.22 to 0.61)
50 mg	0.44 (0.03 to 0.86)
100 mg	0.01 (-0.58 to 0.60)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.11 (-0.55 to 0.77)
50 mg vs 0 mg	0.36 (-0.33 to 1.04)
100 mg vs 0 mg	-0.08 (-0.92 to 0.77)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.15 (-0.31 to 0.62)

Mean relative vortex area

A total of 27 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.03 (-0.02 to 0.08)
25 mg	-0.02 (-0.07 to 0.03)
50 mg	0.00 (-0.05 to 0.05)
100 mg	0.03 (-0.05 to 0.10)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.06 (-0.14 to 0.02)
50 mg vs 0 mg	-0.03 (-0.11 to 0.05)
100 mg vs 0 mg	-0.01 (-0.11 to 0.10)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.04 (-0.09 to 0.02)

Mean tissue elasticity

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.04 (0.00 to 0.09)
25 mg	0.02 (-0.03 to 0.06)
50 mg	-0.01 (-0.06 to 0.04)
100 mg	-0.01 (-0.06 to 0.04)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.03 (-0.09 to 0.04)
50 mg vs 0 mg	-0.05 (-0.13 to 0.02)
100 mg vs 0 mg	-0.05 (-0.12 to 0.03)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.04 (-0.09 to 0.01)

Mean tissue relative viscosity

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.01 (-0.06 to 0.05)
25 mg	0.02 (-0.03 to 0.07)
50 mg	-0.07 (-0.13 to -0.01)
100 mg	-0.01 (-0.06 to 0.05)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.02 (-0.06 to 0.11)
50 mg vs 0 mg	-0.07 (-0.17 to 0.03)
100 mg vs 0 mg	0.00 (-0.09 to 0.09)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.00 (-0.07 to 0.06)

Mean tissue stiffness

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.03 (-0.01 to 0.08)
25 mg	0.02 (-0.03 to 0.06)
50 mg	-0.04 (-0.09 to 0.01)
100 mg	0.01 (-0.04 to 0.06)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.02 (-0.08 to 0.04)
50 mg vs 0 mg	-0.08 (-0.15 to 0.00)
100 mg vs 0 mg	-0.03 (-0.10 to 0.04)

Plot




Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.04 (-0.08 to 0.01)

Mean tissue viscosity

A total of 40 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	0.04 (-0.04 to 0.11)
25 mg	0.05 (-0.03 to 0.13)
50 mg	-0.09 (-0.18 to 0.00)
100 mg	0.05 (-0.04 to 0.14)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.01 (-0.11 to 0.13)
50 mg vs 0 mg	-0.13 (-0.26 to 0.01)
100 mg vs 0 mg	0.02 (-0.11 to 0.15)

Plot



Supplementary analysis: any dose vs 0 mg


Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.02 (-0.11 to 0.06)

Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start

Mean Response over Time for Total volume of contrast agent enhancing lesion




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	-0.06 (-1.56 to 1.43)
25 mg	0.35 (-1.35 to 2.06)
50 mg	-0.50 (-2.23 to 1.24)
100 mg	-0.58 (-2.32 to 1.16)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	0.42 (-1.69 to 2.52)
50 mg vs 0 mg	-0.43 (-2.60 to 1.74)
100 mg vs 0 mg	-0.52 (-2.66 to 1.62)

Plot



Supplementary analysis: any dose vs 0 mg


Comparison	Difference (95% CI)
Any dose vs 0 mg	-0.16 (-1.48 to 1.16)

Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3

A total of 53 subjects were included in the analysis for this endpoint.

Line plot by treatment start

Mean Response over Time for Total volume of pathologic signal on T2/FLAIR




Means by dose

Dose	Estimated change from baseline (95% CI)
0 mg	6.23 (1.20 to 11.25)
25 mg	6.04 (0.25 to 11.83)
50 mg	6.19 (0.32 to 12.06)
100 mg	8.10 (2.21 to 13.98)

Differences vs 0 mg

Comparison	Difference (95% CI)
25 mg vs 0 mg	-0.18 (-7.64 to 7.27)
50 mg vs 0 mg	-0.03 (-7.65 to 7.58)
100 mg vs 0 mg	1.87 (-5.64 to 9.39)

Plot



Supplementary analysis: any dose vs 0 mg

Comparison	Difference (95% CI)
Any dose vs 0 mg	0.54 (-4.15 to 5.23)

MRI Endpoints at Cycle 11 (Day 141)

For all MRI endpoints, we report estimated change from baseline at Cycle 11 (day 141), grouped by the extended treatment variable. The summaries include mean estimates, contrasts versus the reference group, and a plot with confidence intervals.

Mean DSC K2 leakage

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.85 (-2.06 to 0.35)
losartan 25 mg	-2.15 (-4.14 to -0.16)
losartan 50 mg	-1.13 (-2.93 to 0.68)
losartan 100 mg	-1.24 (-2.63 to 0.14)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-1.30 (-4.10 to 1.51)
losartan 50 mg - losartan 0 mg	-0.28 (-2.91 to 2.36)
losartan 100 mg - losartan 0 mg	-0.39 (-2.63 to 1.84)

Plot



Mean relative DSC extravasation

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.01 (-0.39 to 0.41)
losartan 25 mg	0.38 (-0.36 to 1.11)
losartan 50 mg	-0.06 (-0.77 to 0.64)
losartan 100 mg	-0.27 (-0.97 to 0.44)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.36 (-0.75 to 1.48)
losartan 50 mg - losartan 0 mg	-0.07 (-1.06 to 0.92)
losartan 100 mg - losartan 0 mg	-0.28 (-1.26 to 0.71)

Plot



Mean relative apparent diffusion coefficient

A total of 43 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.23 (0.13 to 0.34)
losartan 25 mg	0.17 (0.01 to 0.34)
losartan 50 mg	0.25 (0.08 to 0.42)
losartan 100 mg	0.30 (0.17 to 0.43)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.06 (-0.30 to 0.18)
losartan 50 mg - losartan 0 mg	0.02 (-0.23 to 0.27)
losartan 100 mg - losartan 0 mg	0.07 (-0.13 to 0.26)

Plot



Mean relative cerebral blood

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.24 (-0.43 to -0.06)
losartan 25 mg	-0.13 (-0.42 to 0.17)
losartan 50 mg	-0.26 (-0.54 to 0.02)
losartan 100 mg	-0.20 (-0.41 to 0.01)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.12 (-0.31 to 0.54)
losartan 50 mg - losartan 0 mg	-0.02 (-0.43 to 0.39)
losartan 100 mg - losartan 0 mg	0.04 (-0.29 to 0.38)

Plot



Mean relative cerebral blood flow (PTA-ROI)

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.18 (-0.31 to -0.04)
losartan 25 mg	-0.11 (-0.33 to 0.11)
losartan 50 mg	-0.21 (-0.42 to 0.00)
losartan 100 mg	-0.13 (-0.28 to 0.03)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.07 (-0.24 to 0.38)
losartan 50 mg - losartan 0 mg	-0.03 (-0.34 to 0.27)
losartan 100 mg - losartan 0 mg	0.05 (-0.20 to 0.30)

Plot



Mean relative cerebral blood flow (ROI2)

A total of 40 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.35 (-0.47 to -0.22)
losartan 25 mg	-0.18 (-0.39 to 0.04)
losartan 50 mg	-0.36 (-0.56 to -0.16)
losartan 100 mg	-0.23 (-0.40 to -0.05)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.17 (-0.13 to 0.47)
losartan 50 mg - losartan 0 mg	-0.01 (-0.30 to 0.27)
losartan 100 mg - losartan 0 mg	0.12 (-0.14 to 0.38)

Plot



Mean relative direction value

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	2.20 (-38.97 to 43.37)
losartan 25 mg	72.20 (7.03 to 137.37)
losartan 50 mg	14.32 (-69.57 to 98.21)
losartan 100 mg	-17.81 (-105.95 to 70.34)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	69.99 (-25.02 to 165.01)
losartan 50 mg - losartan 0 mg	12.12 (-102.56 to 126.79)
losartan 100 mg - losartan 0 mg	-20.01 (-140.71 to 100.69)

Plot



Mean relative fractional anisotropy

A total of 40 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.00 (-0.07 to 0.06)
losartan 25 mg	-0.04 (-0.15 to 0.06)
losartan 50 mg	-0.09 (-0.21 to 0.04)
losartan 100 mg	-0.12 (-0.20 to -0.03)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.04 (-0.19 to 0.11)
losartan 50 mg - losartan 0 mg	-0.08 (-0.25 to 0.09)
losartan 100 mg - losartan 0 mg	-0.12 (-0.25 to 0.02)

Plot



Mean relative restricted spectrum imaging

A total of 40 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.16 (-0.22 to -0.09)
losartan 25 mg	-0.14 (-0.25 to -0.03)
losartan 50 mg	-0.23 (-0.33 to -0.13)
losartan 100 mg	-0.20 (-0.29 to -0.11)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.02 (-0.13 to 0.17)
losartan 50 mg - losartan 0 mg	-0.07 (-0.22 to 0.08)
losartan 100 mg - losartan 0 mg	-0.04 (-0.17 to 0.09)

Plot



Mean relative solid stress (PTA-ROI)

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.02 (-0.07 to 0.11)
losartan 25 mg	0.03 (-0.12 to 0.18)
losartan 50 mg	0.04 (-0.11 to 0.19)
losartan 100 mg	-0.07 (-0.20 to 0.07)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.01 (-0.21 to 0.22)
losartan 50 mg - losartan 0 mg	0.02 (-0.19 to 0.23)
losartan 100 mg - losartan 0 mg	-0.09 (-0.28 to 0.11)

Plot



Mean relative solid stress (ROI2)

A total of 23 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.02 (-0.09 to 0.05)
losartan 25 mg	-0.03 (-0.17 to 0.10)
losartan 50 mg	0.07 (-0.05 to 0.19)
losartan 100 mg	-0.14 (-0.26 to -0.03)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.01 (-0.20 to 0.17)
losartan 50 mg - losartan 0 mg	0.09 (-0.07 to 0.26)
losartan 100 mg - losartan 0 mg	-0.12 (-0.29 to 0.04)

Plot



Mean relative transit time

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	-0.03 (-0.12 to 0.05)
losartan 25 mg	-0.07 (-0.21 to 0.06)
losartan 50 mg	-0.04 (-0.17 to 0.09)
losartan 100 mg	-0.16 (-0.26 to -0.06)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.04 (-0.24 to 0.16)
losartan 50 mg - losartan 0 mg	-0.01 (-0.20 to 0.18)
losartan 100 mg - losartan 0 mg	-0.12 (-0.28 to 0.04)

Plot



Mean relative vessel caliber

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.16 (-0.08 to 0.40)
losartan 25 mg	0.17 (-0.22 to 0.56)
losartan 50 mg	-0.12 (-0.56 to 0.31)
losartan 100 mg	-0.05 (-0.48 to 0.38)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.01 (-0.56 to 0.59)
losartan 50 mg - losartan 0 mg	-0.28 (-0.89 to 0.32)
losartan 100 mg - losartan 0 mg	-0.21 (-0.82 to 0.40)

Plot



Mean relative vessel size index

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.34 (-0.10 to 0.78)
losartan 25 mg	0.17 (-0.54 to 0.88)
losartan 50 mg	-0.21 (-1.01 to 0.59)
losartan 100 mg	0.07 (-0.72 to 0.87)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.17 (-1.20 to 0.86)
losartan 50 mg - losartan 0 mg	-0.55 (-1.68 to 0.57)
losartan 100 mg - losartan 0 mg	-0.26 (-1.39 to 0.86)

Plot



Mean relative vortex area

A total of 20 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.03 (-0.03 to 0.08)
losartan 25 mg	-0.01 (-0.10 to 0.08)
losartan 50 mg	0.06 (-0.05 to 0.17)
losartan 100 mg	0.05 (-0.06 to 0.15)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.04 (-0.17 to 0.09)
losartan 50 mg - losartan 0 mg	0.04 (-0.12 to 0.19)
losartan 100 mg - losartan 0 mg	0.02 (-0.13 to 0.17)

Plot



Mean tissue elasticity

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.03 (-0.05 to 0.11)
losartan 25 mg	0.06 (-0.07 to 0.19)
losartan 50 mg	0.08 (-0.05 to 0.21)
losartan 100 mg	-0.06 (-0.17 to 0.06)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.03 (-0.16 to 0.21)
losartan 50 mg - losartan 0 mg	0.05 (-0.13 to 0.23)
losartan 100 mg - losartan 0 mg	-0.09 (-0.26 to 0.08)

Plot



Mean tissue relative viscosity

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.03 (-0.04 to 0.10)
losartan 25 mg	-0.03 (-0.14 to 0.09)
losartan 50 mg	-0.05 (-0.16 to 0.07)
losartan 100 mg	-0.02 (-0.12 to 0.09)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.06 (-0.22 to 0.11)
losartan 50 mg - losartan 0 mg	-0.08 (-0.24 to 0.08)
losartan 100 mg - losartan 0 mg	-0.05 (-0.20 to 0.10)

Plot



Mean tissue stiffness

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.02 (-0.07 to 0.11)
losartan 25 mg	0.03 (-0.12 to 0.18)
losartan 50 mg	0.04 (-0.11 to 0.19)
losartan 100 mg	-0.07 (-0.20 to 0.07)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	0.01 (-0.21 to 0.22)
losartan 50 mg - losartan 0 mg	0.02 (-0.19 to 0.23)
losartan 100 mg - losartan 0 mg	-0.09 (-0.28 to 0.11)

Plot



Mean tissue viscosity

A total of 29 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	0.06 (-0.06 to 0.18)
losartan 25 mg	0.04 (-0.16 to 0.24)
losartan 50 mg	0.03 (-0.18 to 0.24)
losartan 100 mg	-0.08 (-0.26 to 0.11)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.02 (-0.31 to 0.27)
losartan 50 mg - losartan 0 mg	-0.03 (-0.32 to 0.26)
losartan 100 mg - losartan 0 mg	-0.14 (-0.41 to 0.13)

Plot



Total volume of contrast agent enhancing lesion on T1-weighted MRI in cm3

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	2.51 (-0.92 to 5.94)
losartan 25 mg	1.14 (-4.30 to 6.57)
losartan 50 mg	1.58 (-3.55 to 6.72)
losartan 100 mg	1.28 (-2.57 to 5.12)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-1.37 (-9.19 to 6.44)
losartan 50 mg - losartan 0 mg	-0.93 (-8.50 to 6.65)
losartan 100 mg - losartan 0 mg	-1.23 (-7.51 to 5.04)

Plot



Total volume of pathologic signal on T2/FLAIR-weighted MRI in cm3

A total of 47 subjects were included in the analysis for this endpoint.


Means by treatment

Treatment	Estimated change from baseline (95% CI)
losartan 0 mg	22.48 (7.88 to 37.07)
losartan 25 mg	21.79 (-1.48 to 45.06)
losartan 50 mg	13.16 (-8.61 to 34.94)
losartan 100 mg	15.51 (-1.04 to 32.06)

Differences vs reference

Comparison	Difference (95% CI)
losartan 25 mg - losartan 0 mg	-0.69 (-34.00 to 32.62)
losartan 50 mg - losartan 0 mg	-9.31 (-41.20 to 22.57)
losartan 100 mg - losartan 0 mg	-6.97 (-33.91 to 19.97)

Plot




Neurologic Performance

Change from baseline in KPS, ECOG, and NANO scores at Days 15/43 and model-based estimates at Days 155/239 are reported below for the newly diagnosed glioblastoma cohort. Note that since the day 29 is not measured for any of the measures, the dose-response analyses are based on Days 15 and 43 only for the early timepoints. There might therefore be a limited ability to detect dose-response relationships at these early visits because we are lacking one contrast.

Karnofsky Performance Scale

A total of 54 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Days 15/43 (change from baseline)

Dose (mg)	Estimated change from baseline (95% CI)
0	-0.63 (-2.41 to 1.14)
25	0.89 (-1.03 to 2.81)
50	1.19 (-0.73 to 3.11)
100	-1.55 (-3.49 to 0.39)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg0	1.53 (-1.59 to 4.65)
losartan 50 mg vs losartan 0 mg0	1.83 (-1.30 to 4.95)
losartan 100 mg vs losartan 0 mg0	-0.91 (-4.03 to 2.21)

Plot




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-0.77 (-3.70 to 2.16)
155	losartan 25 mg	-0.23 (-4.66 to 4.20)
155	losartan 50 mg	4.23 (0.00 to 8.46)
155	losartan 100 mg	-3.51 (-6.73 to -0.29)
239	losartan 0 mg	1.63 (-1.56 to 4.83)
239	losartan 25 mg	-0.84 (-5.82 to 4.14)
239	losartan 50 mg	3.76 (-0.66 to 8.17)
239	losartan 100 mg	-0.28 (-3.59 to 3.02)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	0.54 (-5.86 to 6.94)
155	losartan 50 mg - losartan 0 mg	5.00 (-1.23 to 11.22)
155	losartan 100 mg - losartan 0 mg	-2.74 (-8.00 to 2.53)
239	losartan 25 mg - losartan 0 mg	-2.47 (-9.60 to 4.65)
239	losartan 50 mg - losartan 0 mg	2.12 (-4.47 to 8.72)
239	losartan 100 mg - losartan 0 mg	-1.92 (-7.48 to 3.64)


Plot



ECOG Score

A total of 54 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Days 15/43 (change from baseline)

Dose (mg)	Estimated change from baseline (95% CI)
0	0.08 (-0.12 to 0.27)
25	-0.05 (-0.25 to 0.14)
50	-0.04 (-0.23 to 0.16)
100	0.11 (-0.09 to 0.31)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	-0.13 (-0.49 to 0.23)
losartan 50 mg vs losartan 0 mg	-0.12 (-0.48 to 0.25)
losartan 100 mg vs losartan 0 mg	0.03 (-0.34 to 0.40)

Plot




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	0.09 (-0.14 to 0.31)
155	losartan 25 mg	-0.27 (-0.62 to 0.09)
155	losartan 50 mg	-0.17 (-0.51 to 0.18)
155	losartan 100 mg	0.29 (0.03 to 0.55)
239	losartan 0 mg	0.03 (-0.24 to 0.29)
239	losartan 25 mg	-0.21 (-0.62 to 0.20)
239	losartan 50 mg	-0.30 (-0.66 to 0.06)
239	losartan 100 mg	0.02 (-0.25 to 0.29)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	-0.35 (-0.86 to 0.16)
155	losartan 50 mg - losartan 0 mg	-0.25 (-0.75 to 0.25)
155	losartan 100 mg - losartan 0 mg	0.21 (-0.21 to 0.62)
239	losartan 25 mg - losartan 0 mg	-0.23 (-0.82 to 0.35)
239	losartan 50 mg - losartan 0 mg	-0.32 (-0.87 to 0.22)
239	losartan 100 mg - losartan 0 mg	0.00 (-0.46 to 0.45)


Plot



Total NANO score

A total of 54 subjects were included in the analysis for this endpoint.

Line plot by treatment start




Days 15/43 (change from baseline)

Dose (mg)	Estimated change from baseline (95% CI)
0	-0.19 (-0.49 to 0.12)
25	-0.01 (-0.34 to 0.32)
50	-0.14 (-0.48 to 0.20)
100	-0.15 (-0.49 to 0.18)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg0	0.18 (-0.36 to 0.72)
losartan 50 mg vs losartan 0 mg0	0.05 (-0.50 to 0.60)
losartan 100 mg vs losartan 0 mg0	0.04 (-0.50 to 0.57)

Plot




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	0.03 (-0.39 to 0.45)
155	losartan 25 mg	0.46 (-0.17 to 1.10)
155	losartan 50 mg	-0.34 (-0.94 to 0.27)
155	losartan 100 mg	-0.25 (-0.69 to 0.20)
239	losartan 0 mg	-0.44 (-0.89 to 0.02)
239	losartan 25 mg	0.16 (-0.54 to 0.86)
239	losartan 50 mg	-0.19 (-0.82 to 0.44)
239	losartan 100 mg	-0.22 (-0.70 to 0.26)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	0.43 (-0.49 to 1.36)
155	losartan 50 mg - losartan 0 mg	-0.37 (-1.26 to 0.53)
155	losartan 100 mg - losartan 0 mg	-0.27 (-1.02 to 0.47)
239	losartan 25 mg - losartan 0 mg	0.60 (-0.41 to 1.61)
239	losartan 50 mg - losartan 0 mg	0.24 (-0.69 to 1.18)
239	losartan 100 mg - losartan 0 mg	0.22 (-0.58 to 1.02)

Plot



Quality of Life Outcomes

Model based estimates of change from baseline in QLQ-C30 and QLQ-BN20 scores at Days 155/239 are reported below for the newly diagnosed glioblastoma cohort. Note that even if the QoL outcomes were collected at Day 43 it is not possible to make a meaningful analysis of the effect of losartan as all participants received the treatment at that time point.

EORTC QLQ-C30 score

A total of 19 subjects were included in the analysis for this endpoint.


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-2.63 (-7.82 to 2.56)
155	losartan 25 mg	-4.63 (-14.69 to 5.42)
155	losartan 50 mg	1.62 (-6.31 to 9.56)
155	losartan 100 mg	0.27 (-6.63 to 7.17)
239	losartan 0 mg	3.55 (-2.03 to 9.13)
239	losartan 25 mg	-4.63 (-14.69 to 5.42)
239	losartan 50 mg	-3.05 (-12.66 to 6.56)
239	losartan 100 mg	-2.73 (-9.63 to 4.17)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	-2.01 (-15.76 to 11.75)
155	losartan 50 mg - losartan 0 mg	4.25 (-7.29 to 15.79)
155	losartan 100 mg - losartan 0 mg	2.90 (-7.61 to 13.41)
239	losartan 25 mg - losartan 0 mg	-8.19 (-22.19 to 5.82)
239	losartan 50 mg - losartan 0 mg	-6.60 (-20.11 to 6.91)
239	losartan 100 mg - losartan 0 mg	-6.28 (-17.07 to 4.52)

Plot



EORTC QLQ-BN20 score

A total of 18 subjects were included in the analysis for this endpoint.


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-0.86 (-4.46 to 2.74)
155	losartan 25 mg	-3.96 (-11.13 to 3.22)
155	losartan 50 mg	3.50 (-2.15 to 9.15)
155	losartan 100 mg	-1.30 (-6.09 to 3.49)
239	losartan 0 mg	0.60 (-3.14 to 4.34)
239	losartan 25 mg	-3.96 (-11.13 to 3.22)
239	losartan 50 mg	-1.27 (-7.51 to 4.97)
239	losartan 100 mg	-4.05 (-8.84 to 0.74)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	-3.09 (-12.94 to 6.75)
155	losartan 50 mg - losartan 0 mg	4.36 (-3.85 to 12.58)
155	losartan 100 mg - losartan 0 mg	-0.44 (-7.77 to 6.89)
239	losartan 25 mg - losartan 0 mg	-4.56 (-14.47 to 5.36)
239	losartan 50 mg - losartan 0 mg	-1.87 (-10.75 to 7.01)
239	losartan 100 mg - losartan 0 mg	-4.65 (-12.08 to 2.78)

Plot



Steroid Use

Binary steroid use is analysed with longitudinal logistic regression and population-averaged marginal risk differences for early (Days 15/29/43) and late (Days 155/239) visits in the newly diagnosed glioblastoma cohort.

Steroid use

A total of 54 subjects were included in the analysis.


Days 15/29/43 (by dose)

Treatment	Risk (95% CI)
losartan 0 mg	0.224 (0.220 to 0.229)
losartan 25 mg	0.226 (0.221 to 0.230)
losartan 50 mg	0.226 (0.221 to 0.231)
losartan 100 mg	0.225 (0.223 to 0.228)

Risk differences vs 0 mg

Comparison	Risk difference (95% CI)
100 - 0	0.001 (-0.004 to 0.006)
25 - 0	0.002 (-0.006 to 0.009)
50 - 0	0.002 (-0.006 to 0.010)

Plot




Days 155/239 (population-averaged risk)

Visit	Treatment	Risk (95% CI)
155	losartan 0 mg	0.295 (0.084 to 0.505)
239	losartan 0 mg	0.227 (0.013 to 0.440)
155	losartan 25 mg	0.000 (0.000 to 0.000)
239	losartan 25 mg	0.000 (0.000 to 0.000)
155	losartan 50 mg	0.265 (-0.054 to 0.584)
239	losartan 50 mg	0.299 (-0.053 to 0.651)
155	losartan 100 mg	0.331 (0.078 to 0.584)
239	losartan 100 mg	0.149 (-0.051 to 0.348)

Risk differences vs reference

Visit	Comparison	Risk difference (95% CI)
155	losartan 100 mg - losartan 0 mg	0.034 (-0.275 to 0.343)
155	losartan 25 mg - losartan 0 mg	-0.300 (-0.499 to -0.102)
155	losartan 50 mg - losartan 0 mg	-0.028 (-0.393 to 0.337)
239	losartan 100 mg - losartan 0 mg	-0.078 (-0.366 to 0.210)
239	losartan 25 mg - losartan 0 mg	-0.241 (-0.446 to -0.036)
239	losartan 50 mg - losartan 0 mg	0.070 (-0.326 to 0.466)

Plot



RANO Responses

Ordinal RANO outcomes are analysed using proportional odds models within the configured cohort. For State of disease according to the RANO criteria, visits at Days 141 and 225 are included; for Best overall radiographic response, only Day 225 is included. Proportional odds ratios and confidence intervals are reported below.

State of disease according to the RANO criteria

Visit 141

Comparison	Propoprtional Odds Ratio (95% CI)
losartan 25 vs 0 mg	1.226 (0.189 to 7.519)
losartan 50 vs 0 mg	0.763 (0.128 to 3.986)
losartan 100 vs 0 mg	2.384 (0.606 to 10.009)

Visit 225

Comparison	Propoprtional Odds Ratio (95% CI)
losartan 25 vs 0 mg	0.545 (0.063 to 5.303)
losartan 50 vs 0 mg	0.485 (0.070 to 3.359)
losartan 100 vs 0 mg	0.727 (0.133 to 3.936)

Best overall radiographic response

Visit 225

Comparison	Propoprtional Odds Ratio (95% CI)
losartan 25 vs 0 mg	2.034 (0.160 to 33.506)
losartan 50 vs 0 mg	0.275 (0.038 to 1.865)
losartan 100 vs 0 mg	1.249 (0.199 to 8.688)

Biomarker Endpoints (Serum Cytokines / Angiogenic Factors)

Serum cytokine and angiogenic-factor concentrations from Bio-Plex 19-plex and 14-plex panels are analysed as exploratory biomarker endpoints. For each analyte, a linear mixed model with random subject intercept is fitted with change from Screening baseline as the response. Where the log-scale model fits better (lower AIC), the analysis is performed on the log-concentration scale and results are labelled accordingly.

Biomarker samples are collected at screening (baseline), Day 15 (Cycle 2), and Day 43 (Cycle 4) for the early analysis, and again at Cycle 17 (Day 225) and later for the long-term analysis. The early analysis uses the stepped-wedge treatment assignment; the long-term analysis uses the assigned long-term dose (`trt02p`). The early analyses results are presented first, then the long term.

A formal model selection and residual diagnostic assessment is available in a separate document (`impress_lb_model_fit.pdf`).

Fractalkine

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.022 (-0.232 to 0.276)
25 mg	-0.128 (-0.384 to 0.128)
50 mg	0.254 (-0.013 to 0.521)
100 mg	0.095 (-0.181 to 0.371)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.15 (-0.61 to 0.31)
losartan 50 mg vs losartan 0 mg0	0.23 (-0.25 to 0.71)
losartan 100 mg vs losartan 0 mg0	0.07 (-0.42 to 0.56)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1458.8	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	219.4	TRUE

Plot



Granzyme B

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.105 (-0.284 to 0.074)
25 mg	-0.416 (-0.607 to -0.225)
50 mg	0.048 (-0.153 to 0.249)
100 mg	-0.045 (-0.249 to 0.160)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.31 (-0.63 to 0.01)
losartan 50 mg vs losartan 0 mg0	0.15 (-0.18 to 0.49)
losartan 100 mg vs losartan 0 mg0	0.06 (-0.28 to 0.40)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	820.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	150.7	TRUE

Plot



IFN-g

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.077 (-0.409 to 0.255)
25 mg	-0.175 (-0.535 to 0.185)
50 mg	-0.221 (-0.607 to 0.165)
100 mg	-0.344 (-0.732 to 0.045)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.10 (-0.68 to 0.49)
losartan 50 mg vs losartan 0 mg0	-0.14 (-0.77 to 0.48)
losartan 100 mg vs losartan 0 mg0	-0.27 (-0.90 to 0.36)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	624.1	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	271.7	TRUE

Plot



IL-1b

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.001 (-0.355 to 0.358)
25 mg	-0.084 (-0.490 to 0.322)
50 mg	-0.156 (-0.582 to 0.271)
100 mg	-0.085 (-0.516 to 0.346)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.09 (-0.68 to 0.51)
losartan 50 mg vs losartan 0 mg0	-0.16 (-0.79 to 0.48)
losartan 100 mg vs losartan 0 mg0	-0.09 (-0.73 to 0.55)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	512.8	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	288.5	TRUE

Plot



IL-6

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.467 (-0.050 to 0.984)
25 mg	-0.671 (-1.207 to -0.134)
50 mg	-0.174 (-0.741 to 0.394)
100 mg	-0.563 (-1.140 to 0.013)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-1.14 (-2.06 to -0.21)
losartan 50 mg vs losartan 0 mg0	-0.64 (-1.62 to 0.34)
losartan 100 mg vs losartan 0 mg0	-1.03 (-2.02 to -0.04)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	387.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	355.0	TRUE

Plot



IL-8

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.076 (-0.190 to 0.037)
25 mg	-0.081 (-0.207 to 0.044)
50 mg	-0.029 (-0.161 to 0.104)
100 mg	-0.114 (-0.249 to 0.021)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.00 (-0.20 to 0.19)
losartan 50 mg vs losartan 0 mg0	0.05 (-0.16 to 0.26)
losartan 100 mg vs losartan 0 mg0	-0.04 (-0.25 to 0.18)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	425.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	64.9	TRUE

Plot



MCP-1

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.040 (-0.180 to 0.260)
25 mg	-0.157 (-0.403 to 0.089)
50 mg	0.020 (-0.240 to 0.279)
100 mg	-0.407 (-0.668 to -0.147)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.20 (-0.58 to 0.19)
losartan 50 mg vs losartan 0 mg0	-0.02 (-0.43 to 0.39)
losartan 100 mg vs losartan 0 mg0	-0.45 (-0.86 to -0.04)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1144.2	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	192.4	TRUE

Plot



MMP-10

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.046 (-0.047 to 0.140)
25 mg	0.024 (-0.078 to 0.126)
50 mg	0.062 (-0.044 to 0.169)
100 mg	0.103 (-0.005 to 0.211)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.02 (-0.19 to 0.14)
losartan 50 mg vs losartan 0 mg0	0.02 (-0.16 to 0.19)
losartan 100 mg vs losartan 0 mg0	0.06 (-0.12 to 0.23)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1254.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	25.5	TRUE

Plot



PDGF-DD

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.205 (-0.427 to 0.018)
25 mg	-0.350 (-0.599 to -0.101)
50 mg	-0.146 (-0.408 to 0.116)
100 mg	-0.434 (-0.699 to -0.168)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.15 (-0.53 to 0.24)
losartan 50 mg vs losartan 0 mg0	0.06 (-0.35 to 0.46)
losartan 100 mg vs losartan 0 mg0	-0.23 (-0.64 to 0.18)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1106.7	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	195.4	TRUE

Plot



PECAM-1

A total of 52 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.172 (-0.481 to 0.137)
25 mg	-0.402 (-0.731 to -0.072)
50 mg	0.020 (-0.313 to 0.353)
100 mg	-0.436 (-0.777 to -0.095)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.23 (-0.79 to 0.33)
losartan 50 mg vs losartan 0 mg0	0.19 (-0.38 to 0.76)
losartan 100 mg vs losartan 0 mg0	-0.26 (-0.85 to 0.32)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	2298.7	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	248.3	TRUE

Plot



TGF-alpha

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.025 (-0.340 to 0.289)
25 mg	-0.059 (-0.397 to 0.280)
50 mg	0.196 (-0.160 to 0.552)
100 mg	0.123 (-0.239 to 0.485)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.03 (-0.59 to 0.53)
losartan 50 mg vs losartan 0 mg0	0.22 (-0.37 to 0.81)
losartan 100 mg vs losartan 0 mg0	0.15 (-0.45 to 0.75)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	539.2	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	261.7	TRUE

Plot



Thrombospondin-2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.058 (-0.183 to 0.066)
25 mg	-0.084 (-0.225 to 0.058)
50 mg	-0.057 (-0.206 to 0.091)
100 mg	-0.175 (-0.325 to -0.025)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.03 (-0.23 to 0.18)
losartan 50 mg vs losartan 0 mg0	0.00 (-0.22 to 0.22)
losartan 100 mg vs losartan 0 mg0	-0.12 (-0.34 to 0.11)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1880.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	83.1	TRUE

Plot



VEGF-A

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.292 (-0.512 to -0.072)
25 mg	-0.494 (-0.735 to -0.253)
50 mg	-0.330 (-0.585 to -0.075)
100 mg	-0.322 (-0.580 to -0.064)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.20 (-0.59 to 0.18)
losartan 50 mg vs losartan 0 mg0	-0.04 (-0.45 to 0.37)
losartan 100 mg vs losartan 0 mg0	-0.03 (-0.44 to 0.38)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1077.7	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	192.6	TRUE

Plot



VEGF-C

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.073 (-0.237 to 0.091)
25 mg	-0.122 (-0.311 to 0.066)
50 mg	-0.033 (-0.230 to 0.165)
100 mg	-0.277 (-0.476 to -0.078)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.05 (-0.32 to 0.22)
losartan 50 mg vs losartan 0 mg0	0.04 (-0.25 to 0.33)
losartan 100 mg vs losartan 0 mg0	-0.20 (-0.49 to 0.09)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1476.2	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	136.8	TRUE

Plot



Angiopoietin-1

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.074 (-0.301 to 0.152)
25 mg	-0.216 (-0.464 to 0.032)
50 mg	-0.170 (-0.427 to 0.088)
100 mg	-0.379 (-0.643 to -0.115)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.14 (-0.55 to 0.26)
losartan 50 mg vs losartan 0 mg0	-0.10 (-0.52 to 0.33)
losartan 100 mg vs losartan 0 mg0	-0.30 (-0.74 to 0.13)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1917.7	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	197.6	TRUE

Plot



Angiopoietin-2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.004 (-0.074 to 0.083)
25 mg	-0.060 (-0.145 to 0.025)
50 mg	-0.057 (-0.147 to 0.033)
100 mg	-0.077 (-0.170 to 0.016)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.06 (-0.20 to 0.07)
losartan 50 mg vs losartan 0 mg0	-0.06 (-0.21 to 0.09)
losartan 100 mg vs losartan 0 mg0	-0.08 (-0.23 to 0.07)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1604.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	-7.0	TRUE

Plot



ANGPTL4

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.070 (-0.166 to 0.026)
25 mg	-0.012 (-0.119 to 0.094)
50 mg	-0.057 (-0.171 to 0.058)
100 mg	-0.027 (-0.143 to 0.089)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.06 (-0.11 to 0.22)
losartan 50 mg vs losartan 0 mg0	0.01 (-0.17 to 0.19)
losartan 100 mg vs losartan 0 mg0	0.04 (-0.14 to 0.22)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	2349.9	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	31.9	TRUE

Plot



CA9

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.129 (-0.139 to 0.396)
25 mg	0.000 (-0.300 to 0.300)
50 mg	0.211 (-0.107 to 0.529)
100 mg	-0.118 (-0.444 to 0.207)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.13 (-0.59 to 0.33)
losartan 50 mg vs losartan 0 mg0	0.08 (-0.40 to 0.57)
losartan 100 mg vs losartan 0 mg0	-0.25 (-0.75 to 0.25)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1075.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	231.5	TRUE

Plot



CD163

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.108 (0.017 to 0.200)
25 mg	0.082 (-0.019 to 0.182)
50 mg	0.135 (0.029 to 0.241)
100 mg	0.049 (-0.058 to 0.156)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.03 (-0.19 to 0.13)
losartan 50 mg vs losartan 0 mg0	0.03 (-0.14 to 0.20)
losartan 100 mg vs losartan 0 mg0	-0.06 (-0.23 to 0.11)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	2660.7	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	22.6	TRUE

Plot



Collagen IV alpha 1

A total of 52 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.099 (0.012 to 0.186)
25 mg	0.039 (-0.058 to 0.136)
50 mg	0.141 (0.041 to 0.240)
100 mg	0.065 (-0.036 to 0.167)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.06 (-0.21 to 0.09)
losartan 50 mg vs losartan 0 mg0	0.04 (-0.12 to 0.20)
losartan 100 mg vs losartan 0 mg0	-0.03 (-0.20 to 0.13)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1328.6	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	11.3	TRUE

Plot



EGF

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.717 (-1.469 to 0.036)
25 mg	-0.774 (-1.604 to 0.056)
50 mg	-0.174 (-1.046 to 0.698)
100 mg	-1.310 (-2.197 to -0.423)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.06 (-1.38 to 1.26)
losartan 50 mg vs losartan 0 mg0	0.54 (-0.85 to 1.93)
losartan 100 mg vs losartan 0 mg0	-0.59 (-2.01 to 0.82)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	869.9	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	430.9	TRUE

Plot



ErbB2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.080 (0.005 to 0.156)
25 mg	0.058 (-0.028 to 0.145)
50 mg	0.032 (-0.058 to 0.122)
100 mg	-0.093 (-0.184 to -0.001)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.02 (-0.15 to 0.10)
losartan 50 mg vs losartan 0 mg0	-0.05 (-0.18 to 0.09)
losartan 100 mg vs losartan 0 mg0	-0.17 (-0.31 to -0.04)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1697	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	-13	TRUE

Plot



FGF Basic

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.188 (-0.428 to 0.052)
25 mg	-0.237 (-0.505 to 0.031)
50 mg	-0.240 (-0.523 to 0.044)
100 mg	-0.291 (-0.577 to -0.004)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.05 (-0.46 to 0.36)
losartan 50 mg vs losartan 0 mg0	-0.05 (-0.49 to 0.39)
losartan 100 mg vs losartan 0 mg0	-0.10 (-0.55 to 0.34)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	649.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	209.9	TRUE

Plot



IL-12 p70

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.199 (-0.095 to 0.493)
25 mg	0.181 (-0.159 to 0.521)
50 mg	0.086 (-0.270 to 0.442)
100 mg	0.099 (-0.257 to 0.456)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.02 (-0.47 to 0.43)
losartan 50 mg vs losartan 0 mg0	-0.11 (-0.60 to 0.37)
losartan 100 mg vs losartan 0 mg0	-0.10 (-0.58 to 0.38)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1279.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	248.8	TRUE

Plot



IL-2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.278 (-0.247 to 0.803)
25 mg	-0.154 (-0.707 to 0.400)
50 mg	-0.354 (-0.936 to 0.229)
100 mg	-0.486 (-1.082 to 0.110)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.43 (-1.37 to 0.51)
losartan 50 mg vs losartan 0 mg0	-0.63 (-1.62 to 0.35)
losartan 100 mg vs losartan 0 mg0	-0.76 (-1.77 to 0.24)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	786.7	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	360.3	TRUE

Plot



IP-10 (CXCL10)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.262 (0.060 to 0.463)
25 mg	0.159 (-0.062 to 0.380)
50 mg	0.202 (-0.029 to 0.434)
100 mg	0.184 (-0.056 to 0.424)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.10 (-0.46 to 0.25)
losartan 50 mg vs losartan 0 mg0	-0.06 (-0.43 to 0.31)
losartan 100 mg vs losartan 0 mg0	-0.08 (-0.46 to 0.31)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1152.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	174.9	TRUE

Plot



I-TAC (CXCL11)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.000 (-0.096 to 0.096)
25 mg	-0.055 (-0.166 to 0.056)
50 mg	0.017 (-0.100 to 0.133)
100 mg	-0.045 (-0.163 to 0.072)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.05 (-0.21 to 0.10)
losartan 50 mg vs losartan 0 mg0	0.02 (-0.15 to 0.18)
losartan 100 mg vs losartan 0 mg0	-0.05 (-0.21 to 0.12)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1171.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	33.0	TRUE

Plot



MIG (CXCL9)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.068 (-0.397 to 0.534)
25 mg	-0.100 (-0.619 to 0.419)
50 mg	0.155 (-0.392 to 0.703)
100 mg	0.151 (-0.403 to 0.705)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.17 (-0.97 to 0.64)
losartan 50 mg vs losartan 0 mg0	0.09 (-0.77 to 0.94)
losartan 100 mg vs losartan 0 mg0	0.08 (-0.78 to 0.94)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	1453.2	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	338.2	TRUE

Plot



PIGF

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.049 (-0.021 to 0.119)
25 mg	0.061 (-0.019 to 0.141)
50 mg	0.044 (-0.040 to 0.128)
100 mg	0.079 (-0.006 to 0.163)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	0.01 (-0.11 to 0.13)
losartan 50 mg vs losartan 0 mg0	0.00 (-0.13 to 0.12)
losartan 100 mg vs losartan 0 mg0	0.03 (-0.10 to 0.16)

Model comparison (AIC)

Model	AIC	Preferred
Linear ($\text{chg} \sim \text{base} + \text{trtcd} + \text{avisitn}$)	789.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trtcd} + \text{avisitn}$)	-25.9	TRUE

Plot



Tenascin C

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.061 (0.016 to 0.105)
25 mg	-0.006 (-0.056 to 0.044)
50 mg	0.045 (-0.008 to 0.098)
100 mg	0.025 (-0.028 to 0.079)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.07 (-0.14 to 0.01)
losartan 50 mg vs losartan 0 mg0	-0.02 (-0.10 to 0.07)
losartan 100 mg vs losartan 0 mg0	-0.04 (-0.12 to 0.05)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1752.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	-115.2	TRUE

Plot



TNFRSF18 (GITR)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	-0.071 (-0.319 to 0.176)
25 mg	-0.366 (-0.647 to -0.086)
50 mg	0.086 (-0.209 to 0.381)
100 mg	0.056 (-0.243 to 0.355)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.30 (-0.71 to 0.12)
losartan 50 mg vs losartan 0 mg0	0.16 (-0.28 to 0.60)
losartan 100 mg vs losartan 0 mg0	0.13 (-0.32 to 0.58)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1513.0	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	217.5	TRUE

Plot



TNFSF6 (FasL)

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.047 (-0.065 to 0.159)
25 mg	-0.082 (-0.208 to 0.044)
50 mg	-0.024 (-0.156 to 0.109)
100 mg	-0.098 (-0.232 to 0.037)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.13 (-0.32 to 0.06)
losartan 50 mg vs losartan 0 mg0	-0.07 (-0.27 to 0.13)
losartan 100 mg vs losartan 0 mg0	-0.14 (-0.35 to 0.06)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	876.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	62.8	TRUE

Plot



VEGFR2

A total of 53 participants contributed data for this analyte.

Estimated change from baseline by dose (early analyses (baseline to Day 43))

Dose (mg)	Change from baseline on the natural logarithmic scale (95% CI)
0 mg	0.058 (0.005 to 0.112)
25 mg	0.020 (-0.042 to 0.081)
50 mg	0.008 (-0.056 to 0.071)
100 mg	0.002 (-0.063 to 0.066)


Differences vs 0 mg (early analyses (baseline to Day 43))

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg vs losartan 0 mg0	-0.04 (-0.13 to 0.05)
losartan 50 mg vs losartan 0 mg0	-0.05 (-0.14 to 0.04)
losartan 100 mg vs losartan 0 mg0	-0.06 (-0.15 to 0.04)

Model comparison (AIC)

Model	AIC	Preferred
Linear (chg ~ base + trtcd + avisitn)	1751.4	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trtcd + avisitn)	-81.7	TRUE

Plot



Biomarker Endpoints — Long-term (Cycle 15+)

Model-based estimates of change from baseline at Cycle 15 and later are reported below for analytes with available long-term samples. The same scale (linear or log-transformed) as selected for the early analysis is used for each analyte.

Fractalkine

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.148 (-0.448 to 0.744)
losartan 25 mg	-0.219 (-1.082 to 0.644)
losartan 50 mg	0.642 (-0.094 to 1.378)
losartan 100 mg	0.439 (-0.142 to 1.021)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.37 (-1.65 to 0.92)
losartan 50 mg - losartan 0 mg	0.49 (-0.68 to 1.67)
losartan 100 mg - losartan 0 mg	0.29 (-0.73 to 1.31)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	520.9	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	99.0	TRUE

Plot



Granzyme B

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.608 (-1.388 to 0.172)
losartan 25 mg	-0.550 (-1.702 to 0.602)
losartan 50 mg	-0.381 (-1.371 to 0.610)
losartan 100 mg	-0.684 (-1.458 to 0.089)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.06 (-1.63 to 1.74)
losartan 50 mg - losartan 0 mg	0.23 (-1.33 to 1.78)
losartan 100 mg - losartan 0 mg	-0.08 (-1.42 to 1.27)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	240.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	118.4	TRUE

Plot



IFN-g

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.083 (-0.559 to 0.393)
losartan 25 mg	-0.200 (-0.910 to 0.511)
losartan 50 mg	-0.344 (-0.948 to 0.260)
losartan 100 mg	-0.405 (-0.884 to 0.074)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.12 (-1.17 to 0.93)
losartan 50 mg - losartan 0 mg	-0.26 (-1.19 to 0.67)
losartan 100 mg - losartan 0 mg	-0.32 (-1.15 to 0.51)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	111.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	85.1	TRUE

Plot



IL-1b

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.117 (-0.305 to 0.539)
losartan 25 mg	0.002 (-0.625 to 0.629)
losartan 50 mg	-0.062 (-0.593 to 0.468)
losartan 100 mg	0.104 (-0.319 to 0.526)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.11 (-1.04 to 0.81)
losartan 50 mg - losartan 0 mg	-0.18 (-1.00 to 0.65)
losartan 100 mg - losartan 0 mg	-0.01 (-0.74 to 0.72)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	103.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	77.3	TRUE

Plot



IL-6

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.621 (-1.365 to 0.123)
losartan 25 mg	-0.644 (-1.751 to 0.464)
losartan 50 mg	-0.044 (-0.977 to 0.889)
losartan 100 mg	-0.130 (-0.876 to 0.615)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.02 (-1.65 to 1.60)
losartan 50 mg - losartan 0 mg	0.58 (-0.88 to 2.03)
losartan 100 mg - losartan 0 mg	0.49 (-0.79 to 1.77)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	106.9	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	115.9	TRUE

Plot



IL-8

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.160 (-0.380 to 0.701)
losartan 25 mg	-0.546 (-1.340 to 0.247)
losartan 50 mg	-0.017 (-0.685 to 0.651)
losartan 100 mg	0.175 (-0.364 to 0.713)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.71 (-1.89 to 0.47)
losartan 50 mg - losartan 0 mg	-0.18 (-1.22 to 0.86)
losartan 100 mg - losartan 0 mg	0.01 (-0.93 to 0.96)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	158.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	93.0	TRUE

Plot



MCP-1

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.049 (-0.152 to 0.250)
losartan 25 mg	-0.143 (-0.431 to 0.146)
losartan 50 mg	-0.139 (-0.397 to 0.119)
losartan 100 mg	0.005 (-0.189 to 0.199)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.19 (-0.62 to 0.23)
losartan 50 mg - losartan 0 mg	-0.19 (-0.60 to 0.23)
losartan 100 mg - losartan 0 mg	-0.04 (-0.39 to 0.30)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	364.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	24.3	TRUE

Plot



MMP-10

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.073 (-0.254 to 0.108)
losartan 25 mg	0.035 (-0.234 to 0.304)
losartan 50 mg	-0.100 (-0.328 to 0.127)
losartan 100 mg	0.176 (-0.004 to 0.356)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.11 (-0.29 to 0.51)
losartan 50 mg - losartan 0 mg	-0.03 (-0.38 to 0.33)
losartan 100 mg - losartan 0 mg	0.25 (-0.06 to 0.56)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	435.6	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	19.1	TRUE

Plot



PDGF-DD

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.336 (-0.693 to 0.021)
losartan 25 mg	-0.464 (-0.976 to 0.048)
losartan 50 mg	0.131 (-0.300 to 0.563)
losartan 100 mg	-0.235 (-0.589 to 0.119)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.13 (-0.90 to 0.64)
losartan 50 mg - losartan 0 mg	0.47 (-0.21 to 1.15)
losartan 100 mg - losartan 0 mg	0.10 (-0.53 to 0.73)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	361.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	63.4	TRUE

Plot



PECAM-1

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.174 (-0.680 to 0.332)
losartan 25 mg	0.256 (-0.498 to 1.009)
losartan 50 mg	0.185 (-0.449 to 0.820)
losartan 100 mg	0.247 (-0.258 to 0.753)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.43 (-0.68 to 1.54)
losartan 50 mg - losartan 0 mg	0.36 (-0.63 to 1.35)
losartan 100 mg - losartan 0 mg	0.42 (-0.45 to 1.29)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	832.5	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	89.6	TRUE

Plot



TGF-alpha

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.305 (-0.615 to 0.005)
losartan 25 mg	-0.251 (-0.708 to 0.206)
losartan 50 mg	0.274 (-0.105 to 0.653)
losartan 100 mg	-0.203 (-0.505 to 0.099)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.05 (-0.63 to 0.74)
losartan 50 mg - losartan 0 mg	0.58 (-0.02 to 1.18)
losartan 100 mg - losartan 0 mg	0.10 (-0.43 to 0.63)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	142.0	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	54.5	TRUE

Plot



Thrombospondin-2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.125 (-0.313 to 0.063)
losartan 25 mg	-0.047 (-0.326 to 0.232)
losartan 50 mg	-0.142 (-0.378 to 0.094)
losartan 100 mg	0.036 (-0.151 to 0.224)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.08 (-0.33 to 0.49)
losartan 50 mg - losartan 0 mg	-0.02 (-0.39 to 0.35)
losartan 100 mg - losartan 0 mg	0.16 (-0.16 to 0.49)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	660.8	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	22.3	TRUE

Plot



VEGF-A

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.453 (-0.825 to -0.081)
losartan 25 mg	-0.419 (-0.975 to 0.137)
losartan 50 mg	0.045 (-0.422 to 0.512)
losartan 100 mg	-0.066 (-0.439 to 0.307)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.03 (-0.78 to 0.85)
losartan 50 mg - losartan 0 mg	0.50 (-0.23 to 1.23)
losartan 100 mg - losartan 0 mg	0.39 (-0.25 to 1.03)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	381.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	68.6	TRUE

Plot



VEGF-C

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.172 (-0.476 to 0.131)
losartan 25 mg	-0.248 (-0.686 to 0.190)
losartan 50 mg	0.137 (-0.235 to 0.509)
losartan 100 mg	-0.348 (-0.645 to -0.050)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.08 (-0.73 to 0.58)
losartan 50 mg - losartan 0 mg	0.31 (-0.28 to 0.90)
losartan 100 mg - losartan 0 mg	-0.18 (-0.70 to 0.35)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	495.6	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	53.0	TRUE

Plot



Angiopoietin-1

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.338 (-0.625 to -0.051)
losartan 25 mg	-0.505 (-0.909 to -0.101)
losartan 50 mg	-0.111 (-0.452 to 0.229)
losartan 100 mg	-0.399 (-0.674 to -0.125)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.17 (-0.78 to 0.45)
losartan 50 mg - losartan 0 mg	0.23 (-0.33 to 0.78)
losartan 100 mg - losartan 0 mg	-0.06 (-0.56 to 0.44)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	629.9	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	46.9	TRUE

Plot



Angiopoietin-2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.003 (-0.128 to 0.122)
losartan 25 mg	0.014 (-0.166 to 0.193)
losartan 50 mg	0.013 (-0.139 to 0.166)
losartan 100 mg	0.067 (-0.060 to 0.195)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.02 (-0.25 to 0.28)
losartan 50 mg - losartan 0 mg	0.02 (-0.22 to 0.25)
losartan 100 mg - losartan 0 mg	0.07 (-0.16 to 0.30)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	553.3	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-7.8	TRUE

Plot



ANGPTL4

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.059 (-0.203 to 0.084)
losartan 25 mg	-0.036 (-0.248 to 0.176)
losartan 50 mg	0.013 (-0.179 to 0.204)
losartan 100 mg	0.015 (-0.137 to 0.167)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.02 (-0.29 to 0.34)
losartan 50 mg - losartan 0 mg	0.07 (-0.21 to 0.36)
losartan 100 mg - losartan 0 mg	0.07 (-0.19 to 0.34)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	823.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	3.2	TRUE

Plot



CA9

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.067 (-0.672 to 0.538)
losartan 25 mg	0.188 (-0.703 to 1.079)
losartan 50 mg	0.492 (-0.360 to 1.344)
losartan 100 mg	-0.113 (-0.737 to 0.511)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.26 (-1.06 to 1.57)
losartan 50 mg - losartan 0 mg	0.56 (-0.75 to 1.87)
losartan 100 mg - losartan 0 mg	-0.05 (-1.09 to 1.00)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	353.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	101.3	TRUE

Plot



CD163

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.069 (-0.129 to 0.268)
losartan 25 mg	0.041 (-0.241 to 0.322)
losartan 50 mg	0.144 (-0.094 to 0.382)
losartan 100 mg	0.121 (-0.069 to 0.311)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.03 (-0.46 to 0.40)
losartan 50 mg - losartan 0 mg	0.07 (-0.31 to 0.46)
losartan 100 mg - losartan 0 mg	0.05 (-0.29 to 0.39)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	972.1	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	22.3	TRUE

Plot



Collagen IV alpha 1

A total of 33 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.087 (-0.072 to 0.245)
losartan 25 mg	0.080 (-0.143 to 0.303)
losartan 50 mg	0.036 (-0.157 to 0.230)
losartan 100 mg	0.170 (0.019 to 0.321)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.01 (-0.34 to 0.33)
losartan 50 mg - losartan 0 mg	-0.05 (-0.36 to 0.26)
losartan 100 mg - losartan 0 mg	0.08 (-0.18 to 0.35)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	460.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	6.8	TRUE

Plot



EGF

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.529 (-1.108 to 0.050)
losartan 25 mg	-0.336 (-1.192 to 0.519)
losartan 50 mg	-0.356 (-1.089 to 0.377)
losartan 100 mg	-0.188 (-0.766 to 0.390)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.19 (-1.07 to 1.45)
losartan 50 mg - losartan 0 mg	0.17 (-0.97 to 1.32)
losartan 100 mg - losartan 0 mg	0.34 (-0.65 to 1.33)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	309.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	98.5	TRUE

Plot



ErbB2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.088 (-0.049 to 0.225)
losartan 25 mg	0.026 (-0.180 to 0.232)
losartan 50 mg	0.050 (-0.123 to 0.222)
losartan 100 mg	0.082 (-0.056 to 0.220)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.06 (-0.36 to 0.24)
losartan 50 mg - losartan 0 mg	-0.04 (-0.31 to 0.23)
losartan 100 mg - losartan 0 mg	-0.01 (-0.24 to 0.23)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	605.8	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	0.8	TRUE

Plot



FGF Basic

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.007 (-0.415 to 0.401)
losartan 25 mg	-0.178 (-0.784 to 0.427)
losartan 50 mg	-0.176 (-0.690 to 0.337)
losartan 100 mg	0.315 (-0.093 to 0.724)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.17 (-1.06 to 0.72)
losartan 50 mg - losartan 0 mg	-0.17 (-0.97 to 0.63)
losartan 100 mg - losartan 0 mg	0.32 (-0.38 to 1.02)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	269.8	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	74.9	TRUE

Plot



IL-12 p70

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.435 (-1.085 to 0.215)
losartan 25 mg	1.012 (0.056 to 1.969)
losartan 50 mg	0.454 (-0.395 to 1.304)
losartan 100 mg	0.020 (-0.623 to 0.663)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	1.45 (0.05 to 2.85)
losartan 50 mg - losartan 0 mg	0.89 (-0.44 to 2.22)
losartan 100 mg - losartan 0 mg	0.46 (-0.66 to 1.57)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	441.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	105.9	TRUE

Plot



IL-2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.640 (-1.484 to 0.204)
losartan 25 mg	0.605 (-0.708 to 1.917)
losartan 50 mg	0.181 (-0.871 to 1.232)
losartan 100 mg	-0.084 (-0.928 to 0.761)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	1.24 (-0.69 to 3.18)
losartan 50 mg - losartan 0 mg	0.82 (-0.82 to 2.46)
losartan 100 mg - losartan 0 mg	0.56 (-0.89 to 2.00)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	349.7	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	124.0	TRUE

Plot



IP-10 (CXCL10)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.173 (-0.048 to 0.394)
losartan 25 mg	0.016 (-0.317 to 0.349)
losartan 50 mg	-0.147 (-0.429 to 0.134)
losartan 100 mg	0.173 (-0.048 to 0.395)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.16 (-0.65 to 0.33)
losartan 50 mg - losartan 0 mg	-0.32 (-0.75 to 0.11)
losartan 100 mg - losartan 0 mg	0.00 (-0.38 to 0.38)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	338.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	33.2	TRUE

Plot



I-TAC (CXCL11)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.266 (-0.464 to -0.068)
losartan 25 mg	-0.104 (-0.398 to 0.190)
losartan 50 mg	0.012 (-0.243 to 0.267)
losartan 100 mg	-0.022 (-0.223 to 0.178)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.16 (-0.27 to 0.60)
losartan 50 mg - losartan 0 mg	0.28 (-0.12 to 0.68)
losartan 100 mg - losartan 0 mg	0.24 (-0.10 to 0.58)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	419.1	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	25.6	TRUE

Plot



MIG (CXCL9)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.390 (-0.919 to 0.139)
losartan 25 mg	0.400 (-0.408 to 1.208)
losartan 50 mg	-0.081 (-0.753 to 0.590)
losartan 100 mg	0.009 (-0.518 to 0.537)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.79 (-0.40 to 1.98)
losartan 50 mg - losartan 0 mg	0.31 (-0.73 to 1.34)
losartan 100 mg - losartan 0 mg	0.40 (-0.51 to 1.31)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	490.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	92.4	TRUE

Plot



PIGF

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.022 (-0.156 to 0.112)
losartan 25 mg	0.127 (-0.075 to 0.328)
losartan 50 mg	0.174 (0.007 to 0.341)
losartan 100 mg	0.062 (-0.072 to 0.195)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.15 (-0.15 to 0.45)
losartan 50 mg - losartan 0 mg	0.20 (-0.06 to 0.46)
losartan 100 mg - losartan 0 mg	0.08 (-0.15 to 0.31)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	275.2	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-1.2	TRUE

Plot



Tenascin C

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.048 (-0.134 to 0.038)
losartan 25 mg	0.020 (-0.107 to 0.147)
losartan 50 mg	0.022 (-0.086 to 0.129)
losartan 100 mg	0.020 (-0.067 to 0.106)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.07 (-0.12 to 0.25)
losartan 50 mg - losartan 0 mg	0.07 (-0.10 to 0.24)
losartan 100 mg - losartan 0 mg	0.07 (-0.08 to 0.22)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	620.8	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-31.1	TRUE

Plot



TNFRSF18 (GITR)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-1.327 (-2.192 to -0.462)
losartan 25 mg	-0.007 (-1.238 to 1.224)
losartan 50 mg	-0.149 (-1.196 to 0.898)
losartan 100 mg	-0.182 (-1.021 to 0.657)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	1.32 (-0.53 to 3.17)
losartan 50 mg - losartan 0 mg	1.18 (-0.51 to 2.86)
losartan 100 mg - losartan 0 mg	1.15 (-0.36 to 2.65)

Model comparison (AIC)

Model	AIC	Used
Linear ($\text{chg} \sim \text{base} + \text{trt02p}$)	500.3	FALSE
Log-transformed ($\log(\text{value}) - \log(\text{base}) \sim \log(\text{base}) + \text{trt02p}$)	123.2	TRUE

Plot



TNFSF6 (FasL)

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	-0.063 (-0.232 to 0.106)
losartan 25 mg	0.008 (-0.243 to 0.259)
losartan 50 mg	-0.130 (-0.345 to 0.085)
losartan 100 mg	-0.091 (-0.260 to 0.079)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	0.07 (-0.30 to 0.44)
losartan 50 mg - losartan 0 mg	-0.07 (-0.40 to 0.27)
losartan 100 mg - losartan 0 mg	-0.03 (-0.32 to 0.26)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	284.9	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	15.1	TRUE

Plot



VEGFR2

A total of 34 participants contributed data for this analyte.

Estimated change from baseline by long-term dose

Long-term dose	Change from baseline on the natural logarithmic scale (95% CI)
losartan 0 mg	0.091 (-0.027 to 0.208)
losartan 25 mg	-0.051 (-0.221 to 0.118)
losartan 50 mg	-0.008 (-0.152 to 0.135)
losartan 100 mg	-0.008 (-0.126 to 0.110)


Differences vs 0 mg by long-term dose

Comparison	Difference of the log-transformed values (95% CI)
losartan 25 mg - losartan 0 mg	-0.14 (-0.39 to 0.11)
losartan 50 mg - losartan 0 mg	-0.10 (-0.33 to 0.13)
losartan 100 mg - losartan 0 mg	-0.10 (-0.31 to 0.11)

Model comparison (AIC)

Model	AIC	Used
Linear (chg ~ base + trt02p)	645.5	FALSE
Log-transformed (log(value) - log(base) ~ log(base) + trt02p)	-11.5	TRUE

Plot




Time-to-Event Endpoints

Kaplan-Meier curves are shown with the at-risk table below. Overall log-rank p-values are reported for each endpoint, followed by Cox regression hazard ratios versus 0 mg.

The proportional-hazards assumption underlying the Cox hazard ratios is assessed with standard diagnostics (scaled Schoenfeld residual tests and plots, and complementary log-log survival plots) in a separate document ([impress_tte_ph_check.pdf](#)).

Progression-free survival



Number at risk

Dose	0	250	500	750	1000	1250
0 mg	18	11	7	2	1	1
25 mg	9	3	2	0	0	0
50 mg	9	5	2	0	0	0
100 mg	18	11	3	2	1	0

Days since randomisation


Log-rank p-value

Overall log-rank p-value: 0.439

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	1.99 (0.82 to 4.85)	0.128
50 mg vs 0 mg	1.52 (0.65 to 3.58)	0.335
100 mg vs 0 mg	1.56 (0.75 to 3.25)	0.239

6-month progression-free survival




Log-rank p-value

Overall log-rank p-value: 0.729

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	2.25 (0.32 to 16.01)	0.417
50 mg vs 0 mg	1.02 (0.09 to 11.26)	0.987
100 mg vs 0 mg	2.19 (0.40 to 11.96)	0.366

Overall survival



Number at risk

Dose	0	250	500	750	1000
0 mg	18	18	8	3	0
25 mg	9	8	2	0	0
50 mg	9	8	5	3	1
100 mg	18	15	10	2	0

Days since randomisation


Log-rank p-value

Overall log-rank p-value: 0.797

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	1.62 (0.61 to 4.26)	0.332
50 mg vs 0 mg	1.13 (0.41 to 3.12)	0.809
100 mg vs 0 mg	1.12 (0.47 to 2.65)	0.793

Overall survival at 12 months




Log-rank p-value

Overall log-rank p-value: 0.772

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	2.34 (0.47 to 11.58)	0.299
50 mg vs 0 mg	1.47 (0.25 to 8.82)	0.671
100 mg vs 0 mg	1.54 (0.34 to 6.87)	0.573

Overall survival at 24 months



Log-rank p-value

Overall log-rank p-value: 0.797

Cox regression (vs 0 mg)

Comparison	HR (95% CI)	P-value
25 mg vs 0 mg	1.62 (0.61 to 4.26)	0.332
50 mg vs 0 mg	1.13 (0.41 to 3.12)	0.809
100 mg vs 0 mg	1.12 (0.47 to 2.65)	0.793

Safety

Safety analyses follow SAP §7 and are descriptive/exploratory only, by treatment dose group and over time, with no formal hypothesis testing. The Safety Set is nominally all participants who received at least one dose of losartan, analysed by treatment actually received. **In this sham mock-up the dosing export (`_EX.csv`) is empty, so the Safety Set is approximated by the Full Analysis Set and all safety outputs are grouped by planned dose.** Assessment by treatment actually received will be added at database lock and unblinding. Conventional laboratory parameters are limited to creatinine, the only safety-lab assay populated in the current export.

Extent of Exposure

N/% exposed per planned dose level, treatment duration overall and by long-term dose, short-term vs continued treatment, dose switches / early discontinuation, and duration categories, for the newly diagnosed glioblastoma cohort.

A total of 54 subjects are included.

Subjects exposed per planned dose level

Dose (mg)	Short-term (randomised)	Long-term (maintenance)
0	0 (0.0%)	18 (33.3%)
25	18 (33.3%)	9 (16.7%)
50	18 (33.3%)	9 (16.7%)
100	18 (33.3%)	18 (33.3%)

Treatment duration (overall)

Group	N	Median (range), days	Mean (SD), days
Overall	54	450 (15-1065)	471.2 (256.8)

Treatment duration by long-term dose

Long-term dose (mg)	N	Median (range), days	Mean (SD), days
0	18	432 (153-786)	516.3 (215.1)
25	9	448 (60-737)	380.3 (255.2)
50	9	665 (100-1065)	569.6 (311.1)
100	18	454 (15-779)	422.4 (262.5)

Treatment continuation and changes

Measure	N (%)
Continued to long-term treatment	52 (96.3%)
Short-term assignment only	2 (3.7%)
Switched dose (short- vs long-term)	18 (33.3%)
Discontinued treatment early	5 (9.3%)

Duration categories

Duration category	N (%)
≤ 14 days	0 (0.0%)
$> 14 - \leq 42$ days	2 (3.7%)
> 42 days	52 (96.3%)

Adverse Events

Treatment-emergent adverse event (TEAE) incidence (subjects with at least one event), coded with MedDRA, split by onset into an **early** phase (onset Day ≤ 44) and a **late** phase (onset Day ≥ 45). Early-phase events are grouped by the stepped-wedge dose in effect at onset (a 0 mg group captures events before a subject's randomised step started; columns are not mutually exclusive, and each denominator is the number of subjects ever at that dose during Days 1-44). Late-phase events are grouped by the long-term planned dose (denominator = subjects per dose in the analysis set).

Early phase (onset Day ≤ 44 , by stepped-wedge dose)

A total of 17 subjects had at least one TEAE in this phase.

Overall summary

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Subjects in analysis set	N=54	N=18	N=18	N=18	N=54
Subjects with ≥ 1 TEAE	7 (13.0%)	2 (11.1%)	5 (27.8%)	3 (16.7%)	17 (31.5%)
with ≥ 1 serious TEAE	6 (11.1%)	0 (0.0%)	3 (16.7%)	2 (11.1%)	11 (20.4%)
with ≥ 1 treatment-related TEAE	1 (1.9%)	0 (0.0%)	1 (5.6%)	1 (5.6%)	3 (5.6%)
with ≥ 1 TEAE leading to discontinuation	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
with ≥ 1 fatal TEAE	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)

Subjects by maximum severity

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Mild	2 (3.7%)	1 (5.6%)	2 (11.1%)	0 (0.0%)	5 (9.3%)
Moderate	2 (3.7%)	0 (0.0%)	1 (5.6%)	1 (5.6%)	4 (7.4%)
Severe	3 (5.6%)	1 (5.6%)	2 (11.1%)	2 (11.1%)	8 (14.8%)
Life-threatening	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Death	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)

Subjects by relationship to treatment

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Unrelated	4 (7.4%)	2 (11.1%)	4 (22.2%)	2 (11.1%)	12 (22.2%)
Probable related	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (5.6%)	2 (3.7%)
Not applicable	2 (3.7%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (3.7%)
Possibly related	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Unlikely related	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)

Incidence by system organ class and preferred term

SOC / Preferred term	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Endocrine disorders	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Inappropriate antidiuretic hormone secretion	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Eye disorders	0 (0.0%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (1.9%)
Retinal detachment	0 (0.0%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (1.9%)
Gastrointestinal disorders	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Lower gastrointestinal haemorrhage	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Infections and infestations	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Wound infection	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Investigations	0 (0.0%)	2 (11.1%)	1 (5.6%)	2 (11.1%)	5 (9.3%)
Blood creatinine increased	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Neutrophil count decreased	0 (0.0%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (1.9%)
Platelet count decreased	0 (0.0%)	2 (11.1%)	0 (0.0%)	2 (11.1%)	4 (7.4%)
Nervous system disorders	3 (5.6%)	0 (0.0%)	2 (11.1%)	1 (5.6%)	6 (11.1%)
Brain oedema	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Dizziness	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Epilepsy	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (5.6%)	2 (3.7%)
Headache	1 (1.9%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	2 (3.7%)
Social circumstances	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Loss of personal independence in daily activities	1 (1.9%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Vascular disorders	0 (0.0%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (1.9%)
Hypotension	0 (0.0%)	0 (0.0%)	1 (5.6%)	0 (0.0%)	1 (1.9%)

Late phase (onset Day $\geq$ 45, by long-term dose)

A total of 21 subjects had at least one TEAE in this phase.

Overall summary

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Subjects in analysis set	N=18	N=9	N=9	N=18	N=54
Subjects with $\geq$ 1 TEAE	7 (38.9%)	4 (44.4%)	4 (44.4%)	6 (33.3%)	21 (38.9%)
with $\geq$ 1 serious TEAE	5 (27.8%)	2 (22.2%)	3 (33.3%)	6 (33.3%)	16 (29.6%)
with $\geq$ 1 treatment-related TEAE	2 (11.1%)	1 (11.1%)	0 (0.0%)	2 (11.1%)	5 (9.3%)
with $\geq$ 1 TEAE leading to discontinuation	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
with $\geq$ 1 fatal TEAE	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)

Subjects by maximum severity

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Mild	2 (11.1%)	2 (22.2%)	2 (22.2%)	1 (5.6%)	7 (13.0%)
Moderate	1 (5.6%)	0 (0.0%)	0 (0.0%)	2 (11.1%)	3 (5.6%)
Severe	4 (22.2%)	2 (22.2%)	1 (11.1%)	3 (16.7%)	10 (18.5%)
Life-threatening	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Death	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)

Subjects by relationship to treatment

Category	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Definite	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Unrelated	5 (27.8%)	3 (33.3%)	4 (44.4%)	6 (33.3%)	18 (33.3%)
Probable related	1 (5.6%)	1 (11.1%)	0 (0.0%)	1 (5.6%)	3 (5.6%)
Possibly related	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Unlikely related	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)

Incidence by system organ class and preferred term

SOC / Preferred term	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
General disorders and administration site conditions	1 (5.6%)	0 (0.0%)	2 (22.2%)	1 (5.6%)	4 (7.4%)
Gait disturbance	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)
Pain	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Pyrexia	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Sudden death	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)
Infections and infestations	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Wound infection	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)

SOC / Preferred term	losartan 0 mg	losartan 25 mg	losartan 50 mg	losartan 100 mg	Total
Investigations	1 (5.6%)	1 (11.1%)	1 (11.1%)	2 (11.1%)	5 (9.3%)
Alanine aminotransferase increased	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Gamma-glutamyltransferase increased	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Neutrophil count decreased	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)
Platelet count decreased	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (5.6%)	2 (3.7%)
Nervous system disorders	6 (33.3%)	2 (22.2%)	2 (22.2%)	5 (27.8%)	15 (27.8%)
Ataxia	1 (5.6%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	2 (3.7%)
Dizziness	1 (5.6%)	0 (0.0%)	0 (0.0%)	2 (11.1%)	3 (5.6%)
Epilepsy	2 (11.1%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	3 (5.6%)
Generalised tonic-clonic seizure	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Haemorrhage intracranial	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Headache	0 (0.0%)	0 (0.0%)	1 (11.1%)	0 (0.0%)	1 (1.9%)
Partial seizures	2 (11.1%)	0 (0.0%)	1 (11.1%)	1 (5.6%)	4 (7.4%)
Seizure	1 (5.6%)	1 (11.1%)	0 (0.0%)	2 (11.1%)	4 (7.4%)
Somnolence	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Syncope	1 (5.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Psychiatric disorders	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Confusional state	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Renal and urinary disorders	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Renal failure	0 (0.0%)	1 (11.1%)	0 (0.0%)	0 (0.0%)	1 (1.9%)
Respiratory, thoracic and mediastinal disorders	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)
Pulmonary embolism	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (5.6%)	1 (1.9%)

Vital Signs

Systolic and diastolic blood pressure and pulse rate are summarised descriptively by dose group and visit (observed and change from baseline) and analysed with a repeated- measures linear mixed model analogous to the efficacy endpoints: change from baseline with baseline value, nominal visit and treatment group as fixed effects and a subject random intercept. Short-term (Days 15/29/43) and long-term (Days 155/239) windows are reported.

Systolic BP

A total of 54 subjects contributed to the model-based analyses.

Descriptive summary by dose and visit

Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	3	128.0 (19.7)	0.0 (0.0)
15	0 mg	36	120.4 (10.6)	-7.1 (12.1)
15	25 mg	6	135.8 (18.0)	-6.2 (14.6)
15	50 mg	6	119.5 (8.3)	-6.0 (12.5)
15	100 mg	6	129.7 (21.7)	-7.8 (15.5)
29	0 mg	18	112.4 (10.1)	-15.2 (11.0)
29	25 mg	12	121.2 (16.3)	-13.0 (11.4)
29	50 mg	12	115.5 (6.1)	-10.9 (13.3)
29	100 mg	11	117.1 (12.6)	-15.4 (16.0)
43	25 mg	18	116.9 (14.6)	-14.4 (11.5)
43	50 mg	18	119.2 (13.2)	-8.7 (13.7)
43	100 mg	16	112.2 (10.1)	-17.3 (20.4)
57	25 mg	1	116.0 (NA)	-19.0 (NA)
71	0 mg	18	123.9 (16.0)	-3.2 (9.6)
71	25 mg	8	119.0 (9.8)	-15.0 (13.9)
71	50 mg	9	121.9 (15.1)	-8.6 (10.9)
71	100 mg	16	119.1 (11.2)	-12.4 (19.4)
99	0 mg	18	119.7 (17.6)	-7.3 (11.5)
99	25 mg	9	124.7 (11.8)	-9.8 (12.1)
99	50 mg	9	118.7 (15.5)	-11.8 (11.6)
99	100 mg	15	117.8 (12.4)	-12.9 (17.4)
127	0 mg	16	119.8 (17.0)	-7.2 (9.2)
127	25 mg	7	132.7 (18.1)	-0.7 (17.8)
127	50 mg	7	122.0 (11.7)	-8.9 (14.3)
127	100 mg	12	118.8 (12.7)	-11.5 (25.5)
141	100 mg	1	124.0 (NA)	-19.0 (NA)
155	0 mg	19	121.0 (14.4)	-6.9 (10.1)
155	25 mg	7	128.4 (12.6)	-5.0 (14.9)
155	50 mg	8	118.6 (14.7)	-11.1 (12.7)
155	100 mg	14	123.3 (12.5)	-6.9 (16.5)
183	0 mg	17	123.9 (15.9)	-2.1 (12.5)
183	25 mg	7	125.0 (13.7)	-8.4 (18.9)
183	50 mg	8	120.9 (13.5)	-8.9 (20.5)
183	100 mg	12	123.8 (11.5)	-6.3 (19.9)
211	0 mg	15	119.2 (13.3)	-5.6 (12.3)
211	25 mg	6	118.3 (14.3)	-17.3 (17.3)
211	50 mg	7	114.6 (11.7)	-16.3 (13.6)
211	100 mg	10	123.1 (10.3)	-5.5 (21.1)
239	0 mg	13	121.2 (17.6)	-4.4 (10.8)


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
239	25 mg	5	125.8 (4.5)	-10.2 (7.7)
239	50 mg	7	128.7 (11.0)	-2.1 (18.3)
239	100 mg	12	121.9 (17.2)	-8.2 (25.2)
253	0 mg	46	125.5 (15.8)	-3.9 (15.4)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	-13.78 (-17.39 to -10.18)
25 mg	-7.63 (-11.89 to -3.37)
50 mg	-8.47 (-12.71 to -4.22)
100 mg	-14.55 (-18.84 to -10.27)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg0	6.16 (-0.33 to 12.64)
losartan 50 mg vs losartan 0 mg0	5.32 (-1.11 to 11.74)
losartan 100 mg vs losartan 0 mg0	-0.77 (-7.20 to 5.66)




Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-7.84 (-13.65 to -2.03)
155	losartan 25 mg	-2.88 (-12.27 to 6.51)
155	losartan 50 mg	-11.04 (-19.79 to -2.28)
155	losartan 100 mg	-6.58 (-13.20 to 0.03)
239	losartan 0 mg	-6.98 (-13.60 to -0.37)
239	losartan 25 mg	-6.57 (-17.25 to 4.11)
239	losartan 50 mg	-2.37 (-11.56 to 6.83)
239	losartan 100 mg	-7.11 (-14.11 to -0.10)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	4.96 (-8.45 to 18.37)
155	losartan 50 mg - losartan 0 mg	-3.19 (-15.91 to 9.53)
155	losartan 100 mg - losartan 0 mg	1.26 (-9.41 to 11.93)
239	losartan 25 mg - losartan 0 mg	0.41 (-14.85 to 15.67)
239	losartan 50 mg - losartan 0 mg	4.62 (-9.10 to 18.33)
239	losartan 100 mg - losartan 0 mg	-0.13 (-11.79 to 11.54)



Diastolic BP

A total of 54 subjects contributed to the model-based analyses.

Descriptive summary by dose and visit


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	3	72.7 (5.1)	0.0 (0.0)
15	0 mg	36	78.6 (7.4)	-2.6 (7.6)
15	25 mg	6	80.5 (8.5)	-0.3 (5.5)
15	50 mg	6	77.5 (6.3)	-4.2 (5.8)
15	100 mg	6	77.0 (7.8)	-9.7 (9.7)
29	0 mg	18	74.2 (6.2)	-6.9 (6.3)
29	25 mg	12	76.2 (8.9)	-5.2 (6.6)
29	50 mg	12	75.2 (6.1)	-4.6 (8.7)
29	100 mg	11	76.7 (7.2)	-8.2 (8.2)
43	25 mg	18	74.4 (7.6)	-7.3 (7.5)
43	50 mg	18	75.7 (7.4)	-4.8 (7.9)
43	100 mg	16	72.9 (5.7)	-9.6 (9.0)
57	25 mg	1	74.0 (NA)	-10.0 (NA)
71	0 mg	18	77.3 (6.8)	-2.3 (6.6)
71	25 mg	8	77.9 (8.0)	-4.6 (7.0)
71	50 mg	9	76.8 (10.3)	-5.7 (6.8)
71	100 mg	16	76.8 (6.9)	-6.6 (9.4)
99	0 mg	18	76.3 (9.6)	-3.3 (9.1)
99	25 mg	9	80.6 (9.2)	-2.6 (11.0)
99	50 mg	9	76.7 (9.0)	-5.8 (9.6)
99	100 mg	15	77.7 (9.0)	-5.4 (9.7)
127	0 mg	16	74.9 (6.9)	-5.1 (5.3)
127	25 mg	7	82.3 (9.3)	0.6 (13.6)
127	50 mg	7	76.1 (6.4)	-8.0 (5.4)
127	100 mg	12	78.8 (9.5)	-4.0 (11.7)
141	100 mg	1	77.0 (NA)	-7.0 (NA)
155	0 mg	19	75.9 (6.1)	-3.4 (6.4)
155	25 mg	7	82.9 (7.5)	1.1 (9.0)
155	50 mg	8	77.9 (9.5)	-4.1 (6.3)
155	100 mg	14	80.1 (8.5)	-3.4 (8.0)
183	0 mg	17	77.8 (7.6)	-2.1 (7.4)
183	25 mg	7	78.3 (7.3)	-3.4 (7.0)
183	50 mg	8	79.4 (6.8)	-2.6 (8.3)
183	100 mg	12	81.7 (7.7)	-1.8 (7.8)
211	0 mg	15	74.1 (6.1)	-5.0 (6.5)
211	25 mg	6	79.5 (7.2)	-5.8 (6.6)
211	50 mg	7	76.0 (7.2)	-8.1 (5.1)
211	100 mg	10	79.4 (6.7)	-3.2 (7.6)
239	0 mg	13	77.1 (5.1)	-2.2 (5.5)
239	25 mg	5	81.2 (8.1)	-1.8 (4.0)
239	50 mg	7	82.0 (3.9)	-2.1 (7.9)
239	100 mg	12	79.2 (10.1)	-4.2 (10.0)
253	0 mg	46	80.5 (9.0)	-1.7 (8.3)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	-5.19 (-7.38 to -3.01)
25 mg	-5.08 (-7.66 to -2.50)
50 mg	-4.57 (-7.16 to -1.99)
100 mg	-8.02 (-10.65 to -5.38)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	0.11 (-3.77 to 4.00)
losartan 50 mg vs losartan 0 mg	0.62 (-3.27 to 4.51)
losartan 100 mg vs losartan 0 mg	-2.82 (-6.73 to 1.09)



Plot


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-4.63 (-7.76 to -1.51)
155	losartan 25 mg	1.18 (-3.79 to 6.15)
155	losartan 50 mg	-3.95 (-8.60 to 0.70)
155	losartan 100 mg	-2.48 (-6.02 to 1.07)
239	losartan 0 mg	-3.35 (-6.86 to 0.16)
239	losartan 25 mg	-1.19 (-6.79 to 4.40)

Visit	Treatment	Estimated change from baseline (95% CI)
239	losartan 50 mg	-1.37 (-6.24 to 3.50)
239	losartan 100 mg	-2.68 (-6.41 to 1.05)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	5.81 (-1.30 to 12.92)
155	losartan 50 mg - losartan 0 mg	0.68 (-6.12 to 7.47)
155	losartan 100 mg - losartan 0 mg	2.15 (-3.64 to 7.94)
239	losartan 25 mg - losartan 0 mg	2.16 (-5.85 to 10.17)
239	losartan 50 mg - losartan 0 mg	1.98 (-5.32 to 9.29)
239	losartan 100 mg - losartan 0 mg	0.67 (-5.59 to 6.94)



Plot

Pulse rate

A total of 54 subjects contributed to the model-based analyses.

Descriptive summary by dose and visit


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	3	71.7 (4.5)	0.0 (0.0)
15	0 mg	36	75.2 (12.3)	-6.2 (14.3)
15	25 mg	6	70.7 (4.2)	-5.3 (8.2)
15	50 mg	6	79.2 (14.7)	-3.2 (13.5)
15	100 mg	6	72.0 (10.4)	-6.0 (9.7)
29	0 mg	18	82.1 (12.5)	-2.9 (11.1)
29	25 mg	12	73.0 (11.6)	-1.3 (12.7)
29	50 mg	12	72.3 (14.2)	-7.8 (14.6)
29	100 mg	11	79.7 (10.5)	0.5 (12.5)
43	25 mg	18	72.6 (12.1)	-7.3 (15.3)
43	50 mg	18	75.8 (14.9)	-4.4 (12.1)
43	100 mg	16	73.3 (10.8)	-7.0 (11.1)
57	25 mg	1	60.0 (NA)	6.0 (NA)
71	0 mg	18	74.0 (14.7)	-2.5 (13.0)
71	25 mg	8	73.6 (12.5)	-15.0 (15.4)
71	50 mg	9	74.6 (8.1)	-7.7 (13.3)
71	100 mg	16	73.9 (8.9)	-7.2 (12.3)
99	0 mg	18	73.4 (12.3)	-3.1 (11.6)
99	25 mg	9	75.0 (11.6)	-12.4 (16.5)
99	50 mg	9	77.2 (8.5)	-5.0 (11.0)
99	100 mg	15	79.3 (11.2)	-1.3 (17.9)
127	0 mg	16	71.9 (12.4)	-5.6 (15.6)
127	25 mg	7	70.6 (10.3)	-19.6 (11.3)
127	50 mg	7	74.3 (10.6)	-8.7 (17.9)
127	100 mg	12	75.9 (12.9)	-4.3 (13.0)
141	100 mg	1	65.0 (NA)	-4.0 (NA)
155	0 mg	19	70.9 (10.6)	-5.4 (12.1)
155	25 mg	7	72.4 (7.3)	-17.7 (11.0)
155	50 mg	8	77.2 (9.4)	-4.4 (12.4)
155	100 mg	14	79.4 (14.9)	-4.2 (14.3)
183	0 mg	17	71.9 (9.2)	-4.9 (15.9)
183	25 mg	7	76.7 (11.6)	-13.4 (11.8)
183	50 mg	8	74.0 (9.4)	-7.6 (11.0)
183	100 mg	12	74.7 (11.2)	-10.6 (14.3)
211	0 mg	15	72.7 (15.3)	-3.3 (17.0)
211	25 mg	6	78.0 (7.9)	-14.8 (10.3)
211	50 mg	7	70.0 (3.6)	-13.0 (17.7)
211	100 mg	10	75.9 (9.8)	-11.7 (13.1)
239	0 mg	13	78.2 (17.3)	0.5 (19.9)
239	25 mg	5	73.0 (12.9)	-14.8 (9.9)
239	50 mg	7	75.4 (13.3)	-7.6 (15.7)
239	100 mg	12	74.3 (10.9)	-10.9 (15.7)
253	0 mg	45	77.1 (12.6)	-5.2 (16.2)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	-4.04 (-7.36 to -0.73)
25 mg	-6.65 (-10.59 to -2.70)
50 mg	-5.22 (-9.16 to -1.28)
100 mg	-4.52 (-8.48 to -0.55)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	-2.61 (-8.33 to 3.12)
losartan 50 mg vs losartan 0 mg	-1.17 (-6.88 to 4.53)
losartan 100 mg vs losartan 0 mg	-0.47 (-6.15 to 5.20)



Plot


Days 155/239 (mixed model)

Visit	Treatment	Estimated change from baseline (95% CI)
155	losartan 0 mg	-8.47 (-13.77 to -3.17)
155	losartan 25 mg	-12.76 (-21.31 to -4.22)
155	losartan 50 mg	-4.51 (-12.35 to 3.34)
155	losartan 100 mg	-3.14 (-9.08 to 2.80)
239	losartan 0 mg	-2.62 (-8.61 to 3.37)
239	losartan 25 mg	-10.78 (-20.44 to -1.13)

Visit	Treatment	Estimated change from baseline (95% CI)
239	losartan 50 mg	-7.36 (-15.61 to 0.89)
239	losartan 100 mg	-9.02 (-15.32 to -2.71)

Differences vs reference

Visit	Comparison	Difference (95% CI)
155	losartan 25 mg - losartan 0 mg	-4.29 (-16.67 to 8.10)
155	losartan 50 mg - losartan 0 mg	3.96 (-7.49 to 15.42)
155	losartan 100 mg - losartan 0 mg	5.33 (-4.37 to 15.03)
239	losartan 25 mg - losartan 0 mg	-8.17 (-22.05 to 5.72)
239	losartan 50 mg - losartan 0 mg	-4.74 (-17.08 to 7.61)
239	losartan 100 mg - losartan 0 mg	-6.40 (-16.99 to 4.19)



Plot

Conventional Laboratory Parameters

Conventional clinical laboratory parameters (creatinine in this export) are summarised descriptively and with the same repeated-measures mixed model as vital signs, plus categorisation against reference ranges (counts/proportions outside the normal range).

Creatinine

Reference-range categorisation by dose

trt	LOW	NORMAL	HIGH
0 mg	3 (2.9%)	100 (97.1%)	0 (0.0%)
25 mg	0 (0.0%)	29 (100.0%)	0 (0.0%)
50 mg	2 (6.5%)	29 (93.5%)	0 (0.0%)
100 mg	0 (0.0%)	32 (94.1%)	2 (5.9%)

Descriptive summary by dose and visit


Visit (day)	Dose	N	Observed mean (SD)	Change mean (SD)
-4	0 mg	54	70.4 (12.9)	0.0 (0.0)
15	0 mg	36	72.0 (10.6)	3.8 (9.6)
15	25 mg	6	77.7 (14.2)	0.3 (6.7)
15	50 mg	6	65.3 (10.8)	4.2 (6.0)
15	100 mg	5	79.8 (11.8)	1.0 (4.1)
43	25 mg	18	72.3 (12.4)	4.1 (6.3)
43	50 mg	18	71.6 (11.8)	6.5 (6.4)
43	100 mg	17	81.7 (20.4)	2.8 (5.7)
225	0 mg	13	76.3 (11.3)	6.8 (10.2)
225	25 mg	5	81.6 (15.4)	4.2 (6.7)
225	50 mg	7	62.4 (6.6)	4.6 (6.6)
225	100 mg	12	77.1 (9.7)	0.5 (8.6)

Days 15/29/43 (change from baseline, by dose)

Dose (mg)	Estimated change from baseline (95% CI)
0 mg	4.27 (0.93 to 7.61)
25 mg	2.69 (-0.67 to 6.05)
50 mg	3.88 (0.11 to 7.66)
100 mg	4.12 (0.44 to 7.80)

Differences vs 0 mg

Comparison	Difference (95% CI)
losartan 25 mg vs losartan 0 mg	-1.58 (-7.67 to 4.52)
losartan 50 mg vs losartan 0 mg	-0.38 (-6.81 to 6.05)
losartan 100 mg vs losartan 0 mg	-0.15 (-6.63 to 6.34)



Deaths

Deaths are summarised by long-term planned dose group (in addition to the overall survival analysis in the time-to-event section) and listed with date and cause.

A total of 34 deaths were observed among 54 subjects.

Deaths by dose group

Long-term dose	N	Deaths n (%)
0 mg	18	10 (55.6%)
25 mg	9	7 (77.8%)
50 mg	9	6 (66.7%)
100 mg	18	11 (61.1%)

Listing of deaths

Subject	Dose	Death date	Days from start	Cause
ImPRESS-ImPR-2005	0 mg	2021-04-20	307	Progressive disease
ImPRESS-ImPR-2012	0 mg	2021-11-03	421	Progressive disease
ImPRESS-ImPR-2015	0 mg	2022-01-11	429	Progressive disease
ImPRESS-ImPR-2023	0 mg	2022-08-28	432	Progressive disease
ImPRESS-ImPR-2031	0 mg	2022-12-06	331	Progressive disease
ImPRESS-ImPR-2035	0 mg	2023-05-07	362	Progressive disease
ImPRESS-ImPR-2037	0 mg	2024-02-02	387	Progressive disease
ImPRESS-ImPR-2045	0 mg	2024-06-15	401	Progressive disease
ImPRESS-ImPR-2044	0 mg	2024-07-14	433	Progressive disease
ImPRESS-ImPR-2050	0 mg	2025-03-28	430	Progressive disease
ImPRESS-ImPR-2008	25 mg	2021-11-10	465	Progressive disease
ImPRESS-ImPR-2019	25 mg	2022-01-01	300	Progressive disease
ImPRESS-ImPR-2013	25 mg	2022-02-01	484	Progressive disease
ImPRESS-ImPR-2030	25 mg	2022-11-20	356	Progressive disease
ImPRESS-ImPR-2029	25 mg	2023-01-13	437	Progressive disease
ImPRESS-ImPR-2046	25 mg	2023-07-29	67	Progressive disease
ImPRESS-ImPR-2038	25 mg	2024-04-24	448	Progressive disease
ImPRESS-ImPR-2004	50 mg	2021-07-29	430	Progressive disease
ImPRESS-ImPR-2026	50 mg	2022-10-30	419	Progressive disease
ImPRESS-ImPR-2040	50 mg	2023-09-05	182	Other cause
ImPRESS-ImPR-2027	50 mg	2023-09-07	717	Progressive disease
ImPRESS-ImPR-2041	50 mg	2024-01-18	303	Progressive disease
ImPRESS-ImPR-2048	50 mg	2025-09-09	665	Progressive disease
ImPRESS-ImPR-2014	100 mg	2021-03-09	134	Progressive disease
ImPRESS-ImPR-2001	100 mg	2021-06-29	527	Progressive disease
ImPRESS-ImPR-2002	100 mg	2021-08-31	583	Progressive disease
ImPRESS-ImPR-2009	100 mg	2022-02-04	551	Progressive disease
ImPRESS-ImPR-2011	100 mg	2022-02-17	540	Progressive disease
ImPRESS-ImPR-2016	100 mg	2022-03-21	456	Progressive disease
ImPRESS-ImPR-2022	100 mg	2022-08-28	455	Progressive disease
ImPRESS-ImPR-2033	100 mg	2023-02-26	370	Progressive disease
ImPRESS-ImPR-2034	100 mg	2023-03-18	347	Progressive disease
ImPRESS-ImPR-2036	100 mg	2023-06-15	205	Progressive disease
ImPRESS-ImPR-2052	100 mg	2025-04-26	312	Progressive disease