



Distinct SU(2) reductions: spectrum and static response


A One-square Jacobi diagnostic



B Static tilted-Haar measure



C Variance retained in the scalar equation



Different obligations

Jacobi: infinite character tower;
finite spatial graph.
N = 24 states do not certify its tail.


Static response: a probability measure;
no physical time generator is identified.

Dropping variance changes the origin
slope from 1/4 to 1/3.


Gap floor: inherited exact Round10 certificate on $0 \leq \lambda/\alpha \leq 10$. Static curves: exact Bessel formula evaluated numerically. Numerical agreement and truncation refinement are diagnostics; neither proves a continuum theorem.

Complete support controls the available time window

A Strict region: $\gamma < 3(1 - \beta)$



B Exact support budget evaluated




Canonical summable family, $\eta = 1/2$, $C = \ell = 1$; $L = \max(1, \text{floor}(\epsilon^{-\beta}))$.


Boundary curves are excluded from the vanishing certificate. A positive or divergent upper bound is not a lower bound on actual correlation error. The $q = 1$ Hamiltonian is not inserted into the summable family.

Filtered correction: contraction and locality are separate

A Origin source: three crossing stars




B Translated family: twelve crossings




Panel A: AB2 original cap $M = 35/1664$. Panel B: AE2 requires $2/(1-25M) < T < 1/(192M)$. Its joint open interval exists exactly when $409M < 1$. The zero-frequency residual is one for both filters. These are sufficient upper certificates; neither a budget floor nor a failed locality estimate proves actual divergence.

Wilson endpoints: one clock, distinguishable probes


A Quadratic real response




B Linear signed response



C More than one reached frequency



D Equal variance does not fix the operator




Actual multiplier endpoints, $0 \leq z \leq 10^{-6}$. The elementary face is a null probe at this clock.

The coherent sum shares the rank-swap endpoint scalar, but its norm squared is 8 and fourth vacuum moment is $5/2$.

Curves evaluate exact limiting formulas. They do not include finite-q remainders, physical clock calibration or measured data.

Heat approximation: physical omission and numerical error



$\lambda \leq 0.01$ for AC2; $\lambda = 0.01$ for AF2. Inputs: normalized original P21 vectors within 0.01 of the vacuum.
Physical bound = 457097/12472250000; denominator $\geq 7127/7200$. The 293-dimensional space is an enrichment.
This companion recomputes scalar certificates. It does not execute the original 293-coordinate heat solver.