

INFINITE TETRADES OF CONGRUENT FACTORIALS

OCTAVIO A. AGUSTÍN-AQUINO AND JOSÉ HERNÁNDEZ SANTIAGO

ABSTRACT. We prove that there exist infinitely many primes p such that there exist k_1, k_2, k_3, k_4 distinct such that

$$k_1! \equiv k_2! \equiv k_3! \equiv k_4! \equiv 1 \pmod{p}.$$

1. INTRODUCTION

Following [1], we write $a \setminus b$ to mean a divides b . The following fact is relatively well-known.

Lemma 1.1 (Wilson's symmetry). *Let p be an odd prime and $0 \leq n \leq p-1$ an odd integer such that $n! \equiv 1 \pmod{p}$. Then*

$$(p-1-n)! \equiv 1 \pmod{p}.$$

Proof. By Wilson's theorem we have $(p-1)! \equiv -1 \pmod{p}$. Then

$$(p-1)! = (p-1-n)! \prod_{k=1}^n (p-k).$$

Thus, modulo p , we have

$$\begin{aligned} (p-1-n)! \prod_{k=1}^n (p-k) &\equiv (p-1-n)! (-1)^n n! \\ &\equiv -(p-1-n)! \equiv -1 \pmod{p}, \end{aligned}$$

hence $(p-1-n)! \equiv 1 \pmod{p}$. □

Wilson's symmetry is crucial for the following construction.

Lemma 1.2 (Valid tetrad). *Let $q \geq 3$ be an odd integer, and p a prime divisor of $q! - 1$. If p is distinct from $q+2$ and $2q+1$, then*

$$k_1 = 1, \quad k_2 = q, \quad k_3 = p-2, \quad k_4 = p-1-q,$$

are four different integers such that $k_i! \equiv 1 \pmod{p}$ for $1 \leq i \leq 4$.

2020 *Mathematics Subject Classification.* 11B65, 11A07, 11A41.

Proof. Since $p \nmid (q! - 1)$, then $p > q$. Otherwise, $q! \equiv 0 \pmod{p}$, which contradicts that $q! \equiv 1 \pmod{p}$. We already have $1!, q! \equiv 1 \pmod{p}$, and by Wilson's symmetry, $(p-2)!, (p-1-q)! \equiv 1 \pmod{p}$ because q is odd. If the k_i were not distinct, then $p-2 = 1$ if and only if $p = 3$, $p-1-q = 1$ if and only if $p = q+2$, $q = p-2$ if and only if $p = q+2$, or $q = p-1-q$ if and only if $p = 2q+1$, or $p-2 = p-1-q$ if and only if $q = 1$, but all of these are blocked by the hypotheses. $\square$

2. MAIN THEOREM

Theorem 2.1. *There exist infinitely many primes p such that*

$$|\{1 \leq n \leq p-1 : n! \equiv 1 \pmod{p}\}| \geq 4.$$

Proof. Let $q = 6t + 1$ for an integer $t \geq 1$. It follows that q is odd, while

$$q+2 = 6t+3 = 3(2t+1) \quad \text{and} \quad 2q+1 = 12t+3 = 3(4t+1),$$

which implies $q+2$ and $2q+1$ are composite. Thus any prime divisor p of $q! - 1$ cannot be equal to $q+2$ or $2q+1$. Now $q! - 1 > 1$, and hence such a prime p exists. By the valid tetrad lemma, there exist k_1, k_2, k_3, k_4 , all distinct, such that their factorials coincide in having residue 1 modulo p .

Let S be a finite set of positive primes with such a property. Choose $q = 6t + 1$ larger than any of those primes. Then $q! - 1$ has a positive prime divisor p that is larger than q , and thus it cannot belong to S , and the result follows. $\square$

Remark 2.2. This result is connected to problem A15 of Guy's compendium [2, p. 54] and the corresponding observation by Mąkowski [4].

Remark 2.3. For $q = at + b$, with a even and b odd, the construction of the proof of the main theorem (cf. [3, Remark 2.8, p. 556]) works whenever

$$\gcd(a, b+2) > 1, \quad \text{and} \quad \gcd(a, 2b+1) > 1,$$

Indeed, these conditions force $q+2$ and $2q+1$ to be composite for every t , so the two possible Wilson-pair collisions are ruled out. The choice $q = 6t + 1$ is a minimal case and makes the aforementioned pair divisible by same prime divisor, namely 3.

3. SOME COMPUTATIONS

3.1. Euclidean Scheme. We can obtain sequences of primes with at least a tetrad of factorials congruent to 1 using the Euclidean scheme of the proof (see [1, Section 4.3]). If we begin with $t = 1$, then we have $q = 6 \cdot 1 + 1 = 7$. But $7! - 1 = 5039$ is prime, and the smallest q of the form $6t + 1$ that is larger than 5039 is 5041. The least prime factor of $5041! - 1$ is 20549. This was found considering that if we write the prime divisor as $s = 5041 + r$, then by Wilson's theorem

$$(s - 1)! \equiv -1 \pmod{s}.$$

By the definition of s , we have

$$(s - 1)! = 5041!5042 \cdot 5043 \cdots (s - 1),$$

and modulo s we have

$$\begin{aligned} 5042 \cdot 5043 \cdots (s - 1) &\equiv (-r + 1)(-r + 2) \cdots (-1) \\ &\equiv -(r - 1)! \pmod{s} \end{aligned}$$

because r is even. Thus

$$-1 \equiv 5041!(-(r - 1)!) \pmod{s}.$$

Hence $5041! \equiv 1 \pmod{s}$ if, and only if, $(r - 1)! \equiv 1 \pmod{s}$. A direct check using the primes greater than 5041 with a Python script yields $r = 15508$. The next q is 20551, and the script yields $r = 4296$ for the prime factor $s = 24847$. After this the algorithm takes more than a few seconds to finish the computation.

If we begin with $t = 2$, then $q = 6 \cdot 2 + 1 = 13$, and the least prime factor of $13! - 1$ is 1733. The next q of the form $6t + 1$ is 1735. After this the algorithm takes more than a few seconds to finish the computation.

If we begin with $t = 3$, then $q = 6 \cdot 3 + 1 = 19$, and the least prime factor of $19! - 1$ is 653. Now the next q of the form $6t + 1$ is 655, and the least prime factor of $655! - 1$ is 5407. After this the algorithm takes more than a few seconds to finish the computation.

3.2. Tetrad-Only Primes. It is interesting to calculate the sequence of primes p that admit exactly one tetrad of factorials congruent to 1 modulo p . Using a simple Python script, we can readily find its first terms:

$$17, 53, 59, 73, 83, 89, 97, 139, 239, \dots$$

ACKNOWLEDGEMENTS

The exploration to find the valid tetrad lemma and the main theorem was done with ChatGPT 5.5 Pro. The write up was done by the authors and we take full responsibility for the manuscript.

SOURCE CODE

3.3. Euclidean Scheme.

```
import sympy as sp

def search_prime_divisor(n):
    # Here n is assumed to be
    # an odd number.
    for s in sp.primerange(n + 1, 10**7):
        r = s - n
        prod = 1
        for u in range(1, r):
            prod = (prod * u) % s
        if prod == 1:
            return s, r
```

3.4. Tetrads-Only Primes.

```
import sympy as sp

for p in sp.primerange(2, 10000):
    restos = []
    fact = 1

    for n in range(1, p):
        fact = (fact * n) % p

        if fact == 1:
            restos.append(n)

        # Optional early stop:
        # we only care about exactly 4
        if len(restos) > 4:
            break

    if len(restos) == 4:
        print(p)
```

REFERENCES

- [1] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik. *Concrete Mathematics*. 2nd ed. Boston, MA: Addison Wesley, 1994.
- [2] Richard K. Guy. *Unsolved Problems in Number Theory*. 3rd ed. Springer, 2004.
- [3] G. E. Hardy and M. V. Subbarao. “A Modified Problem of Pillai and Some Related Questions”. In: *The American Mathematical Monthly* 109.6 (2002), pp. 554–559.
- [4] Andrzej Mąkowski. “On a number-theoretical problem of Erdős”. In: *Elemente der Mathematik* 38 (1983), pp. 101–102.

INSTITUTO DE FÍSICA Y MATEMÁTICAS, UNIVERSIDAD TECNOLÓGICA DE LA MIXTECA, HUAJUAPAN DE LEÓN, OAXACA, MÉXICO

Email address: `octavioalberto@mixteco.utm.mx`

URL: `www.utm.mx/~octavioalberto`

ESCUELA SUPERIOR DE MATEMÁTICAS NO. 2, UNIVERSIDAD AUTÓNOMA DE GUERRERO, CIUDAD ALTAMIRANO, GUERRERO, MÉXICO

Email address: `15620@uagro.mx`

URL: `https://euclid.uagro.mx/`