

# The State of Comon's Conjecture

**Lael JOHN**

Supervisor: Prof. N.  
Vannieuwenhoven  
[KU Leuven](#)

Co-supervisor: Dr. D. Taufer  
[Max Planck Institute \(MIS Leipzig\)](#)

Thesis presented in  
fulfillment of the requirements  
for the degree of Master of Science  
in Mathematics

Academic year 2025 -2026

---

© Copyright by KU Leuven

Without prior written permission from both the supervisor(s) and the author(s), copying, reproducing, using, or realizing this publication or parts thereof is prohibited. For requests or information regarding the copying and/or use and/or realization of parts of this publication, please contact KU Leuven, Faculty of Science, Celestijnenlaan 200H - box 2100, 3001 Leuven (Heverlee), Telephone +32 16 32 14 01.

Prior written permission from the supervisor(s) is also required for using the (original) methods, products, circuits, and programs described in this work for industrial or commercial purposes, and for submitting this publication for scientific awards or contests.

# Foreword

This thesis aims to document the mathematical progress made around Comon's Conjecture (a question about tensor rank and symmetry). Covered within this text are some attempts made to prove the conjecture in various contexts, published after it was first posited in 2008. The main focus of this work centers around the tools and methodology proposed by Shitov (both from published works [22] and preprints [23–25]) that aim to construct a valid counterexample to the conjecture. The culmination of this thesis is an examination of Wang and Seigal's paper [27] detailing the construction of a valid counterexample in  $\mathbb{R}$ . While the existence of a counterexample in  $\mathbb{C}$  is still an open question, this thesis concludes that the ideas detailed within the papers cited provide a valid pathway to one.

## Acknowledgements and Contribution Statement

This thesis was written in consultation with Prof. N. Vannieuwenhoven and Dr. D. Taufer. Prof. Vannieuwenhoven provided the initial idea and shaped the overall direction this thesis took, and Dr. Taufer's guidance over the year helped in developing this project on a solid foundation. I am grateful to both for the structure and feedback they have provided over the course of my interaction with them. Personal contributions to this thesis are detailed in the introduction. Generative AI tools were not used in the writing of this thesis.

This thesis would not exist without friends, family, coworkers and mentors, both in Belgium and from around the world — I would not be in any position to write this work without them.

# Contents

|                                                          |            |
|----------------------------------------------------------|------------|
| <b>Foreword</b>                                          | <b>ii</b>  |
| <b>List of Notation</b>                                  | <b>iv</b>  |
| <b>List of Figures</b>                                   | <b>vii</b> |
| <b>1 Introduction</b>                                    | <b>1</b>   |
| <b>2 Preliminaries Regarding Tensors</b>                 | <b>4</b>   |
| 2.1 Tensors in an Algebraic Setting . . . . .            | 4          |
| 2.2 Rank . . . . .                                       | 6          |
| 2.3 Symmetric Tensors . . . . .                          | 8          |
| <b>3 Literature Review</b>                               | <b>11</b>  |
| 3.1 Attempts to Prove the Conjecture . . . . .           | 11         |
| 3.2 Attempts to Disprove the Conjecture . . . . .        | 16         |
| 3.3 Overview of Valid Cases of the Conjecture . . . . .  | 16         |
| <b>4 Constructions required for a Counterexample</b>     | <b>17</b>  |
| 4.1 Background to the Counterexample . . . . .           | 17         |
| 4.1.1 Shitov's First Paper in 2018 . . . . .             | 17         |
| 4.1.2 Shitov's Subsequent Preprints . . . . .            | 18         |
| 4.1.3 Slicing a Tensor . . . . .                         | 19         |
| 4.1.4 Flattening a Tensor . . . . .                      | 21         |
| 4.2 Major New Constructions from Shitov . . . . .        | 22         |
| 4.2.1 Background to Adjoining . . . . .                  | 23         |
| 4.2.2 Adjoining Slices to a Tensor . . . . .             | 25         |
| 4.2.3 Generating Tensor Families modulo Slices . . . . . | 25         |
| 4.2.4 Cloning Tensors . . . . .                          | 28         |
| 4.3 The Counterexample over $\mathbb{R}$ . . . . .       | 31         |
| 4.3.1 Details from the Paper . . . . .                   | 31         |
| 4.3.2 Emulating Family Construction . . . . .            | 35         |
| <b>5 Conclusion</b>                                      | <b>37</b>  |
| 5.1 Future Work . . . . .                                | 37         |

# List of Notation

## General Objects

|                                 |                                                                                                               |
|---------------------------------|---------------------------------------------------------------------------------------------------------------|
| $[n]$                           | The set $\{1, 2, \dots, n\}$                                                                                  |
| $\mathbb{F}$                    | A general field, in this thesis either $\mathbb{R}$ or $\mathbb{C}$ unless specified                          |
| $\mathbb{F}[x_1, \dots, x_n]_d$ | Subring of polynomial ring in $n$ variables over $\mathbb{F}$ , consisting of polynomials of fixed degree $d$ |
| $\sigma$                        | A permutation of a finite collection of elements (the length of the collection is specified in context)       |

## Vector Spaces

|                               |                                                                                                                                                       |
|-------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\hat{V}_i$                   | The tensor product $V_1 \otimes \dots \otimes V_{i-1} \otimes (\bigoplus_{\lambda \in \Lambda} \mathbb{F}) \otimes V_{i+1} \otimes \dots \otimes V_d$ |
| $\langle v_i \rangle_{i=1}^n$ | The linear span of $n$ vectors                                                                                                                        |
| $S^d(\mathbb{F}^n)$           | Symmetric tensors (forms) of order $d$ over $\mathbb{F}^n$                                                                                            |
| $V_i$                         | A vector space over $\mathbb{F}$ having dimension $n_i$                                                                                               |
| $V_i^*$                       | The dual vector space to $V_i$                                                                                                                        |
| $\hat{V}_i$                   | The tensor product of vector spaces excluding $V_i$ , defined as $V_1 \otimes \dots \otimes V_{i-1} \otimes V_{i+1} \otimes \dots \otimes V_d$        |

## Vectors

|                     |                                                                                                             |
|---------------------|-------------------------------------------------------------------------------------------------------------|
| $\mathbf{1}_n$      | The vector of all 1's lying in $\mathbb{F}^n$                                                               |
| $\alpha$            | A vector of non-negative integers $(a_1, \dots, a_n)$                                                       |
| $\mathbf{x}^\alpha$ | $\prod_{i=1}^n x_i^{a_i}$ where $n$ is taken from context                                                   |
| $\mathbf{x}$        | A vector of indeterminates $(x_1, \dots, x_n)$                                                              |
| $e_{i,j}$           | The $j$ -th standard ordered basis vector in $V_i$ . The $i$ may not be dropped if it is clear from context |
| $v(k)$              | The $k$ th entry in a vector $v$                                                                            |
| $v_i^k$             | The $k$ -th vector lying in $V_i$                                                                           |

**Tensors**

|                      |                                                                                                                                                          |
|----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\bar{\mathcal{M}}$  | The tensor constructed by taking elements of $\mathcal{M}$ as slices                                                                                     |
| $\mathcal{M}$        | A finite family of tensors                                                                                                                               |
| $T(k_1, \dots, k_d)$ | The element located at the position $(k_1, \dots, k_d)$ in a tensor $T$                                                                                  |
| $T_j^i$              | The $j$ -th slice along the $i$ -th mode of a tensor $T$ (see Definition 4.1)                                                                            |
| $T^{(i)}$            | The flattening of a tensor $T$ along the $i$ th mode (see Definition 4.2)                                                                                |
| $T^\emptyset$        | The unravelling of a tensor $T$ into a vector, with elements indexed by $\times_{i=1}^d [n_i]$ in lexicographic order (see remarks after Definition 4.2) |
| $T^{\otimes d}$      | The tensor product of $T$ with itself $d$ times, $\underbrace{T \otimes T \otimes \dots \otimes T}_{d \text{ times}}$                                    |

**General Functions**

|                                               |                                                                                                                                                                                          |
|-----------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\pi_N^i(T)$                                  | The mapping denoting a projection down to the vector space $\mathbb{F}^N$ along the $i$ -th mode of a tensor (see Definition 4.6)                                                        |
| $Adj(T, \mathcal{M}_1, \dots, \mathcal{M}_d)$ | The mapping denoting the general adjoining of collections of slices $\mathcal{M}_i$ along all $i$ modes of a tensor $T$                                                                  |
| $Adj^i(T, \bar{\mathcal{M}})$                 | The mapping denoting the adjoining of $\mathcal{M}$ as slices along the $i$ -th mode of a tensor $T$ (see Definition 4.4)                                                                |
| $C_n(T)$                                      | The mapping that constructs the $n$ -th clone of a tensor $T$                                                                                                                            |
| $F^{(i)}(T)$                                  | The mapping denoting the flattening of a tensor $T$ along the $i$ -th mode                                                                                                               |
| $Mod^i(T, \bar{\mathcal{M}})$                 | The mapping generating a family of tensors by slicing $T$ with elements from $\mathcal{M}$ along the $i$ th mode (see Definition 4.9)                                                    |
| $pad^i(A, T)$                                 | The mapping denoting the padding of a lower order tensor $A$ into the same format as $T$ by inserting 0's along the $i$ th mode (see Definition 4.7)                                     |
| $Pad_j^i(A, T)$                               | The mapping taking a lower order tensor $A$ into the same format as a higher order tensor $T$ , now seen as the only non-zero $j$ -th slice along the $i$ th mode (see Definition 4.8)   |
| $S_j^i(T)$                                    | The mapping that selects the $j$ -th slice along the $i$ -th mode of a tensor $T$                                                                                                        |
| $SAdj(T, \mathcal{M})$                        | The mapping denoting the symmetric adjoining operation defined for a symmetric tensor $T$ and a collection of symmetric slices $\mathcal{M}$ , $Adj(T, \mathcal{M}, \dots, \mathcal{M})$ |

$T \bmod (\mathcal{M}_1, \dots, \mathcal{M}_d)$  The mapping denoting the family generated by slicing  $T$  along each  $i$ -th mode with slices from  $\mathcal{M}_i$

### Scalar-Valued Functions

**Rk**( $T$ ) The rank of a tensor  $T$

**SRk**( $T$ ) The symmetric rank of a symmetric tensor  $T$

# List of Figures

|   |                                                                                                                                         |    |
|---|-----------------------------------------------------------------------------------------------------------------------------------------|----|
| 1 | Example of a tensor $T \in \mathbb{F}^4 \otimes \mathbb{F}^3 \otimes \mathbb{F}^2$ , represented in the form of a 3-way array . . . . . | 1  |
| 2 | Illustrating a tensor product of $R$ -modules . . . . .                                                                                 | 4  |
| 3 | Slicing a tensor $T$ along different modes . . . . .                                                                                    | 20 |
| 4 | Flattening a tensor $T$ . . . . .                                                                                                       | 22 |
| 5 | Adjoining a slice $A$ to a tensor $T$ . . . . .                                                                                         | 26 |
| 6 | Family generated from a tensor $T$ by a slice $A$ . . . . .                                                                             | 28 |
| 7 | 2 clone of a tensor $T$ . . . . .                                                                                                       | 29 |



# Chapter 1

## Introduction

The title of this thesis centers around an open question about tensors. Tensors are mathematical tools, used to describe multiple linear relationships that might exist between mathematical objects. This ability of tensors to describe such inter-related information has spurred the development the study of *multi*-linear algebra. The focus of this thesis is threefold: how are tensors constructed, where do they live when seen as mathematical objects, and how they can be broken down?

The current state of Comon's conjecture (an open problem for the past 20 years) is hard to describe succinctly. While mathematicians have steadily increased the number of cases for which the conjecture remains valid, an answer to the question in full generality for  $\mathbb{R}$ - or  $\mathbb{C}$ -tensors remains unknown. The hope of this thesis is that at least some of the work done in recent years can be highlighted as useful, and paves the way towards either a definitive counterexample, or a proof of the conjecture.

This question about symmetric tensors merits study because of the powerful nature of tensors as describing of multilinear relationships. One of the more important problems that has come to light in recent years is the computational complexity of matrix multiplication ([18]), and tensors and their ranks help to quantify exactly how efficient the algorithms for matrix multiplication are. Tensors also find their use in quantum mechanics (describing the entanglement of particles), general relativity (as tools to describe the curvature of space-time [9]), chemometrics, psychometrics (representing combinations of chemicals or traits that are interrelated [28]), in data science and




Figure 1: Example of a tensor  $T \in \mathbb{F}^4 \otimes \mathbb{F}^3 \otimes \mathbb{F}^2$ , represented in the form of a 3-way array

computer vision [20] and in signal processing (the source selection problem [18]) — to name some more examples.

## Comon's Conjecture

At its heart, Comon's Conjecture centers around two central themes: the *rank* and the *symmetry* of tensors. A tensor's rank describes how it can be decomposed into much simpler constituent tensors, and its symmetry describes how the tensor behaves under permutation actions.

The conjecture was first stated openly in July 2008 at "Geometry and Representation Theory of Tensors for Computer Science, Statistics and Other Areas" - a workshop organised at the American Institute for Mathematics [19] in the group focusing on "Multilinear Techniques in Data Analysis and Signal Processing". The conjecture as stated in the report is as follows:

**Conjecture** (Comon, (Oeding 2008 [19])). *Do the rank and symmetric rank of  $T$  always coincide, where  $T$  is a symmetric, real or complex tensor?*

This report also covered work done previously by Comon et al. [13] where they also showed that the conjecture held for conditions bounding the symmetric rank or order of a tensor in question. Since 2008, the mathematical community has further proved the conjecture true under various other assumptions (see Chapter 3).

In 2018, Shitov [22] proposed the construction of a counterexample living in  $(\mathbb{C}^{800})^{\otimes 3}$  with symmetric rank at least 904, and a rank of at most 903. In 2024 however, an erratum published [14] pointed out an error in the original paper, rendering the conjecture still unsolved in the complex case. In the intervening time, a paper published by Wang and Seigal [27] demonstrated the construction of a real, order 6, symmetric tensor having real rank 30913 and symmetric rank 30914, thus disproving the conjecture for the real case. This paper built on a method of construction developed in a preprint from Shitov in 2020 [23]. These ideas have also been developed further in subsequent preprints - [24] and [25], again from Shitov.

The following chapters and sections highlight important terminology required to state and describe the setting of Comon's conjecture, highlight the work done to prove the conjecture in the past, and they finally describe the major constructions required to construct the real counterexample.

**Chapter 2** covers some preliminary terminology and definitions required to understand the history of attempts to prove Comon's conjecture. Personal contributions in this chapter include the proof of a theorem showing the isomorphism between symmetric tensors of fixed order and homogenous polynomials of fixed degree. [ADD ANY OTHER PERSONAL CONTRIBUTIONS HERE]

**Chapter 3** goes over the literature that currently exists since the statement of the problem in 2008. The papers discussed highlight the valid attempts at proving Comon's conjecture over  $\mathbb{R}$  and  $\mathbb{C}$ . Personal contributions to this chapter are the compilation of all these results in a singular location, highlighting the state of the art currently. [PROOFS INCLUDED IN THIS CHAPTER ARE NOT PERSONAL, OTHERWISE SPECIFY HERE]

**Chapter 4** focuses on understanding Shitov's methodology and approach to constructing a counterexample to Comon's conjecture. The culmination of this chapter is an examination of the paper from Wang and Seigal [27], a published real-valued counterexample to the conjecture. Personal contributions to this chapter include explicit proofs for the various operations discussed, most of which do not exist in the existing literature. [PROVIDE EXAMPLES]

**Chapter 5** highlights the gaps present in the literature so far [WHAT ARE THESE], illustrates problems with the current approach to constructing counterexamples, and seeks to highlight potential future avenues to solving this problem once and for all. [COPY PASTE THIS TO THE FINAL CHAPTER TO KNOW WHAT TO FOCUS ON WHEN FINISHING WRITING THIS THESIS]

# Chapter 2

## Preliminaries Regarding Tensors

### 2.1 Tensors in an Algebraic Setting

The most abstract way of thinking about tensors is when describing them as elements of a tensor product of modules. Consider  $A, B$  and  $P$  as being modules over a ring  $R$ . Then any bi-linear mapping,  $t : A \times B \rightarrow P$ , is described by acting on elements from  $A \times B$  in the following manner:

$$t(\lambda a, b) = t(a, \lambda b) = \lambda t(a, b), a \in A, b \in B, \lambda \in R, t(a, b) \in P$$




Figure 2: Illustrating a tensor product of  $R$ -modules

Such a bilinear (or even generalising to a multilinear mapping) between  $R$ -modules can be represented by an element lying in their tensor product, such that Figure 2 commutes [26] (illustrated specifically for the case mentioned above). Such a tensor product of  $R$ -modules is unique upto unique isomorphisms, and this is referred to in algebraic literature as a *universal property* of tensor products.

While such an abstract description is useful when showing that any general elements lying in the specified spaces  $A$  and  $B$  can be combined to form a tensor, it does not specify details required for showing representations of this tensor.

A more useful approach to understanding tensors is to begin from a foundation in linear algebra. As matrices represent linear transformations between vector spaces  $V$  and  $W$ , so  $d$ -way arrays can be used to represent multilinear relationships between vector spaces  $V_1, V_2 \dots V_d$ , where each  $V_i$  has a dimension  $n_i$ . Diving deeper into this analogy, a matrix of the form  $n_i \times n_j$  behaves either as a bilinear map from the product  $V_i^* \times V_j \rightarrow \mathbb{F}$ , or as linear map  $V_j \rightarrow V_i$  or as its dual  $V_i^* \rightarrow V_j^*$ . This hybrid nature of matrices is a reflection of the underlying fact that matrices represent order-2 tensors. Since computing elements  $A_{ij}$  of the matrix  $A$  depends on a choice of bases for both

vector spaces  $V$  and  $W$ , an analogous choice of bases must be made to represent the  $d$ -linear tensor  $T \in V_1 \otimes V_2 \otimes \cdots \otimes V_d$  as a  $d$ -way array.

For the purposes of this thesis, unless specified, each vector space will be considered over a field  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{C}$ . Here each vector space  $V_i$  is equipped with its standard basis  $\{e_{i,j}\}_{j=1}^{n_i}$  where  $e_{i,j}(k) = \delta_{j,k}$ . Each different choice of basis changes how the tensor  $T$  is represented as a  $d$ -way array.

**Definition 2.1** (Order of a Tensor). An element lying in the tensor product of  $d$  vector spaces,  $T \in V_1 \otimes \cdots \otimes V_d$ , is said to be an **order- $d$  tensor**.

Consider  $T := v_1 \otimes \cdots \otimes v_d$ , a specific order- $d$  tensor, with each component vector  $v_i \in V_i$ . Then, in coordinates

$$v_i = \sum_{j=1}^{n_i} v_i(j) e_{i,j}, \quad v_i(j) \in \mathbb{F} \quad \forall j = 1, \dots, n_i.$$

The elements of a  $d$ -way array that corresponds to  $T$  then can be given in the following manner:

$$T(k_1, k_2, \dots, k_d) = \prod_{i=1}^d v_i(k_i),$$

where  $0 \leq k_i \leq n_i$ .

In a similar fashion, given two tensors in  $V_1 \otimes \cdots \otimes V_d$  of the specific form  $T_1 := v_1 \otimes \cdots \otimes v_d$  and  $T_2 := w_1 \otimes \cdots \otimes w_d$ , then one can describe the  $d$ -way array corresponding to the tensor  $T_1 + T_2$  in coordinates as follows

$$(T_1 + T_2)(k_1, k_2, \dots, k_d) = \left( \prod_{i=1}^d v_i(k_i) \right) + \left( \prod_{i=1}^d w_i(k_i) \right),$$

indicating simply a componentwise addition in the corresponding  $d$ -way arrays. A similar description can be made when scaling a tensor by any factor  $\lambda \in \mathbb{F} \rightarrow \lambda T$ .

Therefore an order  $d$  tensor  $T \in V_1 \otimes \cdots \otimes V_d$  that can be expressed as a sum of these specific *elementary* forms (as above) can also be represented by a  $d$ -way array. This property is satisfied by any general order- $d$  tensor (see Definition 2.4).

**Definition 2.2** (Modes of a Tensor). Given an order- $d$  tensor,  $T \in V_1 \otimes \cdots \otimes V_d$ , the modes of  $T$  refer to specifying the  $d$  distinct ways that  $T$  as a tensor can be oriented. Thus a tensor of order  $n$  has exactly  $n$  modes.

**Remark.** Modes of are useful because they provide a specific orientation along which operations can be done on a given tensor. When working with tensors as  $d$ -way arrays, the modes can be seen as the distinct indices in the overall tuples that index the elements of the array.

**Example 2.1.** Consider

$$T := (a, b, c, d) \otimes (x, y, z) \otimes (p, q) \in \mathbb{F}^4 \otimes \mathbb{F}^3 \otimes \mathbb{F}^2.$$

Then  $T$  is an order-3 tensor having 3 modes. each entry in the 3-way array representing  $T$  is indexed by a tuple  $(i_1, i_2, i_3) \in [4] \times [3] \times [2]$ .

## Viewing Tensors

A tensor  $T := v_1 \otimes \cdots \otimes v_d \in V_1 \otimes \cdots \otimes V_d$  can be viewed in the following different ways [18]:

1. As a multilinear mapping between  $V_1^* \times V_2^* \times \cdots \times V_d^* \rightarrow \mathbb{F}$  (extending the analogy of looking at matrices as order-2 tensors)

$$(f_1, f_2 \dots f_d) \mapsto \prod_{i=1}^d f_i(v_i).$$

Here  $f_i \in V_i^*$ , the dual space to each vector space  $V_i$ .

2. For every choice of  $i \in \{1, \dots, d\}$ ,  $T$  can be viewed as a linear mapping from  $V_i^* \rightarrow V_i := V_1 \otimes \cdots \otimes V_{i-1} \otimes V_{i+1} \otimes \cdots \otimes V_d$ . Viewing  $T$  this way takes a vector (an order-1 tensor) to an order- $(d-1)$  tensor.

$$f_j \mapsto f_j(v_i)(v_1 \otimes \cdots \otimes v_{i-1} \otimes v_{i+1} \otimes \cdots \otimes v_d). \quad (2.1)$$

This viewpoint can be seen when defining an operation that selects a specific "slice" of a tensor (or even a linear combination of them), after fixing the mode  $i$ .

3. Finally, as the dual of the defined linear map (see Equation (2.1)), now going from  $V_i^* \rightarrow V_i$ . This point of view of transforms an order- $(d-1)$  tensor to a rescaling of the vector  $v_i$ .

$$f_1 \otimes \cdots \otimes f_{i-1} \otimes f_{i+1} \otimes \cdots \otimes f_d \mapsto \left( \prod_{j \neq i} f_j(v_j) \right) v_i.$$

Here note that the dual operation and the tensor product operation commute.

Each of these different viewpoints holds for order- $d$  tensors in general, not just for these specific elementary tensors. This becomes evident in the following section on tensor rank and decompositions.

## 2.2 Rank

The first key focus of Comon's conjecture involves figuring out how tensors can be built. The following example shows that there exist tensors that cannot be written as simply the tensor product of vectors analogously to how not all vectors in a vector space can be written as rescaled basis vectors.

**Example 2.2.** When dealing with  $2 \times 2$  matrices  $\in \mathbb{F}^2 \otimes \mathbb{F}^2$ , consider  $e_1 \otimes e_1 + e_2 \otimes e_2$  (the identity matrix). Such a matrix cannot be written as a simple tensor product of two vectors  $v_1 \otimes v_2$ , where  $v_1, v_2 \in \mathbb{F}^2$ .

Making this preliminary observation leads to the following definition —

**Definition 2.3** (Elementary Tensor). Consider an order- $d$  tensor,  $T$ , of the specific form  $T := v_1 \otimes \cdots \otimes v_d \in V_1 \otimes \cdots \otimes V_d$ . Since  $T$  can be written explicitly as the tensor product of  $d$  vectors, it is called an **elementary tensor**. Equivalent names for such tensors in the literature reviewed are *rank-1* tensors and *decomposable* tensors.

Using these elementary tensors as building blocks provides a way to describe the construction of any general order- $d$  tensors. (In the most naive way, this can be done by considering all elementary tensors of the form  $e_{1,j_1} \otimes \cdots \otimes e_{d,j_d}$ ). This is why any general order- $d$  tensor can be represented by a  $d$ -way array.

**Definition 2.4** (Tensor Decomposition [28]). Consider any order- $d$  tensor  $T \in V_1 \otimes \cdots \otimes V_d$ . A **decomposition** of  $T$  consists of writing  $T$  as a sum of elementary tensors

$$T = \sum_{k=1}^s v_1^k \otimes \cdots \otimes v_d^k,$$

with each of the  $v_i^k \in V_i$ .

**Remark.** Here note that there exist multiple names for describing the decomposition of a tensor into the sum of elementary tensors - polyadic forms, canonical decomposition (CANDECOMP), parallel factors (PARAFAC) etc. The above definition agrees with what is known in the literature as CP-decomposition (a combination of CANDECOMP/PARAFAC). [16]

Thus, for any given tensor  $T$ , there can exist multiple decompositions, not just depending on the choice of basis vectors for each  $V_i$ . Two questions naturally follow this definition

- Q1.** For a given order- $d$  tensor  $T$ , does there exist a shortest decomposition?
- Q2.** If such a shortest decomposition exists, is it unique (upto the choice of basis vectors)?

The answer to the first question can be given in terms of the following definition:

**Definition 2.5** (Tensor Rank). The **rank** of a tensor  $T$ , denoted  $\mathbf{Rk}(T)$ , is the minimal length of any possible decomposition of  $T$ .

A rank decomposition of a tensor always exists, because tensors can always be written in terms of elementary tensors. Computing the rank of a given tensor is not an easy problem, but there exist multiple approaches in the literature that rely on both the algebraic properties of tensors and associated transformations, and the geometric properties of the varieties on which these tensors lie. Tensor rank also does not depend on which elementary tensors are involved in a decomposition. Thus Q1 as given above has an affirmative answer, but Q2 does not — there can exist multiple rank decompositions.

## 2.3 Symmetric Tensors

The other focus of Comon's Conjecture revolves around tensors that exhibit the properties of symmetry. These tensors become valuable in applications simply because the symmetry means that they require "less" information to be constructed. What makes a tensor symmetric is the next logical question to tackle, and makes it necessary to define how permutations act on a given tensor.

Given  $\sigma \in S_d$  (the symmetry group of  $\{1, 2, \dots, d\}$ ), one can define the permutation action of  $\sigma$  on a tensor  $T$  as follows:

**Definition 2.6** (Permutation of a Tensor). Let  $T \in V := V_1 \otimes \dots \otimes V_d$  be an order- $d$  tensor. Then  $\sigma \in S_d$  acts on all tensors in  $V$  by linearly extending the following mapping:

$$\sigma : V_1 \otimes \dots \otimes V_d \rightarrow V_{\sigma_1} \otimes \dots \otimes V_{\sigma_d}, v_1 \otimes \dots \otimes v_d \rightarrow v_{\sigma_1} \otimes \dots \otimes v_{\sigma_d}.$$

This induces a corresponding transformation from a general  $T \mapsto \sigma \cdot T := \sigma \cdot (\sum_{i=1}^r T_i)$ , where each  $T_i$  is an elementary tensor.

To talk about tensors being invariant under permutation actions, it is necessary to work with tensors that retain the same format after permutation. This means they must exist in the space  $V^{\otimes d} := \underbrace{V \otimes \dots \otimes V}_{d \text{ times}}$ . These tensors are sometimes referred to in the

literature as being *cubical*.

The definition of a symmetric tensor now follows:

**Definition 2.7** (Symmetric Tensor [13]). Consider a tensor  $T \in V^{\otimes d}$ .  $T$  is said to be **symmetric** if the following is true:

$$T = \sigma \cdot T \quad \forall \sigma \in S_d.$$

**Remark.** As a consequence of this definition, the following equivalent equality condition also holds for symmetric tensors

$$T = \frac{1}{d!} \sum_{\sigma \in S_d} \sigma \cdot T.$$

This gives rise to what can be called the **symmetrisation operator**  $\mathcal{S} : V \rightarrow V$  that takes  $T \mapsto \mathcal{S}(T) := \frac{1}{d!} \sum_{\sigma \in S_d} \sigma \cdot T$ . This resulting tensor  $\mathcal{S}(T)$  is always symmetric.

Since every tensor can be expressed as a sum of elementary tensors (see Definition 2.4),  $\mathbf{Rk}(T)$  is defined for symmetric tensors  $T$  as well. However, one can define *elementary symmetric tensors* as tensors in  $V^{\otimes d}$  of the form  $v^{\otimes d}$  where  $v \in V$ . Establishing this definition allows for symmetric tensors to be written as a sum of these elementary symmetric tensors, resulting in the definition of **symmetric decompositions** and the analogous notion of **symmetric rank**, denoted  $\mathbf{SRk}(T)$ .

**Lemma 2.1** ([13]). Let  $T$  be a symmetric, order- $d$  tensor lying in  $(\mathbb{F}^n)^{\otimes d}$ . Then there exist vectors  $v_1, \dots, v_s \in \mathbb{F}^n$  such that

$$T = \sum_{i=1}^s v_i^{\otimes d}$$



*Proof.* Suppose not. Then there must exist a symmetric, order- $d$  tensor  $T$  that cannot be expressed in the form  $\sum_{i=1}^s v_i^{\otimes d}$  for any  $s > 0$ . Then either  $T$  is asymmetric (a violation of the initial assumption that the lemma was false, leading to a contradiction), or that  $T$  can only be expressed as an infinite sum of symmetric tensors. But since the space  $V_1 \otimes \cdots \otimes V_d$  is finite dimensional, such a sum can be expressed in a finite manner, again leading to a contradiction to our initial assumption. Therefore, a symmetric decomposition must always exist for  $T$ , a symmetric order- $d$  tensor.  $\square$

**Remark.** Comon's proof involves showing that the space generated by all  $k$ -th tensor powers of linear forms in the symmetric tensors cannot be contained in any non-zero non-trivial hyperplane. Their proof also involves defining a bilinear form on the space of the symmetric tensors.

**Example 2.3.** Consider the tensor

$$\begin{aligned} T &:= (1, 0, 0) \otimes (1, 0, 0) \otimes (1, 0, 0) + (0, 0, 1) \otimes (0, 0, 1) \otimes (0, 0, 1) \\ &\quad + (0, 1, 0) \otimes (0, 1, 0) \otimes (0, 1, 0) \\ &= (1, 0, 0)^{\otimes 3} + (0, 0, 1)^{\otimes 3} + (0, 1, 0)^{\otimes 3}. \end{aligned}$$

This is definitely a symmetric decomposition, and  $T$  is a symmetric tensor.

It is clear that not every tensor can be seen as symmetric, and therefore the subset of all order- $d$  tensors that are symmetric is denoted  $S^d(\mathbb{F}^n) \subseteq (\mathbb{F}^n)^{\otimes d}$  — notation usually reserved for homogenous polynomials in  $n$  variables of fixed degree  $d$ . In fact it can be shown that these spaces,  $S^d(\mathbb{F}^n)$  and  $\mathbb{F}[x_1, \dots, x_n]_d$ , are isomorphic to each other. This isomorphism is a powerful tool that allows for an interchange of polynomial and tensor notation (in an appropriate manner) when dealing with symmetric tensors.

## Symmetric Tensors and Homogenous Polynomials of Fixed Degree

A homogenous polynomial  $f \in \mathbb{F}[x_1, \dots, x_n]_d$  is of the form

$$f(x_1, \dots, x_n) = \sum_{a_i \geq 0, \sum_i a_i = d} \lambda_{(a_1, \dots, a_n)} \prod_{i=1}^n x_i^{a_i},$$

where all the coefficients  $\lambda_{(a_1, \dots, a_n)} \in \mathbb{F}$ . The following notation is also standard in the literature:  $\mathbf{x} := (x_1, x_2, \dots, x_n)$  and  $\alpha := (a_1, a_2, \dots, a_n) \in (\mathbb{N} \cup \{0\})^n$ . Then one has  $\mathbf{x}^\alpha := \prod_{i=1}^n x_i^{a_i}$ . Further  $|\alpha| = \sum_i a_i$ , so the homogenous polynomial as stated above can be rewritten as

$$f(\mathbf{x}) = \sum_{|\alpha|=d} \lambda_\alpha \mathbf{x}^\alpha.$$

The following proposition is standard in tensor decomposition literature, but is stated again for convenience.

**Proposition 2.1.** *For fixed values of  $d, n > 0$ , the map*

$$\Phi_d : \mathbb{F}[x_1, \dots, x_n]_d \rightarrow S^d(\mathbb{F}^n),$$

defined by linearly extending the following mapping of the degree- $d$  monomials

$$\mathbf{x}^\alpha := \prod_{i=1}^n x_i^{a_i} \mapsto \frac{1}{d!} \sum_{\sigma \in S_d} \sigma \cdot \left( \bigotimes_{i=1}^n e_i^{\otimes a_i} \right),$$

is an isomorphism of vector spaces over  $\mathbb{F}$ .

*Proof.* Each degree- $d$  monomial can be seen as a basis vector in the vector space  $\mathbb{F}[x_1, \dots, x_n]_d \cong \mathbb{F}^{\binom{n+d-1}{d}}$ . These monomials are mapped to symmetric tensors (by the construction above, and by the definition of symmetric tensors).  $\Phi_d$  is thus well defined mapping, since all other homogenous degree- $d$  polynomials can be constructed as linear combinations of these basis vectors. Therefore for  $\lambda \in \mathbb{F}$ ,

$$\Phi(\lambda \mathbf{x}^\alpha + \mathbf{x}^\beta) = \lambda \left( \frac{1}{d!} \sum_{\sigma \in S_d} \sigma \cdot \left( \bigotimes_{i=1}^n e_i^{\otimes a_i} \right) \right) + \left( \frac{1}{d!} \sum_{\sigma \in S_d} \sigma \cdot \left( \bigotimes_{i=1}^n e_i^{\otimes b_i} \right) \right) = \lambda \Phi(\mathbf{x}^\alpha) + \Phi(\mathbf{x}^\beta).$$

Further, every elementary symmetric tensor  $v^{\otimes d} = (\sum_{i=1}^n v_i e_i)^{\otimes d}$  can be seen as the image of the  $d$ -th power of the linear form  $(\sum_{i=1}^n v_i x_i)^d$ . Since every symmetric tensor has a symmetric tensor decomposition,  $\Phi_d$  must also be surjective.

[CHECK INJECTIVITY ARGUMENT HERE]

Suppose  $\Phi_d$  is not injective. Then there exist  $f, g \in \mathbb{F}[x_1, \dots, x_n]_d$  such that  $\Phi_d(f) = \Phi_d(g)$  but  $f \neq g$ . Now if  $f \neq g$ , then there is at least one coefficient  $f_{\alpha'} \neq g_{\alpha'}$ . Without a loss of generality, let this be the only index where the functions are not equal. Now we definitely have that  $\Phi_d(f) - \Phi_d(g) = 0$ , by our assumption. This is equivalent to the following

$$\sum_{|\alpha|=d} \frac{f_\alpha - g_\alpha}{d!} \sum_{\sigma \in S_d} \sigma \cdot \left( \bigotimes_{i=1}^n e_i^{\otimes a_i} \right) = 0.$$

But this leads to the following expression

$$\frac{f_{\alpha'} - g_{\alpha'}}{d!} \left( \sum_{\sigma \in S_d} \sigma \cdot \left( \bigotimes_{i=1}^n e_i^{\otimes a_i} \right) \right) = 0,$$

a contradiction, because none of the tensors on the left side of the multiplication are 0, and by the initial assumption, the fraction must also be non-zero, and this multiplication is occurring in an integral domain. Thus the initial assumption must be false, and  $\Phi_d$  must be injective.  $\square$

In fact  $\Phi_d$  can be seen as a component of a larger isomorphism between graded vector spaces,  $\Phi : \mathbb{F}[x_1, \dots, x_n] \rightarrow \bigoplus_{d \geq 0} S^d(\mathbb{F}^n)$  [13].

# Chapter 3

## Literature Review

Since the statement of the conjecture in 2008, there has been a substantial amount of work done to arrive at a conclusive final resolution to the problem. Some of this body of work relies on the specific geometric construction of secant varieties and how they relate to the question of computing rank and symmetric rank. This connection to geometry is not surprising because tensors can also be seen as points lying in some topological space (much like vectors and matrices can be seen as representing points, as well as algebraic objects.) This chapter goes over some of the work on proving and disproving Comon, with the final section of this chapter compiling all the conditions currently known for which the conjecture holds or doesn't hold.

### 3.1 Attempts to Prove the Conjecture

The statement of Comon's Conjecture can now be stated in more formal terminology, as introduced in the previous chapter.

**Conjecture** (Comon et al. [13]). *Given a symmetric, order- $d$  tensor  $T \in S^d(\mathbb{F}^n) \subset (\mathbb{F}^n)^{\otimes d}$ , then  $\mathbf{Rk}(T) = \mathbf{SRk}(T)$ .*

The ability to rewrite symmetric tensors of fixed order both as homogenous polynomials and as a sum of regular tensors was the key factor that seems to have inspired this conjecture, as stated in subsection 4.4 from Comon et al. [13]. This section details some of the papers that prove that the conjecture holds under certain constraints.

As mentioned in the introduction Comon et al. [13] showed that the conjecture was true for tensors of "large enough" order — a notion that became concrete in later literature. [COMON ET AL. 2008 PROPOSITION 5.4 - PROOF OF SUFFICIENTLY LARGE (include proposition 4.3 here) CAUSING REWRITE OF THE FOLLOWING PARAGRAPH] To establish such a proof, the authors first proved the following lemma (stated as a proposition), highlighting how a finite collection of linear polynomials, under pairwise independence constraints could be used to generate a linearly independent set of degree- $d$  homogenous polynomials. Doing so requires the following definition:

**Definition 3.1** (Bilinear form on the space of homogenous polynomials [13]). Let  $F, G \in \mathbb{F}[x_1, \dots, x_n]_d$ , where

$$F(\mathbf{x}) = \sum_{|\alpha|=d} \binom{d}{a_1, \dots, a_n} \lambda_{\alpha} \mathbf{x}^{\alpha}, \quad G(\mathbf{x}) = \sum_{|\alpha|=d} \binom{d}{a_1, \dots, a_n} \mu_{\alpha} \mathbf{x}^{\alpha}.$$

Here  $\alpha = (a_1, \dots, a_n)$  and  $\lambda_\alpha, \mu_\alpha \in \mathbb{F}$  for all  $\alpha$ . Then the form  $\langle \cdot, \cdot \rangle : \mathbb{F}[x_1, \dots, x_n]_d \times \mathbb{F}[x_1, \dots, x_n]_d \rightarrow \mathbb{F}$  defined as

$$\langle F, G \rangle := \sum_{|\alpha|=d} \binom{d}{a_1, \dots, a_n} \lambda_\alpha \mu_\alpha,$$

is a symmetric, non-degenerate bilinear form.

**Lemma 3.1** ([13]). *Let  $L_1, \dots, L_r \in \mathbb{F}[x_1, \dots, x_n]_1$ , be linear forms in  $n$  variables. Suppose for all  $i \neq j$ ,  $L_i \neq \alpha L_j$  (where  $\alpha \in \mathbb{F}$ ). Then for any  $d \geq r - 1$ , the polynomials  $L_1^d, \dots, L_r^d$  are linearly independent in  $\mathbb{F}[x_1, \dots, x_n]$ .*

*Proof.* Fix  $d \geq r - 1$ . Suppose if possible, that there exist coefficients  $\alpha_1, \dots, \alpha_r \in \mathbb{F}$  such that  $\sum_{i=1}^r \alpha_i L_i^d = 0$ . Since the bilinear form is non-degenerate, the expression

$$\langle F, \sum_{i=1}^r \alpha_i L_i^d \rangle = 0$$

for any  $F \in \mathbb{F}[x_1, \dots, x_n]_d$ . This simplifies to

$$\sum_{i=1}^r \alpha_i \langle F, L_i^d \rangle = \sum_{i=1}^r \alpha_i F(L_i) = 0$$

where the expression  $F(L_i) = F(\beta_1, \dots, \beta_n)$  when  $L_i = \sum_{i=1}^n \beta_i x_i$ . The next step is to show that a particular degree- $d$  homogenous polynomial  $F$  exists, such that it vanishes on all except one of the  $L_i$ . Consider  $F$ , a degree- $d$  homogenous polynomial, to be the multiple of a product of  $r - 1$  linear forms, such that  $F(L_r) \neq 0$  but  $F(L_i) = 0$  for all  $i < r$ . Then one can determine that  $\alpha_r = 0$ , by the bilinear form expansion derived above. In a similar manner, similar manner, it can be shown that all the other  $\alpha_i$  are also zero, leading to the conclusion that the forms  $L_i^d$  are linearly independent.  $\square$

This lemma also allows for determining when symmetric, order- $d$  tensors can be considered as linearly independent, using the isomorphism in Proposition 2.1.

**Proposition 3.1** ([13]). *Let  $v_1, \dots, v_s \in \mathbb{F}^n$  be pairwise linearly independent. If  $d$  is sufficiently large, then the symmetric tensor, defined as*

$$T := \sum_{i=1}^s v_i^{\otimes d}$$

*satisfies  $\mathbf{Rk}(T) = \mathbf{SRk}(T)$  generically.*

*Proof.* To begin with, for any symmetric tensor  $T$ , it is known that  $\mathbf{Rk}(T) \leq \mathbf{SRk}(T)$ . It is enough to show that in this specific case, for a tensor  $T$  as defined in the statement of this proposition, the reverse inequality i.e.  $\mathbf{Rk}(T) \geq \mathbf{SRk}(T)$  holds.

Suppose  $\mathbf{Rk}(T) = r$  and  $\mathbf{SRk}(T) = s$ . Then there exist decompositions

$$\sum_{i=1}^r x_1^i \otimes \dots \otimes x_d^i = T = \sum_{j=1}^s v_j^{\otimes d}$$

. Without a loss of generality, assume that  $d$  is even. Then  $T$  can be written instead as  $T = \sum_{i=1}^s v_j^{\otimes d/2} \otimes v_j^{\otimes d/2}$ . By the previous lemma, for  $d/2 \geq s - 1$ , the tensors  $v_j^{\otimes d/2}$  are generically linearly independent. Now therefore, one can find functionals  $\Phi_1, \dots, \Phi_s \in S^{d/2}(\mathbb{F}^n)^*$  that are duals to the vectors  $v_j^{\otimes d/2}$ . i.e.  $\Phi_i(v_j^{\otimes d/2}) = \delta_{i,j}$ , the Kronecker delta. Then the initial equality of decompositions can be transformed (after contracting along the first  $d/2$  modes) into the following expressions

$$\sum_{i=1}^r \alpha_{ij} x_{d/2+1}^i \otimes \dots \otimes x_d^i = v_j^{\otimes d/2},$$

for each of the  $v_j$ , where  $\alpha_{ij} = \Phi_j(x_1^i \otimes \dots \otimes x_{d/2}^i)$  where  $i = 1, \dots, r$ . This then implies that the  $s$  linearly independent vectors  $v_j^{\otimes d/2}$  lie in the span of the  $r$  tensors of the form  $\{x_{d/2+1}^i \otimes \dots \otimes x_d^i\}_{i=1}^r$ . This then implies that  $r \leq s$ , and the proof is complete.  $\square$

The authors noted that, as a result of the Veronese embedding (a canonical construction in algebraic geometry), the rank of a symmetric tensor  $T \in S^d(\mathbb{C}^n)$  cannot be more than  $\binom{n+d-1}{d}$ . Also, since  $S^d(\mathbb{C}^n) \subseteq (\mathbb{C}^n)^{\otimes d}$ , they pointed out that the following was true for all  $T \in S^d(\mathbb{C}^n)$ : that the  $\mathbf{Rk}(T) \leq \mathbf{SRk}(T)$ . Their major result was that the conjecture holds "generically" (i.e in a dense subset of order- $d$  symmetric tensors) whenever  $\mathbf{SRk}(T) \leq n$ , the dimension of the underlying vector space.

In particular, for every  $T \in S^d(\mathbb{C}^n)$ , they showed that the conjecture held if the  $\mathbf{SRk}(T) = 1$  or 2.

A concurrent approach to tackling Comon's conjecture arises from the transposition of a classical problem from number theory — the Waring decomposition of a number — to the language of polynomials. This famous problem then gets restated as follows:

---

*"What is the minimum number of linear polynomials required such that a linear combination of them to the  $d$ -th power is equal to a homogenous degree- $d$  polynomial?"*

---

In mathematical notation, take  $f \in \mathbb{F}[x_1, \dots, x_n]_d$ , a homogenous polynomial of degree  $d$  and  $\mathbf{a} := (a_1, \dots, a_n)$  denoting a vector containing coefficients such that  $\mathbf{a} \cdot \mathbf{x} := a_1 x_1 + \dots + a_n x_n$ . Then the Waring problem for polynomials is the following minimisation problem

$$f := \sum_{|\alpha|=d} f_{\alpha} \mathbf{x}^{\alpha} = \min_{s \geq 0} \sum_{i=1}^s (\mathbf{a}_i \cdot \mathbf{x})^d \quad (3.1)$$

A solution to the polynomial Waring problem solves the issue of computing symmetric rank for a given symmetric tensor (see Section 2.3). In fact this relationship is so significant that in the literature, the term **Waring Rank** is used synonymously with symmetric rank when discussing symmetric tensors. An equivalent formulation for the Waring problem is in terms of predicting the dimensions of higher secant varieties and relating these predictions to the symmetric (or Waring) decompositions for symmetric tensors [4].

IS THIS PARAGRAPH REALLY NECESSARY TO THE REST OF THE THESIS?  
DOES IT SHOW THE STATE OF THE ART? HOW CAN IT BE EXPANDED IF IT HAS

TO BE INCLUDED? [The polynomial Waring problem for generic forms was solved by Alexander and Hirschowitz in 1995, by using this secant variety formulation, building on previous work from Campbell, Palatini and Terracini, and using the novel "differential Horace's method" [4]. They showed that  $\sigma_r(\nu_d(\mathbb{P}\mathbb{F}^{n+1}))$  - the  $r$ -th secant variety to the  $d$ -th Veronese embedding of  $\mathbb{F}^{n+1}$  - is of expected dimension with 5 exceptional cases. Brambilla and Ottaviani [6] in 2008 gave a self-contained proof of the theorem, with some simplifications, particularly in the case of order 3 symmetric tensors. The existence of concrete methods for computing rank and symmetric rank meant that for any given symmetric tensor, Comon's conjecture could be quickly verified or disproved.]

Ballico and Bernardi [2] in 2013 showed that that a tensor  $T$  lying on the tangent variety to a Veronese variety must always have  $\mathbf{Rk}(T) = \mathbf{SRk}(T)$ . Their paper highlighted explicit algorithms for the computation of the rank of a tensor that could be expressed as the limit of a linear combination of elementary tensors. The paper showed an application to Comon's conjecture occurred when working specifically with symmetric tensors. [BALLICO BERNARDI 2013 COROLLARY 2 (SECTION 5)]

IF NOT INCLUDING SECANT VARIETIES, WHY INCLUDE THIS? JUSTIFY.[While secant varieties are convenient geometric tools to use for computing rank and symmetric rank via the Alexander-Hirschowitz theorem, Buczyńska and Buczyński in 2013 [7] demonstrated how the commonly used practice of using the  $(r+1) \times (r+1)$  minors of the *catalecticant matrix* to compute the generating equations for the secant variety are not sufficient for higher order Veronese varieties (order  $\gg 2$ ). This paper resulted in the development of *cactus varieties*.]

Computing decompositions covers the first question about symmetric rank — the existence of a minimal length symmetric decomposition. The other question, concerning the uniqueness of a decomposition, requires the definition of when a tensor is *identifiability*.

**Definition 3.2** (Identifiability). A tensor  $T \in V_1 \otimes \cdots \otimes V_d$  is said to be **identifiable** if it admits a unique rank decomposition (modulo scaling by  $d$ -th roots of unity and permutations of summands)

**Example 3.1.** The tensor  $(3, 4) \otimes (1, 1) + (4, 5) \otimes (1, 0)$  can be written as  $(7, 9) \otimes (1, 0) + (3, 4) \otimes (0, 1)$ , and is thus not identifiable, since it can be decomposed in distinct ways, not involving simply a permutation of summands or a direct rescaling.

Identifiability is a useful property when working with tensors because no further computation is required once a rank decomposition has been found for a given identifiable tensor.

When dealing with questions of tensor rank specifically, given an identifiable tensor, it suffices to find only one rank decomposition. This then becomes useful for the purposes of looking for a counterexample to Comon's conjecture. The likely candidate symmetric tensors must either have a symmetric rank greater than the generic rank, or must lie in the complement of a dense subset. [REFINE THIS SENTENCE]

Generic identifiability is the idea that almost every tensor (i.e. a tensor in a dense open subset of tensors), is identifiable. Specific identifiability poses the opposite question - given a specific tensor  $T$ , does it admit a unique decomposition? Chiantini, Ottaviani and Vannieuwenhoven [11, 12] in two subsequent papers tackled both of these questions. Their earlier paper showed that a general symmetric tensor in  $S^d(\mathbb{C}^{n+1})$  of

sub-generic rank  $r$  was identifiable, provided  $r \leq \binom{n+d}{d}/(n+1)$  (with all exceptions again being covered by the Alexander-Hirschowitz theorem). Their second paper focused on showing what criteria could be called "effective" to determine if a given tensor was specifically identifiable. Criteria were designated effective if their conditions were satisfied by generic points (i.e. the criteria were satisfied in an open, dense subset). One of the conclusions of this last paper was that Kruskal's criterion (a condition on collections of order- $d$  tensors, seen as points in a vector space) was *effective*. [READ BOTH PAPERS AGAIN THOROUGHLY. IDENTIFY MAIN QUESTION BEING ANSWERED, MAIN STEPS TO ANSWERING THEM. DO THEY RELATE TO COMON'S CONJECTURE?]

[PROVIDE FULL STATEMENT FOR FRIEDLAND THEOREM 1.1. COMBINE SECTIONS 3,5,6.] Friedland [15] in 2016 focused not on the computation of the symmetric rank, but on the properties of symmetric tensors of a given symmetric rank. In particular the paper focused on showing that when a symmetric tensor  $T$  can be flattened into a matrix  $T^{(i)}$  (see Section 4.1.4), then Comon's Conjecture held if the rank of the tensor,  $\mathbf{Rk}(T) = \mathbf{Rk}(T^{(i)})$  or if  $\mathbf{Rk}(T) = \mathbf{Rk}(T^{(i)})$ . [INCLUDE THEOREM 3.8 AND PROOF] Zhang et al. [28] conclusively showed that Comon's conjecture held for all tensors where the rank of the tensor under consideration is not more than its order.

[REMARK 3.11 AFTER COROLLARY 3.10] Ballico et al [3] at the end of 2018, refocused their efforts on the question of identifiability, and worked on refining Kruskal's criterion (sharp and algebraic in nature) into a geometric criterion that can be verified more easily. They concluded that Comon's conjecture held for tensors whose symmetric rank  $r$  is strictly bounded above by  $\binom{n+e}{e}$  for a specific choice of  $e$ . This choice depended upon splitting the Segre product (another classical construction from algebraic geometry) into two disjoint components and then imposing conditions on a set of points on the Segre variety that exhibited distinct behaviour when projected down to the first component versus the second. The authors of this paper clearly pointed out that even though they had increased the scope of validity of Comon's conjecture, Shitov's proposed counterexample [22] lay outside this scope and thus might have still been a plausible counterexample.

[GO BACK TO QUESTION ON SECANT VARIETIES] Bernardi et al. [4] in 2018 comprehensively covered the literature and algorithms required for the application of secant varieties to questions about (generic) tensor decomposition, with a detailed explanation of the case of symmetric tensor decomposition and its relationship to the AH theorem.

[PROPOSITION 1. MAYBE THEOREM 1] Casarotti et al. [10] in 2018 proposed to improve the bounds for Comon's conjecture when working with flattenings, provided that equations for the secant varieties associated to the Veronese variety could be generated. Their result was for a tensor  $T$  having rank  $r < \binom{n+\lfloor \frac{d}{2} \rfloor}{n}$ , the  $\mathbf{SRk}(T) = r$ , in cases where such generating determinantal equations were known.

[THEOREM 1.3 - MAY NEED TO COVER ALL SECTIONS?] Finally in a paper from 2020, Seigal [21] demonstrated that for all "cubic surfaces", Comon's conjecture does hold. Here cubic surfaces were used to refer to the fact that under the identification with homogenous polynomials, order-3 symmetric tensors correspond to cubic polynomials, and thus their zero sets could be called cubic surfaces. The author also used flattenings of tensors (see Section 4.1.4), dividing their proof into cases depending on the number of singularities on the cubic surfaces in question.



## 3.2 Attempts to Disprove the Conjecture

## 3.3 Overview of Valid Cases of the Conjecture

The current overview of cases where Comon's Conjecture is known to hold is as follows:

- For  $T = \sum_{i=1}^s y_i^{\otimes d}$  and  $d$  large enough [13]. The large enough argument in the paper hinges on a classical result from algebraic geometry, that  $d$  points in  $\mathbb{C}^n$  form the solution set of polynomial equations of degree  $\leq d$ .
- For  $T \in S^d(\mathbb{C}^n)$  and  $\mathbf{SRk}(T) = 1, 2$  [13].
- For  $T \in S^d \mathbb{F}^n$ , and  $\mathbf{Rk}(T) \leq \frac{dn-d+1}{2}$  where  $\mathbb{F} = \mathbb{R}, \mathbb{C}$  [19].
- For  $T \in S^d(\mathbb{C}^n)$  for  $(d, n) \in \{(3, 2), (4, 2), (3, 3)\}$  [15].
- In the case  $d \geq 3$ ,  $\text{char}(\mathbb{F}) \geq 3$ ,  $T \in S^d(\mathbb{F}^n)$  and  $\mathbf{Rk}(T) = \mathbf{Rk}(T^{(i)})$  or  $\mathbf{Rk}(T^{(i)}) + 1$  (Here  $T^{(i)}$  denotes the flattening (see Section 4.1.4) of  $T$  along any mode. Since  $T$  is symmetric all directions are equivalent.) [15]
- When  $T \in S^3(\mathbb{C}^n)$  and  $\mathbf{Rk}(T) \leq 5$  or  $\mathbf{SRk}(T) \leq 6$  [15].
- For  $T \in S^d(\mathbb{F}^n)$  where  $\mathbf{Rk}(T) \leq d$  where  $\mathbb{F} = \mathbb{R}, \mathbb{C}$  [28].
- If  $T$  is a tensor lying on a tangential variety to a Veronese variety [2].
- If  $T \in \mathbb{C}^2 \otimes \mathbb{C}^n \otimes \mathbb{C}^n$  [8].
- For  $T$  being a Coppersmith-Winograd tensor [17].
- Whenever a tensor  $T$  is cubic surface (real or complex) i.e.  $T$  can be represented by a homogenous cubic polynomial  $\sum_{|\alpha|=3} \lambda_\alpha \mathbf{x}^\alpha$  [21].



# Chapter 4

## Constructions required for a Counterexample

### 4.1 Background to the Counterexample

In spite of all of these preliminary results indicating positive answers towards Comon's Conjecture, Yaroslav Shitov seemed to arrive at a method of constructing a potential counterexample, as early as 2018 [22]. An error found in this paper led to the invalidation [REPLACE THIS WORD] of this specific counterexample in 2024, but the period of six years in the interim led to multiple other preprints from Shitov [23–25].

These preprints started with a newer, alternative methodology for the construction of counterexamples. While there is now no material carrying the stamp of peer-review and Shitov's name, his initial contribution and subsequent preprints seem to indicate a viable approach to constructing a counterexample to Comon's conjecture over  $\mathbb{R}$ , if not  $\mathbb{C}$ . [VALUED TENSORS RATHER THAN FIELD]

#### 4.1.1 Shitov's First Paper in 2018

The 2018 paper began with taking the ground field as  $\mathbb{C}$ , and fixing the notation to be used. The main objects of interest in the paper were order-3 tensors, and the introductory section of the paper highlighted the behaviour of such tensors under elementary transformations with respect to their constituent order-2 "slices". The construction of the counterexample began with the description of a  $100 \times 100$  matrix, partitioned into 20 submatrices [INCLUDE GRAPHIC ILLUSTRATING THIS]. Each submatrix was then mapped to a corresponding  $100 \times 100$  matrix, now with entries having the same indices as the submatrix set to 1 and all other entries as 0. Taking the 1– and 2– supports of each of these, and pre- and post-multiplying them with a column-shift operator with parameter  $a$  led to another set of matrices of the same size (still  $100 \times 100$ ).

It is at this point that the paper made a concerning [CHANGE WORD?] construction. A set  $\mathcal{B}$  was constructed consisting of matrices of distinct formats (viz.  $200 \times 200$  and  $400 \times 400$ ). While there is no issue in defining such a set, the paper then constructed "the  $\mathbb{C}$ -linear span" of the elements of this set. Addition of quantities that do not live in the same dimensional vector space, without a specified or implied embedding map (at least from this author's knowledge) presented a major problem with following this

paper.

Later in the paper a tensor decomposition was described (Eq. 5.6 in [22].) This statement was the subject of the erratum published in 2024 [14]. The erratum indicated such a statement was true only in specific cases, but could not be said to hold in general otherwise without proof. Proving the statement has been a non-trivial challenge, and since no answer has been arrived at, the current consensus is that the paper [22] does not lead to a valid counterexample. [PARAGRAPH IS CLUNKY, TRY REWRITING]

While certain choices were made during this process that led to unanswered questions, the overall tools for later establishing a valid counterexample were first laid out in this paper. These included the two minor operations - slicing and flattening, and the three major operations - adjoining slices to tensors, generating linear families modulo slices and cloning.

### 4.1.2 Shitov's Subsequent Preprints

In the following years, the preprints [23–25] from Shitov either tackled Comon's conjecture in a restricted setting, or developed tools to construct more general counterexamples. These might also be used to construct a counterexample to Strassen's conjecture (another tensor rank conjecture concerning the additivity of rank). The preprint from 2020 [23] tackled the conjecture over  $\mathbb{R}$  and it was this preprint that formed the basis for the paper from Wang and Seigal [27].

The constructions made in each preprint followed the same pattern, indicating a common toolset and structure of approach. Each approach saw an order- $d$  tensor as composed of lower order- $(d - 1)$  tensors, which allowed for tensors to be transformed by "slice" along a specified mode. Viewing tensors in this manner opened a way to describe tensors modulo these slices — a useful operation, [seen in both number theory when working with integers modulo primes, and algebraic geometry as a tool to study problems in a more localised setting]. Such a family, when generated, gave an idea of the general properties of the original tensor (how many linearly independent slices along a specified mode, maximum slice rank along each mode etc.) Flattenings assisted with the computation of rank for these slices (because computing the rank of matrices is a known and easier problem to solve than doing so for higher order tensors) and the flattenings lent themselves well to describing rank information that might change in the relationship between a tensor and its slices.

These tools allowed for an analysis of any particular tensor, but the construction aspect of this approach became evident with the adjoining of slices to a tensor and the cloning operation. The former operation changed the format of the original tensor with respect to certain chosen slices, and allowed the rank information of the family generated modulo the slices to be preserved in a new tensor containing both the original and the slices that were required.

Cloning was a way to maintain rank information by simply making a higher format copy, and allowed for more freedom when modifying slices — simply because there were more to choose from. Cloning started to play a role because perturbing such an adjoined tensor (usually by the modification of one of the constituent slices) caused a violation of the properties that were supposed to be preserved.

While each of these preprints started with this general structure, and [25] proposed the construction of a general counterexample over any  $\mathbb{F}$  with  $\text{char}(\mathbb{F}) \neq 3$ , an interme-

diated step involved in all of these was the construction of an *emulating family*. Such a family, if it existed, would allow for the rest of the constructions to proceed. However, proving the existence of such a family was still not trivial. The latest preprint attempted to do so, highlighting a similar construction from [24], but tracing the construction back led to the definition of a function dependent on an arbitrary constant - in the same way as the paper [22] relied on the existence of an arbitrarily partitioned  $100 \times 100$  matrix. Such choices, while perhaps valid, did not appear to be explained at any point in their corresponding documents, raising some skepticism and prompting further investigation into better documented methods that could explain, or proofs that could deny it. [IT HERE UNCLEAR].

### 4.1.3 Slicing a Tensor

Viewing the columns or rows of a matrix as linearly dependent or independent vectors and how they affect the overall linear transformation described by the matrix is the exact low-order analogue for describing tensors as composed of lower-order slices. Conventionally the columns of a matrix would be the 1-slices, since they occur along the "first" mode, and the rows would therefore be considered 2-slices.

**Definition 4.1** (Slices of a Tensor). Let  $T := v_1 \otimes \cdots \otimes v_d \in V_1 \otimes \cdots \otimes V_d$  be an elementary, order- $d$  tensor. Here each  $v_i \in V_i$ , with  $V_i$  a vector space with dimension  $n_i$ .

Then, the  $j$ -th  $i$ -slice of  $T$  is given by

$$T_j^i := (v_i)(j)(v_1 \otimes \cdots \otimes v_{i-1} \otimes v_{i+1} \otimes \cdots \otimes v_d) \in V_i.$$

Here  $1 \leq j \leq n_i$ . For each *mode*  $1 \leq i \leq d$ , there are  $n_i$  order- $(d-1)$  slices that make up  $T$ .

This definition is equivalent to selecting the  $j$ -th  $i$ -slice of the  $d$ -way array that represents  $T$ . For a general tensor  $T \in V_1 \otimes \cdots \otimes V_d$ , the slice  $T_j^i$  is exactly the sum of the  $j$ -th  $i$ -slices of elementary tensors that make up a rank decomposition of  $T$ .


**Observation.** Through the rest of this thesis, the superscript accompanying a tensor variable indicates the chosen mode along which a specified operation must occur.

**Example 4.1.** Consider a tensor  $T \in \mathbb{F}^4 \otimes \mathbb{F}^3 \otimes \mathbb{F}^2$ , where  $T := (1, 2, 3, 4) \otimes (1, 2, 3) \otimes (1, 2)$ . Then slices along different modes are computed as follows (visualised in Figure 3)

- $T_3^1 = 3((1, 2, 3) \otimes (1, 2))$
- $T_2^2 = 2((1, 2, 3, 4) \otimes (1, 2))$
- $T_2^3 = 2((1, 2, 3, 4) \otimes (1, 2, 3))$

**Observation.** The operation defined above for taking the  $j$ -th  $i$ -slice of an order- $d$  tensor,  $S_j^i : V := V_1 \otimes \cdots \otimes V_d \rightarrow V_i$  defined as  $T \mapsto T_j^i$  is a well-defined linear mapping between vector spaces, by Definition 4.1.

[ILLUSTRATE/HAVE A BETTER EXAMPLE?]


 Figure 3: Slicing a tensor  $T$  along different modes

**Lemma 4.1.** *Given a tensor  $T$  such that  $\mathbf{Rk}(T) = r$ , then there is at least one mode  $1 \leq i \leq d$  having  $r$  linearly independent slices  $T_j^i$  where  $1 \leq j \leq n_i$ .*

*REVIEW PROOF.* Suppose the claim is false i.e. there is no mode  $i$  such that the number of linearly independent slices  $T_j^i$  is  $r$ . Since we are only dealing with a finite order  $d$ , there is a mode  $i$  with the largest number of linearly independent slices  $T_j^i$ . Let this number be  $\rho$ . Suppose

$\rho < r$ : Then each of the slices  $S_j^i(T)$  can be written as a linear combination of exactly  $\rho$  slices of order  $(d-1)$ . One can therefore rewrite the decomposition of  $T$  in terms of vectors in  $V_i$  and the linearly independent set of  $S_j^i(T)$ , a decomposition of length less than  $r$ , contradicting the definition of rank.

$\rho > r$ : Then any decomposition of length  $r$  must miss out at least one of the linearly independent slices that make up  $T$ , a contradiction to the definition of a tensor decomposition.

Thus the assumption of the claim being false leads to a contradiction.  $\square$

A major claim from the literature on Comon's conjecture is the following theorem about the number of linearly independent slices along a chosen mode, when working with rank decompositions. The theorem as stated below first appeared as Lemma 3.1 in [28], but in a more geometric formulation it was proved earlier as Proposition 2.4 in [8].

**Theorem 4.1** ([28], [8]). *Let  $T$  be an order- $d$  tensor of rank  $r$ . Then*

$$T := \sum_{k=1}^r v_1^k \otimes \cdots \otimes v_d^k.$$

For any mode  $i$ , the set of slices of the decomposition

$$\{v_1^k \otimes \cdots \otimes v_{i-1}^k \otimes v_{i+1}^k \otimes \cdots \otimes v_d^k \in V_i \mid 1 \leq k \leq r\}$$

is linearly independent.

[INCLUDE PROOF DON'T LEAVE THIS UNPROVEN] When working with symmetric tensors, slicing along any mode results in a lower order tensor in the same *format* (referring to the dimensions of the underlying vector space). In this case  $S_j^i : (V)^{\otimes d} \rightarrow (V)^{\otimes d-1}$  for all modes  $i$  and all slices  $1 \leq j \leq n_i$ . [CITE REFERENCE HERE FRIEDLAND I THINK]

#### 4.1.4 Flattening a Tensor

Flattening, while not as important in Shitov's constructions[OCCURS IN THE WANG-SEIGAL PAPER MAYBE INCLUDE THOSE DEFINITIONS AT THE END OF THIS SUBSECTION], is an useful operation to verify rank information for tensors. Such an operation can also be used as a quick check when performing operations or constructions on tensors to detect changes in rank. This is due to the fact that computing the rank of a matrix is a much easier problem than computing the rank of an order  $d$  tensor, especially as  $d$  increases.

**Definition 4.2** (Flattening a Tensor). [INCORRECT DEFINITION] Let  $T \in V_1 \otimes \cdots \otimes V_d$ , be an order- $d$  tensor. Define  $T^{(i)} \in \left(\bigoplus_{j \neq i} V_j\right) \otimes V_i$ , the  $i$ -th flattening as follows

$$T^{(i)}((k_1, \dots, k_{i-1}, k_{i+1}, \dots, k_d), k_i) := T(k_1, k_2, \dots, k_d)$$

As with slicing, this definition is equivalent to flattening the  $d$ -way array representing  $T$  into a matrix with the elements in a row indexed by  $[n_i]$  and elements in a column indexed by  $\times_{j \neq i} [n_j]$ , in lexicographic order.


[REORDER EXAMPLES AND OBSERVATIONS, REWRITE TO MAKE CLEARER FIRST WHAT THE NOTATION IS, WHY THE EMPTY SET GETS USED, AND THEN SHOW WITH AN EXAMPLE]

**Example 4.2.** Consider a tensor  $T \in \mathbb{F}^4 \otimes \mathbb{F}^3 \otimes \mathbb{F}^2$ , where  $T := (1, 2, 3, 4) \otimes (1, 2, 3) \otimes (1, 2)$ . Then flattening the tensor along the 2-mode can be computed as

$$T^{(2)} = ((1, 2, 3, 4) \otimes (1, 2))^{\emptyset} \otimes (1, 2, 3).$$

While this looks strange, the vector required to be tensored with  $(1, 2, 3)$  is formed by flattening the matrix  $(1, 2, 3, 4) \otimes (1, 2)$  along the 1-mode. This is equivalent to the vector  $(1, 2, 3, 4, 2, 4, 6, 8)$ . Figure 4 provides an illustration of this example, also highlighting how the column and row indices are generated.

**Observation.** Flattening  $T$  to  $T^{(i)}$  is the operation by which  $T$  could be seen as a linear map between vector spaces instead (a modification of the illustration in Section 2.1). Here  $T^{(i)} \in V_i \otimes V_i$ , seen as a matrix, representing a mapping from  $V_i \rightarrow V_i$ . Each column  $T_j^{(i)} := T^{(i)}(\dots | j)$  represents the *unfolding* of the  $j$ -th  $i$  slice of  $T$ , denoted  $(T_j^i)^{\emptyset}$ , into a vector.

Figure 4: Flattening a tensor  $T$ 

**Observation.** The flattening operation  $F^{(i)} : V := V_1 \otimes \cdots \otimes V_d \rightarrow W \otimes V_i, T \mapsto T^{(i)}$  is a well defined linear mapping between vector spaces where  $W = \mathbb{F}^{\prod_{j \neq i} n_j}$ .

**Remark.** In the literature surveyed, flattening can be done not just along one mode, but for a collection of modes. This gives rise to the empty-set notation denoting the unravelling of a vector. Since this is less important for the following constructions, this more general definition of flattening is not covered. However, in this document, the spirit of such a general operation is retained by also describing the unfolding of an order- $d$  tensor, seen in the following remark.

**Remark.** For a tensor  $T \in V_1 \otimes \cdots \otimes V_d$ , the unfolding  $T^\emptyset$  denotes a vector lying in  $\mathbb{F}^{\prod_{i=1}^d n_i}$ . Here each component of the vector is indexed by the the set  $\times_{i=1}^d [n_i]$ , and  $T^\emptyset((k_1, \dots, k_d)) = T(k_1, \dots, k_d)$ .

When dealing with symmetric tensors, flattening along any mode is equivalent (since the tensor is symmetric), and so  $T^{(i)} = T^{(j)}$  for all  $i \neq j$ . The resulting matrix is an element of  $\mathbb{F}^{(d-1)n} \otimes \mathbb{F}^n$ .

## 4.2 Major New Constructions from Shitov

While the preceding constructions are relatively "minor" in the grand scheme of Shitov's proposed counterexample to Comon, the following three major operations on tensors are integral to the final construction proposed (and indeed implemented by Wang and Seigal [27]). [CLEARER WRITING] The following subsections deal with constructions involving the operations on a specific tensor parametrised by families of tensors of

lower order and appropriate format. It is convenient to represent such families of slices in terms of an appropriate order- $d$  tensor, and then reframe definitions accordingly in terms of this newly created order- $d$  tensor.

**Definition 4.3** (Creating a Tensor from a Family of Slices). Let  $d > 0$  and  $1 \leq i \leq d$ . Take  $\mathcal{M} := \{M_1, \dots, M_N\} \subset V_i$  as a finite collection of order- $(d-1)$  tensors. Since  $\mathcal{M}$  is finite, it can be rewritten as a tensor of order  $d$ . This newly constructed tensor is denoted by  $\bar{\mathcal{M}} \in V_1 \otimes \dots \otimes V_{i-1} \otimes \mathbb{F}^N \otimes V_{i+1} \otimes \dots \otimes V_d$ , defined in terms of each of its  $i$ -slices  $\bar{\mathcal{M}}_j^i$ , where  $1 \leq j \leq N$ ,

$$\bar{\mathcal{M}}(k_1, \dots, k_{i-1}, j, k_{i+1}, \dots, k_d) := M_j(k_1, \dots, k_{i-1}, k_{i+1}, \dots, k_d).$$

[COORDINATE INDEPENDENT DEFINITION POSSIBLE?]

### 4.2.1 Background to Adjoining

Once the notion of a tensor of order  $d$  being composed of various order- $(d-1)$  slices has been established, one natural operation that can be performed is the *adjoining* of slices along the  $d$  modes of the tensor, provided that each slice is of appropriate format. To perform this operation, one needs two pieces of information — the tensor to be transformed, and the collections of slices (of appropriate format) that are to be adjoined. This operation seems to have been first formally defined by Shitov in his 2018 paper [22], and further developed into the language below in his preprint from 2020[23].

[MAYBE REPHRASE? OR SHOW, NOT TELL] In this paper, the operation involving only one mode is discussed, before reframing the operation involving all the modes as the result of a series of successive operations. Doing so requires the tensor to be first embedded into the tensor product of  $(d-1)$  finite-dimensional vector spaces and the following direct sum of the field  $\mathbb{F}$  (denoted  $(\bigoplus_{\lambda \in \Lambda} \mathbb{F})$ , where  $\Lambda$  is an indexing set). To work with this embedding that captures all the information required, it is then necessary to project back down to a tensor product of finite-dimensional vector spaces. Finally when moving between modes, since the lower-order tensors chosen were appropriately sized for the original tensor, a padding function is required to continue the adjoining process, without adding any extraneous information.

**Definition 4.4** (Adjoining Slices to a Tensor along a specific Mode). Let  $T \in V := V_1 \otimes \dots \otimes V_d$  be an order- $d$  tensor, and  $\mathcal{M} = \{M_1, \dots, M_N\} \subset V_i$  be a finite collection of order- $(d-1)$  slices. Define  $\hat{V}_i := V_1 \otimes \dots \otimes V_{i-1} \otimes (\bigoplus_{\lambda \in \Lambda} \mathbb{F}) \otimes V_{i+1} \otimes \dots \otimes V_d$ . Defining  $\hat{V}_i$  in this manner allows for the following definition of the adjoining operation, now in terms of a well-defined function:

$$Adj^i : V \times \hat{V}_i \rightarrow \hat{V}_i,$$

taking  $(T, \bar{\mathcal{M}}) \mapsto \tau$  as:

$$\tau(k_1, \dots, k_d) = \begin{cases} T(k_1, \dots, k_d) & (k_i)_{i=1}^d \in \times_{i=1}^d [n_i] \\ \bar{\mathcal{M}}(k_1, \dots, k_d) & (k_j)_{j \neq i} \in \times_{j \neq i} [n_j] \text{ and } n_i \leq k_i \leq n_i + N \\ 0 & k_i > N + n_i \end{cases}$$



It becomes necessary to define an appropriate projection mapping, to allow the image of this map to be viewed as a tensor in finite dimensions along every mode, and the following definition can be given.

**Definition 4.5** (Tensor of Finite Format). A tensor  $T \in V_1 \otimes \cdots \otimes V_d$  where all the  $V_i$  are finite-dimensional vector spaces is called a tensor of **finite format**

Post the adjoining operation,  $Adj^i(T, \bar{\mathcal{M}})$  lies in a tensor product of vector spaces over  $\mathbb{F}$ , but cannot be said to be of finite format, given the infinite direct sum present in the definition of  $\hat{V}_i$ . To continue working with this object, it is necessary to also develop a method of returning to a finite format context. [DEFINITION BELOW CAN BE GIVEN WITHOUT COORDINATES NO?]

**Definition 4.6** (Projecting to obtain a tensor of finite format). Given an element  $T \in \hat{V}_i$ , define the projection mapping

$$\pi_N^i : \hat{V}_i \rightarrow V_1 \otimes \cdots \otimes V_{i-1} \otimes \mathbb{F}^N \otimes V_{i+1} \otimes \cdots \otimes V_d$$

as restricting  $T$  to an order- $d$  tensor of finite format in a natural way. In coordinates

$$\pi_N^i(T)(k_1, \dots, k_d) = T(k_1, \dots, k_d), (k_i)_{i=1}^d \in [n_1] \times \cdots \times [n_{i-1}] \times [N] \times [n_{i+1}] \times \cdots [n_d]$$

**Remark.** The above two definitions allow for obtaining a tensor of finite format when adjoining a family of slices along only one mode. The next major step to be taken is adjoining slices along multiple directions. Doing so requires the definition of a *padding* operation, that allows for order- $(d-1)$  tensors to be adjusted into appropriate format, depending on the original order- $d$  tensor, before being adjoined in the required direction.

[CAN BE DESCRIBED VIA CANONICAL INJECTION? NO NEED FOR COORDINATE SPECIFICATION]

**Definition 4.7** (Padding an order- $(d-1)$  tensor without increasing order). Given  $d > 0$   $1 \leq i \leq d$  as fixed. Let  $T \in V_1 \otimes \cdots \otimes V_d$  be an order- $d$  tensor, and  $A \in W_1 \otimes \cdots \otimes W_{i-1} \otimes W_{i+1} \otimes \cdots \otimes W_d =: W_i$  be an order  $d-1$  tensor, such that  $\dim(W_j) \geq \dim(V_j)$  for all  $j \neq i$ . This condition allows for  $A$  to be padded according to the  $i$  slices of  $T$

Then

$$pad^i : W_i \times V \rightarrow V_i$$

as a function can be defined in coordinates as

$$pad^i(A, T)(k_1 \dots k_{i-1}, k_{i+1} \dots k_d) = \begin{cases} A(k_1 \dots k_{i-1}, k_{i+1} \dots k_d) & 1 \leq k_j \leq \dim(W_j) \forall j \neq i \\ 0 & \text{otherwise} \end{cases}$$

**Example 4.3.** Consider  $T := (1, 2, 3, 4) \otimes (5, 6, 7) \otimes (8, 9) \in \mathbb{F}^4 \otimes \mathbb{F}^3 \otimes \mathbb{F}^2$  and  $A := (1, 1) \otimes (2, 2) \in (\mathbb{F}^2)^{\otimes 2}$ . Then  $pad^1(A, T) = (1, 1, 0) \otimes (2, 2) \in (\mathbb{F}^3)^{\otimes 2}$ , the same format as the 1 slices of  $T$ . Likewise  $pad^2(A, T) = (1, 1, 0, 0) \otimes (2, 2)$  and  $pad^3(A, T) = (1, 1, 0, 0) \otimes (2, 2, 0)$ .



### 4.2.2 Adjoining Slices to a Tensor

[CHECK ALGORITHM - SYMBOLS ARE OFF. THIS OPERATION REQUIRES A CO-ORDINATE DEFINITION AT LEAST] Combining all of these functions ( $Adj^i$ ,  $\pi_N^i$  and  $pad^i$ ) allows for adjoining families of order- $(d-1)$  tensors along every mode of a given order  $d$  tensor. Given an order- $d$  tensor  $T \in V := V_1 \otimes \cdots \otimes V_d$ , and a collection of finite subsets of order- $(d-1)$  tensors  $\mathcal{M}_i \subset V_i$  for  $1 \leq i \leq d$ , the adjoin of  $T$  with all of the  $\mathcal{M}_i$ , denoted  $Adj(T, \mathcal{M}_1, \dots, \mathcal{M}_d)$ , can be constructed in the following steps:

**Step 1**  $T_1 := \pi_{|\mathcal{M}_1|+n_1}^1 \circ Adj^1(T, \bar{\mathcal{M}}_1)$  resulting in a new intermediate tensor with slices adjoined along the 1-mode, of finite format.

**Step 2a** Let  $\mathcal{M}_2 := \{pad^2(A, T_1) | A \in \mathcal{M}_2\}$ , creating a set of appropriately padded slices, now to be adjoined along the 2-mode.

**Step 2b**  $T_2 := \pi_{|\mathcal{M}_2|+n_2}^2 \circ Adj^2(T_1, \bar{\mathcal{M}}_2)$

$\vdots$

**Step da**  $\mathcal{M}_d = \{pad^d(A, T_{d-1}) | A \in \mathcal{M}_d\}$

**Step db**  $T_d = \pi_{|\mathcal{M}_d|+n_d}^d \circ Adj^d(T_{d-1}, \bar{\mathcal{M}}_d)$

The tensor  $T_d \in (V_1 \oplus \mathbb{F}^{|\mathcal{M}_1|}) \otimes \cdots \otimes (V_d \oplus \mathbb{F}^{|\mathcal{M}_d|})$  is the required result of the process, and can be denoted  $Adj(T, \mathcal{M}_1, \dots, \mathcal{M}_d)$ .

**Observation.** Figure 5 illustrates the above procedure for the case when only one slice is being adjoined in one direction. According to the above definition  $\tau$  can be seen as an order- $d$  tensor still, but now lying in  $(V_1 \oplus \langle e_{1,n_1+1} \rangle) \otimes \cdots \otimes V_d$  where  $e_{1,n_1+1}$  indicates a "new" basis vector added onto the original vector space  $V_1$ .


The **symmetric adjoin** can be denoted  $SAdj(T, \mathcal{M})$  where  $\mathcal{M}$  is adjoined along every mode, following the same process as given in Section 4.2.2.

The adjoining process as described creates a tensor of larger format, but the rank of such a tensor may be appropriately constrained depending on the slices being adjoined. It is this constraining property that Shitov [25] and Wang and Seigal [27] use to construct an intermediate tensor with adjoined slices that has a differing rank and symmetric rank.[IS THERE A PROOF? ILLUSTRATE/DEMONSTRATE]

**Remark.** It may be trivial to note that while the order of the slices in  $\mathcal{M}_i$  matters when constructing the final tensor, the rank information of the constructed tensor is independent of such changes in ordering. Further, the adjoining procedure itself can be modified, to follow any ordering of modes, as long as the slices required have been padded appropriately.

### 4.2.3 Generating Tensor Families modulo Slices

Given that a tensor can admit many non-unique decompositions, one way to analyse the structure of a tensor is via an analogy to Gaussian elimination for matrices. The substitution method as described in [1, 17] has been a computational tool for analysing


 Figure 5: Adjoining a slice  $A$  to a tensor  $T$ 

tensors since the 1960's. Rather than arriving at one possible "lowest" optimal tensor, generating a family of tensors modulo a set of slices is equivalent to looking at all the possible ways a tensor could be transformed, based on transforming its constituent slices along different modes.

Defining such an operation that generates a linear span necessitates one additional, alternate description of padding a tensor into an appropriate format, now in terms of seeing the lower order slice as embedded into an empty higher order tensor. In doing so, it is also necessary to take a candidate higher order tensor as a "template" that the slice can be made to resemble. [REWRITE, CAN MAKE THIS DEFINITION SIMPLER. SIMPLY TENSOR WITH AN APPROPRIATE VECTOR ALONG THE MISSING MODE]

**Definition 4.8** (Padding an order- $(d - 1)$  tensor into higher order). Let  $d > 0$ , and  $1 \leq i \leq d$ . Let  $T \in V_1 \otimes \cdots \otimes V_d$  be the candidate higher order tensor to be used as a template, and  $A \in V_i$  the order  $d - 1$  tensor to be padded into order  $d$ .

Then define  $\text{Pad}_j^i : V_i \times V \rightarrow V$  in coordinates as follows

$$\text{Pad}_j^i(A, T)(k_1, \dots, k_d) = \begin{cases} A(k_1, \dots, k_{i-1}, k_{i+1}, \dots, k_d) & k_i = j \\ 0 & \text{otherwise} \end{cases}$$

Equipped with this method of padding one can now define slicing a tensor  $T$  along one particular mode.

**Definition 4.9** (Tensor Family Generated by Slices along a Mode). Let  $d > 0$  and  $1 \leq i \leq d$ . Let  $T \in V := V_1 \otimes \cdots \otimes V_d$  be an order- $d$  tensor, and  $\mathcal{M} := \{M_1, \dots, M_N\} \subset V_i$  be a finite collection of order- $(d - 1)$  slices of the appropriate format.

Define  $\text{Mod}^i(T, \bar{\mathcal{M}}) : V \times V_1 \otimes \cdots \otimes V_{i-1} \otimes \mathbb{F}^N \otimes V_{i+1} \otimes \cdots \otimes V_d \rightarrow 2^V$  in terms of coordinates as follows ( $2^P$  denoting the power set of a set  $P$ ):

$$\text{Mod}^i(T, \bar{\mathcal{M}}) = \left\{ T + \sum_{k=1}^{n_i} \sum_{j=1}^N \lambda_{j,k} \text{Pad}_k^i(S_j^i(\bar{\mathcal{M}})) \mid \lambda_{j,k} \in \mathbb{F} \right\}.$$

Here slices are first selected from the constructed order- $d$  tensor from the family of lower-order tensors, then these are padded appropriately to remain of the same format as  $T$ , and then scaled and added to  $T$ .

This definition can now be extended to performing this operation along all the  $d$  modes. The only difference now notationally is writing  $\text{Mod}^j(\text{Mod}^i(T, \bar{\mathcal{M}}_i), \bar{\mathcal{M}}_j)$  to denote the  $\text{Mod}^j(S, \bar{\mathcal{M}}_j)$  for every  $S \in \text{Mod}^i(T, \bar{\mathcal{M}}_i)$  and then taking the union of all the resulting sets. Thus the final family  $F := T \text{ mod } (\mathcal{M}_1, \dots, \mathcal{M}_d)$  is a result of the  $d$ -fold composition of the distinct  $\text{Mod}^i$  operations.

**Proposition 4.1.** *Let  $T$  be an order- $d$  tensor  $\in V_1 \otimes \dots \otimes V_d$ . Consider  $\mathcal{M}_i \subset V_i$  and  $\mathcal{M}_j \subset V_j$  as collections of lower-order tensors of format appropriate to the  $i$  and  $j$  slices of  $T$  respectively. Then the family generated modulo  $\mathcal{M}_i$  and  $\mathcal{M}_j$  is the same, regardless of which modulo operation is performed first, i.e.*

$$\text{Mod}^j(\text{Mod}^i(T, \bar{\mathcal{M}}_i), \bar{\mathcal{M}}_j) = \text{Mod}^i(\text{Mod}^j(T, \bar{\mathcal{M}}_j), \bar{\mathcal{M}}_i)$$

*Proof.*

$$\begin{aligned} & \text{Mod}^j(\text{Mod}^i(T, \bar{\mathcal{M}}_i), \bar{\mathcal{M}}_j) \\ &= \{ \text{Mod}^j(S, \bar{\mathcal{M}}_j) \mid S \in \text{Mod}^i(T, \bar{\mathcal{M}}_i) \} \\ &= \left\{ S + \sum_{a=1}^{n_j} \sum_{b=1}^{N_j} \lambda_{a,b} \text{Pad}_a^j(S_b^j(\bar{\mathcal{M}}_j)) \mid \lambda_{a,b} \in \mathbb{F}, S \in \text{Mod}^i(T, \bar{\mathcal{M}}_i) \right\} \\ &= \left\{ T + \sum_{c=1}^{n_i} \sum_{d=1}^{N_i} \lambda_{c,d} \text{Pad}_c^i(S_d^i(\bar{\mathcal{M}}_i)) + \sum_{b=1}^{n_j} \sum_{a=1}^{N_j} \lambda_{a,b} \text{Pad}_a^j(S_b^j(\bar{\mathcal{M}}_j)) \mid \lambda_{a,b}, \lambda_{c,d} \in \mathbb{F} \right\} \\ &= \left\{ S + \sum_{c=1}^{n_i} \sum_{d=1}^{N_i} \lambda_{c,d} \text{Pad}_c^i(S_d^i(\bar{\mathcal{M}}_i)) \mid \lambda_{c,d} \in \mathbb{F}, S \in \text{Mod}^j(T, \bar{\mathcal{M}}_j) \right\} \\ &= \{ \text{Mod}^i(S, \bar{\mathcal{M}}_i) \mid S \in \text{Mod}^j(T, \bar{\mathcal{M}}_j) \} \\ &= \text{Mod}^i(\text{Mod}^j(T, \bar{\mathcal{M}}_j), \bar{\mathcal{M}}_i) \end{aligned}$$

□

The above proof shows that the order of applying modulo operations on a given tensor does not affect the final resultant linear span.

**Observation.** Here, the family  $F \subseteq V_1 \otimes \dots \otimes V_d$ , i.e. all its elements remain of the same order as the original tensor  $T$ . Figure 6 illustrates how one such element from such a family can be generated. [HOW/PROVIDE MORE DETAIL]

**Remark.** As with the adjoining operation for symmetric tensors, generating a family of symmetric tensors modulo symmetric slices also follows immediately from the definitions provided above, but in this case it is the same family of order- $(d-1)$  slices,  $\mathcal{M}$ , being used along every mode. Therefore  $T \text{ mod } \mathcal{M} := T \text{ mod } (\mathcal{M}, \dots, \mathcal{M})$ .

$$T \in \mathbb{F}^4 \otimes \mathbb{F}^3 \otimes \mathbb{F}^2$$




Figure 6: Family generated from a tensor  $T$  by a slice  $A$

Since performing the modulo operation on a tensor gives a family instead of a single output tensor, one can talk about how the maximal or minimal rank of the tensor *family* is affected.

## Combining Adjoining and Modulo

As seen in the previous two subsections, rank information of a tensor does change when adjoining slices, and the following theorem becomes important in Wang and Seigal's 2023 paper [21].

**Theorem 4.2** ([27]). *Suppose  $T \in V := V_1 \otimes \cdots \otimes V_d$  is an order- $d$  tensor, and  $\mathcal{M}_i \subset V_i$  for every  $1 \leq i \leq d$ . Then*

$$\mathbf{Rk}(\text{Adj}(T, \mathcal{M}_1, \dots, \mathcal{M}_d)) \geq \min \mathbf{Rk}(T \bmod (\mathcal{M}_1, \dots, \mathcal{M}_d)) + \sum_{j=1}^d \dim \langle \mathcal{M}_j \rangle$$


*Equality holds if each of the  $\mathcal{M}_i$  consist of only rank-one tensors.*

[ADD PROOF]

### 4.2.4 Cloning Tensors

The idea of cloning a tensor is born out of necessity. When constructing an emulating family (see Section 4.3.2), difficulties arise when working with tensors of a fixed, finite format. Cloning a tensor, as the name suggests, seeks to preserve the properties of the original, while embedding it in a much "larger space".

**Definition 4.10** (Cloning a Tensor). Fix  $d > 0$ . Let  $T \in V := V_1 \otimes \cdots \otimes V_d$  be an order- $d$  tensor.

Figure 7: 2 clone of a tensor  $T$ 

The  $n$ -th cloning function  $C_n : V \rightarrow (\bigoplus_{i=1}^n V_1) \otimes \cdots \otimes (\bigoplus_{i=1}^n V_d)$  can be defined in coordinates in the following manner:

$$C_n(T)(k_{1,j_1}, \dots, k_{d,j_d}) = T(k_1, \dots, k_d).$$

Here, the notation  $k_{i,j_i} := k_{(j_i-1)n+i}$  where  $1 \leq j_i \leq n$ .

**Observation.** Since the definition of cloning depends on a non-zero  $d$ , one can clone tensors of any order, in particular vectors, as well. This leads to the description of the clone of a standard ordered basis vector  $e_i \in V$ , an  $\mathbb{F}$  vector space as:  $C_n(e_i) = \sum_{j=1}^n e_{i,j}$ .

**Lemma 4.2.** *The cloning operation as defined above distributes over the tensor product i.e.  $C_n(A \otimes B) = C_n(A) \otimes C_n(B)$  where  $A, B$  are vectors in an arbitrary vector space.*

*Proof.* Consider two arbitrary vectors,  $A := (a_1, \dots, a_k), B := (b_1, \dots, b_k) \in V$ , an  $\mathbb{F}$  vector space. Then  $A \otimes B$  is an order 2 tensor (a matrix) lying in  $V \otimes V$ , and  $(A \otimes B)(i, j) = a_i b_j$ . Then according to Definition 4.10,

$$C_n(A \otimes B)(k_1, k_2) = (A \otimes B)(k_1 \pmod n, k_2 \pmod n) = A(k_1 \pmod n)B(k_2 \pmod n),$$

here using the regular modulo operation from number theory. This tensor lies in  $(\bigoplus_{i=1}^n V) \otimes (\bigoplus_{i=1}^n V)$ .

Now taking the vectors  $C_n(A), C_n(B) \in (\bigoplus_{i=1}^n V)$ . Here

$$C_n(A) = (\underbrace{a_1, \dots, a_1}_{n \text{ times}}, \dots, \underbrace{a_k, \dots, a_k}_{n \text{ times}}),$$

and likewise for  $C_n(B)$ . Their tensor product is also a matrix lying in  $(\bigoplus_{i=1}^n V) \otimes (\bigoplus_{i=1}^n V)$  where now

$$(C_n(A) \otimes C_n(B))(k_1, k_2) = C_n(A)(k_1)C_n(B)(k_2) = A(k_1 \pmod n)B(k_2 \pmod n).$$

Thus, since each component of both tensors is equal, they are the same tensor.  $\square$

**Remark.** For two tensors  $A, B$  to show that cloning distributes over the tensor product  $A \otimes B$  is a little more involved. However this problem simplifies to asking whether  $C_n(a \otimes b \otimes c) = C_n(a) \otimes C_n(b) \otimes C_n(c)$  where  $a, b, c$  are vectors. This simplified problem can be solved in the same way as the proof above, by working in coordinates and using the modulo function from number theory.

Another description of cloning is by first transforming the tensor under consideration in the following manner. Denote  $\mathbf{1}_n$  as the vector of 1's lying in  $\mathbb{F}^n$ . Then the  $n$ -clone of a simple vector  $v$  can be described as  $(v \otimes \mathbf{1}_n)^\emptyset$  or the flattening of the matrix created by tensoring  $v$  with the vector of ones. Thus another way to look at  $C_n(T)$  is with the following procedure:

$$\begin{aligned} C_n(T) &= C_n\left(\sum_{i=1}^r T_i\right) \\ &= \sum_{i=1}^r C_n(T_i) \\ &= \sum_{i=1}^r C_n(v_1^i \otimes \dots \otimes v_d^i) \\ &= \sum_{i=1}^r C_n(v_1^i) \otimes \dots \otimes C_n(v_d^i) \\ &= \sum_{i=1}^r (v_1^i \otimes \mathbf{1}_n)^\emptyset \otimes \dots \otimes (v_d^i \otimes \mathbf{1}_n)^\emptyset \end{aligned}$$

**Remark.** This property, along with the observation that cloning is a linear operation, allows for the proof that cloning a tensor does preserve its rank.

**Proposition 4.2.**  $\mathbf{Rk}(T) = \mathbf{Rk}(C_n(T))$  for  $C_n$  the cloning operation as defined in Definition 4.10, and  $T \in V := V_1 \otimes \dots \otimes V_d$ .

*Proof.* The proof is straightforward, and is as follows

$\leq$ : Since  $T$  can be seen as a sub-tensor lying in  $C_n(T)$ , this inequality naturally exists. The rank of a subtensor is always at most the rank of the tensor, but never more.

$\geq$ : Since cloning distributes over addition (Lemma 4.2), if one has the rank decomposition  $T = \sum_{i=1}^r T_i$ , then  $C_n(T) = \sum_{i=1}^r C_n(T_i)$  is a decomposition of  $C_n(T)$ . Thus the rank of  $C_n(T)$  is at most that of  $T$ .

$\square$

## 4.3 The Counterexample over $\mathbb{R}$

Wang and Seigal [27] in 2023 published a paper that demonstrated the construction of a distinct counterexample to Comon's conjecture, following Shitov [23]. Their paper first examined the constructions made by Shitov, established new notation and terminology to prove results on rank pre- and post-operations, and finally they used MATLAB to construct a distinct order-6 example with real rank 30913 and real symmetric rank 30914.

As the title of the paper suggests, such a counterexample relies on certain lower bounds and inequalities remaining strict, particularly when making constructions in a field (namely  $\mathbb{R}$ ) that is not algebraically complete.

### 4.3.1 Details from the Paper

The paper in 2023 [27] works over  $\mathbb{R}$ , specifically in  $\mathbb{R}[x, y]$  as the underlying ring of polynomials from which the homogenous polynomials can be selected. For simplicity's sake, there are only 2 variables, corresponding to an underlying real vector space of dimension 2.

The particular tensor under consideration is represented by the homogenous polynomial  $x^d - ky^d$  where  $d > 0$  is a fixed, even integer, and  $k > 0$  (also fixed). Such a homogenous polynomial corresponds to the following tensor

$$T := (e_1)^{\otimes d} - k(e_2)^{\otimes d}, k > 0, d = 2n, n \in \mathbb{Z}_+ \quad (4.1)$$

Visualised as a multiway array, this is a tensor that has non-zero entries at the two extreme "corners" and 0's everywhere else. Such a tensor clearly has rank and symmetric rank as 2.

Since the stated goal is to construct a counterexample to Comon's conjecture, this initial "seed" tensor,  $T$ , is first modified by the using a specific family of slices to generate a linear span of tensors (see Section 4.2.3). Such a family when constructed has elements having minimal rank and symmetric rank. The elements in this family achieving these minimal values do not have to be the same.

The family chosen in this case is

$$\mathcal{M} := \{x^{d-1-i}y^i\}_{i=1}^{d-2} \quad (4.2)$$

This is a collection of monomials of degree  $(d - 1)$  that are of *mixed* form i.e. they are monomials not purely in either variable. When describing such a family in terms of symmetric tensors (see Section 2.3), the family can be written as

$$\mathcal{M} := \left\{ \frac{1}{(d-1)!} \sum_{\sigma \in S_{d-1}} \sigma((e_1)^{\otimes d-1-i} \otimes (e_2)^{\otimes i}) \right\}_{i=1}^{d-2} \quad (4.3)$$

These order- $(d - 1)$  tensors, since they are not formed of a single standard ordered basis vector, they do not affect the initial tensor  $T$ , when performing the operation of generating a family modulo slices:  $T \bmod \mathcal{M}$ . (Note here since they are working with symmetric tensors, it is enough to specify only one family.)

## Establishing A Difference in Rank and Symmetric Rank

Within the constructed family  $T \bmod \mathcal{M}$ , a tensor  $P \in T \bmod \mathcal{M}$  corresponds to the following polynomial

$$P(x, y) := x^d - ky^d + \sum_{i=1}^{d-2} (\alpha_i x + \beta_i y) x^{d-1-i} y^i, \alpha_i, \beta_i \in \mathbb{R} \forall i = 1, \dots, d-2, k > 0 \quad (4.4)$$

First, the symmetric rank decomposition is computed, and shown to be at least 2 or higher.

**Proposition 4.3** ([27]). *A polynomial of the form  $P(x, y)$  (as Equation (4.4)) has symmetric rank at least 2.*

*Proof.* Suppose not. Then one can rewrite  $P(x, y)$  in terms of a linear polynomial raised to the  $d$ th power i.e.

$$P(x, y) = (ax + by)^d$$

for some choices of  $a, b \in \mathbb{R}$ .

Expanding the RHS would then lead to the following

$$\begin{aligned} P(x, y) &= (ax + by)^d \\ &= \sum_{j=0}^d \binom{d}{j} a^{d-j} b^j x^{d-j} y^j. \end{aligned}$$

This leads to the following equality of polynomial expressions:

$$x^d - ky^d + \sum_{i=1}^{d-2} (\alpha_i x + \beta_i y) x^{d-1-i} y^i = \sum_{j=0}^d \binom{d}{j} a^{d-j} b^j x^{d-j} y^j.$$

This leads to a contradiction simply by looking at the coefficient of the  $y^d$  monomial on either side of the equality. One must have  $b^d = -k$  for some choice of  $b \in \mathbb{R}$ . Since such an equation has no real solutions, the initial assumption of being able to write  $P$  in terms of the  $d$ -th power of a single linear form was false, and therefore the Waring rank (equivalent to the symmetric rank) of  $P \geq 2$ .  $\square$

This establishes  $\min_{P \in T \bmod \mathcal{M}} \mathbf{SRk}(P) = 2$ . The next step in the paper is the establishing of the fact that the minimum *rank* element of such a family has rank 1.

**Proposition 4.4** ([27]). *There is a tensor of the form  $P(x, y)$  (see Equation (4.4)) that is an elementary tensor*

*Proof.* Consider the original tensor  $T := x^d - ky^d$  (see Equation (4.1)). This is a tensor of the form as in Equation (4.4). Now summing all the elements in  $\mathcal{M}$  as defined earlier (see Equation (4.2)) gives an order  $d - 1$  tensor with non-zero entries everywhere except for two "corners" — corresponding to  $x^{d-1}$  and  $y^{d-1}$ . These can be added along the  $d$ -th mode to the first slice, and after scaling by a factor of  $-k$ , to the last slice.



Formally the tensor  $T$  is now modified to

$$x^d - ky^d + \underbrace{\sum_{i=1}^{d-2} x^{d-i}y^i}_{\text{modifying the first } d\text{-slice}} + \underbrace{-k \left( \sum_{j=1}^{d-2} x^{d-1-j}y^{j+1} \right)}_{\text{scaling and modifying the last } d\text{-slice}}$$

An important point to note here is that changing the degree of the polynomial changes the order, and since the tensors all lie in  $(\mathbb{R}^2)^{\otimes d}$ , changes affecting the "first" slice have an extra factor of  $x$ , and changes affecting the "last" or second have an extra factor of  $y$ .

The final step of construction is adding the order- $(d-1)$  tensor corresponding to the monomial  $x^{d-2}y$  (scaled by an appropriate  $a_i$ ) to the first  $i$ -slice for all  $d$  modes of the tensor  $T$ . Likewise for the tensor corresponding to the monomial  $xy^{d-2}$  (with appropriate  $b_i$  added to the last  $i$ -slices). The choices of these scaling coefficients are such that the following conditions are satisfied:

$$\left( \sum_{i=1}^d a_i \right) - a_j = 0, \forall j \neq d \quad (4.5)$$

$$\sum_{i=1}^{d-1} a_i = -k \quad (4.6)$$

$$\left( \sum_{i=1}^d b_i \right) - b_j = 0, \forall j \neq d \quad (4.7)$$

$$\sum_{i=1}^{d-1} b_i = 1 \quad (4.8)$$

These choices for  $a_i$  are made because modifying the tensor by adding the scaled tensors modifies only one entry in last  $d$  slice. The last addition of  $x^{d-1}y$  along the  $d$ -mode only affects the first  $d$ -slice. A similar line of reasoning follows the choices for the  $b_i$ . Making such choices is always possible, and non-unique.

Performing these operations results in a tensor still lying in the family  $T \bmod \mathcal{M}$ , now of the form  $(\mathbf{1}_2)^{\otimes d-1} \otimes (1, -k)$ , an elementary order  $d$  tensor. Here again  $\mathbf{1}_n$  denotes the vector of all 1's lying in  $\mathbb{F}^n$   $\square$

**Remark.** It is useful to note that proving such an elementary tensor exists is not a function of the underlying field. (assuming characteristic 0). Such a proof could also be modified for any number of variables (moving from working in  $\mathbb{F}^2$  to  $\mathbb{F}^n$ ), and for tensors of any even order  $d$ .

The following statement holds for the family  $T \bmod \mathcal{M}$

$$\min \mathbf{Rk}(T \bmod \mathcal{M}) < \min \mathbf{SRk}(T \bmod \mathcal{M}) \quad (4.9)$$

## Moving from Families to a Specific Tensor

Now that the existence of a family having elements with differing minimal rank and symmetric rank has been established, the authors construct  $SAdj(T, \mathcal{M})$ , as a symmetric tensor that is a candidate counterexample to Comon's conjecture. The problem

of computing the rank of such a construction now arises, as it relates to the original  $T$  and candidate family  $\mathcal{M}$ .

The solution proposed originally by Shitov [23], is a proposed modification of  $\mathcal{M}$  to a family of *elementary* order  $(d - 1)$  tensors that span the same linear space [WHY? SPECIFIED EARLIER?]. Unfortunately, in the format  $(\mathbb{R}^2)^{\otimes 4}$ , doing so also violates the vital inequality required for rank and symmetric rank of the family  $T \bmod \mathcal{M}$  (see Equation (4.9)).

Such a modification is required for the original computation of rank (see Theorem 4.2) to be an equality (because equality only holds when the family consists of elementary tensors).

[REWORK THIS WHOLE SECTION + PROOF] The following proposition as adapted from [27], shows that for higher order examples (viz. order  $2n$  where  $n > 2$ ), simply modifying Shitov's argument from the order 4 case does not work i.e. it may be possible to modify the family  $\mathcal{M}$  to elementary tensors that preserves the inequality in Equation (4.9).

**Proposition 4.5.** *Consider  $T$  as in Equation (4.1), and  $\mathcal{M}$  as Equation (4.2). Let  $\mathcal{W} \subset (\mathbb{R}^2)^{\otimes d-1}$ , a finite set of elementary symmetric tensors such that  $\mathcal{W}$  spans  $\mathcal{M}$ . Then the argument that  $0 \in T \bmod \mathcal{W}$  when  $d = 2n, n > 2$  from [27] does not hold.*

*Proof.* Consider  $\mathcal{W}$ , a collection of elementary symmetric tensors of appropriate order. Showing  $0 \in T \bmod \mathcal{W}$  is equivalent to asking that  $\mathcal{W}$  spans the set  $\{x^i y^{d-i}\}_{i=0}^{d-1} \supseteq \mathcal{M}$ . This is due to the fact that every slice of  $T$  (seen as an order- $(d-1)$  tensor and therefore represented by a degree- $(d-1)$  polynomial) can then be expressed as lying in the span of  $\mathcal{W}$ . In particular, if one can show that the dimension of  $\text{span } \mathcal{W}$  is  $d$  (the number of degree  $d-1$  monomials in 2 variables), this completes the proof that would show that the inequality in Equation (4.9) is violated.

Suppose the dimension of the span of  $\mathcal{W}$  is less than  $d$ . Consider the tensor represented by  $x^{d-2}y$ . This monomial has rank  $d-1$  since it can be written minimally as sum of  $d-1$  elementary tensors. Thus the dimension of  $\text{span } \mathcal{W}$  is at least  $d-1$ . Since  $\mathcal{W}$  consists of symmetric tensors, its basis vectors must be a linear forms raised to the  $d-1$ th power:  $(\lambda_i x + \mu_i y)^{d-1}$ . These forms must also correspond to elementary tensors, by virtue of being elements of  $\mathcal{W}$ .

Now since this set,  $\{(\lambda_i x + \mu_i y)^{d-1}\}_{i=1}^{d-1}$  is a basis for  $\text{span } \mathcal{W} \supseteq \mathcal{M}$ , one must have that the tensors  $x^{d-2}y$  and  $xy^{d-2}$  can be written as linear combinations of the elements of the set. Doing so imposes conditions on  $\lambda_i, \mu_i > 0$  such that

$$\sum_{i=1}^d \frac{\lambda_i}{\mu_i} = 0 = \sum_{i=1}^d \frac{\mu_i}{\lambda_i}.$$

This set of equalities results in the statement

$$1 = \frac{\lambda_1}{\mu_1} \frac{\mu_1}{\lambda_1} = \left( \sum_{i=2}^d \frac{\lambda_i}{\mu_i} \right) \left( \sum_{i=2}^d \frac{\mu_i}{\lambda_i} \right) = (d-1) + \sum_{i=2}^d \sum_{j \neq i} \frac{\lambda_i \mu_j}{\mu_i \lambda_j}.$$

If one can show that

$$\sum_{i=2}^d \sum_{j < i} t_{ij} + t_{ij}^{-1} + d - 2 \neq 0,$$

where  $t_{ij} := \frac{\lambda_i \mu_j}{\mu_i \lambda_j}$ , then the statement arrived at is false. This would mean that the span  $\mathcal{W}$  being of a dimension less than  $d$  cannot be true, since this leads to a contradictory statement.

However, the range of this function is the entirety of  $\mathbb{R}$  whenever dealing with a value of  $d > 4$  since the function generated in terms of  $\{t_{ij}, t_{ij}^{-1}\}$  is no longer univariate.

Therefore there are choices of  $\lambda_i, \mu_i$  that can be made such that the span of  $\mathcal{W}$  has a dimension less than  $d$ . Therefore it cannot be guaranteed, by the above modified argument, that  $0 \in T \bmod \mathcal{W}$ .

□

### 4.3.2 Emulating Family Construction

[REWRITE/REJUSTIFY] While for the order-4 case from Shitov, the issue of Equation (4.9) being violated when modifying the family  $\mathcal{M}$  arose, as seen in the previous proposition, such issues do not arise when working with the order-6 or higher cases. However, the authors of [27] continue to use the cloning methodology later in their paper, and doing so ensures that there are no further complications [SUCH AS?] that could arise when modifying the initial family of slices  $\mathcal{M}$ .

As demonstrated in Section 4.2.4, since cloning preserves rank, then inequality is preserved after taking the  $k$ -th clones, and transforms into

$$\min \mathbf{Rk} C_k(T) \bmod C_k(\mathcal{M}) < \min \mathbf{SRk} C_k(T) \bmod C_k(\mathcal{M}) \quad (4.10)$$

In this situation, modifying  $C_k(M)$  (a family of order- $(d-1)$  tensors lying in  $C_k(V^{\otimes d-1})$ ) into  $\mathcal{W}$  (a family of order- $(d-1)$  elementary tensors) can occur successfully. This results in working with a symmetric tensor  $SAdj(C_k(T), \mathcal{W})$  - the final symmetric tensor having a definite distinction between rank and symmetric rank. [HOW DOES ONE CHOOSE  $K$ ]

It is past this point that both Shitov [23], and Wang and Seigal [27] go forward with the construction of a family  $\mathcal{W}$  to replace  $C_k(\mathcal{M})$  that can be described algorithmically, but the justification for why such choices are made remains opaque. [RETRY TO FIGURE OUT WHAT THE ALGORITHM IS]

The later paper [27] constructs this family for their candidate counterexample of order 6 as described below. In personal correspondence, the authors indicated that this falls in step with the construction in the preprint [23]:

They begin with vectors lying in  $\mathbb{R}^{2n}$  of the form  $a_i$ , where  $i \in [n]$  and each vector is such that  $a_i(2i) = a_i(2i-1) = 1$  and all the other entries are 0. Then the new vectors constructed lie in  $\mathbb{R}^{4n}$  and are one of the following forms (where the  $|$  indicates the distinction between the first  $2n$  entries and the last  $2n$  entries.)

1.  $(a_{i_1} + a_{i_2} + a_{i_3} + a_{i_4} \mid a_{k_1})$
2.  $(a_{i_1} + a_{i_2} + a_{i_3} + a_{i_4} \mid 0)$
3.  $(a_{i_1} + a_{i_2} + a_{i_3} \mid \frac{n-2}{n-1} a_{k_1})$
4.  $(a_{i_1} + a_{i_2} + a_{i_3} \mid \frac{n-3}{n-4} a_{k_1})$
5.  $(a_{i_1} + a_{i_2} + a_{i_3} \mid a_{k_1} + a_{k_2})$

6.  $(a_{i_1} + a_{i_2} + a_{i_3} \mid 0)$
7.  $(a_{i_1} + a_{i_2} \mid \frac{(n-2)^2}{(n-3)(n-1)}a_{k_1})$
8.  $(a_{i_1} + a_{i_2} \mid \frac{n-2}{n-4}a_{k_1})$
9.  $(a_{i_1} + a_{i_2} \mid \frac{n-2}{n-3}(a_{k_1} + a_{k_2}))$
10.  $(a_{i_1} + a_{i_2} \mid 0)$
11.  $(a_{i_1} \mid \frac{n-2}{n-3}a_{k_1})$
12.  $(a_{i_1} \mid \frac{n-1}{n-4}a_{k_1})$
13.  $(a_{i_1} \mid \frac{n-1}{n-3}(a_{k_1} + a_{k_2}))$
14.  $(a_{i_1} \mid 0)$
15.  $(0 \mid a_{k_1})$
16.  $(0 \mid a_{k_1} + a_{k_2})$

Here  $1 \leq i_1 < i_2 < i_3 < i_4 \leq n$  and  $1 \leq k_1 < k_2 \leq n$ . The collection of all the 16 vectors in  $\mathbb{R}^{4n}$  described above, for all choices of indices  $i_1, i_2, i_3, i_4$  and  $k_1, k_2$  forms the set  $\mathcal{W}_1$ . This is only half of their construction.

Under the permutation operation

$$\pi : (i_1, i_2, \dots, i_{2n}, k_1, k_2, \dots, k_{2n}) \mapsto (k_2, \dots, k_{2n}, k_1, i_2, \dots, i_{2n}, i_1),$$

the same set  $\mathcal{W}_1$  can be transformed into a new set  $\mathcal{W}_2$ . The final emulating family they use for their order-6 counterexample is  $\mathcal{W} := \mathcal{W}_1 \cup \mathcal{W}_2$

This emulating family proves to be the last piece in their construction of  $SAdj(C_k(T), \mathcal{W})$ , an order-6 symmetric tensor having real symmetric rank 30914 and real rank 30913.

Extending this counterexample to higher orders requires a clearer understanding of the choices made in this construction of  $\mathcal{W}$ . [TOOTHLESS]

# Chapter 5

## Conclusion

While this thesis only condenses the literature surrounding Comon’s conjecture and attempts to generate counterexamples to it, the fact that a computed, real counterexample does exist provides some hope for a conclusive answer in the future. As mentioned in Chapter 4, since the real counterexample requires the discrepancy from Equation (4.9), an equivalent condition in the complex case seems to be the pathway forward.

To be more precise: if a symmetric, complex-valued, order- $d$  tensor  $T$  exists, along with a family of symmetric, complex-valued, order- $(d - 1)$  tensors  $\mathcal{M}$ , such that the statement

$$\min \mathbf{Rk}(T \bmod \mathcal{M}) < \min \mathbf{SRk}(T \bmod \mathcal{M})$$

still holds, then a viable candidate counterexample can be constructed, paralleling the constructions in [23, 27], since the only part of these constructions dependent on the underlying field is the choice of candidate real tensor  $T_{\mathbb{R}}$  represented by the homogeneous even order  $d$  polynomial  $x^d - ky^d$ .

### 5.1 Future Work

The work in this thesis establishes a baseline for exploring constructions of candidate counterexamples to Comon’s Conjecture.

- One way this work can be extended is to focus more on a geometric description of the constructions made in this paper, particularly for Equation (4.9). In doing so, the hope is that one would uncover geometric conditions that indicate a candidate counterexample to the conjecture, and these are hopefully independent of the base field chosen. [ILLUSTRATE/ELABORATE?]
- A more explicit understanding of the choices of vectors made in Section 4.3.2. The choices made so far stem from [23], and while they have proved to be effective, have no indication as to *why* they are. Studying the construction of such a family more carefully might indicate algebraic or geometric conditions that are being met by the collection of vectors or slices, allowing for the existence of an emulating family of elementary tensors. [MAY NEED TO REMOVE DEPENDING ON HOW THE REST OF THIS GOES]

- Comon's conjecture as originally stated in [19] was only for  $\mathbb{R}$ - or  $\mathbb{C}$ -valued tensors. A conclusive understanding of whether Comon's conjecture can be extended to finite fields, or if such an extension is non-trivial can be one path forward. [MAYBE LOOK AT ZHENG 2020]
- For the real, even-order case, the development of explicit algorithms to compute rank decomposition of length shorter than the symmetric rank for a given symmetric tensor, in cases when such a discrepancy can be shown to exist. [MAY NEED TO REMOVE]
- Adjusting the work in this paper to understand the smallest possible cloning operation required to satisfy the conditions of the emulating family could be another potential extension of this work. The paper from Borovik et al. in 2025 [5], while not explicitly related to Comon's conjecture, does establish an upper bound for how large a cloning operation may be required. This paper builds on [24], a preprint from Shitov more in line with constructing a counterexample to Strassen's conjecture, but relies on similar methodology.

# Bibliography

- [1] Boris Alexeev, Michael A. Forbes, and Jacob Tsimerman, *Tensor Rank: Some Lower and Upper Bounds*, 2011 IEEE 26th Annual Conference on Computational Complexity, June 2011, pp. 283–291.
- [2] E. Ballico and A. Bernardi, *Tensor ranks on tangent developable of Segre varieties*, Linear and Multilinear Algebra **61** (July 2013), no. 7, 881–894 (en).
- [3] Edoardo Ballico, Alessandra Bernardi, Luca Chiantini, and Elena Guardo, *Bounds on the tensor rank*, Annali di Matematica Pura ed Applicata (1923 -) **197** (December 2018), no. 6, 1771–1785 (en).
- [4] Alessandra Bernardi, Enrico Carlini, Maria Virginia Catalisano, Alessandro Gimigliano, and Alessandro Oneto, *The Hitchhiker Guide to: Secant Varieties and Tensor Decomposition*, Mathematics **6** (December 2018), no. 12, 314 (en).
- [5] Viktoriia Borovik, Cosimo Flavi, Paweł Pielasa, Anatoli Shatsila, and Jeyoung Song, *On the construction of a counterexample to Strassen’s rank additivity conjecture*, arXiv, 2025 (en). arXiv:2507.17890 [math].
- [6] Maria Chiara Brambilla and Giorgio Ottaviani, *On the Alexander–Hirschowitz theorem*, Journal of Pure and Applied Algebra **212** (May 2008), no. 5, 1229–1251 (en).
- [7] Weronika Buczyńska and Jarosław Buczyński, *Secant varieties to high degree Veronese reembeddings, catalecticant matrices and smoothable Gorenstein schemes*, Journal of Algebraic Geometry **23** (September 2013), no. 1, 63–90 (en).
- [8] Jarosław Buczyński and J.M. Landsberg, *Ranks of tensors and a generalization of secant varieties*, Linear Algebra and its Applications **438** (January 2013), no. 2, 668–689 (en).
- [9] Sean Carroll, *Spacetime and geometry: an introduction to general relativity*, Cambridge university press, Cambridge, 2019 (eng).
- [10] Alex Casarotti, Alex Massarenti, and Massimiliano Mella, *On Comon’s and Strassen’s Conjectures*, Mathematics **6** (October 2018), no. 11, 217 (en).
- [11] Luca Chiantini, Giorgio Ottaviani, and Nick Vannieuwenhoven, *On generic identifiability of symmetric tensors of subgeneric rank*, Transactions of the American Mathematical Society **369** (November 2016), no. 6, 4021–4042 (en).
- [12] ———, *Effective Criteria for Specific Identifiability of Tensors and Forms*, SIAM Journal on Matrix Analysis and Applications **38** (January 2017), no. 2, 656–681 (en).
- [13] Pierre Comon, Gene Golub, Lek-Heng Lim, and Bernard Mourrain, *Symmetric Tensors and Symmetric Tensor Rank*, SIAM Journal on Matrix Analysis and Applications **30** (January 2008), no. 3, 1254–1279 (en).
- [14] Jan Draisma, *Erratum: A Counterexample to Comon’s Conjecture*, SIAM Journal on Applied Algebra and Geometry **8** (March 2024), no. 1, 225–225 (en).
- [15] Shmuel Friedland, *Remarks on the Symmetric Rank of Symmetric Tensors*, SIAM Journal on Matrix Analysis and Applications **37** (January 2016), no. 1, 320–337 (en).
- [16] Tamara G. Kolda and Brett W. Bader, *Tensor Decompositions and Applications*, SIAM Review **51** (August 2009), no. 3, 455–500 (en).

- [17] J.M. Landsberg and Mateusz Michałek, *Abelian tensors*, Journal de Mathématiques Pures et Appliquées **108** (September 2017), no. 3, 333–371 (en).
- [18] Joseph M. Landsberg, *Tensors: geometry and applications*, Graduate studies in mathematics, American mathematical society, Providence (R.I.), 2012 (eng).
- [19] Luke Oeding, *Report on "Geometry and representation theory of tensors for computer science, statistics and other areas."*, arXiv, 2008 (en). arXiv:0810.3940 [math].
- [20] Yannis Panagakis, Jean Kossaifi, Grigorios G. Chrysos, James Oldfield, Mihalios A. Nicolaou, Anima Anandkumar, and Stefanos Zafeiriou, *Tensor Methods in Computer Vision and Deep Learning*, Proceedings of the IEEE **109** (May 2021), no. 5, 863–890 (en).
- [21] Anna Seigal, *Ranks and symmetric ranks of cubic surfaces*, Journal of Symbolic Computation **101** (November 2020), 304–317 (en).
- [22] Yaroslav Shitov, *A Counterexample to Comon's Conjecture*, SIAM Journal on Applied Algebra and Geometry **2** (January 2018), no. 3, 428–443 (en).
- [23] ———, *Comon's conjecture over the reals* (2020), available at <https://vixra.org/abs/2009.0134> (en).
- [24] ———, *Higher rank substitutions for tensor decompositions. I. Direct sum conjectures* (2024), available at <https://rgdoi.net/10.13140/RG.2.2.21768.43528> (en).
- [25] ———, *Higher rank substitutions for tensor decompositions. II. Comon's conjecture* (2024), available at <https://rgdoi.net/10.13140/RG.2.2.17574.13125> (en).
- [26] Ravi Vakil, *The rising sea: foundations of algebraic geometry*, Princeton University Press, Princeton, 2025 (eng).
- [27] Kexin Wang and Anna Seigal, *Lower bounds on the rank and symmetric rank of real tensors*, Journal of Symbolic Computation **118** (September 2023), 69–92 (en).
- [28] Xinzheng Zhang, Zheng-Hai Huang, and Liqun Qi, *Comon's Conjecture, Rank Decomposition, and Symmetric Rank Decomposition of Symmetric Tensors*, SIAM Journal on Matrix Analysis and Applications **37** (January 2016), no. 4, 1719–1728 (en).



**Department of Mathematics**  
Celestijnenlaan 200B  
3001 LEUVEN (HEVERLEE), BELGIË  
tel. + 32 16 32 70 07  
fax + 32 16 32 79 98  
wis.kuleuven.be

