

# LHCb MC group meeting: $B^0 \rightarrow K^{*0} \mu \mu$

Oliver Dahme

University of Zurich

*o.dahme@cern.ch*

October 4, 2017

$$B^0 \rightarrow K^{*0} \mu \mu$$

- Decay is a FCNC
- four charged particles in final state:
- $K^+$  and  $\pi^-$  from the  $K^{*0}$  decay
- two leptons from loop or box diagrams:




Figure: Feynman diagrams for decay  $B_d^0 \rightarrow \mu^+ \mu^-$  at lowest order

$$B^0 \rightarrow K^{*0} \mu \mu$$

three angles define the kinematics of the decay:

- ThetaK or  $\theta_K$
- ThetaL or  $\theta_L$
- Phi or  $\phi$





Figure: kinematic variables of the decay  $B^0 \rightarrow K^{*0} \mu \mu$

# classification with Machine Learning (BDT)


- to eliminate combinatorial background, Boost Decision Trees are used as a black box to classify data into signal and background.
- They assign a 'probability' to each event
- sk bdt from the scikit-learn package is used, and uBoost from the hep ml package
- average of both gives the BDT response
- training has been performed on the Linux Cluster in Zurich

# Region to train

- you want the bdt to train in the background region
- J/Psi has two resonance which will be cut off



(a) Before Cut



(b) After Cut

# training of the BDTs

8 train parameters:


- B0 P
- B0 PT
- B0 ENDVERTEX CHI2
- B0 IP OWNPV
- B0 IPCHI2 OWNPV
- B0 FD OWNPV
- B0 FDCHI2 OWNPV
- B0 relinfo VTXISOBDTHARDFIRSTVALUE

4 uniform parameters:


- B0 M
- B0 ThetaK
- B0 ThetaL
- B0 Phi

uniform parameters have to be uncorrelated since we know that from physics. But BDT might find a correlation which is bad.

# BDT correlation check




# BDT correlation check





# BDT correlation check



# BDT correlation check

