



# Numerical Methods

## Exercise Sheet 6

HS 16  
M. Chrzaszcz  
D. van Dyk

I. Bezshyiko, A. Pattori

<http://www.physik.uzh.ch/en/teaching/PHY233/>

HS2016.html

Issued: 16.11.2016

Due: 23.11.2016

### Exercise 1: Explicit Runge-Kutta methods (120 Pts.)

In this exercise you will be asked to find numerically the solution to the following ODE:

$$\frac{dy}{dx} = f(x, y) = 8(1 - 2x)y, \quad y(0) = e^{-2}. \quad (1)$$

You will do it using the so-called Bogacki-Shampine method, which makes use of the following two Butcher tableaus:

Tableau A (2 <sup>nd</sup> order):	Tableau B (3 <sup>rd</sup> order):																																													
<table style="border-collapse: collapse; margin: auto;"> <tr><td style="border-right: 1px solid black; padding: 5px 10px;">0</td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"><math>\frac{1}{2}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{2}</math></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"><math>\frac{3}{4}</math></td><td style="padding: 5px 10px;">0</td><td style="padding: 5px 10px;"><math>\frac{3}{4}</math></td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;">1</td><td style="padding: 5px 10px;"><math>\frac{2}{9}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{3}</math></td><td style="padding: 5px 10px;"><math>\frac{4}{9}</math></td></tr> <tr style="border-top: 1px solid black;"><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;"><math>\frac{2}{9}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{3}</math></td><td style="padding: 5px 10px;"><math>\frac{4}{9}</math>   0</td></tr> </table>	0				$\frac{1}{2}$	$\frac{1}{2}$			$\frac{3}{4}$	0	$\frac{3}{4}$		1	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$		$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$ 0	<table style="border-collapse: collapse; margin: auto;"> <tr><td style="border-right: 1px solid black; padding: 5px 10px;">0</td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"><math>\frac{1}{2}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{2}</math></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"><math>\frac{3}{4}</math></td><td style="padding: 5px 10px;">0</td><td style="padding: 5px 10px;"><math>\frac{3}{4}</math></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;">1</td><td style="padding: 5px 10px;"><math>\frac{2}{9}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{3}</math></td><td style="padding: 5px 10px;"><math>\frac{4}{9}</math></td><td style="padding: 5px 10px;"></td></tr> <tr style="border-top: 1px solid black;"><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;"><math>\frac{7}{24}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{4}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{3}</math></td><td style="padding: 5px 10px;"><math>\frac{1}{8}</math></td></tr> </table>	0					$\frac{1}{2}$	$\frac{1}{2}$				$\frac{3}{4}$	0	$\frac{3}{4}$			1	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$			$\frac{7}{24}$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{8}$
0																																														
$\frac{1}{2}$	$\frac{1}{2}$																																													
$\frac{3}{4}$	0	$\frac{3}{4}$																																												
1	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$																																											
	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$ 0																																											
0																																														
$\frac{1}{2}$	$\frac{1}{2}$																																													
$\frac{3}{4}$	0	$\frac{3}{4}$																																												
1	$\frac{2}{9}$	$\frac{1}{3}$	$\frac{4}{9}$																																											
	$\frac{7}{24}$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{8}$																																										

Tableau B will be used to extract  $y_{n+1} \equiv y_{n+1}^B$ , while tableau A will be used to estimate the error on  $y_{n+1}$  through the difference  $\delta y_{n+1} = y_{n+1}^B - y_{n+1}^A$ .

Please notice that tableau A and B differ only by the last row.

- a) Implement the two tableaus A and B in a function

$$\Phi(x_n, y_n, h) \quad (2)$$

which returns:

- The point  $\{x_{n+1}, y_{n+1}\}^B$ , computed using the tableau B (3<sup>rd</sup> order).
- The value  $\delta y_{n+1} = y_{n+1}^B - y_{n+1}^A$ , defined as the difference in the prediction for  $y_{n+1}$  between the tableaus B and A.

(20 points)

Optional: make the ODE function  $f(x, y)$  and the Butcher tableaus themselves input parameters of  $\Phi$ , in order to give more flexibility to your code.

(20 points)

- b) As explained,  $\delta y_{n+1}$  gives you an estimate of the error with which  $y_{n+1}^B$  is extracted. Compute  $\delta y_1(h)$  for  $\{x_0, y_0\} = \{0, e^{-2}\}$  and different values of  $h$  (e.g.  $h = 0.01, 0.02, \dots, 0.1$ ). Use these results to show that your error indeed scales as  $h^3$ , as expected by the order of the method.

(20 points)

c) *Fixed step size method.* We want to solve eq. (1) in the range  $x \in [0, 2]$ . We will first do it using a fixed step size. Set  $h = 0.1$  and solve ODE (1) using your function  $\Phi(x_n, y_n, h)$ .

Plot the result and the errors  $\delta y_{n+1}$  of each step (you may do it in two different plots). (20 points)

d) *Adaptive step size method.* Suppose we want to impose  $\delta y_n < 10^{-4}$  at every step. A possibility could be to reduce uniformly the step size  $h$  until we reach such goal. This is not an efficient solution and sometimes it is overwhelmingly time-consuming.

A more interesting solution is the adaptive step size method, in which you adjust  $h$  at each step accordingly to the estimated error  $\delta y_{n+1}$ .

Implement a simple adaptive step size method in this way:

1- Start with  $h = 0.1$

2- At every step  $n \rightarrow n + 1$ , evaluate  $\delta y_{n+1}$ :

- If  $\delta y_{n+1} \in (10^{-5}, 10^{-4})$ , keep the result and go to next step
- If  $\delta y_{n+1} > 10^{-4}$ , reduce  $h$  accordingly:

$$h \rightarrow 0.9 \sqrt[3]{\frac{10^{-4}}{\delta y_{n+1}}} h$$

then compute again step  $n \rightarrow n + 1$

- If  $\delta y_{n+1} < 10^{-5}$ , increase  $h$  accordingly:

$$h \rightarrow 1.1 \sqrt[3]{\frac{10^{-5}}{\delta y_{n+1}}} h$$

and go to the next step

Solve ODE (1) for  $x \in [0, 2]$  with this method. (20 points)

Compare the result and the errors  $\delta y_n$  with what obtained in task (1c). (10 points)

Also, keep tracks of the adapted  $h_n$  of each step, plot and comment them. (10 points)

**Maximum number of points for mandatory tasks on 23.11 : 100**

**Maximum possible number of points for tasks on 23.11 : 120**