

Exoplanet Atmospheres

O. Daniel O'Carroll

22337259

oocarrol@tcd.ie

June 3, 2025(45²)

CONTENTS

1	ELLIPTICAL ORBITS	2
1.1	KEPLER'S LAWS	2
1.2	LAGRANGE POINTS	2
2	KEYWORDS & DEFINITIONS	3
2.1	HOT JUPITERS	3
2.2	SUPER-EARTHS/SMALL NEPTUNES	3
2.3	RADIAL VELOCITY SEMI-AMPLITUDE	3
2.4	EMISSION SPECTROSCOPY	3
2.5	TRANSMISSION SPECTROSCOPY	3
2.6	HIGH VS LOW RESOLUTION SPECTROSCOPY	4
3	CODE AND COMPUTER MODELLING	5
3.1	CROSS CORRELATION FUNCTIONS	5
3.2	KERNEL	5
3.3	RETRIEVALS	5
3.4	PETITRADTRANS PYTHON PACKAGE	5
3.5	BEER'S LAW	6
3.6	BEER'S LAW APPLIED TO PLANETARY TRANSITS	6
	REFERENCES	7

1 ELLIPTICAL ORBITS

Definiton 1.1: Kepler's Laws

Kepler's First Law: Equation of Elliptical Orbits

$$r(t) = \frac{a(1 - e^2)}{1 + e \cos(\theta(t))}. \quad (1.1.1)$$

with a being the semi-major axis $l = \frac{b^2}{a}$ the semi-latus rectum

Kepler's Second Law: Area/Angular Momentum Conservation

$$A = \frac{1}{2m} \int_{t_0}^{t_1} J(t) dt \quad (1.1.2)$$

with $J(t) = mr^2\dot{\theta}$ a constant.

Kepler's Third Law: Orbital Period

$$P^2 = \frac{4\pi^2}{G^2(M_\star + M_p)} a^3 \quad (1.1.3)$$

Definiton 1.2: Lagrange Points

In the orbit of two bodies where one is much more massive than the other, the *Lagrange points* are the points in the system which have the same orbital period as the smaller object.




Figure 1.2.1: Diagram of the JWST in the L2 point around Earth.

2 KEYWORDS & DEFINITIONS

Definiton 2.1: Hot Jupiters

Jupiter sized planet close to its host star.

Definiton 2.2: Super-Earths/Small Neptunes

Large Earth-sized planet or small Neptune sized planet.

Definiton 2.3: Radial Velocity Semi-Amplitude

$$K = v \sin(i). \quad (2.3.1)$$

Definiton 2.4: Emission Spectroscopy

Measuring the spectrum of the planets atmosphere using its own radiation.

Definiton 2.5: Transmission Spectroscopy

Measuring the spectrum of a planets atmosphere using the starlight of its host star when it transits.




Figure 2.5.1: Planet in transit and its absorbtion spectrum.

Definiton 2.6: High vs Low Resolution Spectroscopy

The formula for the resolution (or resolving power) of a spectrometer is

$$R = \frac{\lambda}{\Delta\lambda}. \quad (2.6.1)$$

High resolution is when $R \propto 10^6$, this is most ground-based telescopes such as the VST (Very Large Telescope). The JWST is a low resolution telescope, $R \propto 10^2$.

High resolution telescopes are more complex but produce spectra with more peaks and as such are better at identifying specific molecules. They are more limited in their range as the light is diffracted more, the uncertainty in the wavelengths become larger and more importantly some of the modes start to overlap. The fix for this is to look at a narrower band of wavelengths.

Low resolution spectroscopy on the other hand can have a broader band of wavelengths to detect but can't resolve individual peaks.

3 CODE AND COMPUTER MODELLING

Definiton 3.1: Cross Correlation Functions

When a spectrum is measured a cross correlation functions is used by taking a model of a spectrum and integrating it across the data to find the best fit.

Definiton 3.2: Kernel

A kernel will be convolved with the model of the cross correlation function to account for things like the light and dark side of the planet, red and blue shifting, and atmospheric broadening.

Definiton 3.3: Retrievals

Retrievals are when the 10 or so model parameters which are statistically correlated are determined modelled and analysed, usually displayed as a 10 x 10 corner plot.

Definiton 3.4: petitRADTRANS Python Package

Python package that can generate example spectra and create retrievals for high and low resolution spectra.




Figure 3.4.1: Tutorial spectrum from petitRADTRANS [3].

Definiton 3.5: Beer's Law

Beer's law describes the intensity fall off due to the increase in optical path length. Below is the equation of electric flux varying with distance assuming that the light is being observed in the x -direction.

$$\frac{d\phi_e}{dx} = -\mu(x)\phi_e(x), \quad (3.5.1)$$

where $\mu(z)$ is the attenuation coefficient [4].

Definiton 3.6: Beer's Law Applied to Planetary Transits

$$R(\lambda) = R_0 + H \left[\gamma + \ln \left(\frac{P_0}{mg} \sqrt{\frac{2\pi R_0}{H}} \right) \right] + H \ln \sum_j \chi_j \sigma_j(\lambda), \quad (3.6.1)$$

where H is the *scale height*, R_0 is the *reference radius*, σ_j is the absorption cross-section as a function of wavelength for species j , χ_j is the volume mixing ratio of species j , P_0 is the pressure at the reference radius, m is the mean molecular weight of the atmosphere, g is the *surface gravity*, and γ is a dimensionless constant ≈ 0.56 .

$$H = \frac{kT}{mg}. \quad (3.6.2)$$

REFERENCES

- [1] Transits and Occultations - Joshua N Winn
- [2] Exoplanetary Atmospheres: Key Insights, Challenges, and Prospects - Nikku Madhusudhan
- [3] petitRADTRANS Python package website & documentation.
- [4] The theory of transmission spectra revisited: a semi-analytical method for interpreting WFC3 data and an unresolved challenge - Kevin Heng, Daniel Kitzmann