

Active Winter Action

$$\begin{cases} \mathrm{d}X_t^i &= \mathrm{Pev}(\theta_t^i)\mathrm{d}t + \sqrt{2\sigma_x}\mathrm{d}W_t^i \\ \mathrm{d}\theta_t^i &= \chi F\left(\left(X_t^i, \theta_t^i\right), \mu_t^N\right)\mathrm{d}t + \sqrt{2}\mathrm{d}B_t^i \end{cases}$$

- Interaction strength $\chi > 0$

- Empirical measure: $\mu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{(X_t^i, \theta_t^i)}$, McKean-Vlasov

- Famous interacting active Brownian particle model:
Vicsek model (Phys. Rev. Lett., 1995) for flocking birds, 6000+ citations



Active Vlasov interaction

$$\begin{cases} dX_t^i &= \text{Pev}(\theta_t^i)dt + \sqrt{2\sigma_x}dW_t^i \\ d\theta_t^i &= \chi F\left((X_t^i, \theta_t^i), \mu_t^N\right)dt + \sqrt{2}dB_t^i \end{cases}$$


- Interaction strength $\chi > 0$
- Empirical measure: $\mu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{(X_t^i, \theta_t^i)}$, McKean-Vlasov
- Famous interacting active Brownian particle model: Vicsek model (Phys. Rev. Lett., 1995) for flocking birds, 6000+ citations



Active Vlasov interaction

$$\begin{cases} dX_t^i &= \text{Pev}(\theta_t^i)dt + \sqrt{2\sigma_x}dW_t^i \\ d\theta_t^i &= \chi F\left((X_t^i, \theta_t^i), \mu_t^N\right)dt + \sqrt{2}dB_t^i \end{cases}$$

- Interaction strength $\chi > 0$
- Empirical measure: $\mu_t^N = \frac{1}{N} \sum_{i=1}^N \delta_{(X_t^i, \theta_t^i)}$, McKean-Vlasov
- Famous interacting active Brownian particle model: Vicsek model (Phys. Rev. Lett., 1995) for flocking birds, 6000+ citations



Vicsek model simulation © F. Turci 2020