

Trigonometry

Contents

1. Trigonometry	2
1.1. Domain/Range of trig functions	2
1.2. Trigonometric Formulas	2
1.3. Value of θ	3
2. Properties of a triangles	4
2.1. Law of Sines:	4
2.2. Law of Cosines:	4
2.3. Projection Rule:	4
2.4. Napier's analogy:	4
3. Inverse Trigonometry	5
3.1. Domain/Range of inverse trig functions	5
3.2. Inverse Trigonometric Formulas	5

1. Trigonometry

1.1. Domain/Range of trig functions

Identities	Domain	Range
$\sin \theta$	$\mathbb{R}$	$[-1, 1]$
$\cos \theta$	$\mathbb{R}$	$[-1, 1]$
$\tan \theta$	$\mathbb{R} - \{(2n + 1)\frac{\pi}{2}\}$	$\mathbb{R}$
$\cot \theta$	$\mathbb{R} - \{n\pi\}$	$\mathbb{R}$
$\csc \theta$	$\mathbb{R} - \{n\pi\}$	$\mathbb{R} - (-1, 1)$
$\sec \theta$	$\mathbb{R} - \{(2n + 1)\frac{\pi}{2}\}$	$\mathbb{R} - (-1, 1)$

1.2. Trigonometric Formulas

1. $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$
2. $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$
3. $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
4. $\cot(A \pm B) = \frac{\cot A \cot B \mp 1}{\cot B \pm \cot A}$
5. $\sin(A + B) + \sin(A - B) = 2 \sin A \cos B$
6. $\sin(A + B) - \sin(A - B) = 2 \cos A \sin B$
7. $\cos(A + B) + \cos(A - B) = 2 \cos A \cos B$
8. $\cos(A + B) - \cos(A - B) = -2 \sin A \sin B$
9. $\sin(A + B) \cdot \sin(A - B) = \sin^2 A - \sin^2 B = \cos^2 B - \cos^2 A$
10. $\cos(A + B) \cdot \cos(A - B) = \cos^2 A - \sin^2 B = \cos^2 B - \sin^2 A$
11. $\sin C + \sin D = 2 \sin\left(\frac{C + D}{2}\right) \cdot \cos\left(\frac{C - D}{2}\right)$
12. $\sin C - \sin D = 2 \cos\left(\frac{C + D}{2}\right) \cdot \sin\left(\frac{C - D}{2}\right)$
13. $\cos C + \cos D = 2 \cos\left(\frac{C + D}{2}\right) \cdot \cos\left(\frac{C - D}{2}\right)$

$$14. \cos C - \cos D = 2 \sin\left(\frac{C+D}{2}\right) \cdot \sin\left(\frac{D-C}{2}\right)$$

$$15. \sin 2A = 2 \sin A \cos A = \frac{2 \tan A}{1 + \tan^2 A}$$

$$16. \cos 2A = \cos^2 A - \sin^2 A = 1 - 2 \sin^2 A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$$

$$17. \tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

$$18. \cot 2A = \frac{\cot^2 A - 1}{2 \cot A}$$

$$19. \sin 3A = 3 \sin A - 4 \sin^3 A$$

$$20. \cos 3A = 4 \cos^3 A - 3 \cos A$$

$$21. \tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$$

$$22. \cot 3A = \frac{\cot^3 A - 3 \cot A}{3 \cot^2 A - 1}$$

$$23. \tan^2 A = \frac{1 - \cos 2A}{1 + \cos 2A}$$

1.3. Value of θ

$$1. \sin \theta = \sin \alpha \implies \theta = n\pi + (-1)^n \alpha$$

$$2. \cos \theta = \cos \alpha \implies \theta = 2n\pi \pm \alpha$$

$$3. \tan \theta = \tan \alpha \implies \theta = n\pi + \alpha$$

$$4. \sin \theta = 0 \implies \theta = n\pi$$


$$5. \cos \theta = 0 \implies \theta = n\pi + \frac{\pi}{2} = (2n+1)\frac{\pi}{2}$$

$$6. \tan \theta = 0 \implies \theta = n\pi$$

$$7. \sin \theta = 1 \implies \theta = 2n\pi + \frac{\pi}{2} = (4n+1)\frac{\pi}{2}$$

$$8. \cos \theta = 1 \implies \theta = 2n\pi$$

2. Properties of a triangles



2.1. Law of Sines:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

2.2. Law of Cosines:

$$1. \ a^2 = b^2 + c^2 - 2bc \cos A \implies \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$2. \ b^2 = a^2 + c^2 - 2ac \cos B \implies \cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$3. \ c^2 = a^2 + b^2 - 2ab \cos C \implies \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

2.3. Projection Rule:

$$1. \ a = b \cos C + c \cos B$$

$$2. \ b = c \cos A + a \cos C$$

$$3. \ c = a \cos B + b \cos A$$

2.4. Napier's analogy:

$$1. \ \tan\left(\frac{B-C}{2}\right) = \frac{b-c}{b+c} \cot \frac{A}{2}$$

$$2. \tan\left(\frac{C-A}{2}\right) = \frac{c-a}{c+a} \cot \frac{B}{2}$$

$$3. \tan\left(\frac{A-B}{2}\right) = \frac{a-b}{a+b} \cot \frac{C}{2}$$

3. Inverse Trigonometry

3.1. Domain/Range of inverse trig functions

Identities	Domain	Range
$\sin^{-1} \theta$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$\cos^{-1} \theta$	$[-1, 1]$	$[0, \pi]$
$\tan^{-1} \theta$	$\mathbb{R}$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$\cot^{-1} \theta$	$\mathbb{R}$	$(0, \pi)$
$\csc^{-1} \theta$	$(-\infty, -1] \cup [1, \infty)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
$\sec^{-1} \theta$	$(-\infty, -1] \cup [1, \infty)$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$

3.2. Inverse Trigonometric Formulas

$$1. \sin^{-1} x = \csc^{-1}\left(\frac{1}{x}\right)$$

$$2. \cos^{-1} x = \sec^{-1}\left(\frac{1}{x}\right)$$

$$3. \tan^{-1} x = \cot^{-1}\left(\frac{1}{x}\right), x > 0$$

$$= \cot^{-1}\left(\frac{1}{x}\right) - \pi, x < 0$$

$$= 0, x = 0$$

$$4. \sin^{-1}(-x) = -\sin^{-1} x$$

$$5. \cos^{-1}(-x) = \pi - \cos^{-1} x$$

6. $\tan^{-1}(-x) = -\tan^{-1} x$
7. $\cot^{-1}(-x) = \pi - \cot^{-1} x$
8. $\csc^{-1}(-x) = -\csc^{-1} x$
9. $\sec^{-1}(-x) = \pi - \sec^{-1} x$
10. $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$
11. $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$
12. $\csc^{-1} x + \sec^{-1} x = \frac{\pi}{2}$
13. $\sin^{-1} x \pm \sin^{-1} y = \sin^{-1}\left(x\sqrt{1-y^2} \pm y\sqrt{1-x^2}\right)$
14. $\cos^{-1} x \pm \cos^{-1} y = \cos^{-1}\left(xy \mp \sqrt{1-x^2}\sqrt{1-y^2}\right)$
15. $\tan^{-1} x + \tan^{-1} y = \tan^{-1}\left(\frac{x+y}{1-xy}\right), xy < 1$
 $= \pi + \tan^{-1}\left(\frac{x+y}{1-xy}\right), xy > 1, x > 0, y > 0$
 $= -\pi + \tan^{-1}\left(\frac{x+y}{1-xy}\right), xy > 1, x < 0, y < 0$
16. $\tan^{-1} x - \tan^{-1} y = \tan^{-1}\left(\frac{x-y}{1+xy}\right)$
17. $2 \sin^{-1} x = \cos^{-1}(1-2x^2) = \sin^{-1}(2x\sqrt{1-x^2})$
18. $2 \cos^{-1} x = \cos^{-1}(2x^2-1)$
19. $2 \tan^{-1} x = \tan^{-1}\left(\frac{2x}{1-x^2}\right) = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$
20. $3 \sin^{-1} x = \sin^{-1}(3x-4x^3)$
21. $3 \cos^{-1} x = \cos^{-1}(4x^3-3x)$