

# Sparsity Tables for Polyhedra Scars and Exact Scars

Camilla Polvara

October 23, 2025

**Polyhedra scars (fixed  $L$ ).** For each polyhedron we list the scar multiplets (including pseudoscars where observed) and their sparsity, defined as  $\text{sparsity} = \frac{\text{number of non-zero components}}{\dim \mathcal{H}}$ . Data are taken from the polyhedra study (icosahedron, dodecahedron, Catalan solids, and the Archimedean solid with a scar).

Table 1: Icosahedron ( $V=12$ ,  $\dim \mathcal{H}=2^{12}=4096$ ). Includes pseudoscars.

Label	Energy $E$	Degeneracy	Non-zero comps.	Sparsity (%)
Scar $S_1$	-6	5	280	6.836
Pseudoscar $S'_1$	-4	1	720	17.578
Pseudoscar $S'_2$	0	1	600	14.648

Table 2: Dodecahedron ( $V=20$ ,  $\dim \mathcal{H}=2^{20}=1,048,576$ ).

Label	Energy $E$	Degeneracy	Non-zero comps.	Sparsity (%)
Scar $S_1$				

Table 3: Catalan solids (fixed  $L$  of each graph).

Solid	Label	Energy $E$	Degeneracy	Non-zero comps.	Sparsity (%)
Rhombic dodecahedron ( $\dim \mathcal{H}=16384$ )	$S_1$	-12	3	432	2.637
Rhombic dodecahedron	$S_2$	-6	18	432	2.637
Rhombic dodecahedron	$S_3$	6	18	432	2.637
Rhombic dodecahedron	$S_4$	12	3	432	2.637
Triakis octahedron ( $\dim \mathcal{H}=16384$ )	$S_1$	-4	1	192	1.172
Triakis octahedron	$S_2$	0	4	368	2.246
Tetrakis hexahedron ( $\dim \mathcal{H}=16384$ )	$S_1$	-4	12	1172	7.153
Tetrakis hexahedron	$S_2$	0	1	192	1.172

Table 4: Archimedean solid with a scar: Truncated tetrahedron ( $\dim \mathcal{H}=4096$ ).

Label	Energy $E$	Degeneracy	Non-zero comps.	Sparsity (%)
$S_1$	0	1	48	1.172

**Exact scars (top-of-tower vs.  $L$ ).** Below we list the sparsity of the *top scar in the tower* as a function of chain length  $L$  for each exactly solvable model.

Table 5: Dimer scar (spin- $\frac{1}{2}$ ).

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
14	252	$2^{14} = 16384$	1.538
16	510	$2^{16} = 65536$	0.778
18	1020	$2^{18} = 262144$	0.389
20	2046	$2^{20} = 1048576$	0.195

Table 6: Domain-wall conserving spin- $\frac{1}{2}$  model — tower  $|S_n\rangle$ .

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
12	54	4096	1.318
14	210	16384	1.282
16	352	65536	0.537
18	1287	262144	0.491

Table 7: Domain-wall conserving spin- $\frac{1}{2}$  model — tower  $|S'_n\rangle$ .

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
12	54	4096	1.318
14	210	16384	1.282
16	352	65536	0.537
18	1287	262144	0.491

Table 8: Hubbard/Hirsch model —  $\eta$  tower.

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
8	70	65536	0.107
10	252	1048576	0.024
12	924	16777216	0.006

Table 9: Hubbard/Hirsch model —  $\tau$  tower.

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
8	120	65536	0.183
10	700	1048576	0.067
12	3780	16777216	0.023

Table 10: Hubbard/Hirsch model —  $\varsigma$  tower.

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
8	320	65536	0.488
10	1400	1048576	0.134
12	6048	16777216	0.036

Table 11: Spin-1 XY model — Tower A.

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
8	70	6561	1.067
10	252	59049	0.427
12	924	531441	0.174
14	3432	4782969	0.072

Table 12: Spin-1 XY model — Tower B.

$L$	Non-zero comps.	$\dim \mathcal{H}$	Sparsity (%)
8	69	6561	1.052
10	250	59049	0.423
12	923	531441	0.174
14	3430	4782969	0.072