
MATH3430: DIFFERENTIAL EQUATIONS

ADITHYA BHASKARA
PROFESSOR: SHEAGAN JOHN

TEXTBOOK: MORRIS TENENBAUM & HARRY POLLARD

UNIVERSITY OF COLORADO BOULDER




University of Colorado
Boulder

Draft: February 8, 2023

EDITION 1

TRUSTWORTHINESS OF INFORMATION BY FILE EXTENSION



Contents

Preface	iii
1 Basic Notions	1
1.1 Lecture 1: January 20, 2023	1
1.1.1 Definition of an Ordinary Differential Equation	1
1.2 Lecture 2: January 23, 2023	2
1.2.1 Explicit and Implicit Solutions	2
1.3 Lecture 3: January 25, 2023	9
1.3.1 General and Particular Solutions	9
1.4 Lecture 4: January 27, 2023	11
1.4.1 Initial Conditions	11
1.5 Lecture 5: January 30, 2023	12
1.5.1 Finding an n th Order Differential Equation Given an n -Parameter Family of Solutions	12
2 First Order Differential Equations	13
2.1 Lecture 6: February 1, 2023	13
2.1.1 An Introduction to Separable Differential Equations	13
2.2 Lecture 7: February 3, 2023	18
2.2.1 Differential Equations with Homogeneous Coefficients	18
2.3 Lecture 8: February 6, 2023	20
2.3.1 Using the Homogeneous Substitution to Solve Differential Equations	20
2.3.2 Exactness	21
2.4 Lecture 9: February 8, 2022	22
2.4.1 Solving Exact Differential Equations	22
3 Higher Order Differential Equations	23
4 Systems of Differential Equations	24
Differential Equations as a Word Cloud	25
List of Theorems and Definitions	26

Draft: February 8, 2023

Preface

To the interested reader,

This document is a compilation of lecture notes taken during the Spring 2023 semester for MATH3430: Ordinary Differential Equations at the University of Colorado Boulder. The course used *Ordinary Differential Equations*¹ by Morris Tenenbaum and Harry Pollard as its primary text. As such, many theorems, definitions, and content may be quoted from the book. This course was taught by Sheagan John, Ph. D.

The author would like to provide the following resources for students currently taking a differential equations course:

1. MIT OpenCourseWare Differential Equations Lectures From Spring 2006.
2. 3Blue1Brown's *Overview of Differential Equations*.

While much effort has been put in to remove typos and mathematical errors, it is very likely that some errors, both small and large, are present. Please keep in mind that the author wrote this resource during his second semester of his undergraduate studies. If an error needs to be resolved, please contact Adithya Bhaskara at adithya.bhaskara@colorado.edu.

Finally, the author would like to dedicate this set of lecture notes to *Aidan Janney*, *Erika Sjöblom*, and *Tate McDonald*, three of the author's closest friends who plan to take Differential Equations in the upcoming semester; Fall 2023, at the time of writing.

Best Regards,
Adithya Bhaskara

REVISED: February 8, 2023

¹Tenenbaum, M., & Pollard, H. (1985). *Ordinary Differential Equations*. Dover Publications.

1

Basic Notions

1.1 Lecture 1: January 20, 2023

1.1.1 Definition of an Ordinary Differential Equation

Consider the following definitions.

Definition 1.1.1: 🌐 Ordinary Differential Equations

An ordinary differential equation is an equation of the form

$$F(x, y(x), y'(x), \dots, y^{(n)}(x)) = 0$$

where x is an independent variable and y is n th order differentiable.

We remark that every ordinary differential equation is valid as an expression only when we specify the values of x for which it is defined.

Definition 1.1.2: 🌐 Order of a Differential Equation

The order of an ordinary differential equation is the highest nontrivial derivative present in the equation.

Consider the following ordinary differential equations. From now on, we may omit the term “ordinary,” as partial differential equations are not considered in this text.

1. $F(x, y, y') = \cos(xy') + y^2 y' + x^2 = 0$ is a first order differential equation.
2. $F(x, y, y', y'') = -\frac{1}{1-x^2} + y'' = 0$ is a second order differential equation.
3. $F(x, y, y', y'', y''', y''') = e^{xy} y'''' - x^2 y'' - \sin x = 0$ is a fourth order differential equation.

1.2 Lecture 2: January 23, 2023

1.2.1 Explicit and Implicit Solutions

We will now consider explicit and implicit solutions to differential equations.

Definition 1.2.1: 🎯 Explicit Solutions to Ordinary Differential Equations

Let $F(x, y, y', \dots, y^{(n)}) = 0$ be a differential equation defined on the interval I . Then, an explicit solution to F is a function

$$f : \mathbb{R} \rightarrow \mathbb{C}$$

for which $f(x)$ is well-defined on some set X such that $I \cap X \neq \emptyset$ and f satisfies the differential equation for all $x \in I$. That is,

$$F(x, f, f', \dots, f^{(n)}) = 0.$$

Definition 1.2.2: 🎯 Implicit Functions

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfying $f(x, y) = 0$. Then, f defines y as an implicit function of x if and only if

1. There exists an explicit function $g(x)$ such that $y = g(x)$ on some interval $I \subseteq \mathbb{R}$.
2. For all $x \in I$, $f(x, g(x)) = 0$.

Consider the following examples.

Example 1.2.1: 🎯 Is it an Implicit Function? 1

Does $f(x, y) = x^2 + y^2 - 25 = 0$ define an implicit function of x on $\{x : x \in \mathbb{R} : -5 \leq x \leq 5\}$?

We consider solving for y . We obtain $y = \pm\sqrt{25 - x^2}$. Therefore we define $y_1(x) = \sqrt{25 - x^2}$ and $y_2(x) = -\sqrt{25 - x^2}$. For $y_1(x)$, we see that

$$f(x, y_1(x)) = x^2 + (\sqrt{25 - x^2})^2 - 25 = x^2 + 25 - x^2 - 25 = 0,$$

as desired. Also, $y_1(x)$ is defined for on $\{x : x \in \mathbb{R} : -5 \leq x \leq 5\}$. Therefore, $f(x, y)$ does define an implicit function of x . Upon further inspection, $f(x, y)$ defines more than one implicit function of x .

Example 1.2.2: ♣♣ Is it an Implicit Function? 2

Does $f(x, y) = \sqrt{x^2 - y^2} + \arccos\left(\frac{x}{y}\right) = 0$, $y \neq 0$ define an implicit function?

Here, it is too difficult to solve for y . Instead, we notice that $\arccos(1) = 0$. Therefore, we try $y(x) = x$. By substitution,

$$f(x, y(x)) = \sqrt{x^2 - x^2} + \arccos\left(\frac{x}{x}\right) = 0,$$

as desired. Also, $y(x)$ is defined on \mathbb{R} . Therefore, $f(x, y)$ defines an implicit function on $\mathbb{R} - \{0\}$.

Consider the following definition.

Definition 1.2.3: ☉ Implicit Solutions to Ordinary Differential Equations

Let $F(x, y, y', \dots, y^{(n)}) = 0$ be a differential equation defined on the interval I . Then, an implicit solution to F is a relation $f(x, y) = 0$ if and only if f defines y as an implicit function of x on I , and if $y(x)$ is an explicit solution to the differential equation.

Consider the following examples.

Example 1.2.3: ♣♣ Verifying Explicit Solutions 1

Show that $y(x)$ is an explicit solution to the differential equation

$$F(x, y, y', y'') = x^2 y'' + xy' = 0.$$

Provide the domain of definition for $y(x)$ along with the solution set. Let $y(x) = \log x$, where $\log x = \log_e x$.

Proof. We first take the derivative of $y(x)$ to obtain $y'(x) = \frac{1}{x}$. The second derivative of $y(x)$ is $y''(x) = -\frac{1}{x^2}$. Then, we have

$$\begin{aligned} x^2 y''(x) + xy'(x) &= x^2 \cdot -\frac{1}{x^2} + x \cdot \frac{1}{x} \\ &= -1 + 1 = 0, \quad x \neq 0, \end{aligned}$$

as desired. The domain of definition for $y(x)$ is $\{x : x \in \mathbb{R} : x > 0\}$. The differential equation is defined on $\{x : x \in \mathbb{R}\}$. Additionally, we have the restriction $x \neq 0$. Therefore, the solution set is

$$\{x : x \in \mathbb{R}\} \cap \{x : x \in \mathbb{R} : x > 0\} \cap \{x : x \in \mathbb{R} : x \neq 0\} = \{x : x \in \mathbb{R} : x > 0\}.$$

We can then state that $y(x) = \log x$ is an explicit solution for F on $\{x : x \in \mathbb{R} : x > 0\}$. \square

Example 1.2.4: 🚩🚩 Verifying Explicit Solutions 2

Show that $y(x)$ is an explicit solution to the differential equation

$$F(x, y, y') = yy' - 4 = 0.$$

Provide the domain of definition for $y(x)$ along with the solution set. Let $y(x) = 2\sqrt{2x}$.

Proof. We first take the derivative of $y(x)$ to obtain $y'(x) = \frac{2}{\sqrt{2x}}$. Then, we have

$$\begin{aligned} y(x)y'(x) - 4 &= 2\sqrt{2x} \cdot \frac{2}{\sqrt{2x}} - 4 \\ &= 4 - 4 = 0, \quad x \neq 0, \end{aligned}$$

as desired. The domain of definition for $y(x)$ is $\{x : x \in \mathbb{R} : x \geq 0\}$. The differential equation is defined on $\{x : x \in \mathbb{R}\}$. Additionally, we have the restriction $x \neq 0$. Therefore, the solution set is

$$\{x : x \in \mathbb{R}\} \cap \{x : x \in \mathbb{R} : x \geq 0\} \cap \{x : x \in \mathbb{R} : x \neq 0\} = \{x : x \in \mathbb{R} : x > 0\}.$$

We can then state that $y(x) = 2\sqrt{2x}$ is an explicit solution for F on $\{x : x \in \mathbb{R} : x > 0\}$. \square

Example 1.2.5: 🚩🚩 Verifying Explicit Solutions 3

Show that $y(x)$ is an explicit solution to the differential equation

$$F(x, y, y', y'') = y''^3 + y'^2 - y - 3x^2 - 8.$$

Provide the domain of definition for $y(x)$ along with the solution set. Let $y(x) = x^2$.

Proof. We first take the derivative of $y(x)$ to obtain $y'(x) = 2x$. The second derivative of $y(x)$ is $y''(x) = 2$. Then, we have

$$\begin{aligned} y''^3 + y'^2 - y - 3x^2 - 8 &= (2)^3 + (2x)^2 - x^2 - 3x^2 - 8 \\ &= 8 + 4x^2 - x^2 - 3x^2 - 8 = 0 \end{aligned}$$

as desired. The domain of definition for $y(x)$ is \mathbb{R} . The differential equation is defined on \mathbb{R} . We have no additional restrictions. Therefore, the solution set is \mathbb{R} . We can then state that $y(x) = x^2$ is an explicit solution for F on \mathbb{R} . \square

Example 1.2.6: 🐼🐼 Verifying Explicit Solutions 4

Show that $y(x)$ is an explicit solution to the differential equation

$$F(x, y, y') = (x + y)^2 - y' = 0.$$

Provide the domain of definition for $y(x)$ along with the solution set. Let $y(x) = \tan x - x$.

Proof. We first take the derivative of $y(x)$ to obtain $y'(x) = \sec^2 x - 1 = \tan^2 x$. Then, we have

$$\begin{aligned} (x + \tan x - x)^2 - \tan^2 x &= (\tan x)^2 - \tan^2 x \\ &= \tan^2 x - \tan^2 x = 0 \end{aligned}$$

as desired. The domain of definition for $y(x)$ is $\{x : x \in \mathbb{R} : x \neq \frac{\pi}{2} + k\pi : k \in \mathbb{Z}\}$. The differential equation is defined on $\{x : x \in \mathbb{R}\}$. We have no additional restrictions. Therefore, the solution set is

$$\{x : x \in \mathbb{R}\} \cap \{x : x \in \mathbb{R} : x \neq \frac{\pi}{2} + k\pi : k \in \mathbb{Z}\} = \{x : x \in \mathbb{R} : x \neq \frac{\pi}{2} + k\pi : k \in \mathbb{Z}\}.$$

We can then state that $y(x) = \tan x - x$ is an explicit solution for F on $\{x : x \in \mathbb{R} : x \neq \frac{\pi}{2} + k\pi : k \in \mathbb{Z}\}$. \square

Example 1.2.7: 🐼🐼 Verifying Implicit Solutions 1

Determine whether $f(x, y) = x^2 + y^2 + 4 = 0$ provides an implicit solution to

$$F(x, y, y') = 2x + 2y'' = 0.$$

Provide the intervals of solution.

First, we determine whether $f(x, y)$ defines y as an implicit function of x . Consider the functions $g_1(x) = \sqrt{-x^2 - 4}$ and $g_2(x) = -\sqrt{-x^2 - 4}$; these functions are defined nowhere on \mathbb{R} . Thus, $f(x, y) = 0$ does not provide an implicit solution to the differential equation.

Example 1.2.8: ♠♠ Verifying Implicit Solutions 2

Determine whether $f(x, y) = xy - y^2 = 0$ provides an implicit solution to

$$F(x, y, y', y'') = \frac{1}{y - x^2} y'' + yy' - y = 0.$$

Provide the intervals of solution.

First, we determine whether $f(x, y) = xy - y^2 = y(x - y)$ defines y as an implicit function of x . Consider the functions $g_1(x) = 0$ and $g_2(x) = x$. Both g_1 and g_2 are defined on \mathbb{R} . Note that F has the restriction $y - x^2 \neq 0$. We see that $f(x, g_1(x)) = 0$ for all $x \in \mathbb{R}$ and $f(x, g_2(x)) = x^2 - x^2 = 0$ for all $x \in \mathbb{R}$. Therefore, f defines y as an implicit function of x . Taking $y = g_1(x)$ gives

$$F(x, g_1(x), g_1'(x), g_1''(x)) = \frac{1}{0 - x^2}(0) + (0)(0) - (0) = -\frac{1}{x^2} = 0, \quad x \neq 0.$$

Then, if we take $y = g_2(x)$, we have

$$F(x, g_2(x), g_2'(x), g_2''(x)) = \frac{1}{x - x^2} y'' + xy' - x = \frac{1}{x(1 - x)}(0) + x(1) - x = 0, \quad x \neq 0, x \neq 1.$$

Therefore, $f(x, y)$ provides an implicit solution to the differential equation. When providing the intervals of solution, we must explicitly pick which solution, g_1 or g_2 , to provide the interval with respect to. For $g_1(x)$, this is

$$\{x : x \in \mathbb{R} : x \neq 0\}.$$

and for $g_2(x)$, it is

$$\{x : x \in \mathbb{R} : x \neq 0, x \neq 1\}.$$

Note that we did not need to consider $y = g_2(x)$ to show that $f(x, y)$ provides an implicit solution.

Note that it is bad practice to immediately differentiate the relation $f(x, y)$. For example, in Example 1.2.7, if we immediately differentiated f , we would indeed obtain a symbolic equivalent to the differential equation, but we would not account for the domain restrictions.

Example 1.2.9: ♣♣ Verifying Implicit Solutions 3

Determine whether $f(x, y) = e^{2y} + e^{2x} - 1 = 0$ provides an implicit solution to

$$F(x, y, y') = e^{x-y} + y'e^{y-x} = 0.$$

Provide the intervals of solution.



Example 1.2.10: ♠♠ Verifying Implicit Solutions 4

Determine whether $f(x, y) = -(x+2)^2 + y^2 - 1 = 0$ provides an implicit solution to

$$F(x, y, y') = y^2 - 1 - y'(2y + xy) = 0.$$

Provide the intervals of solution.

First, we determine whether $f(x, y) = -(x+2)^2 + y^2 - 1 = 0 = 0$ defines y as an implicit function of x . Consider the functions $g_1(x) = \sqrt{1 + (x+2)^2}$ and $g_2(x) = -\sqrt{1 + (x+2)^2}$. We see that $g_1(x)$ and $g_2(x)$ are defined on \mathbb{R} . We see that $f(x, g_1(x)) = f(x, g_2(x)) = 0$ on \mathbb{R} . Therefore, f defines g_1 and g_2 as implicit functions of x . If we take $y = g_1(x)$, we have

$$\begin{aligned} F(x, g_1(x), g_1'(x)) &= 1 + (x+2)^2 - 1 - \frac{2(x+2)}{2\sqrt{1+(x+2)^2}}(2\sqrt{1+(x+2)^2} + x\sqrt{1+(x+2)^2}) \\ &= (x+2)^2 - \frac{x+2}{\sqrt{1+(x+2)^2}}(x+2)(\sqrt{1+(x+2)^2}) \\ &= (x+2)^2 - (x+2)^2 = 0. \end{aligned}$$

Then, if we take $y = g_2(x)$, we have

$$\begin{aligned} F(x, g_2(x), g_2''(x)) &= 1 + (x+2)^2 - 1 + \frac{2(x+2)}{2\sqrt{1+(x+2)^2}}(-2\sqrt{1+(x+2)^2} - x\sqrt{1+(x+2)^2}) \\ &= (x+2)^2 - \frac{x+2}{\sqrt{1+(x+2)^2}}(x+2)(\sqrt{1+(x+2)^2}) \\ &= (x+2)^2 - (x+2)^2 = 0. \end{aligned}$$

Note that F has no additional restrictions, so the interval of solution for both $g_1(x)$ and $g_2(x)$ is \mathbb{R} . Note that we did not to consider $y = g_2(x)$ to show that $f(x, y)$ provides an implicit solution.

1.3 Lecture 3: January 25, 2023

1.3.1 General and Particular Solutions

Consider the following definitions.

Definition 1.3.1: n -Parameter Families of Solutions

A differential equation $F(x, y, y', \dots, y^{(n)}) = 0$ possesses an n -parameter family of solutions $y(x, c_1, \dots, c_n)$ if and only if y is a solution for any choice of $c_1, \dots, c_n \in \mathbb{F}$.

Definition 1.3.2: Particular Solutions of Differential Equations

Let $y(x, c_1, \dots, c_n)$ be an n -parameter family of solutions to $F(x, y, y', \dots, y^{(n)}) = 0$. Then, for each choice of c_1, \dots, c_n , we obtain one particular solution.

Consider the following example.

Example 1.3.1: Finding an n -Parameter Family of Solutions

Consider $F(x, y, y', y'') = y'' = 0$. Note that F has solutions $y(x) = x$ and $y(x) = \pi$ on \mathbb{R} . Both these solutions are particular, as they contain no arbitrary constants. If we take the linear combination of the solutions to obtain

$$y(x, c_1, c_2) = c_1x + c_2,$$

as our 2-parameter family of solutions.

Note that we will often rewrite $y(x, c_1, \dots, c_n)$ as $y(x)$ even though this is an abuse of notation.

Definition 1.3.3: General Solutions of Differential Equations

Let $y(x, c_1, \dots, c_n)$ be an n -parameter family of solutions to $F(x, y, y', \dots, y^{(n)}) = 0$. Then, y is a general solution if and only if every solution to F can be obtained from some choice of c_1, \dots, c_n .

In various engineering applications, the terms defined in Definition 1.3.1 and Definition 1.3.3 are equivalent; however, this construction can break. Consider the following examples.

1. The differential equation $F(x, y, y', y'') = y'' = 0$ has the general solution $y(x, c_1, c_2) = c_1x + c_2$.
2. The differential equation $F(x, y, y') = y'^2 + y^2 = 0$ has only one particular solution $y(x) = 0$.
3. Examples 1.3.2 and 1.3.3 demonstrate where the differential equations has an n -parameter family of solutions but no general solution.

Consider the following examples.

Example 1.3.2: 🍄🍄 n -Parameter Families and General Solutions 1

Show that $F(x, y, y') = y'^2 - 3y' = 0$ has a 1-parameter family of solutions but no general solution.

Note that $y'^2 - 3y' = y'(y' - 3) = 0$. Therefore, either $y' = 0$ or $y' = 3$. We have that $y' = 0$ implies

$$y(x, c_1) = c_1.$$

For $y' = 3$, we have that $y(x) = 3x$. Both $y(x)$ and $y(x, c_1)$ are valid on \mathbb{R} . The particular solution $y(x)$ cannot be obtained from $y(x, c_1)$. But, we can take the linear combination of both solutions $y_2(x, c_1) = 3x + c_1$ because, then, we have

$$y_2'^2 - 3y_2' = (3)^2 - 3(3) = 0$$

on \mathbb{R} . Therefore, we redefine $y(x, c_1) = y_2(x, c_1)$. Still, there is no choice of c_1 which produces the particular solution $y(x) = 5$ for $y(x, c_1) = 3x + c_1$. Therefore, we have found that not all solutions to F can be obtained from $y(x, c_1)$, so y is not a general solution.

Example 1.3.3: 🍄🍄 n -Parameter Families and General Solutions 2

Show that $F(x, y, y') = y'^2 + (y - 2)y' - 2y = 0$ has two distinct 1-parameter families of solutions. Does F have a general solution?

Note that $y'^2 + (y - 2)y' - 2y = (y' + y)(y' - 2) = 0$. Therefore, $y' = 2$ or $y' = -y$. For $y' = 2$, we have the 1-parameter family $y_1(x, c_{1_1}) = 2x + c_{1_1}$. For $y' = -y$, we have $y_2(x, c_{1_2}) = c_{1_2}e^{-x}$. Both 1-parameter families are valid on \mathbb{R} . Note that both n -parameter families are distinct; they cannot be obtained from each other. They cannot be combined into a single general solution.

1.4 Lecture 4: January 27, 2023

1.4.1 Initial Conditions

Consider the following definition.

Definition 1.4.1: Initial Conditions

Let $F(x, y, y', \dots, y^{(n)}) = 0$ possess an n -parameter family of solutions. Any system of n equations which determine unique values for the arbitrary constants is called a set of initial conditions.

Consider the following example.

Example 1.4.1: Finding a Particular Solution Given Initial Conditions 1

Recall that $F(x, y, y', y'') = y'' = 0$ has a general solution $y(x, c_1, c_2) = c_1x + c_2$. Find the particular solution satisfying $y(0) = 5$ and $y'(1) = 3$.

We have that $y(0) = 5$ implies that $c_2 = 5$ and $y'(1) = 3$ implies that $c_1 = 3$. Our particular solution is then

$$y(x) = 3x + 5.$$

Example 1.4.2: Finding a Particular Solution Given Initial Conditions 2

Recall that $F(x, y, y', y'') = y'' = 0$ has a general solution $y(x, c_1, c_2) = c_1x + c_2$. Find the particular solution satisfying $y(2) = 2$ and $y(1) = 3$.

Now, we have

$$\left[\begin{array}{cc|c} 2 & 1 & 2 \\ 1 & 1 & 3 \end{array} \right] \xrightarrow{\text{RREF}} \left[\begin{array}{cc|c} 1 & 0 & -1 \\ 0 & 1 & 4 \end{array} \right].$$

Thus $c_1 = -1$ and $c_2 = 4$. Here, our particular solution is

$$y(x) = -x + 4.$$

1.5 Lecture 5: January 30, 2023

1.5.1 Finding an n th Order Differential Equation Given an n -Parameter Family of Solutions

We now direct our attention to finding an n th order differential equation when given an n -parameter family of solutions. To this, we will look at $y(x, c_1, \dots, c_n)$ and its derivatives and find relations between them. Consider the following examples.

Example 1.5.1: Find a Differential Equation With an n -Parameter Family of Solutions 1

Find an n th order differential equation given the n -parameter family $y(x, c_1) = c_1x$.

Note that $y(x)$ is a 1-parameter family of solutions. If $y(x) = c_1x$, $y'(x) = c_1$. Note that $y(x) = xy'(x)$, which gives the differential equation $F(x, y, y') = xy' - y = 0$.

Example 1.5.2: Find a Differential Equation With an n -Parameter Family of Solutions 2

Find an n th order differential equation given the n -parameter family $y(x, c_1, c_2) = c_1x + c_2e^x$.

Note that $y(x)$ is a 2-parameter family of solutions. If $y(x) = c_1x + c_2e^x$, $y'(x) = c_1 + c_2e^x$, and $y''(x) = c_2e^x$. Consider the system

$$\left[\begin{array}{cc|c} x & e^x & y(x) \\ 1 & e^x & y'(x) \\ 0 & e^x & y''(x) \end{array} \right] \xrightarrow{\text{RREF}} \left[\begin{array}{cc|c} 1 & 0 & -y''(x) + y'(x) \\ 0 & 1 & \frac{-xy'(x) + y(x)}{-xe^x + e^x} \\ 0 & 0 & \frac{-xy'(x) + y(x)}{x-1} + y''(x) \end{array} \right]$$

Thus, we have the differential equation $F(x, y, y'') = \frac{-xy'(x) + y(x)}{x-1} + y''(x) = 0$.

Example 1.5.3: Find a Differential Equation With an n -Parameter Family of Solutions 3

Find an n th order differential equation given the n -parameter family $y(x, c_1, c_2) = c_1e^{3x} + c_2e^{-2x} + x$.

Note that $y(x)$ is a 2-parameter family of solutions. If $y(x) = c_1e^{3x} + c_2e^{-2x} + x$, $y'(x) = 3c_1e^{3x} - 2c_2e^{-2x} + 1$, and $y''(x) = 9c_1e^{3x} + 4c_2e^{-2x}$. Consider the system

$$\left[\begin{array}{cc|c} e^{3x} & e^{-2x} & -x + y(x) \\ 3e^{3x} & -2e^{-2x} & -1 + y'(x) \\ 9e^{3x} & 4e^{-2x} & y''(x) \end{array} \right] \xrightarrow{\text{RREF}} \left[\begin{array}{cc|c} 1 & \frac{e^{-2x}}{e^{3x}} & \frac{-x + y(x)}{e^{3x}} \\ 0 & 1 & \frac{-3(-x + y(x)) - 1 + y'(x)}{-5e^{-2x}} \\ 0 & 0 & -\frac{3(-x + y(x)) - 1 + y'(x)}{-5e^{-2x}} + \frac{-9(-x + y(x)) + y''(x)}{e^{-2x}(4 - 9e^{3x})} \end{array} \right]$$



2

First Order Differential Equations

2.1 Lecture 6: February 1, 2023

2.1.1 An Introduction to Separable Differential Equations

We wish to solve differential equations of the form $F(x, y, y') = f(y)\frac{dy}{dx} + g(x) = 0$ where f and g are continuous on a common interval I . But, what is dx , and what is dy ? Consider the following definition.

Definition 2.1.1: Differentials

Let $y(x)$ be a differentiable function. Then, if Δx represents any small change in x , we define

$$dy(x, \Delta x) = y'(x)\Delta x$$

We wish to apply Definition 2.1.1 to the function $y(x) = x$. Consider the following theorem.

Theorem 2.1.1: A Useful Lemma for a Property of Differentials

If $y(x) = x$, $dy(x, \Delta x) = dx(x, \Delta x) = \Delta x$.

Proof. If $y(x) = x$, $y'(x) = 1$, so by Definition 2.1.1, $dy(x, \Delta x) = \Delta x$. □

Draft: February 8, 2023

Theorem 2.1.2: A Useful Property of Differentials

Let $y(x)$ be a differentiable function. Then, $dy(x, \Delta x) = y'(x) dx(x, \Delta x)$.

Proof. By Definition 2.1.1, $dy(x, \Delta x) = y'(x) \Delta x$. But, by Theorem 2.1.1, $dx(x, \Delta x)$, so

$$dy(x, \Delta x) = y'(x) \Delta x = y'(x) dx(x, \Delta x),$$

as desired. □

More familiarly, $dy = y'(x) dx$.

We may use Theorem 2.1.2 to prove a version of the familiar chain rule.

Theorem 2.1.3: The Chain Rule for Differentials

Let $y = f(x)$ be a differentiable function, and $x(t) = g(t)$. Therefore, $y(x) = f(g(t))$. Then,

$$dy(t, \Delta t) = f'(x(t)) dx(t, \Delta t).$$

Proof. Since $x = g(t)$, we have $dx(t, \Delta t) = g'(t) dt(t, \Delta t)$. Then, using the chain rule for derivatives, we obtain the chain rule for differentials below

$$dy(t, \Delta t) = f'(x(t)) g'(t) dt(t, \Delta t),$$

as desired. □

Note that if $z(x, y)$ is a function differentiable with respect to both x and y , we obtain

$$dz(x, y, \Delta x, \Delta y) = \frac{\partial z}{\partial y} dy + \frac{\partial z}{\partial x} dx.$$

Suppose we have a first order differential equation in the form $F(x, y, y') = f(y) \frac{dy}{dx} + g(x) = 0$. We use that fact that dy and dx are differential functions to write $F(x, y, y') = f(y) dy + g(x) dx = 0$. Then, we obtain the 1-parameter family of solutions

$$\int f(y) dy + \int g(x) dx = c.$$

Consider the following examples.

Example 2.1.1: 🐼🐼 Separation of Variables 1

Consider $F(x, y, y') = \frac{dy}{dx} + y = 0$, therefore, $dy = -y dx$. Assuming that $y(x) \neq 0$, we can divide by y to give

$$\frac{1}{y} dy = -dx.$$

We have a separable differential equation, and by integration, we have

$$\int \frac{1}{y} dy = \int -dx$$

meaning that

$$\log |y| = -x + c_1.$$

This provides an implicit solution. Note that it is a 1-parameter family of solutions, but is not a general function since $y(x) = 0$ is a particular solution to the differential equation, not obtainable from the family. We seek to find an explicit solution. Consider

$$e^{\log |y|} = e^{-x+c_1} = e^{-x} e^{c_1}$$

Then, let $c_2 = e^{c_1}$. Note that $c_2 > 0$. Then, we have $|y| = c_2 e^{-x}$. This 1-parameter family can be made a general solution by allowing $c_2 = 0$. Our general solution is then

$$|y| = c_2 e^{-x}, \quad c_2 \geq 0, \quad x \in \mathbb{R}.$$

If we allow $c_2 \in \mathbb{R}$, we can also write that a general solution is

$$y = c_2 e^{-x}, \quad c_2 \in \mathbb{R}, \quad x \in \mathbb{R}.$$

This method does not always work.

Example 2.1.2: ♠♠♠ Separation of Variables 2

Consider $F(x, y, y') = x^2(y - 2)\frac{dy}{dx} - y^3 = 0$. Assuming that $x \neq 0$ and $y(x) \neq 0$, we can write

$$\frac{1}{x^2} dx = \frac{y - 2}{y^3} dy.$$

We have a separable differential equation, and by integration, we have

$$\int \frac{1}{x^2} dx = \int \frac{y - 2}{y^3} dy$$

meaning that

$$\begin{aligned} -\frac{1}{x} &= \int \left(\frac{1}{y^2} - \frac{2}{y^3} \right) dy \\ &= -\frac{1}{y} + \frac{1}{y^2} + c_1. \end{aligned}$$

We may multiply by y^2 to obtain

$$-\frac{y^2}{x} = -y + 1 + c_1 y^2,$$

and by x to obtain

$$-y^2 = -xy + x + c_1 xy^2,$$

so

$$y^2(c_1 x + 1) = x(y - 1), \quad y(x) \neq 0, \quad x \neq 0$$

is a 1-parameter family of solutions, but it is not general because $y(x) = 0$ is a valid particular solution of F on \mathbb{R} .

Example 2.1.3: ♠♠♠ Separation of Variables 3

Consider $F(x, y, y') = x\sqrt{1-y} - y'\sqrt{1-x^2} = 0$. Assuming that $x^2 \neq 1$ and $y \neq 1$, we can write

$$\frac{x}{\sqrt{1-x^2}} dx = \frac{1}{\sqrt{1-y}} dy.$$

We have a separable differential equation, and by integration, we have

$$\int \frac{x}{\sqrt{1-x^2}} dx = \int \frac{1}{\sqrt{1-y}} dy.$$

meaning that

$$-\sqrt{1-x^2} = -2\sqrt{1-y} + c_1$$

is our 1-parameter family of solutions. We must restrict our solution for $|x| < 1$ and $y < 1$ due to the square roots. Our 1-parameter family of solutions is not general because $y(x) = 1$ is a valid particular solution of F on the interval.

2.2 Lecture 7: February 3, 2023

2.2.1 Differential Equations with Homogeneous Coefficients

Consider the following definitions.

Definition 2.2.1: 🎧 Homogeneous Functions

The function $f(x, y)$ is homogeneous, of order n , on some region $B \subseteq \mathbb{R}^2$ if and only if for all $x, y \in B$, either of the below hold.

1. The function $f(tx, ty) = t^n f(x, y)$ for some $n \in \mathbb{N}$.
2. The function $f(x, y) = x^n g(u)$ for some $u = \frac{y}{x}$ and $n \in \mathbb{N}$.
3. The function $f(x, y) = y^n h(u)$ for some $u = \frac{x}{y}$ and $n \in \mathbb{N}$.

Definition 2.2.2: 🎧 Differential Equations With Homogeneous Coefficients

A first order differential equation $F(x, y, y') = Q(x, y) \frac{dy}{dx} + P(x, y) = 0$ has homogeneous coefficients if and only if both $P(x, y)$ and $Q(x, y)$ are both homogeneous functions of equal order.

Consider the following examples.

Example 2.2.1: 🎧 Is it Homogeneous? 1

Determine whether $f(x, y) = 3x^2y - y^3$ is homogeneous on its domain.

Consider

$$\begin{aligned} f(tx, ty) &= 3(tx)^2(ty) - (ty)^3 \\ &= 3t^3x^2y - t^3y^3 \\ &= t^3(3x^2y - y^3) \\ &= t^3f(x, y). \end{aligned}$$

Therefore, $f(x, y)$ is homogeneous of order 3.

Example 2.2.2: 🎧 Is it Homogeneous? 2

Determine whether $f(x, y) = xy \sin(xy)$ is homogeneous on its domain.

Consider $f(tx, ty) = t^2xy \sin(t^2xy)$. There is no way in which this can be reduced to satisfy Definition 2.2.1.

Example 2.2.3: 🎲🎲 Is it Homogeneous? 3

Determine whether $f(x, y) = xy \sin\left(\frac{x}{y}\right) - x^2$ is homogeneous on its domain.

Consider

$$\begin{aligned} f(tx, ty) &= (tx)(ty) \sin\left(\frac{tx}{ty}\right) - (tx)^2 \\ &= t^2 xy \sin\left(\frac{x}{y}\right) - t^2 x^2 \\ &= t^2 \left(xy \sin\left(\frac{x}{y}\right) - x^2\right) \\ &= t^2 f(x, y). \end{aligned}$$

Therefore, $f(x, y)$ is homogeneous of order 2.

While the first condition is, usually, easiest to use to show that a function is homogeneous, the other conditions are very useful in edge cases and for proofs. Consider the following theorem.

Theorem 2.2.1: 🎯 Differential Equations With Homogeneous Coefficients are Separable

If $F(x, y, y') = Q(x, y)y' + P(x, y) = 0$ has homogeneous coefficients, F can be solved using separation of variables.

Proof. If $F(x, y, y')$ has homogeneous coefficients, $Q(x, y) = x^n g_1(u)$ and $P(x, y) = x^n g_2(u)$. We may make the substitution

$$F(x, y, y') = x^n g_1(u) \frac{dy}{dx} + x^n g_2(u) = 0.$$

Then, since $u = \frac{y}{x}$, $y = ux$, so $y' = u + x \frac{du}{dx}$. Therefore,

$$\begin{aligned} F(x, y, y') &= x^n g_1(u) \left(u + x \frac{du}{dx}\right) + x^n g_2(u) = 0 \\ &= ux^n g_1(u) + x^{n+1} g_1(u) \frac{du}{dx} + x^n g_2(u) = 0 \\ &= ux^n g_1(u) dx + x^{n+1} g_1(u) du + x^n g_2(u) dx = 0 \\ &= \frac{g_1(u)}{x} dx + \frac{g_1(u)}{u} du + \frac{g_2(u)}{ux} dx = 0, \quad ux^{n+1} \neq 0 \\ &= \frac{dx}{x} \left(g_1(u) + \frac{g_2(u)}{u}\right) + \frac{g_1(u)}{u} du = 0 \\ &= \frac{1}{x} dx + \frac{g_1(u)}{ug_1(u) + g_2(u)} du = 0, \quad g_1(u) + \frac{g_2(u)}{u} \neq 0, \quad u \neq 0 \end{aligned}$$

so $F(x, y, y')$ is separable. □

2.3 Lecture 8: February 6, 2023

2.3.1 Using the Homogeneous Substitution to Solve Differential Equations

Consider the following example.

Example 2.3.1: 🌟🌟 A Homogeneous Substitution

Find a 1-parameter family of solutions for

$$F(x, y, y') = 2xy \frac{dy}{dx} - (x^2 + y^2) = 0.$$

We may rewrite the above as

$$2xy \, dy - (x^2 + y^2) \, dx = 0.$$

We see that $2(tx)(ty) = t^2(2xy)$ and $-((tx)^2 + (ty)^2) = -(t^2x^2 + t^2y^2) = -t^2(x^2 + y^2)$ so our differential equation has homogeneous coefficients of order 2. Let $y = ux$, meaning $\frac{dy}{dx} = u + x \frac{du}{dx}$. Then, our differential equation is

$$\begin{aligned} 0 &= 2ux^2 \left(u + x \frac{du}{dx} \right) - (x^2 + u^2x^2) \\ &= 2u^2x^2 + 2ux^3 \frac{du}{dx} - x^2 - u^2x^2 \\ &= x^2(u^2 - 1) \, dx + 2ux^3 \, du \\ &= \frac{1}{x} \, dx + \frac{2u}{u^2 - 1}, \quad x \neq 0, \quad u^2 - 1 \neq 0. \end{aligned}$$

Then, we may integrate to obtain

$$c_1 = \log |x| + \log |u^2 - 1|.$$

If we let $c_2 = e^{c_1}$, we further obtain

$$c_2 = |x| |u^2 - 1|$$

By our earlier substitution, we have

$$c_2 = |x| \left| \left(\frac{y}{x} \right)^2 - 1 \right|.$$

Our restrictions are $x \neq 0$, $y(x) \neq x$; note that $y(x) = x$ is a particular solution, so our 1-parameter solution is not general.

2.3.2 Exactness

Consider the following definitions.

Definition 2.3.1: ☞ Simply Connected Regions

A set $B \subseteq \mathbb{R}^2$ is simply connected if and only if every non-intersecting closed curve lying in B contains only points of B .

Definition 2.3.2: ☞ Exact Differential Equations

The differential equation $P(x, y) dx + Q(x, y) dy = 0$ is exact if and only if $P(x, y)$, $\frac{\partial P}{\partial y}$, $Q(x, y)$, and $\frac{\partial Q}{\partial x}$ are all continuous on a common simply connected region B and

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}.$$

Moreover, there exists $f(x, y)$ such that $\frac{\partial f}{\partial x} = P(x, y)$ and $\frac{\partial f}{\partial y} = Q(x, y)$.

As motivation, let $z = f(x, y)$. Recall that

$$dz = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy.$$

If we take $\frac{\partial f}{\partial x} = P(x, y)$ and $\frac{\partial f}{\partial y} = Q(x, y)$, as stipulated in Definition 2.3.2, the right hand side of the above becomes

$$P(x, y) dx + Q(x, y) dy.$$

2.4 Lecture 9: February 8, 2022

2.4.1 Solving Exact Differential Equations

Recall the Fundamental Theorem of Calculus. That is, for some function $H(x)$, we have

$$H(x) = \int_{t_0}^x H'(t) dt.$$

Then, if $H(x) = x$, we have $x = \int_{t_0}^x 1 dt$. Recall that for $z = f(x, y)$,

$$dz = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy,$$

and if we take $\frac{\partial f}{\partial x} = P(x, y)$ and $\frac{\partial f}{\partial y} = Q(x, y)$,

$$dz = P(x, y) dx + Q(x, y) dy.$$

Notice that $P(x, y) dx + Q(x, y) dy = 0$ is exact. Let the associated simply connected region be B . From the total differential of z , we have

$$f(x, y) = \int_B dz = \int_B P(x, y) dx + \int_B Q(x, y) dy.$$

Note that all the above integrals exist, as by Definition 2.3.2, $P(x, y)$ and $Q(x, y)$ are continuous. Consider the following examples.

Example 2.4.1: 🐾🐾 Exact Differential Equation 1

Solve $F(x, y, y') = 2y \frac{dy}{dx} - \cos x \frac{dy}{dx} + y \sin x = 0$.

We must first write the differential equation in differential form to produce

$$(2y - \cos x) dy + y \sin x dx = 0.$$

Then, note that $\frac{\partial}{\partial y} y \sin x = \sin x$ and $\frac{\partial}{\partial x} (2y - \cos x) = \sin x$. Since the partial derivatives are equal, and P , Q , $\frac{\partial Q}{\partial x}$, and $\frac{\partial P}{\partial y}$ are all continuous on the simply connected region \mathbb{R}^2 , the differential equation is exact. We must now find some $z = f(x, y)$ such that $\frac{\partial f}{\partial x} = y \sin x$ and $\frac{\partial f}{\partial y} = 2y - \cos x$. Then,

$$\begin{aligned} f(x, y) &= \int y \sin x dx \\ &= -y \cos x + c(y). \end{aligned}$$

Note that, now, we have $\frac{\partial f}{\partial y} = -\cos x + c'(y) = 2y - \cos x$. Therefore, $c'(y) = 2y$, and

$$\begin{aligned} f(x, y) &= \int c'(y) dy - y \cos x \\ &= y^2 - y \cos x + c_1. \end{aligned}$$

Then, the 1-parameter family of solutions $f(x, y, c_1) = y^2 - y \cos x + c_1 = 0$ defines an implicit solution to $F(x, y, y')$ on \mathbb{R} .

3

Higher Order Differential Equations

Draft: February 8, 2023

4

Systems of Differential Equations

Draft: February 8, 2023

Chapter 1, PAGE 1

DEFINITION 1.1.1, PAGE 1 *Ordinary Differential Equations*

DEFINITION 1.1.2, PAGE 1 *Order of a Differential Equation*

DEFINITION 1.2.1, PAGE 2 *Explicit Solutions to Ordinary Differential Equations*

DEFINITION 1.2.2, PAGE 2 *Implicit Functions*

DEFINITION 1.2.3, PAGE 3 *Implicit Solutions to Ordinary Differential Equations*

DEFINITION 1.3.1, PAGE 9 *n-Parameter Families of Solutions*

DEFINITION 1.3.2, PAGE 9 *Particular Solutions of Differential Equations*

DEFINITION 1.3.3, PAGE 9 *General Solutions of Differential Equations*

DEFINITION 1.4.1, PAGE 11 *Initial Conditions*

Chapter 2, PAGE 13

DEFINITION 2.1.1, PAGE 13 *Differentials*

THEOREM 2.1.1, PAGE 13 *A Useful Lemma for a Property of Differentials*

THEOREM 2.1.2, PAGE 14 *A Useful Property of Differentials*

THEOREM 2.1.3, PAGE 14 *The Chain Rule for Differentials*

DEFINITION 2.2.1, PAGE 18 *Homogeneous Functions*

DEFINITION 2.2.2, PAGE 18 *Differential Equations With Homogeneous Coefficients*

THEOREM 2.2.1, PAGE 19 *Differential Equations With Homogeneous Coefficients are Separable*

Chapter 3, PAGE 23

Chapter 4, PAGE 24

Draft: February 8, 2023