# MATH2135: LINEAR ALGEBRA

# ADITHYA BHASKARA PROFESSOR: ROBIN DEELEY

TEXTBOOK: STEPHEN ANDRILLI & DAVID HECKER

UNIVERSITY OF COLORADO BOULDER





# Contents

1		Vectors and Matrices			
	1.1	Lecture 1: August 22, 2022			
		1.1.1 Notations, Definitions, and Conventions			
	1.2	Lecture 2: August 24, 2022			
		1.2.1 Vector Operations			
	1.3	Lecture 3: August 26, 2022			
		1.3.1 Matrices			
	1.4	Lecture 4: August 29, 2022			
		1.4.1 The Dot Product			
	1.5	Lecture 5: August 31, 2022			
		1.5.1 Matrix Multiplication			
		·			
2	-	tems of Linear Equations 22			
	2.1	Lecture 6: September 2, 2022			
		2.1.1 Systems of Linear Equations			
	2.2	Lecture 7: September 7, 2022			
		2.2.1 Systems of Linear Equations as Matrices			
		2.2.2 Matrix Row Operations			
	2.3	Lecture 8: September 9, 2022			
		2.3.1 Solving Linear Systems			
	2.4	Lecture 9: September 12, 2022			
		2.4.1 Formalizing Previous Notions Part I			
	2.5	Lecture 10: September 14, 2022			
		2.5.1 Formalizing Previous Notions Part II			
	2.6	Lecture 11: September 16, 2022			
		2.6.1 Linear Maps			
	2.7	Lecture 12: September 19, 2022			
		2.7.1 Inverses of Matrices			
	2.8	Lecture 13: September 21, 2022			
		2.8.1 Finding Inverses of Matrices			
_	_				
3		erminants and Eigenvalues 48			
	3.1	Lecture 13: September 21, 2022			
		3.1.1 Defining the Determinant			
	3.2	Lecture 14: September 23, 2022			
		3.2.1 Determinants of Upper Triangular Matrices			
	3.3	Lecture 15: September 26, 2022			
		3.3.1 Further Properties of Determinants			
		3.3.2 Figenvectors Figenvalues and Diagonalization 55			

CONTENTS	•••
CONTENTS	
CONTENTS	11

4	Finite Dimensional Vector Spaces	56
5	Linear Transformations	57
6	Orthogonality	58
Αp	ppendices	
Α	Introduction to Proofs	60
	A.1 Introduction to Proofs	60
	A.2 Proof by Induction	60
	A.3 Direct Proofs	68
	A.4 Proof by Contrapositive	69
	A.5 Proof by Contradiction	
	A.6 Combinatorial Proofs	

I go by the name of Vector. It's a mathematical term, represented by an arrow with both direction and magnitude. Vector! That's me, because I commit crimes with both direction and magnitude. Oh yeah!

Victor (Vector) Perkins



# 1.1 Lecture 1: August 22, 2022

#### 1.1.1 Notations, Definitions, and Conventions

Consider the following table for a very basic review of fundamental sets.

Symbol	Explanation
$\mathbb{N}$	Natural Numbers $(\mathbb{N}=\{1,2,\})$
$\mathbb Z$	Integers
$\mathbb{Q}$	Rationals
$\mathbb{R}$	Reals
$\mathbb{R}^n$	$\{[v_1, v_2,, v_n] : v_1,, v_n \in \mathbb{R}\}$

We will also note that there is an important distinction between vectors and points. Vectors describe "movement," whereas points describe location. For example, the vector [1,2] starting at the point (1,1) ends at the point (2,3). Consider the following diagram.



Consider the following definitions and statements.

#### **Definition 1.1.1:** The Zero Vector

We define

$$\vec{0} = [0, 0, \dots, 0].$$

#### **Definition 1.1.2: Vector Equality**

Two vectors  $\vec{v} = [v_1, v_2, ..., v_n]$  and  $\vec{w} = [w_1, w_2, ..., w_n]$  are equal if and only if

$$v_1 = w_1, v_2 = w_2, ..., v_n = w_n.$$

#### **Definition 1.1.3: Vector Magnitude**

Given a  $\vec{v} = [v_1, ..., v_n] \in \mathbb{R}^n$ , we define  $||\vec{v}||$ , the magnitude, or norm, of  $\vec{v}$ , as

$$||\vec{v}|| = \sqrt{v_1^2 + \dots + v_n^2}.$$

## Definition 1.1.4: Scalar Multiplication

Given a  $\vec{v} = [v_1, ... v_n] \in \mathbb{R}^n$ , and a scalar  $a \in \mathbb{R}$ , we define scalar multiplication as

$$\overrightarrow{av} = [av_1, \dots, av_n].$$

Consider the following theorem.

#### Theorem 1.1.1: Scalar Multiplication and Magnitude

Given a scalar  $a \in \mathbb{R}$  and  $\vec{v} \in \mathbb{R}^n$ ,

$$||a\overrightarrow{v}|| = |a|||\overrightarrow{v}||.$$

Proof.

$$||a\vec{v}|| = \sqrt{(av_1)^2 + \dots + (av_n)^2}$$

$$= \sqrt{a^2v_1^2 + \dots + a^2v_n^2}$$

$$= \sqrt{a^2}\sqrt{v_1^2 + \dots + v_n^2}$$

$$= |a|||\vec{v}||.$$

The theorem is hence proved.

Consider the following definitions.

#### **Definition 1.1.5:** • **Vector Direction**

Two nonzero vectors  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$  are

- 1. in the same direction if there exists c > 0 such that  $\vec{v} = c\vec{w}$ .
- 2. in opposite directions if there exists c < 0 such that  $\vec{v} = c\vec{w}$ .

#### **Definition 1.1.6:** • Unit Vectors

The vector  $\vec{v} \in \mathbb{R}^n$  is a unit vector if and only if  $||\vec{v}|| = 1$ .

Consider the following theorem.

#### **Theorem 1.1.2:** • Unit Vectors Represent Direction

Given a nonzero  $\vec{v} \in \mathbb{R}^n$ , there exists a unique unit vector in the same direction as  $\vec{v}$ .

*Proof.* Since  $\vec{v} \neq \vec{0}$ ,  $||\vec{v}|| \neq 0$ . Let  $\vec{u} = \frac{1}{||\vec{v}||} \vec{v}$ , meaning that

$$||\vec{u}|| = \left| \left| \frac{1}{||\vec{v}||} \vec{v} \right| \right|$$
$$= \left| \frac{1}{||\vec{v}||} \right| ||\vec{v}||$$
$$= 1$$

This means that  $\vec{u}$  is a unit vector and, because  $\frac{1}{||\vec{v}||}$  is a scalar, is in the same direction as  $\vec{v}$ . We have now shown existence. For uniqueness, assume  $\vec{w}$  is a unit vector in the same direction as  $\vec{v}$ . Then, for a positive  $c \in \mathbb{R}$ ,  $\vec{w} = c\vec{v}$  and  $||\vec{w}|| = 1$ . That is,

$$||\vec{w}|| = |c|||\vec{v}|| = 1$$
,

meaning that  $|c|=rac{1}{||\overrightarrow{v}||}.$  Because  $c>0,\ |c|=c,$  so  $\overrightarrow{w}=\overrightarrow{u}.$ 

In Theorem 1.1.2, we used the fact that for some  $\vec{v} \in \mathbb{R}^n$ ,  $\vec{v} \neq \vec{0} \implies ||\vec{v}|| \neq 0$ . We will now provide a proof.

#### Theorem 1.1.3: Nonzero Vector Implies Nonzero Magnitudes

For some  $\vec{v} = [v_1, ..., v_n] \in \mathbb{R}^n$ ,

$$\vec{v} \neq \vec{0} \implies ||\vec{v}|| \neq 0.$$

*Proof.* We will proceed by proving the contrapositive. Suppose that  $||\vec{v}|| = 0$ , meaning that

$$||\vec{v}|| = \sqrt{v_1^2 + \dots + v_n^2} = 0.$$

The only way this is true is if all of  $v_1, \dots, v_n$  are zero, which, by definition, means that  $\vec{v} = \vec{0}$ .  $\Box$ 

# 1.2 Lecture 2: August 24, 2022

#### 1.2.1 Vector Operations

As a review from the previous lecture, consider the following exercises.

#### Exercise 1.2.1: \*\* Vector Representing Movement

Find the vector representing the movement from (3, 1) to (-1, 2).

We simply find the change in the y coordinates and the change in the x coordinates to find the vector [-4, 1], starting from (3, 1). This is illustrated below.



#### Exercise 1.2.2: \* Finding a Unit Vector

Find the unit vector in the direction  $[3, -1, -\pi]$ .

We construct the unit vector by normalization. This produces  $\frac{[3,-1,-\pi]}{\sqrt{3^2+(-1)^2+(-\pi)^2}}$ .

We will now define addition and subtraction with vectors and provide a few properties.

#### **Definition 1.2.1:** Addition and Subtraction With Vectors

Let  $\vec{v} = [v_1, ..., v_n]$  and  $\vec{w} = [w_1, ..., w_n]$  be vectors in  $\mathbb{R}^n$ . Then,

$$\vec{v} \pm \vec{w} = [v_1 \pm w_1, \dots, v_n \pm w_n].$$

Geometrically, we may visualize vector addition as



#### Theorem 1.2.1: Properties of Vector Addition and Scalar Multiplication

Let  $\vec{u} = [u_1, ..., u_n]$ ,  $\vec{v} = [v_1, ..., v_n]$ , and  $\vec{w} = [w_1, ..., w_n]$  be vectors in  $\mathbb{R}^n$ . Let c and d be scalars. Then,

1. 
$$\vec{u} + \vec{v} = \vec{v} + \vec{u}$$

2. 
$$\vec{u} + (\vec{v} + \vec{w}) = (\vec{u} + \vec{v}) + \vec{w}$$

3. 
$$\vec{0} + \vec{u} = \vec{u} + \vec{0} = \vec{u}$$

4. 
$$\vec{u} + (-\vec{u}) = (-\vec{u}) + \vec{u} = \vec{0}$$

5. 
$$c(\vec{u} + \vec{v}) = c\vec{u} + c\vec{v}$$

6. 
$$(c+d)\vec{u} = c\vec{u} + d\vec{u}$$

7. 
$$(cd)\vec{u} = c(d\vec{u})$$

8. 
$$1\vec{u} = \vec{u}$$

Note that  $\vec{0}$  is called the *identity element* for addition and  $-\vec{u}$  is the *additive inverse element of*  $\vec{u}$ .

We present the proof of one of the components of Theorem 1.2.1.

#### **Example 1.2.1:** \* Commutativity of Addition

Let  $\vec{v} = [v_1, ..., v_n]$  and  $\vec{w} = [w_1, ..., w_n]$  be vectors in  $\mathbb{R}^n$ . Prove that

$$\vec{v} + \vec{w} = \vec{w} + \vec{v}$$
.

Proof. Consider the following.

$$\vec{v} + \vec{w} = [v_1, ..., v_n] + [w_1, ..., w_n]$$
  
=  $[v_1 + w_1, ..., v_n + w_n]$   
=  $[w_1 + v_1, ..., w_n + v_n]$   
=  $\vec{w} + \vec{v}$ .

The proposition is hence proved.

#### Theorem 1.2.2: Scalar Multiplication Producing the Zero Vector

Let  $\vec{v} \in \mathbb{R}^n$  and let c be a scalar. Then,

$$(c = 0 \lor \vec{v} = \vec{0}) \iff c\vec{v} = \vec{0}.$$

Proof. First, we wish to show that

$$(c = 0 \lor \vec{v} = \vec{0}) \implies c\vec{v} = \vec{0}.$$

Suppose c = 0. We have

$$\vec{0} = c\vec{v} = [cv_1, \dots, cv_n].$$

We wish to show that all of  $cv_1, ..., cv_n$  must be zero, no matter the components of  $\vec{v}$ . By basic arithmetic, zero multiplied by any other number is also zero, so if c=0,  $c\vec{v}$  must be  $\vec{0}$ , as all the components of  $c\vec{v}$  are zero. Suppose  $\vec{v}=\vec{0}$ . We again have

$$\vec{0} = c\vec{v} = [cv_1, \dots, cv_n]$$

and wish to show that all of  $cv_1, \ldots, cv_n$  must be zero, no matter the value of c. If  $\vec{v}=0$ , all components of  $\vec{v}$  are zero, and again, zero multiplied by any other number is zero, so all components of  $c\vec{v}$  are zero. We have now shown that, indeed,  $(c=0 \lor \vec{v}=\vec{0}) \implies c\vec{v}=\vec{0}$ . Now, we wish to show that

$$c\vec{v} = \vec{0} \implies (c = 0 \lor \vec{v} = \vec{0}).$$

Suppose that  $\vec{v} \neq \vec{0}$ . Again, we have

$$\vec{0} = c\vec{v} = [cv_1, \dots, cv_n].$$

There must be some nonzero  $v_1, \dots, v_n$ . Let this nonzero number be n. The only way cn = 0 is if c = 0.

# 1.3 Lecture 3: August 26, 2022

#### 1.3.1 Matrices

Consider the following warm-up exercise.

#### Exercise 1.3.1: \*\* General Vector Movement

What is the formula for the vector  $\vec{v}$  representing the movement from  $A=(a_1,\ldots,a_n)$  to  $B=(b_1,\ldots,b_n)$ . Then, find the magnitude of  $\vec{v}$ .

We see that

$$\vec{v} = [b_1 - a_1, \dots, b_n - a_n],$$

starting from A. Then,

$$||\vec{v}|| = \sqrt{(b_1 - a_1)^2 + \dots + (b_n - a_n)^2}.$$

Consider the following definitions.

#### **Definition 1.3.1:** Matrices

Let  $m, n \in \mathbb{N}$ . An  $m \times n$  matrix is a rectangular array of real numbers with m rows and n columns. Matrices are often denoted with capital letters. The elements are called entries, and are usually written as

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}.$$

The main diagonal of A consists of  $a_{11}$ ,  $a_{22}$ ,  $a_{33}$ , ....

#### **Definition 1.3.2:** Square Matrices

A matrix is square if and only if m = n.

#### **Definition 1.3.3:** Diagonal Matrices

A matrix is diagonal if and only if it is square and  $a_{ij}=0$  whenever  $i\neq j$ . That is, all elements not on the main diagonal are zero.

#### **Definition 1.3.4:** The Identity Matrix

The identity matrix A is an  $n \times n$  matrix where

$$a_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}.$$

#### **Definition 1.3.5:** The Zero Matrix

The zero matrix A is an  $m \times n$  matrix where  $a_{ij} = 0$  for all i and j.

#### **Definition 1.3.6:** • Upper Triangular Matrices

An upper triangular matrix is a matrix such that  $a_{ij} = 0$  for all i > j.

#### **Definition 1.3.7:** • Lower Triangular Matrices

An lower triangular matrix is a matrix such that  $a_{ij} = 0$  for all i < j.

#### **Definition 1.3.8:** $\blacksquare$ **The Set of All** $m \times n$ **Matrices**

The set  $\mathcal{M}_{mn}$  is the set of all  $m \times n$  matrices.

#### **Definition 1.3.9:** Matrix Addition

Given  $A, B \in \mathcal{M}_{mn}$ , we define  $A \pm B$  to be the matrix in  $\mathcal{M}_{mn}$  with entries  $a_{ij} \pm b_{ij}$ .

Consider the following example.

#### Example 1.3.1: \* Matrix Addition

Consider the following addition.

$$\begin{bmatrix} 3 & 1 & e \\ 0 & \pi & 5 \end{bmatrix} + \begin{bmatrix} -1 & 3 & e \\ 5 & 0 & -2 \end{bmatrix} = \begin{bmatrix} 2 & 4 & 2e \\ 5 & \pi & 3 \end{bmatrix}.$$

Consider the following definition.

#### **Definition 1.3.10:** Scalar Multiplication With Matrices

Given  $A \in \mathcal{M}_{mn}$  and  $c \in \mathbb{R}$ , we define cA as the matrix with elements  $ca_{ij}$ .

Consider the following example.

#### Example 1.3.2: \* Scalar Multiplication With Matrices

Consider the following scalar multiplication.

$$5\begin{bmatrix}5&1\\-1&0\end{bmatrix}=\begin{bmatrix}25&5\\-5&0\end{bmatrix}.$$

Consider the following definition.

#### **Definition 1.3.11:** Matrix Transpose

Given  $A \in \mathcal{M}_{mn}$ , we define  $A^T \in \mathcal{M}_{nm}$  to be the matrix with the (i,j) entry equal to the (j,i) entry of A.

Consider the following example.

#### Example 1.3.3: \* Matrix Transpose

Consider the following transpose.

$$\begin{bmatrix} 2 & -5 \\ 3 & -1 \\ \pi & -\pi \end{bmatrix}^T = \begin{bmatrix} 2 & 3 & \pi \\ -5 & -1 & -\pi \end{bmatrix}.$$

Consider the following definition.

#### **Definition 1.3.12:** Symmetric and Skew-Symmetric Matrices

Suppose A is a matrix. Then,

- 1. A is symmetric if and only if  $A = A^T$
- 2. A is skew-symmetric if and only if  $A = -A^T$ .

Note that A being square is necessary for the above conditions, but this condition is not sufficient.

#### Example 1.3.4: \* A Symmetric Matrix

Let 
$$A = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$$
. Is  $A$  symmetric?

We see that

$$A^T = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$$
 ,

so A is symmetric.

#### Example 1.3.5: \* A Skew-Symmetric Matrix

Let 
$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$
. Is  $A$  symmetric?

We see that

$$A^T = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = -A,$$

so A is skew-symmetric.

Consider the following theorems.

#### **Theorem 1.3.1:** Transpose Properties

Suppose  $A, B \in \mathcal{M}_{mn}$  and  $c \in \mathbb{R}$ . Then,

1. 
$$(A^T)^T = A$$

2. 
$$(A+B)^T = A^T + B^T$$

3. 
$$(cA)^T = cA^T$$
.

Proof. Consider the matrix

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}.$$

We see that

$$A^T = \begin{bmatrix} a_{11} & \cdots & a_{m1} \\ \vdots & \ddots & \vdots \\ a_{1n} & \cdots & a_{mn} \end{bmatrix}.$$

We will take the transpose of  $A^T$ , that is  $(A^T)^T$ , to yield

$$(A^{T})^{T} = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}$$
$$= A.$$

We have proved the first property. Let the matrix

$$B = \begin{bmatrix} b_{11} & \cdots & b_{1n} \\ \vdots & \ddots & \vdots \\ b_{m1} & \cdots & b_{mn} \end{bmatrix},$$

and therefore,

$$A + B = \begin{bmatrix} a_{11} + b_{11} & \cdots & a_{1n} + b_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & \cdots & a_{mn} + b_{mn} \end{bmatrix}.$$

Then,

$$(A+B)^{T} = \begin{bmatrix} a_{11} + b_{11} & \cdots & a_{m1} + b_{m1} \\ \vdots & \ddots & \vdots \\ a_{1n} + b_{1n} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$
$$= \begin{bmatrix} a_{11} & \cdots & a_{m1} \\ \vdots & \ddots & \vdots \\ a_{1n} & \cdots & a_{mn} \end{bmatrix} + \begin{bmatrix} b_{11} & \cdots & b_{m1} \\ \vdots & \ddots & \vdots \\ b_{1n} & \cdots & b_{mn} \end{bmatrix}$$
$$= A^{T} + B^{T}.$$

Thus, we have proved the second property. Finally, consider the matrix

$$cA = \begin{bmatrix} ca_{11} & \cdots & ca_{1n} \\ \vdots & \ddots & \vdots \\ ca_{m1} & \cdots & ca_{mn} \end{bmatrix}.$$

Then,

$$(cA)^{T} = \begin{bmatrix} ca_{11} & \cdots & ca_{m1} \\ \vdots & \ddots & \vdots \\ ca_{1n} & \cdots & ca_{mn} \end{bmatrix}$$
$$= c \begin{bmatrix} a_{11} & \cdots & a_{m1} \\ \vdots & \ddots & \vdots \\ a_{1n} & \cdots & a_{mn} \end{bmatrix}$$
$$= cA^{T}.$$

The final property is hence proved.

Consider the following definition.

#### **Definition 1.3.13:** © Trace

Let  $A \in \mathcal{M}_{nn}$ . Then,

$$\operatorname{trace} A = \sum_{i=1}^{n} a_{ii}.$$

That is, the sum of the elements on the main diagonal.

#### Theorem 1.3.2: Sum and Difference of Matrices and Their Transpose

Suppose A is an  $n \times n$  matrix. Then,

- 1.  $A + A^T$  is symmetric.
- 2.  $A A^T$  is skew-symmetric.

*Proof.* Consider the transpose of  $A + A^T$ . That is,

$$(A + A^{T})^{T} = A^{T} + (A^{T})^{T}$$
$$= A^{T} + A$$
$$= A + A^{T}.$$

This means that  $A + A^T$  is symmetric. Similarly, Consider the transpose of  $A - A^T$ . That is,

$$(A - A^T)^T = A^T - (A^T)^T$$
$$= A^T - A$$
$$= -(A - A^T).$$

This means that  $A - A^T$  is skew-symmetric.

#### Theorem 1.3.3: The Relation Between Square, Symmetric, and Skew-Symmetric

Suppose A is a square matrix. There exists a symmetric matrix S and a skew-symmetric matrix R such that

$$A = S + R$$
.

Proof. Note that

$$2A = (A + A^T) + (A - A^T).$$

Dividing both sides by 2 produces

$$A = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T),$$

meaning  $S=\frac{1}{2}(A+A^T)$  and  $R=\frac{1}{2}(A-A^T)$ . By Theorem 1.3.2, and basic properties of the transpose operation, S is symmetric and R is skew-symmetric, as desired. We have now shown existence, and will now show uniqueness. Suppose P is a symmetric matrix and Q is a skew-symmetric matrix, and A=P+Q. Then,  $A^T=P^T+Q^T=P-Q$ . This must mean that  $P=\frac{1}{2}(A+A^T)$  and  $Q=\frac{1}{2}(A-A^T)$ .

# 1.4 Lecture 4: August 29, 2022

#### 1.4.1 The Dot Product

We will now define another vector operation: the dot product. We will also provide some important properties.

### Definition 1.4.1: The Dot Product

Let  $\vec{v} = [v_1, ..., v_n]$  and  $\vec{w} = [w_1, ..., w_n]$  be vectors in  $\mathbb{R}^n$ . The dot product, or inner product, of  $\vec{v}$  and  $\vec{w}$  is given by

$$\vec{v} \cdot \vec{w} = \sum_{k=1}^{n} v_k w_k.$$

Note that  $\vec{v}$  and  $\vec{w}$  are orthogonal if and only if  $\vec{v} \cdot \vec{w} = 0$ . Consider the following examples.

#### Example 1.4.1: \* Dot Product 1

Find  $[1, \pi] \cdot [-1, \pi]$ .

We see that  $[1, \pi] \cdot [-1, \pi] = -1 + \pi^2$ .

#### Example 1.4.2: \* Dot Product 2

Find  $[1, 0] \cdot [0, 1]$ .

We see that  $[1, 0] \cdot [0, 1] = 0$ .

#### **Theorem 1.4.1:** Properties of the Dot Product

Let  $\vec{u} = [u_1, \dots, u_n], \ \vec{v} = [v_1, \dots, v_n], \ \text{and} \ \vec{w} = [w_1, \dots, w_n] \ \text{be vectors in } \mathbb{R}^n$ . Let c be a scalar. Then,

1. 
$$\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$$

$$2. \ \overrightarrow{u} \cdot \overrightarrow{u} = ||\overrightarrow{u}||^2 \ge 0$$

3. 
$$\vec{u} \cdot \vec{u} = 0 \iff \vec{u} = \vec{0}$$

4. 
$$c(\vec{u} \cdot \vec{v}) = (c\vec{u}) \cdot \vec{v} = \vec{u} \cdot (c\vec{v})$$

5. 
$$\vec{u} \cdot (\vec{v} + \vec{w}) = (\vec{u} \cdot \vec{v}) + (\vec{u} \cdot \vec{w})$$

6. 
$$(\vec{u} + \vec{v}) \cdot \vec{w} = (\vec{u} \cdot \vec{w}) + (\vec{v} \cdot \vec{w})$$

Consider the following example.

#### Example 1.4.3: \*\* Proving a Dot Product Property

Let  $\vec{v} = [v_1, ..., v_n]$  and  $\vec{w} = [w_1, ..., w_n]$  be vectors in  $\mathbb{R}^n$ . Let c be a scalar. Prove that

$$c(\vec{v} \cdot \vec{w}) = (c\vec{v}) \cdot \vec{w}.$$

Proof. Consider the following transitive chain of equality:

$$c(\vec{v} \cdot \vec{w}) = c(v_1 w_1 + \dots + v_n w_n)$$

$$= (cv_1 w_1 + \dots + cv_n w_n)$$

$$= ((cv_1)w_1 + \dots + (cv_n)w_n)$$

$$= (c\vec{v}) \cdot \vec{w}.$$

The proposition is hence proved.

Consider the following theorem regarding the angle between two vectors.

#### Theorem 1.4.2: The Angle Between Two Vectors

Given nonzero  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$ ,

$$\vec{v} \cdot \vec{w} = ||\vec{v}||||\vec{w}|| \cos \theta,$$

where  $\theta$  is the angle between the two vectors.

*Proof.* Consider two unit vectors  $\hat{v}$  and  $\hat{w}$  where

$$\hat{\mathbf{v}} = [\cos \alpha, \sin \alpha], \quad \hat{\mathbf{w}} = [\cos \beta, \sin \beta].$$

The angle between the two vectors is  $\theta = \beta - \alpha$ . The dot product between the two vectors is

$$\hat{\mathbf{v}} \cdot \hat{\mathbf{w}} = \cos \alpha \cos \beta + \sin \beta \sin \alpha$$
$$= \cos(\beta - \alpha)$$
$$= \cos \theta.$$

Now, if we consider  $\vec{v}=c_1\hat{v}$  and  $\vec{w}=c_2\hat{w}$ , meaning that  $||\vec{v}||=c_1$  and  $||\vec{w}||=c_2$ , we simply scale the above result with  $\hat{v}$  and  $\hat{v}$  by the magnitudes of  $\vec{v}$  and  $\vec{w}$  to produce

$$\vec{v} \cdot \vec{w} = ||\vec{v}||||\vec{w}|| \cos \theta,$$

The proposition is hence proved, but it is also of note to realize that the same proposition can be proved with the law of cosines.  $\Box$ 

We will now state the famous Cauchy-Schwarz Inequality.

#### Theorem 1.4.3: The Cauchy-Schwarz Inequality

If  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$ ,

$$|\vec{v} \cdot \vec{w}| \leq ||\vec{v}||||\vec{w}||.$$

*Proof.* Consider the following lemma. If  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$ , with  $||\vec{v}|| = ||\vec{w}|| = 1$ ,

$$-1 < \vec{v} \cdot \vec{w} < 1$$
.

*Proof.* We will start by showing that  $-1 \le \vec{v} \cdot \vec{w}$ . Consider  $||\vec{v} + \vec{w}||^2 \ge 0$ . We have that

$$\begin{aligned} ||\vec{v} + \vec{w}||^2 &= (\vec{v} + \vec{w}) \cdot (\vec{v} + \vec{w}) \\ &= \vec{v} \cdot \vec{v} + \vec{v} \cdot \vec{w} + \vec{w} \cdot \vec{v} + \vec{w} \cdot \vec{w} \\ &= ||\vec{v}||^2 + 2\vec{v} \cdot \vec{w} + ||\vec{w}||^2 \\ &= 1 + 2\vec{v} \cdot \vec{w} + 1 \\ &= 2 + 2\vec{v} \cdot \vec{w} \\ &\geq 0. \end{aligned}$$

We then solve the resulting inequality which yields

$$2\vec{v}\cdot\vec{w} \ge -2 \implies \vec{v}\cdot\vec{w} \ge -1.$$

Similarly, consider that  $||\vec{v} - \vec{w}||^2 \ge 0$ . We then have

$$||\vec{v} - \vec{w}||^2 = (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w})$$

$$= ||\vec{v}||^2 - 2\vec{v} \cdot \vec{w} + ||\vec{w}||^2$$

$$= 2 - 2\vec{v} \cdot \vec{w}$$

$$\geq 0.$$

Then, we have

$$2-2\vec{v}\cdot\vec{w}\geq 0 \implies \vec{v}\cdot\vec{w}\leq 1$$

as desired.

Consider the following cases. If  $\vec{v}=\vec{0}$  or  $\vec{w}=\vec{0}$ , then,  $\vec{v}\cdot\vec{w}=0$  and  $||\vec{v}||\vec{w}||=0$ . Thus, in this case,  $|\vec{v}\cdot\vec{w}|=||\vec{v}||||\vec{w}||=0$ . If  $\vec{v}\neq\vec{0}$  and  $\vec{w}\neq\vec{0}$ , Consider the unit vectors  $\frac{1}{||\vec{v}||}\vec{v}$  and  $\frac{1}{||\vec{w}||}\vec{w}$ . Hence, by the lemma,

$$-1 \leq \frac{1}{||\vec{v}||}\vec{v} \cdot \frac{1}{||\vec{w}||}\vec{w} \leq 1.$$

This implies that

$$-||\vec{v}||||\vec{w}|| \le \vec{v} \cdot \vec{w} \le ||\vec{v}||||\vec{w}||.$$

That is,

$$|\vec{v}\cdot\vec{w}| \leq ||\vec{v}||||\vec{w}||,$$

which is precisely the statement of the theorem.

We will now state the famous Triangle Inequality, or Minkowski's Inequality.

#### **Theorem 1.4.4:** • The Triangle Inequality

If  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$ ,

$$||\vec{v} + \vec{w}|| \le ||\vec{v}|| + ||\vec{w}||.$$

*Proof.* Since  $f(x) = \sqrt{x}$  is always increasing, we need only prove

$$||\vec{v} + \vec{w}||^2 \le (||\vec{v}|| + ||\vec{w}||)^2.$$

We have

$$\begin{aligned} ||\vec{v} + \vec{w}||^2 &= ||\vec{v}||^2 + 2\vec{v} \cdot \vec{w} + ||\vec{w}||^2 \\ &\leq ||\vec{v}||^2 + 2|\vec{v} \cdot \vec{w}| + ||\vec{w}||^2 \\ &\leq ||\vec{v}||^2 + 2||\vec{v}||||\vec{w}|| + ||\vec{w}||^2 \\ &= (||\vec{v}|| + ||\vec{w}||)^2. \end{aligned}$$

The theorem is hence proved.

We will now define the projection onto a vector.

#### **Definition 1.4.2:** The Projection Onto a Vector

Given  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$ , where  $\vec{w} \neq \vec{0}$ , we conclude that the projection of  $\vec{v}$  onto  $\vec{w}$  is

$$\operatorname{proj}_{\overrightarrow{w}} \overrightarrow{v} = ||\overrightarrow{v}|| \cos \theta \frac{\overrightarrow{w}}{||\overrightarrow{w}||}$$

$$= ||\overrightarrow{v}|| \frac{\overrightarrow{v} \cdot \overrightarrow{w}}{||\overrightarrow{v}|| ||\overrightarrow{w}||} \frac{\overrightarrow{w}}{||\overrightarrow{w}||}$$

$$= \frac{\overrightarrow{v} \cdot \overrightarrow{w}}{||\overrightarrow{w}||^2} \overrightarrow{w}$$

$$= \frac{\overrightarrow{v} \cdot \overrightarrow{w}}{||\overrightarrow{w}||^2} \overrightarrow{w}.$$

Consider the following example.

#### **Example 1.4.4: \* Projections**

Given  $\vec{v} = [x, y]$ , find proj<sub>[1,0]</sub>  $\vec{v}$  and proj<sub>[0,1]</sub>  $\vec{v}$ .

We see that

$$\text{proj}_{[1,0]} \vec{v} = [x, 0], \quad \text{proj}_{[0,1]} \vec{v} = [0, y].$$

Consider the following theorems.

#### Theorem 1.4.5: Sum of Parallel and Perpendicular Projections

Given  $\vec{w} \neq 0$  where  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$ ,

$$\vec{v} = \operatorname{proj}_{\vec{w}} \vec{v} + (\vec{v} - \operatorname{proj}_{\vec{w}} \vec{v}).$$

The first addend returns a vector parallel to  $\vec{w}$ , and the second added returns a vector perpendicular to  $\vec{w}$ 

Proof. Consider the expansion of the right hand side, that is,

$$\operatorname{proj}_{\overrightarrow{w}} \overrightarrow{v} + (\overrightarrow{v} - \operatorname{proj}_{\overrightarrow{w}} \overrightarrow{v}) = \frac{\overrightarrow{v} \cdot \overrightarrow{w}}{\overrightarrow{w} \cdot \overrightarrow{w}} \overrightarrow{w} + \left( \overrightarrow{v} - \frac{\overrightarrow{v} \cdot \overrightarrow{w}}{\overrightarrow{w} \cdot \overrightarrow{w}} \overrightarrow{w} \right)$$
$$= \overrightarrow{v}.$$

We have arrived at the desired result.

#### **Theorem 1.4.6:** Projections Depend on Lines

Given  $\vec{w} \neq 0$  where  $\vec{v}$ ,  $\vec{w} \in \mathbb{R}^n$ , and  $c \in \mathbb{R}$  where  $c \neq 0$ ,

$$\operatorname{proj}_{\overrightarrow{w}}\overrightarrow{v} = \operatorname{proj}_{\overrightarrow{cw}}\overrightarrow{v}.$$

Proof. Consider the expansion of the left hand side, that is,

$$\operatorname{proj}_{c\overrightarrow{w}}\overrightarrow{\overrightarrow{v}} = \frac{\overrightarrow{\overrightarrow{v}} \cdot c\overrightarrow{w}}{c\overrightarrow{w} \cdot c\overrightarrow{w}} c\overrightarrow{w}$$
$$= \frac{\overrightarrow{\overrightarrow{v}} \cdot \overrightarrow{w}}{\overrightarrow{w} \cdot \overrightarrow{w}} \overrightarrow{w}$$
$$= \operatorname{proj}_{\overrightarrow{w}} \overrightarrow{v}.$$

The proposition is hence proved.

# 1.5 Lecture 5: August 31, 2022

#### 1.5.1 Matrix Multiplication

Consider the following definition of matrix multiplication.

#### **Definition 1.5.1:** Matrix Multiplication

For  $A \in \mathcal{M}_{mn}$  and  $B \in \mathcal{M}_{np}$ , such that

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1p} \\ b_{21} & b_{22} & \cdots & b_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{np} \end{bmatrix},$$

the matrix product C=AB is defined such that  $C\in\mathcal{M}_{\mathit{mp}}$  and is given by

$$C = \begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1p} \\ c_{21} & c_{22} & \cdots & c_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ c_{m1} & c_{m2} & \cdots & c_{mp} \end{bmatrix}$$

such that  $c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj} = \sum_{k=1}^{n} a_{ik}b_{kj}$ . The matrix product is only defined if the number of columns of the matrix on the left is equal to the number of the rows of the matrix on the right. Note that matrix multiplication is not commutative. We may also say that the entry in  $c_{ij}$  is the dot product of the *i*th row of A and the *j*th column of B.

Consider the following examples.

#### Example 1.5.1: \* Matrix Multiplication 1

Find 
$$\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ -1 \end{bmatrix}$$
.

We see that

$$\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ -1 \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}.$$

#### Example 1.5.2: \* Matrix Multiplication 2

Find 
$$\begin{bmatrix} 3 & 1 & -1 \end{bmatrix} \begin{bmatrix} \pi \\ -3 \\ 7 \end{bmatrix}$$

We see that

$$\begin{bmatrix} 3 & 1 & -1 \end{bmatrix} \begin{bmatrix} \pi \\ -3 \\ 7 \end{bmatrix} = \begin{bmatrix} 3\pi - 10 \end{bmatrix}.$$

#### Example 1.5.3: \* Matrix Multiplication 3

Find 
$$\begin{bmatrix} 3 & 1 & -1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 7 \\ 1 & 3 \end{bmatrix}$$
.

We see that

$$\begin{bmatrix} 3 & 1 & -1 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 7 \\ 1 & -3 \end{bmatrix} = \begin{bmatrix} 6 & 13 \end{bmatrix}.$$

#### Example 1.5.4: \* Matrix Multiplication 4

Find 
$$\begin{bmatrix} 3 & 1 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

We see that

$$\begin{bmatrix} 3 & 1 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ -1 & 5 \end{bmatrix}.$$

#### Example 1.5.5: \* Matrix Multiplication 5

Find 
$$\begin{bmatrix} 3 & 1 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

We see that

$$\begin{bmatrix} 3 & 1 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Consider the following properties of matrix multiplication.

#### **Theorem 1.5.1:** Properties of Matrix Multiplication

Suppose A, B, and C are matrices such that matrix multipication is well-defined. Then,

1. 
$$A(BC) = (AB)C$$

$$2. \ A(B+C) = AB + AC$$

3. 
$$(A + B)C = AC + BC$$

4. 
$$c(AB) = (cA)B = A(cB)$$
.

We now ponder two questions.

- 1. Does AB = BA? No.
- 2. If AB = 0, does A = 0 or B = 0? No.

Consider the following definition.

# Definition 1.5.2: Raising a Matrix to a Power

Consider an  $n \times n$  matrix A. Then,

$$A^k = \underbrace{A \cdot A \cdot A \cdot \cdots \cdot A}_{k \text{ times}}$$

and  $A^k = I$ .



# Systems of Linear Equations

# 2.1 Lecture 6: September 2, 2022

#### 2.1.1 Systems of Linear Equations

Consider the following definitions.

#### **Definition 2.1.1:** • Linear Equations

A linear equation is an equation of the form

$$a_1x_1+\cdots+a_nx_n=b.$$

#### **Definition 2.1.2:** Systems of Linear Equations

A system of linear equations is a system of the form

$$a_{11}x_1+\cdots+a_{1n}x_n=b_1$$

:

$$a_{m1}x_1+\cdots+a_{mn}x_n=b_m.$$

# 2.2 Lecture 7: September 7, 2022

#### 2.2.1 Systems of Linear Equations as Matrices

We may write systems of linear equations in terms of matrices as

$$A\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix},$$

where A is the matrix with entries  $a_{ij}$ . Consider the following theorem.

#### Theorem 2.2.1: Characterizing Solutions of Linear Systems

A system of linear equations can either have

- 1. No solution.
- 2. One unique solution.
- 3. Infinitely many solutions.

#### 2.2.2 Matrix Row Operations

Consider the following operations.

- 1. Multiplication of a row by a nonzero scalar. Notated as  $c\langle r_1 
  angle o \langle r_1 
  angle$ .
- 2. Addition of a scalar multiple of one row to another. Notated as  $\langle r_1 \rangle + (c) \langle r_2 \rangle \to \langle r_1 \rangle$
- 3. Switching the elements of two rows. Notated as  $\langle r_1 \rangle \leftrightarrow \langle r_2 \rangle$ .

Consider the following examples.

#### Example 2.2.1: \* Row Operation 1

Consider the matrix 
$$\begin{bmatrix} 3 & 1 & -1 \\ 1 & 0 & 1 \\ -1 & 1 & 5 \end{bmatrix}. \text{ Find } 4\langle 3 \rangle \to \langle 3 \rangle.$$

We obtain

$$\begin{bmatrix} 3 & 1 & -1 \\ 1 & 0 & 1 \\ -4 & 4 & 20 \end{bmatrix}.$$

# Example 2.2.2: \* Row Operation 2

Consider the matrix  $\begin{bmatrix} 3 & 1 & -1 \\ 1 & 0 & 1 \\ -1 & 1 & 5 \end{bmatrix}$ . Find  $\langle 1 \rangle + (-3)\langle 2 \rangle \rightarrow \langle 1 \rangle$ .

We obtain

$$\begin{bmatrix} 0 & 1 & -4 \\ 1 & 0 & 1 \\ -1 & 1 & 5 \end{bmatrix}.$$

#### Example 2.2.3: \*\* Row Operation 3

 $\text{Consider the matrix} \begin{bmatrix} 3 & 1 & -1 \\ 1 & 0 & 1 \\ -1 & 1 & 5 \end{bmatrix}. \text{ Find } \langle 2 \rangle \leftrightarrow \langle 3 \rangle.$ 

We obtain

$$\begin{bmatrix} 3 & 1 & -1 \\ -1 & 1 & 5 \\ 1 & 0 & 1 \end{bmatrix}.$$

# 2.3 Lecture 8: September 9, 2022

#### 2.3.1 Solving Linear Systems

Given a linear system of equations, we solve by the following steps.

- 1. Convert the linear system into the matrix equation AX = B, written [A|B].
- 2. Use the three row operations to reduce [A|B] to one with "lots of zeroes and ones."
- 3. Perform back substitution and analyze the solution set.

Consider the following examples.

#### Example 2.3.1: \* No Solution

Consider the matrix

$$\begin{bmatrix} 3 & -6 & 0 & 3 & | & 9 \\ -2 & 4 & 2 & -1 & | & -11 \\ 4 & -8 & 6 & 7 & | & -5 \end{bmatrix}.$$

By row operations, we obtain

$$\begin{bmatrix} 1 & -2 & 0 & 1 & | & 3 \\ 0 & 0 & 1 & \frac{1}{2} & | & -\frac{5}{2} \\ 0 & 0 & 0 & 0 & | & -2 \end{bmatrix}.$$

Looking at the last row, we see the equation 0 = -2, which is not true. Hence, the system has no solution.

#### **Example 2.3.2:** \* Infinitely Many Solutions

Consider the matrix

$$\begin{bmatrix} 3 & 1 & 7 & 2 & | & 13 \\ 2 & -4 & 14 & -1 & | & -10 \\ 5 & 11 & -7 & 8 & | & 59 \\ 2 & 5 & -4 & -3 & | & 39 \end{bmatrix}$$

By row operations, we obtain

$$\begin{bmatrix} 1 & \frac{1}{3} & \frac{7}{3} & \frac{2}{3} & | & \frac{13}{3} \\ 0 & 1 & -2 & \frac{1}{2} & | & 4 \\ 0 & 0 & 0 & 1 & | & 59 \\ 0 & 0 & 0 & 0 & | & 0 \end{bmatrix}.$$

We see that  $x_1$ ,  $x_2$ , and  $x_4$  are determined because their respective column has a 1 in the correct position. In contrast,  $x_3$  is a free variable. To find the solution set, let  $x_3=c\in\mathbb{R}$  and solve for  $x_1$ ,  $x_2$ , and  $x_4$  in terms of c. We have  $x_4=-2$ . Then to find  $x_2$  we have

$$x_2 - 2x_3 + \frac{1}{2}x_4 = 4 \implies x_2 - 2 + \frac{1}{2}(-2) = 4 \implies x_2 = 2c + 5.$$

For  $x_1$ , we have

$$x_1 + \frac{1}{3}x_2 + \frac{7}{3}x_3 + \frac{2}{3}x_4 = \frac{13}{3} \implies x_1 = -3c + 4.$$

The solution set is then  $\{(-3c+4, 2c+5, c, -2) : c \in \mathbb{R}\}.$ 

We generally agree that back substitution is not much fun. Consider the following example.

#### Example 2.3.3: \* No More Back Substitution

Note that the matrix

$$\begin{bmatrix} 3 & -3 & -2 & | & 23 \\ -6 & 4 & 3 & | & -40 \\ -2 & 1 & 1 & | & -12 \end{bmatrix}$$

reduces into

$$\begin{bmatrix} 1 & -1 & -\frac{2}{3} & | & \frac{23}{3} \\ 0 & 1 & \frac{1}{3} & | & -\frac{10}{3} \\ 0 & 0 & 1 & | & 2 \end{bmatrix}.$$

We may now "get rid of" -1,  $-\frac{2}{3}$ , and  $\frac{1}{3}$ . We perform  $-\frac{1}{3}\langle 3\rangle+\langle 2\rangle\to\langle 2\rangle$  which produces

$$\begin{bmatrix} 1 & -1 & -\frac{2}{3} & | & \frac{23}{3} \\ 0 & 1 & 0 & | & -4 \\ 0 & 0 & 1 & | & 2 \end{bmatrix}.$$

Then, we will perform  $\langle 1 \rangle + \langle 2 \rangle \rightarrow \langle 1 \rangle$ , yielding

$$\begin{bmatrix} 1 & 0 & -\frac{2}{3} & | & \frac{23}{3} - 4 \\ 0 & 1 & 0 & | & -4 \\ 0 & 0 & 1 & | & 2 \end{bmatrix}.$$

Finally, we will perform  $\frac{2}{3}\langle 3\rangle + \langle 1\rangle \rightarrow 1$ , obtaining

$$\begin{bmatrix} 1 & 0 & 0 & | & 5 \\ 0 & 1 & 0 & | & -4 \\ 0 & 0 & 1 & | & 2 \end{bmatrix}$$

which means  $x_1 = 5$ ,  $x_2 = -4$ , and  $x_3 = 2$ .

Consider the following theorem.

#### **Theorem 2.3.1:** Row Operations

Suppose  $A \in \mathcal{M}_{mn}$  and  $B \in \mathcal{M}_{np}$ . Then,

- 1. If R is a row operation, R(AB) = (R(A))B.
- 2. If  $R_1, ..., R_n$  are row operations,  $R_n(...(R_2(R_1(AB)))...) = (R_n(...(R_2(R_1(A)))...))B$ .

Note that this result follows from the associativity of matrix multiplication, as any row operation can be represented by a multiplication of two matrices.

Consider the following examples of solving linear systems.

#### Example 2.3.4: \*\* Linear System 1

Solve the following system:

$$\begin{bmatrix} 2 & -1 & 1 & | & 0 \\ 1 & 3 & 4 & | & 0 \end{bmatrix}$$

We first perform the row operation  $\frac{1}{2}\langle 1 \rangle \to \langle 1 \rangle$ , which produces

$$\begin{bmatrix} 1 & -\frac{1}{2} & \frac{1}{2} & | & 0 \\ 1 & 3 & 4 & | & 0 \end{bmatrix}.$$

Then, we perform  $\langle 1 \rangle - \langle 2 \rangle \rightarrow \langle 2 \rangle.$  We obtain

$$\begin{bmatrix} 1 & -\frac{1}{2} & \frac{1}{2} & | & 0 \\ 0 & -\frac{7}{2} & -\frac{7}{2} & | & 0 \end{bmatrix}.$$

Next, we have  $-\frac{2}{7}\langle 2\rangle \rightarrow \langle 2\rangle.$  This yields

$$\begin{bmatrix} 1 & -\frac{1}{2} & \frac{1}{2} & | & 0 \\ 0 & 1 & 1 & | & 0 \end{bmatrix}.$$

From here, let  $x_3 = c$ . Then,  $x_2 = -c$ . To find  $x_1$ , we use the equation

$$x_1 - \frac{1}{2}(-c) + \frac{1}{2}c = 0,$$

which implies that  $x_1 = -c$ . Thus, the solution set is  $\{(-c, -c, c) : c \in \mathbb{R}\}$ .

#### Example 2.3.5: \* Linear System 2

Solve the following system:

$$\begin{bmatrix} 1 & -2 & 1 & 2 & | & 1 \\ 1 & 1 & -1 & 1 & | & 2 \\ 1 & 7 & -5 & -1 & | & 3 \end{bmatrix}$$

First, we perform  $\langle 1 \rangle - \langle 2 \rangle \rightarrow \langle 2 \rangle$ , yielding

$$\begin{bmatrix} 1 & -2 & 1 & 2 & | & 1 \\ 0 & -3 & 2 & 1 & | & -1 \\ 1 & 7 & -5 & -1 & | & 3 \end{bmatrix}.$$

Then, we perform  $\langle 1 \rangle - \langle 3 \rangle \rightarrow \langle 3 \rangle.$  We obtain

$$\begin{bmatrix} 1 & -2 & 1 & 2 & | & 1 \\ 0 & -3 & 2 & 1 & | & -1 \\ 0 & -9 & 6 & 3 & | & -2 \end{bmatrix}.$$

Then, we have  $-\frac{1}{3}\langle 2 \rangle \to \langle 2 \rangle.$  This produces

$$\begin{bmatrix} 1 & -2 & 1 & 2 & | & 1 \\ 0 & 1 & -\frac{2}{3} & -\frac{1}{3} & | & \frac{1}{3} \\ 0 & -9 & 6 & 3 & | & -2 \end{bmatrix}.$$

Our final row operation is  $\langle 3 \rangle + 9 \langle 2 \rangle \rightarrow \langle 3 \rangle.$  This provides us with

$$\begin{bmatrix} 1 & -2 & 1 & 2 & | & 1 \\ 0 & 1 & -\frac{2}{3} & -\frac{1}{3} & | & \frac{1}{3} \\ 0 & 0 & 0 & 0 & | & 1 \end{bmatrix},$$

meaning that there is no solution to the system.

#### Example 2.3.6: \*\* Linear System 3

Solve the following system:

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 \\ 2 & 0 & 2 & | & 1 \\ 1 & -3 & 4 & | & 2 \end{bmatrix}$$

Our first row operation is  $2\langle 1\rangle - \langle 2\rangle \to \langle 2\rangle.$  This produces

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 \\ 0 & -2 & 2 & | & 1 \\ 1 & -3 & 4 & | & 2 \end{bmatrix}.$$

Then, we have  $\langle 1 \rangle - \langle 3 \rangle \rightarrow \langle 3 \rangle$ , providing

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 \\ 0 & -2 & 2 & | & 1 \\ 0 & 2 & -2 & | & -1 \end{bmatrix}.$$

Next, we perform  $\langle 2 \rangle + \langle 3 \rangle \rightarrow \langle 3 \rangle.$  We obtain

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 \\ 0 & -2 & 2 & | & 1 \\ 0 & 0 & 0 & | & 0 \end{bmatrix}.$$

We perform another row operation,  $-\frac{1}{2}\langle 2\rangle \rightarrow \langle 2\rangle$ . This yields

$$\begin{bmatrix} 1 & -1 & 2 & | & 1 \\ 0 & 1 & -1 & | & -\frac{1}{2} \\ 0 & 0 & 0 & | & 0 \end{bmatrix}.$$

Next, we have  $\langle 2 \rangle + \langle 1 \rangle \rightarrow \langle 1 \rangle$ , which gives

$$\begin{bmatrix} 1 & 0 & 1 & | & \frac{1}{2} \\ 0 & 1 & -1 & | & -\frac{1}{2} \\ 0 & 0 & 0 & | & 0 \end{bmatrix}.$$

Let  $x_3=c$ . Then,  $x_2=-\frac{1}{2}+c$  and  $x_1=\frac{1}{2}-c$ . This means the solution set is  $\{\left(\frac{1}{2}-c,-\frac{1}{2}+c,c\right):c\in\mathbb{R}\}$ .

## 2.4 Lecture 9: September 12, 2022

#### 2.4.1 Formalizing Previous Notions Part I

Consider the following definitions.

#### **Definition 2.4.1:** Row Echelon Form

A matrix A is in row echelon form if and only if

- 1. All rows consisting of only zeroes are at the bottom.
- 2. The leading coefficient, or the pivot, of a nonzero row is always strictly to the right of the leading coefficient of the row above it.

#### **Definition 2.4.2:** Reduced Row Echelon Form

A matrix A is in reduced row echelon form if and only if

- 1. The first nonzero entry in each row is one.
- 2. Each successive row has its first nonzero entry in a later column.
- 3. All entries above and below the first nonzero entry are zero.
- 4. All rows consisting of only zeroes are at the bottom.

Note that every matrix has a unique reduced row echelon form.

Consider the following theorems and definitions.

#### Theorem 2.4.1: Number of Solutions to a Linear System

Let AX = B be a system of linear equations. Let C be the reduced row echelon form augmented matrix obtained by row reducing [A|B]. Then,

- 1. If there is a row of C having all zeroes to the left of the augmentation bar but with its last entry nonzero, AX = B has no solution.
- 2. If not, and if one of the columns of C to the left of the augmentation bar has no nonzero pivot entry, AX = B has an infinite number of solutions. The nonpivot columns correspond to (independent) variables that can take on any value, and the values of the remaining (dependent) variables are determined from those.
- 3. Otherwise AX = B has a unique solution.

Consider the following definitions.

#### **Definition 2.4.3:** • Homogeneous Systems

Given  $A \in \mathcal{M}_{mn}$ , the homogeneous system associated with A is

$$AX = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$
.

#### Theorem 2.4.2: Solutions to Homogeneous Systems

Given  $A \in \mathcal{M}_{mn}$ , the homogeneous system always has at least one solution, called the *trivial solution*. Namely,

$$x_1 = 0$$
,  $x_2 = 0$ , ...,  $x_n = 0$ .

Also, consider the following.

1. If m < n, the solution set is infinite.

2. If 
$$X = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$
 and  $X_{\sim} = \begin{bmatrix} x_{\sim 1} \\ \vdots \\ x_{\sim n} \end{bmatrix}$  are solutions,

$$cX + X_{\circ}$$

is a solution for any  $c \in \mathbb{R}$ .

3. If 
$$AX = \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix}$$
 and  $A\hat{X} = B = \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$ ,

$$cX \perp \hat{X}$$

is a solution to [A|B]. Notice that (2) is a special case of (3).

*Proof.* Consider  $A(cX + \hat{X})$ . We wish to show that  $A(cX + \hat{X}) = B$ . We see that

$$A(cX + \hat{X}) = A(cX) + A\hat{X}$$

$$= cAX + A\hat{X}$$

$$= \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} + \begin{bmatrix} b_1 \\ \vdots \\ b_m \end{bmatrix}$$

$$= B,$$

as desired.

This process is analogous to solving homogeneous differential equations to solve nonhomogeneous differential equations.

#### **Definition 2.4.4:** Equivalence of Linear Systems

Two systems [A|B] and  $[A_{\sim}|B_{\sim}]$  are equivalent if and only if

$$AX = B \wedge A_{\sim}X = B_{\sim}.$$

That is, if they have the same solution sets.

#### **Definition 2.4.5:** Row Equivalence

A matrix A is row equivalent to a matrix B if B can be obtained by a finite number of row operations conducted on A.

For example, Gaussian Elimination and Gauss-Jordan Elimination produce matrices that are row equivalent to the original matrix.

One may ask: What is the relationship between these relations? We see that row equivalence implies system equivalence. But, two systems can have the same solution set, but have different sizes, making row equivalence impossible. For the latter case, consider two matrices of different sizes, but with an empty solution set. Recall from discrete mathematics,

#### **Definition 2.4.6: Equivalence Relations**

A relation  $\sim$  on a set S is an equivalence relation on S if and only if  $\sim$  is reflexive, symmetric, and transitive. That is, if

- 1.  $\forall a \in S, a \sim a$ .
- 2.  $\forall a, b \in S, a \sim b \implies b \sim a$
- 3.  $\forall a, b, c \in S, a \sim b \land b \sim c \implies a \sim c$ .

Consider the following theorem.

#### Theorem 2.4.3: System Equivalence and Row Equivalence are Equivalence Relations

First, consider the following table.

Row Operation	Reverse Operation
$ \frac{c\langle i\rangle \to \langle i\rangle}{c\langle i\rangle + \langle j\rangle \to \langle j\rangle} $	$ \frac{\frac{1}{c}\langle i\rangle \to \langle i\rangle}{-c\langle i\rangle + \langle j\rangle \to \langle j\rangle} $
$\langle i \rangle \leftrightarrow \langle j \rangle$	$\langle i \rangle \leftrightarrow \langle j \rangle$

*Proof.* We will consider row equivalence first, and wish to show that row equivalence is reflexive, symmetric, and transitive. Reflexivity is trivial. We can simply not perform any row operations on a matrix A, and we are left with A. The above table can be used to show that row equivalence is symmetric. If a sequence of row operations is carried out on A and produces a matrix B, we can simply carry out the reverse operations on B to lead us back to A. For transitivity, if a sequence of row operations is carried out on A and leads to B, and a second sequence of row operations is performed on B and leads to C, we simply carry out the sequences, in sequence, on A to get us to C.

Now, we consider system equivalence. The system [A|B], of course, has the same solution set as itself. If the system [A|B] has the same solution set as [C|D], [C|D] has the same solution set as [A|B]. If [A|B] has the same solution set as [C|D] and [C|D] has the same solution set as the system [E|F], [A|B] has the same solution set as [E|F].  $\Box$ 

#### 2.5 Lecture 10: September 14, 2022

#### 2.5.1 Formalizing Previous Notions Part II

Consider the following formalization of our last discoveries.

#### Theorem 2.5.1: Row Equivalence Implies System Equivalence

If [A|B] is row equivalent to [C|D], [A|B] is equivalent to [C|D].

#### Theorem 2.5.2: Uniqueness of Reduced Row Echelon Form

Every matrix is row equivalent to a unique matrix in reduced row echelon form. Two matrices are row equivalent if and only if they have the same reduced row echelon form.

#### **Definition 2.5.1:** Rank

Given  $A \in \mathcal{M}_{mn}$ , rank A is the number of nonzero rows in the unique matrix that is row equivalent to A and is in reduced row echelon form.

Consider the following example.

#### Example 2.5.1: \* Rank 1

Consider

$$A = \begin{bmatrix} 3 & 1 & 0 & 1 \\ 0 & -2 & 12 & -5 \\ 2 & -3 & 22 & -14 \end{bmatrix}.$$

By row reduction, we have the matrix

$$\begin{bmatrix} 1 & 0 & 2 & -1 \\ 0 & 1 & -6 & 4 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

and see that rank A = 2.

Consider the following theorem.

#### **Theorem 2.5.3:** Number of Solutions to Homogeneous Systems

If  $A \in \mathcal{M}_{mn}$ ,

- 1. If rank A < n, AX = 0 has an infinite solution set.
- 2. If rank A = n, AX = 0 has only the trivial solution.

We will now define linear combinations of vectors.

#### **Definition 2.5.2:** • Linear Combinations

Let  $\vec{v}_1, \vec{v}_2, ..., \vec{v}_k \in \mathbb{R}^n$ . The vector  $\vec{v}$  is a linear combination of  $\vec{v}_1, \vec{v}_2, ..., \vec{v}_k$  if and only if there are scalars  $c_1, c_2, ..., c_k$  such that

$$\vec{v} = c_1 \vec{v}_1 + \cdots + c_k \vec{v}_k.$$

In general,  $\{c\vec{v}:c\in\mathbb{R}\}$  is a line unless  $\vec{v}=\vec{0}$ . Also,  $\{c_1\vec{v}_1+c_2\vec{v}_2:c_1,c_2\in\mathbb{R}\}$  is usually a plane, but could be either a point or a line. This pattern is an introduction to the concept of linear dependence, which will be elaborated on later in the text.

Consider the following example.

#### Example 2.5.2: \*\* Is a Vector a Linear Combination of Others? 1

Let  $\vec{v}=[1,0], \ \vec{v}_1=[\pi,1], \ \text{and} \ \vec{v}_2=[2,1].$  Is  $\vec{v}$  a linear combination of  $\vec{v}_1$  and  $\vec{v}_2$ ?

Notice that  $\vec{v}_1 - \vec{v}_2 = [\pi - 2, 0]$ . Then,

$$\frac{1}{\pi-2}[\pi-2,0]=[1,0]=\vec{\nu}.$$

We then have that

$$\vec{v} = \frac{1}{\pi - 2} [\pi, 1] - \frac{1}{\pi - 2} [2, 1].$$

Therefore,  $\vec{v}$  is a linear combination of  $\vec{v}_1$  and  $\vec{v}_2$ .

The above solution used a bit of trickery. Instead, given  $\vec{v}_1, \dots, \vec{v}_k$ , we form the equation

$$\begin{bmatrix} \vec{v}_1, \dots, \vec{v}_k \end{bmatrix} \begin{bmatrix} c_1 \\ \vdots \\ c_k \end{bmatrix} = \begin{bmatrix} \vec{v} \end{bmatrix}$$

and solve for the necessary constants.

Consider the following examples.

#### Example 2.5.3: \*\* Is a Vector a Linear Combination of Others? 2

Let  $\vec{v} = [1, 0, 0]$ ,  $\vec{v}_1 = [-4, 2, 0]$ , and  $\vec{v}_2 = [2, 1, 1]$ . Is  $\vec{v}$  a linear combination of  $\vec{v}_1$  and  $\vec{v}_2$ ?

Consider the system

$$\begin{bmatrix} -4 & 2 & | & 1 \\ 2 & 1 & | & 0 \\ 0 & 1 & | & 0 \end{bmatrix}.$$

We first perform the row operation  $-\frac{1}{4}\langle 1\rangle \to \langle 1\rangle$  to obtain

$$\begin{bmatrix} 1 & -\frac{1}{2} & | & -\frac{1}{4} \\ 2 & 1 & | & 0 \\ 0 & 1 & | & 0 \end{bmatrix}.$$

Then, we have  $2\langle 1\rangle - \langle 2\rangle \rightarrow \langle 2\rangle$ , producing

$$\begin{bmatrix} 1 & -\frac{1}{2} & | & -\frac{1}{4} \\ 0 & -2 & | & -\frac{1}{2} \\ 0 & 1 & | & 0 \end{bmatrix}.$$

Next, we will carry out  $-\frac{1}{2}\langle 2\rangle \rightarrow \langle 2\rangle$  to yield

$$\begin{bmatrix} 1 & -\frac{1}{2} & | & -\frac{1}{4} \\ 0 & 1 & | & \frac{1}{4} \\ 0 & 1 & | & 0 \end{bmatrix}.$$

We will then compute  $\langle 2 \rangle - \langle 3 \rangle \rightarrow \langle 3 \rangle$ ; we have

$$\begin{bmatrix} 1 & -\frac{1}{2} & | & -\frac{1}{4} \\ 0 & 1 & | & \frac{1}{4} \\ 0 & 0 & | & \frac{1}{4} \end{bmatrix}.$$

There is no solution, so  $\vec{v}$  is not a linear combination of  $\vec{v}_1$  and  $\vec{v}_2$ .

#### Example 2.5.4: \*\* Is a Vector a Linear Combination of Others? 3

Let  $\vec{v} = [14, -21, 7]$ ,  $\vec{v}_1 = [2, -3, 1]$ , and  $\vec{v}_2 = [-4, 6, 2]$ . Is  $\vec{v}$  a linear combination of  $\vec{v}_1$  and  $\vec{v}_2$ ?

Consider the system

$$\begin{bmatrix} 2 & -4 & | & 14 \\ -3 & 6 & | & -21 \\ 1 & 2 & | & 7 \end{bmatrix}.$$

We first perform the row operation  $\frac{1}{2}\langle 1 \rangle \rightarrow \langle 1 \rangle$  to obtain

$$\begin{bmatrix} 1 & -2 & | & 7 \\ -3 & 6 & | & -21 \\ 1 & 2 & | & 7 \end{bmatrix}.$$

Then, we have  $3\langle 1 \rangle + \langle 2 \rangle \rightarrow \langle 2 \rangle$ , producing

$$\begin{bmatrix} 1 & -2 & | & 7 \\ 0 & 0 & | & 0 \\ 1 & 2 & | & 7 \end{bmatrix}.$$

Next, we will carry out  $\langle 2 \rangle \leftrightarrow \langle 3 \rangle$  to yield

$$\begin{bmatrix} 1 & -2 & | & 7 \\ 1 & 2 & | & 7 \\ 0 & 0 & | & 0 \end{bmatrix}.$$

We will then compute  $\langle 1 \rangle + \langle 2 \rangle \rightarrow \langle 2 \rangle$ ; we have

$$\begin{bmatrix} 1 & -2 & | & 7 \\ 2 & 0 & | & 14 \\ 0 & 0 & | & 0 \end{bmatrix}.$$

Then, we will execute  $2\langle 1 \rangle - \langle 2 \rangle \rightarrow \langle 2 \rangle$ , and we obtain

$$\begin{bmatrix} 1 & -2 & | & 7 \\ 0 & -4 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}.$$

We then have, by  $-\frac{1}{4}\langle 2\rangle \rightarrow \langle 2\rangle$ ,

$$\begin{bmatrix} 1 & -2 & | & 7 \\ 0 & 1 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}.$$

Finally, we have the operation  $\langle 1 \rangle + 2 \langle 2 \rangle \rightarrow \langle 1 \rangle$ , which produces

$$\begin{bmatrix} 1 & 0 & | & 7 \\ 0 & 1 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}.$$

Here, we see that  $\vec{v} = 7\vec{v}_1$ . We note that it would have been simple to conclude this based on the problem statement, but the method shown is the systematic algorithm for answering such questions.

#### Example 2.5.5: \*\* \* Is a Vector a Linear Combination of Others? 4

Let  $\vec{v} = [14, -21, 7]$ ,  $\vec{v}_1 = [2, -3, 1]$ , and  $\vec{v}_2 = [-4, 6, -2]$ . Is  $\vec{v}$  a linear combination of  $\vec{v}_1$  and  $\vec{v}_2$ ?

Consider the system

$$\begin{bmatrix} 2 & -4 & | & 14 \\ -3 & 6 & | & -21 \\ 1 & -2 & | & 7 \end{bmatrix}.$$

By row reduction, we finally obtain

$$\begin{bmatrix} 1 & -2 & | & 7 \\ 0 & 0 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}.$$

Here, the solution set is  $\{(2c+7,c):c\in\mathbb{R}\}$ . There are thus infinitely many ways to express  $\vec{v}$  as a linear combination of  $\vec{v}_1$  and  $\vec{v}_2$ .

Consider the following definition.

#### **Definition 2.5.3:** Row Space

Suppose  $A \in \mathcal{M}_{mn}$ . The row space of A is the subset of  $\mathbb{R}^n$  consisting of the linear combinations of the rows of A.

Consider the following examples.

#### Example 2.5.6: \* Row Space 1

Consider

$$A = \begin{bmatrix} 1 & 2 \\ 5 & 10 \end{bmatrix}.$$

The row space of A is

$$\{c_1[1,2]+c_2[5,10]:c_1,c_2\in\mathbb{R}\}.$$

In this case, the row space of A is a line. Generally, though, with two vectors, the row space will be a plane.

#### Example 2.5.7: \*\* Row Space 2

Consider

$$A = \begin{bmatrix} 1 & 3 \\ 5 & 10 \end{bmatrix}.$$

The row space of A is

$$\{c_1[1,3]+c_2[5,10]:c_1,c_2\in\mathbb{R}\}.$$

In this case, the row space of A is a plane.

To determine if a vector is in the row space of a matrix A, we consider the system  $[A^T|X]$ . One may ask: why? Well, considering A instead of  $A^T$  would provide the wrong system of equations to solve. All we are doing when determining if a vector is in the row space of A is asking if the vector can be written as a linear combination of the rows of A. Consider the following example.

#### Example 2.5.8: \* Are Vectors in the Row Space?

Consider

$$A = \begin{bmatrix} 1 & 2 \\ 5 & 10 \end{bmatrix}$$

and recall that the row space of A is

$$\{c_1[1,2]+c_2[5,10]:c_1,c_2\in\mathbb{R}\}.$$

Is [3, 6] in the row space of A? Is [1, 0] in the row space of A? We consider

$$\begin{bmatrix} 1 & 2 \\ 5 & 10 \end{bmatrix}^T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

For [3, 6], we have

$$\begin{bmatrix} 1 & 5 & | & 3 \\ 2 & 10 & | & 6 \end{bmatrix},$$

which reduces to

$$\begin{bmatrix} 1 & 5 & | & 3 \\ 0 & 0 & | & 0 \end{bmatrix}.$$

The system has (infinitely many) solutions, so [3, 6] is in the row space of A. For [1, 0], we have

$$\begin{bmatrix} 1 & 5 & | & 1 \\ 2 & 10 & | & 0 \end{bmatrix},$$

which reduces to

$$\begin{bmatrix} 1 & 5 & | & 1 \\ 0 & 0 & | & -2 \end{bmatrix}.$$

The system has no solution, so [1,0] is not in the row space of A.

Consider the following theorems.

#### Theorem 2.5.4: Transitivity of Linear Combinations

Suppose that  $\vec{x}$  is a linear combination of  $\vec{q}_1, \dots, \vec{q}_k$ , and suppose also that each of  $\vec{q}_1, \dots, \vec{q}_k$  is itself a linear combination of  $\vec{r}_1, \dots, \vec{r}_l$ . Then,  $\vec{x}$  is a linear combination of  $\vec{r}_1, \dots, \vec{r}_l$ .

*Proof.* Because  $\vec{x}$  is a linear combination of  $\vec{q}_1, \dots, \vec{q}_k$ ,

$$\vec{x} = c_1 \vec{q}_1 + \dots + c_k \vec{q}_k$$

for  $c_1, \ldots, c_k \in \mathbb{R}$ . Then, since each of  $\vec{q}_1, \ldots, \vec{q}_k$  can be written as a linear combination of  $\vec{r}_1, \ldots, \vec{r}_l$ , there exist scalars  $d_{11}, \ldots, d_{kl}$  such that

$$\vec{q}_{1} = d_{11}\vec{r}_{1} + d_{12}\vec{r}_{2} + \dots + d_{1l}\vec{r}_{l}$$

$$\vec{q}_{2} = d_{21}\vec{r}_{1} + d_{22}\vec{r}_{2} + \dots + d_{2l}\vec{r}_{l}$$

$$\vdots$$

$$\vec{q}_{k} = d_{k1}\vec{r}_{1} + d_{k2}\vec{r}_{2} + \dots + d_{kl}\vec{r}_{l}$$

Then,

$$\vec{x} = c_1(d_{11}\vec{r}_1 + d_{12}\vec{r}_2 + \dots + d_{1l}\vec{r}_l) + c_2(d_{21}\vec{r}_1 + d_{22}\vec{r}_2 + \dots + d_{2l}\vec{r}_l) \vdots + c_k(d_{k1}\vec{r}_1 + d_{k2}\vec{r}_2 + \dots + d_{kl}\vec{r}_l) = (c_1d_{11} + c_2d_{21} + \dots + c_kd_{k1})\vec{r}_1 + (c_1d_{12} + c_2d_{22} + \dots + c_kd_{k2})\vec{r}_2 \vdots + (c_1d_{1l} + c_2d_{2l} + \dots + c_kd_{kl})\vec{r}_l.$$

We have just written  $\vec{x}$  as a linear combination of  $\vec{r}_1, \dots, \vec{r}_l$ .

Note that this theorem may be rephrased as follows: If  $\vec{x}$  is in the row space of a matrix Q and each row of Q is in the row space of a matrix R,  $\vec{x}$  is in the row space of R.

#### Theorem 2.5.5: Row Equivalence Implies Equal Row Space

Suppose A and B are row equivalent. Then, the row space of A is equal to the row space of B.

#### 2.6 Lecture 11: September 16, 2022

#### 2.6.1 Linear Maps

Consider the following definition.

#### **Definition 2.6.1:** • Linear Maps

Given  $A \in \mathcal{M}_{mn}$ , we define

$$T_A: \mathbb{R}^n \to \mathbb{R}^m$$

$$: \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \mapsto A \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

Consider the following example.

#### Example 2.6.1: \* Some Special Maps in $\mathbb{R}^2$

Consider the following maps, and name them.

$$1. \ \ \mathsf{If} \ A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \! , \ T_A : \mathbb{R}^2 \to \mathbb{R}^2 \ \mathsf{is the zero map}.$$

2. If 
$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
,  $T_A : \mathbb{R}^2 \to \mathbb{R}^2$  is the identity map.

3. If 
$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$
,  $T_A : \mathbb{R}^2 \to \mathbb{R}^2$  is the projection onto the  $x$  axis.

4. If 
$$A=\begin{bmatrix}0&0\\0&1\end{bmatrix}$$
,  $T_A:\mathbb{R}^2\to\mathbb{R}^2$  is the projection onto the  $y$  axis.

#### 2.7 Lecture 12: September 19, 2022

#### 2.7.1 Inverses of Matrices

Consider the following definitions and theorems.

#### **Definition 2.7.1:** Multiplicative Inverse of a Matrix

Let  $A \in \mathcal{M}_{nn}$ . Then,  $B \in \mathcal{M}_{nn}$  is a multiplicative inverse of A if and only if

$$AB = BA = I_n$$
.

Consider the following examples.

#### Example 2.7.1: \* The Inverse of the Identity

Let  $A = I_n$ . Find the inverse of A.

*Proof.* Since  $I_nI_n=I_nI_n=I_n$ ,  $I_n$  is the inverse of A.

#### Example 2.7.2: \* The Inverse of the Zero

Let  $A = 0_n$ . Show that A does not have an inverse.

*Proof.* For all  $B \in \mathcal{M}_{nn}$ ,  $AB = 0_n B = 0_n \neq I_n$ .

#### Theorem 2.7.1: Inverse Commutativity

Let  $A, B \in \mathcal{M}_{nn}$ . If either AB or BA equals  $I_n$ , the other product also equals  $I_n$ , and A and B are inverses of each other.

#### **Definition 2.7.2:** Singularity

A matrix is *singular* if and only if it is square and does not have an inverse. A matrix is *nonsingular* if and only if it is square and has an inverse.

#### Theorem 2.7.2: Uniqueness of the Inverse

If B and C are both inverses of  $A \in \mathcal{M}_{nn}$ , B = C.

Proof. 
$$B = BI_n = B(AC) = (BA)C = I_nC = C$$
.

We denote the unique inverse of A as  $A^{-1}$ . We can use the inverse to define negative integral powers of a nonsingular matrix A. Consider the following definition.

#### **Definition 2.7.3:** Negative Integral Powers of a Nonsingular Matrices

Let A be a nonsingular matrix. Then, the negative integral powers of A are given as follows:  $A^{-1}$  is the unique inverse of A. For  $k \ge 2$ ,  $A^{-k} = (A^{-1})^k$ .

#### **Theorem 2.7.3:** Properties of Nonsingular Matrices

Let A and B be nonsingular  $n \times n$  matrices. Then,

1.  $A^{-1}$  is nonsingular, and  $(A^{-1})^{-1} = A$ .

*Proof.* We have that  $A^{-1}A = AA^{-1} = I_n$  since  $A^{-1}$  is the inverse of A, hence  $A^{-1}$  is nonsingular and  $(A^{-1})^{-1} = A$ .

2.  $A^k$  is nonsingular, and  $(A^k)^{-1} = (A^{-1})^k = A^{-k}$ , for  $k \in \mathbb{Z}$ .

*Proof.* We have that  $A^kA^{-k}=A^{-k+k}=A^0=I_n$ . Hence,  $A^k$  is nonsingular and  $(A^k)^{-1}=A^{-k}$ .

3. *AB* is nonsingular, and  $(AB)^{-1} = B^{-1}A^{-1}$ .

*Proof.* We have that  $(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AI_nA^{-1} = AA^{-1} = I_n$ . Hence, AB is nonsingular and  $(AB)^{-1} = B^{-1}A^{-1}$ .

4.  $A^T$  is nonsingular, and  $(A^T)^{-1} = (A^{-1})^T$ .

*Proof.* We have that  $(A^{T})(A^{-1})^{T} = (A^{-1}A)^{T} = I_{n}^{T} = I_{n}$ .

Now, we will fully provide statements of matrix exponent laws.

#### **Theorem 2.7.4:** Matrix Exponent Laws

If A is nonsingular and  $p, q \in \mathbb{Z}$ ,

- 1.  $A^{p+q} = (A^p)(A^q)$ .
- 2.  $(A^p)^q = A^{pq} = (A^q)^p$ .

Consider the following theorem.

#### Theorem 2.7.5: $2 \times 2$ Inverse

Suppose  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then, A is nonsingular if and only if  $ad - bc \neq 0$ . In this case,

$$A = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

*Proof.* Suppose  $ad - bc \neq 0$ . We consider

$$\frac{1}{ad-bc}\begin{bmatrix} d & -b \\ -c & a \end{bmatrix}\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Thus,  $A^{-1}$  exists and the formula holds. Now, suppose that ad - bc = 0. We wish to show that A is singular. Consider

$$\begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Suppose A has an inverse. Then,

$$\begin{bmatrix} d & -b \\ -c & a \end{bmatrix} (AA^{-1}) = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

But,

$$\left(\begin{bmatrix} d & -b \\ -c & a \end{bmatrix}A\right)A^{-1} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}A^{-1} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

This implies a=b=c=d=0, meaning  $A=0_2$ . From Example 2.7.2, we know that the matrix  $0_n$  is singular.

#### 2.8 Lecture 13: September 21, 2022

#### 2.8.1 Finding Inverses of Matrices

We have provided a lot of theory about the inverse of a matrix, but given  $A \in \mathcal{M}_{nn}$ , how do we determine if A is invertible. That is, how do we determine if A is nonsingular? If it is, how do we find  $A^{-1}$ ? Consider the matrix equation

$$A\begin{bmatrix} x_{11} & \cdots & x_{1n} \\ \vdots & \ddots & \vdots \\ x_{n1} & \cdots & x_{nn} \end{bmatrix} = \begin{bmatrix} 1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & 1 \end{bmatrix} = I_n.$$

Our goal is to solve for the elements  $x_{ij}$ . We thus have n linear systems to solve. That is,

$$A\begin{bmatrix} x_{11} \\ \vdots \\ x_{n1} \end{bmatrix} = \begin{bmatrix} 1 \\ \vdots \\ 0 \end{bmatrix} \quad , \dots, \quad A\begin{bmatrix} x_{1n} \\ \vdots \\ x_{nn} \end{bmatrix} = \begin{bmatrix} 0 \\ \vdots \\ 1 \end{bmatrix}.$$

To solve equations of the sort, consider the following steps.

- 1. Form  $[A|I_n]$ .
- 2. Row reduce  $[A|I_n]$  to [C|D] until we have reduced row echelon form.
- 3. If  $C = I_n$ ,  $A^{-1}$  exists and  $A^{-1} = D$ . Otherwise,  $A^{-1}$  does not exist.

Consider the following example.

#### **Example 2.8.1:** \* Finding a $2 \times 2$ Inverse

Consider the matrix

$$\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$$

First we form

$$\begin{bmatrix} 2 & 1 & | & 1 & 0 \\ 1 & 1 & | & 0 & 1 \end{bmatrix}.$$

We perform the row operation  $\langle 2 \rangle \leftrightarrow \langle 1 \rangle$ , yielding

$$\begin{bmatrix} 1 & 1 & | & 0 & 1 \\ 2 & 1 & | & 1 & 0 \end{bmatrix}.$$

Then, we execute  $2\langle 1\rangle - \langle 2\rangle \rightarrow \langle 2\rangle,$  which yields

$$\begin{bmatrix} 1 & 1 & | & 0 & 1 \\ 0 & 1 & | & -1 & 2 \end{bmatrix}.$$

We then compute  $\langle 2 \rangle - \langle 1 \rangle \rightarrow \langle 1 \rangle$ , producing

$$\begin{bmatrix} -1 & 0 & | & -1 & 1 \\ 0 & 1 & | & -1 & 2 \end{bmatrix}.$$

Finally, we simply perform  $-\langle 1 \rangle \rightarrow \langle 1 \rangle,$  which obtains

$$\begin{bmatrix} 1 & 0 & | & 1 & -1 \\ 0 & 1 & | & -1 & 2 \end{bmatrix}.$$

Therefore,

$$\begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$$

is the inverse of

$$\begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}.$$

#### **Example 2.8.2:** \* Finding a $3 \times 3$ Inverse

Consider the matrix

$$\begin{bmatrix} 2 & -6 & 5 \\ -4 & 12 & -9 \\ 2 & -9 & 8 \end{bmatrix}.$$

First, we form

$$\begin{bmatrix} 2 & -6 & 5 & | & 1 & 0 & 0 \\ -4 & 12 & -9 & | & 0 & 1 & 0 \\ 2 & -9 & 8 & | & 0 & 0 & 1 \end{bmatrix}.$$

After row reduction, we have

$$\begin{bmatrix} 1 & 0 & 0 & | & \frac{5}{2} & \frac{1}{2} & -1 \\ 0 & 1 & 0 & | & \frac{7}{3} & 1 & -\frac{1}{3} \\ 0 & 0 & 1 & | & 2 & 1 & 0 \end{bmatrix}.$$

Thus,

$$\begin{bmatrix} 2 & -6 & 5 \\ -4 & 12 & -9 \\ 2 & -9 & 8 \end{bmatrix} \begin{bmatrix} \frac{5}{2} & \frac{1}{2} & -1 \\ \frac{7}{3} & 1 & -\frac{1}{3} \\ 2 & 1 & 0 \end{bmatrix} = I_3.$$



#### Determinants and Eigenvalues

#### 3.1 Lecture 13: September 21, 2022

#### 3.1.1 Defining the Determinant

Consider the following theorems and definitions.

#### Theorem 3.1.1: $\ \ \$ The Determinant Determines the Area in $\mathbb{R}^2$

Consider  $\vec{x} = [x_1, x_2]$  and  $\vec{y} = [y_1, y_2]$ . If we form

$$A = \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix},$$

$$|\det A| = ||\vec{x} \times \vec{y}||.$$

That is,  $|\det A|$  provides the area of the parallelogram determined by  $\vec{x}$  and  $\vec{y}$ .

#### Theorem 3.1.2: $\blacksquare$ The Determinant Determines the Volume in $\mathbb{R}^3$

Consider  $\vec{x} = [x_1, x_2, x_3], \ \vec{y} = [y_1, y_2, y_3], \ \text{and} \ \vec{z} = [z_1, z_2, z_3].$  If we form

$$A = \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix},$$

$$|\det A| = \vec{x} \cdot (\vec{y} \times \vec{z}).$$

That is,  $|\det A|$  provides the volume of the parallelepiped determined by  $\vec{x}$ ,  $\vec{y}$ , and  $\vec{z}$ .

#### **Definition 3.1.1:** $\bigcirc$ **The** (i, j) **Submatrix**

Suppose  $A \in \mathcal{M}_{nn}$ . The (i,j) submatrix of A is the  $(n-1) \times (n-1)$  matrix obtained by removing the ith row and the jth column. We denote this by  $A_{(i,j)}$ 

#### **Definition 3.1.2:** $\bigcirc$ **The** (i, j) **Minor**

Suppose  $A \in \mathcal{M}_{nn}$ . The (i, j) minor of A is the determinant of the (i, j) submatrix of A.

#### **Definition 3.1.3:** $\bigcirc$ **The** (i, j) **Cofactor**

Suppose  $A \in \mathcal{M}_{nn}$ . The (i, j) cofactor of  $\overline{A}$  is

$$A_{ij} = (-1)^{i+j} \det(A_{(i,j)}).$$

#### **Definition 3.1.4:** The Determinant

Suppose  $A \in \mathcal{M}_{nn}$ . Then,

1. If 
$$n = 1$$
, and  $A = [a_{11}]$ , det  $A = a_{11}$ .

2. If 
$$n = 2$$
, and  $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$ ,  $\det A = a_{11}a_{22} - a_{12}a_{21}$ .

3. If 
$$n > 2$$
, and  $A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix}$ ,  $\det A = (a_{11}A_{11} + \cdots + a_{1n}A_{1n}) + \cdots + (a_{n1}A_{n1} + \cdots + a_{nn}A_{nn})$ .

For fun, consider the following Python 3 implementation of computing the determinant of any  $n \times n$  matrix.

```
1 # Delete nth Column for Cofactor
def delete_nth_column_helper(matrix, n):
      return [m[:n]+m[n+1:] for m in matrix[1:]]
6 # Calculate Matrix Determinant
7 def determinant(matrix):
      # Base Cases
     if len(matrix) == 1: return matrix[0][0]
     if len(matrix) == 2: return matrix[0][0]*matrix[1][1]-matrix[0][1]*matrix[1][0]
11
12
    # Calculate Determinant Parts
13
     for _ in range(len(matrix[0])):
14
15
          determinant_components = [(-1)**i*matrix[0][i]*determinant(
16
      delete_nth_column_helper(matrix, i)) for i in range(len(matrix[0]))]
17
     # Return Sum
18
     return sum(determinant_components)
19
21 if __name__ == "__main__":
23
      matrix = []
24
      # Get Matrix
     for _ in range(int(input())): matrix.append([int(i) for i in input().split()])
26
27
    # Output
print(determinant(matrix))
```

#### 3.2 Lecture 14: September 23, 2022

#### 3.2.1 Determinants of Upper Triangular Matrices

Consider the following theorems.

#### Theorem 3.2.1: ■ The Determinant of an Upper Triangular Matrix

If  $A \in \mathcal{M}_{nn}$  is upper triangular,

$$\det A = a_{11}a_{22}\cdots a_{nn}.$$

Recall that A is upper triangular if and only if all elements below the main diagonal are zero.

*Proof.* We proceed by induction on n. For n=1,  $A=\begin{bmatrix} a_{11} \end{bmatrix}$ , so  $\det A=a_{11}$ . Suppose for all  $k\in\mathbb{N}$ , and some upper triangular  $A\in\mathcal{M}_{kk}$ ,

$$a_{11}a_{22}\cdots a_{kk}$$
.

Consider

$$B = \begin{bmatrix} b_{11} & \cdots & b_{1(k+1)} \\ \vdots & \ddots & \vdots \\ 0 & \cdots & b_{(k+1)(k+1)} \end{bmatrix}.$$

Then, we compute det B using the last row as our "first row."

$$\det B = (-1)^{1+1}b_{11} \det \begin{bmatrix} b_{22} & b_{23} & \cdots & a_{2(k+1)} \\ 0 & b_{33} & \cdots & b_{3(k+1)} \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & b_{(k+1)(k+1)} \end{bmatrix}$$

$$= b_{11} \underbrace{b_{22} \cdots b_{(k+1)(k+1)}}_{\text{By the inductive hypothesis.}}.$$

This is precisely the stipulation of the theorem when n = k + 1.

#### **Theorem 3.2.2:** Determinants and Row Operations

Suppose  $A \in \mathcal{M}_{nn}$  and let R be a row operation. Then,

1. If R is  $c\langle i \rangle \to \langle i \rangle$  for some  $c \in \mathbb{R}$ ,

$$\det R(A) = c \det A$$
.

Note that if c = 0, R is not a valid row operation.

2. If *R* is  $c\langle i \rangle + \langle j \rangle \rightarrow \langle j \rangle$  for some  $c \in \mathbb{R}$ ,

$$\det R(A) = \det A$$
.

Note that if c = 0, R is not a valid row operation.

3. If R is  $\langle i \rangle \leftrightarrow \langle j \rangle$ ,

$$\det R(A) = -\det A.$$

Note that if det  $A \neq 0$  and R is a row operation, det  $R(A) \neq 0$ .

We can use Theorem 3.2.2 in conjunction with row operations to compute the determinant of a matrix easily. We simply use row operations to create an upper triangular matrix, while keeping track of how the determinant changes. We then apply Theorem 3.2.1. Consider the following examples.

#### **Example 3.2.1:** \*\* Computing a Determinant by Row Reduction

Compute det  $\begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & -2 \\ 4 & 9 & 4 \end{bmatrix}$ .

Let

$$A_0 = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & -2 \\ 4 & 9 & 4 \end{bmatrix}.$$

Consider the following table.

Row Operation	Resultant Matrix			atrix	Effect on the Determinant
	$A_1 = $	1 0 4	1 1 9	1 -4 4	$\det A_1 = \det A_0$
$-4\langle 1\rangle + \langle 3\rangle \rightarrow \langle 3\rangle$	$A_2 =$	1 0 0	1 1 5	1 -4 0	$\det A_2 = \det A_1$
$-5\langle 2\rangle + \langle 3\rangle \rightarrow \langle 3\rangle$	$A_3 =$	1 0 0	1 1 0	1 -4 20	$\det A_3 = \det A_2$

Thus,  $\det A_0 = 20$ .

#### Example 3.2.2: \*\* Computing a Determinant by Row Reduction

Compute det 
$$\begin{bmatrix} 0 & -14 & -8 \\ 1 & 3 & 2 \\ -2 & 0 & 6 \end{bmatrix}$$
.

Let

$$A_0 = \begin{bmatrix} 0 & -14 & -8 \\ 1 & 3 & 2 \\ -2 & 0 & 6 \end{bmatrix}.$$

Consider the following table.

Row Operation	Resultant Matrix	Effect on the Determinant
$\langle 2 \rangle \leftrightarrow \langle 1 \rangle$	$A_1 = \begin{bmatrix} 1 & 3 & 2 \\ 0 & -14 & -8 \\ -2 & 0 & 6 \end{bmatrix}$	$\det A_1 = -\det A_0$
$2\langle 1\rangle + \langle 3\rangle \rightarrow \langle 3\rangle$	$A_2 = egin{bmatrix} 1 & 3 & 2 \ 0 & -14 & -8 \ 0 & 6 & 10 \end{bmatrix}$	$\det A_2 = \det A_1$
$-\frac{1}{14}\langle 2\rangle \to \langle 2\rangle$	$A_3 = \begin{bmatrix} 1 & 3 & 2 \\ 0 & 1 & \frac{4}{7} \\ 0 & 6 & 10 \end{bmatrix}$	$\det A_3 = -\tfrac{1}{14} \det A_2$
$-6\langle 2\rangle + \langle 3\rangle \rightarrow \langle 3\rangle$	$A_3 = \begin{bmatrix} 1 & 3 & 2 \\ 0 & 1 & \frac{4}{7} \\ 0 & 0 & \frac{46}{7} \end{bmatrix}$	$\det A_4 = \det A_3$

Thus, det  $A_0 = \frac{46}{7}(-14)(-1) = 92$ .

Consider the following theorem.

#### Theorem 3.2.3: Inverses and Determinants

Suppose that  $A \in \mathcal{M}_{nn}$ . Then, A is nonsingular if and only if  $\det A \neq 0$ .

*Proof.* If A is nonsingular, we can row reduce A to produce  $I_n$ . Since  $\det I_n \neq 0$ , by Theorem 3.2.2,  $\det A \neq 0$ . If  $\det A \neq 0$ , we form the system [A|B] and reduce it to [C|D]. We know that  $\det C \neq 0$ . Because C is in reduced row echelon form, and square since A is square, it is upper triangular. That means all main diagonal entries are nonzero, meaning they are all 1. Meaning  $C = I_n$ . This means we were able to row reduce A to  $I_n$ , so A is nonsingular.

Consider the following table summarizing various results about singularity, determinants, rank, existence and uniqueness of linear systems, and row equivalence. Statements in each column are equivalent.

$A \in \mathcal{M}_{nn}$ is Nonsingular	$A\in\mathcal{M}_{nn}$ is Singular
- rank $A = n$	rank A < n
$\det A  eq 0$	$\det A = 0$
A is row equivalent to $I_n$ .	A is not row equivalent to $I_n$ .
AX = 0 has only the trivial solution for $X$ .	AX = 0 has a nontrivial solution for $X$ .
,	AX = B does not have a unique solution.

#### 3.3 Lecture 15: September 26, 2022

#### 3.3.1 Further Properties of Determinants

#### **Theorem 3.3.1:** Properties of Determinants

Suppose  $A, B \in \mathcal{M}_{nn}$ . Then,

1. det(AB) = det A det B.

*Proof.* If A or B is singular,  $\det A \det B = 0$ . For now, let B be singular. We don't make any assumptions about A, for now. Sine B is singular, there exists some  $X \neq 0$  such that BX = 0. We consider (AB)X = A(BX) = A(0) = 0. Thus, X is a nontrivial solution to the homogeneous equation associated with AB. Thus AB is singular and  $\det(AB) = 0 = \det A \det B$ . Now, suppose that A is singular and B is nonsingular. There exists a nontrivial solution for Y in the system AY = 0. Since B is nonsingular,  $B^{-1}$  exists. We define  $X = B^{-1}Y$  where  $X \neq 0$ . Then  $ABX = AB(B^{-1}Y) = AY = 0$ . Thus, ABX and  $X \neq 0$  implies that AB is singular, so  $\det(AB) = 0$ . Now, we consider the case where A and B are both nonsingular. There exists row operations  $B_1, \ldots, B_k$  such that  $A = B_1(\ldots(B_k(I_n)\ldots))$  and

$$det(AB) = det(R_1(...(R_k(I_n)...))B)$$

$$= c_1 \cdots c_k \det(I_nB)$$

$$= c_1 \cdots c_k \det B$$

$$= c_1 \cdots c_k \det I_n \det B$$

$$= det(R_1(...(R_k(I_n)...))) \det B$$

$$= det A \det B,$$

hence proving the proposition.

- 2.  $det(A^T) = det A$ .
- 3.  $\det(A^{-1}) = \frac{1}{\det A}$ Suppose that A is nonsingular.

*Proof.* We know that A is nonsingular. We have  $\det I_n = \det(AA^{-1}) = \det A \det(A^{-1}) = 1$ . By simple algebra,  $\det(A^{-1}) = \frac{1}{\det A}$ .

#### 3.3.2 Eigenvectors, Eigenvalues, and Diagonalization

Consider the following definitions and theorems.

#### **Definition 3.3.1:** Similarity

Suppose  $A, B \in \mathcal{M}_{nn}$ . Then, A and B are similar if and only if there exists nonsingular P such that

$$A = PBP^{-1}$$
.

#### **Definition 3.3.2:** Diagonalizability

The matrix  $A \in \mathcal{M}_{nn}$  is diagonalizable if A is similar to a diagonal matrix.

#### **Definition 3.3.3:** © **Eigenvalues and Eigenvectors**

For  $A \in \mathcal{M}_{nn}$ ,  $\lambda \in \mathbb{R}$  is an eigenvalue of A if and only if there exists  $X \neq 0$  such that

$$AX = \lambda X$$
.

If  $\lambda$  is an eigenvalue of A, X is an eigenvector of A with eigenvalue  $\lambda$ .

#### **Theorem 3.3.2:** Finding Eigenvectors and Eigenvalues

Consider a matrix  $A \in \mathcal{M}_{nn}$ . By definition, X is an eigenvector of A with eigenvalue  $\lambda \in \mathbb{R}$  when

$$AX = \lambda X = \lambda IX$$
.

That is, when

$$(A - \lambda I)X = 0.$$

The above equation has a nonzero solution for X if and only if

$$\det(A - \lambda I) = 0.$$

Therefore, the scalar  $\lambda$  is an eigenvalue of A if and only if  $\lambda$  satisfies the above equation.

Note that, for now, we will only consider real eigenvalues; however, complex eigenvalues are incredibly useful and have numerous applications.

Finite Dimensional Vector Spaces

$$\begin{bmatrix} \cos 90^{\circ} & \sin 90^{\circ} \\ -\sin 90^{\circ} & \cos 90^{\circ} \end{bmatrix} \begin{bmatrix} \alpha_{1} \\ \alpha_{2} \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$$

# Linear Transformations

## Orthogonality

### **Appendices**



#### A.1 Introduction to Proofs

Before we delve into techniques to write proofs, let us first define what a proof is.

#### **Definition A.1.1:** Proofs

Mathematical proofs are logical arguments to show that stated premises guarantee that a mathematical statement must be true.

There are multiple techniques to write proofs, but here, we will explore the Proof by Induction, the Direct Proof, the Proof by Contrapositive, the Proof by Contradiction, the Proof by Cases, and the Combinatorial Proof

#### A.2 Proof by Induction

We will use quantifiers to state induction.

Let P(n) be a statement with  $n \in \mathbb{N}$ . Consider the following Rule of Inference.

$$P(0) \Rightarrow P(1)$$

$$P(1) \Rightarrow P(2)$$

$$\vdots$$

$$P(n) \Rightarrow P(n+1)$$

$$\therefore \forall n \in \mathbb{N}, P(n).$$

This may be further collapsed into

$$\frac{P(0)}{\forall k \in \mathbb{N}, P(k) \implies P(k+1)}$$

$$\therefore \forall n \in \mathbb{N}, P(n).$$

Generally, in Proofs by Induction, we follow the following steps.

- Start with an iterative proposition that depends on some  $n \in \mathbb{N}$ , or P(n).
- Prove that the proposition is true for some base case  $n = n_0$ . That is, show that the proposition is true for the smallest fixed number that the proposition makes sense for.
- Prove the inductive step. Suppose that the proposition holds true for n = k, and then prove that the proposition holds for n = k + 1. Essentially, suppose P(k) is true, and prove that P(k + 1) is true.
- Then, the proposition is proved  $\forall n \in \mathbb{N}$ , where  $n \geq n_0$ .

Consider the following examples and exercises.

#### Example A.2.1: \* Gauss' Formula

Prove that the sum of consecutive integers starting at 1 can be found by Gauss' formula. That is,

$$1+2+3\cdots+n=\frac{n(n+1)}{2}$$
.

*Proof.* Consider the base case n=1. Then the left hand side is 1, and the right hand side is

$$\frac{1(1+1)}{2} = 1.$$

Therefore, the left hand side is equal to the right hand side, proving the case base.

We suppose that the relationship is true for n=k where  $k\in\mathbb{N}.$  That is, we suppose that

$$1+2+3+\cdots+k=\frac{k(k+1)}{2}.$$

If we add k + 1 to both sides, we obtain

$$1+2+3+\cdots+k+k+1 = \frac{k(k+1)}{2}+k+1$$

$$= \frac{k(k+1)+2k+2}{2}$$

$$= \frac{k^2+3k+2}{2}$$

$$= \frac{(k+2)(k+1)}{2}$$

$$= \frac{(k+1)((k+1)+1)}{2}.$$

This result is the proposition where n = k+1. Therefore, the inductive step is true. Therefore, Gauss' formula is true for all  $n \in \mathbb{N}$  where  $n \geq 1$ .

#### Example A.2.2: \* Sum of Consequtive Squares

Prove that for all  $n \in \mathbb{N}$ ,  $n \ge 1$ ,

$$1^2 + 2^2 + 3^2 \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

*Proof.* Consider the base case n = 1. Then the left hand side is 1, and the right hand side is

$$\frac{1(1+1)(2+1)}{6} = 1.$$

Therefore, the left hand side is equal to the right hand side, proving the base case.

We suppose that the relationship is true for n=k where  $k\in\mathbb{N}$ . That is, we suppose that

$$1^2 + 2^2 + 3^2 \cdots + k^2 = \frac{k(k+1)(2k+1)}{6}.$$

If we add k + 1 to both sides, we obtain

$$1^{2} + 2^{2} + 3^{2} + k^{2} + (k+1)^{2} = \frac{k(k+1)(2k+1)}{6} + (k+1)^{2}$$

$$= \frac{k(k+1)(2k+1)}{6} + k^{2} + 2k + 1$$

$$= \frac{k(k+1)(2k+1) + 6k^{2} + 12k + 6}{6}$$

$$= \frac{2k^{3} + 9k^{2} + 13k + 6}{6}$$

$$= \frac{(k+1)(k+2)(2k+3)}{6}$$

$$= \frac{(k+1)((k+1) + 1)(2(k+1) + 1)}{6}.$$

This result is the proposition where n=k+1. Therefore, the inductive step is true. Therefore, the above formula is true for all  $n \in \mathbb{N}$  where  $n \ge 1$ .

#### Exercise A.2.1: \* The Power Rule for Derivatives

Prove that for all  $n \in \mathbb{N}$ ,  $n \ge 0$ ,

$$\frac{\mathrm{d}}{\mathrm{d}x}x^n = nx^{n-1}.$$

*Proof.* Consider the base case n = 0. Then the left hand side is 0, as the derivative of any constant is zero, and the right hand side is

$$0x^{0-1} = 0.$$

Therefore, the left hand side is equal to the right hand side, proving the base case.

We suppose that the relationship is true for n=k where  $k\in\mathbb{N}$ . That is, we suppose that

$$\frac{\mathsf{d}}{\mathsf{d}x}x^k = kx^{k-1}.$$

Consider  $\frac{d}{dx}[x^{k+1}]$ , or  $\frac{d}{dx}[xx^k]$ . Then we have

$$\frac{d}{dx}[x^{k+1}] = \frac{d}{dx}[xx^k]$$

$$= x^k + x(kx^{k-1})$$

$$= x^k + kx^k$$

$$= (k+1)x^k.$$

This result is the proposition where n=k+1. Therefore, the inductive step is true. Therefore, the power rule for derivatives is true for all  $n \in \mathbb{N}$  where  $n \geq 0$ .

#### Exercise A.2.2: \* \* nth Derivative

Prove that the *n*th Derivative of  $f(x) = \frac{1}{x}$  is

$$f^{(n)}(x) = \frac{(-1)^n n!}{x^{n+1}}.$$

*Proof.* Consider the base case n = 0. The zeroth derivative of f(x) is f(x) itself. Using the formula, we have

$$f^{(0)}(x) = \frac{(-1)^0 0!}{x^{0+1}}$$
$$= \frac{1}{x}$$
$$= f(x).$$

Therefore, the base case is true. We suppose that the relationship is true for n = k where  $k \in \mathbb{N}$ . That is, we suppose that

$$f^{(k)}(x) = \frac{(-1)^k k!}{x^{k+1}}.$$

To find the (k+1)th derivative, we differentiate  $f^{(k)}$ , producing

$$f^{(k+1)}(x) = \frac{d}{dx} \frac{(-1)^k k!}{x^{k+1}}$$

$$= \frac{(-1)^k k!}{x^{k+1+1}} (-(k+1))$$

$$= \frac{(-1)^{(k+1)} (k+1)!}{x^{(k+1)+1}}.$$

This result is the proposition where n=k+1. Therefore, the inductive step is true. Therefore, the proposition is proved for all  $n \in \mathbb{N}$  where  $n \geq 0$ .

#### Exercise A.2.3: \*\* \* Reduction

Prove that for all  $n \in \mathbb{N}$ ,  $n \ge 0$ ,

$$\int x^n e^{-x} dx = -e^{-x} \left( x^n + n x^{n-1} + n(n-1) x^{n-2} + n(n-1)(n-2) x^{n-3} + \dots + n! \right) + C.$$

*Proof.* Consider the base case n = 0. Then, the left hand side is equal to

$$\int e^{-x} \mathrm{d}x = -e^{-x} + C.$$

The right hand side is equal to  $-e^{-x} + C$ . Therefore, the left hand side is equal to the right hand side, proving the base case.

We assume that the relationship is true for n = k. That is, we assume that

$$\int x^k e^{-x} dx = -e^{-x} (x^k + kx^{k-1} + k(k-1)x^{k-2} + k(k-1)(k-2)x^{k-3} + \dots + k!) + C.$$

Let

$$u = e^{-x}(x^k + kx^{k-1} + k(k-1)x^{k-2} + k(k-1)(k-2)x^{k-3} + \dots + k!).$$

Then,

$$\int x^{k+1}e^{-x}dx = -x^{k+1}e^{-x} - \int -e^{-x}x^{k}(k+1)dx$$

$$= -x^{k+1}e^{-x} - (k+1)\int -x^{k}e^{-x}dx$$

$$= -x^{k+1}e^{-x} + (k+1)\int x^{k}e^{-x}dx$$

$$= -x^{k+1}e^{-x} - e^{-x}(k+1)\frac{u}{e^{-x}} + C$$

$$= -e^{-x}\left(x^{k+1} + \frac{u(k+1)}{e^{-x}}\right) + C$$

$$= -e^{-x}\left(x^{k+1} + (k+1)x^{k} + k(k+1)x^{k-1} + \dots + (k+1)!\right) + C.$$

This result is the proposition where n=k+1. Therefore, the inductive step is true. Therefore, the above formula is true for all  $n \in \mathbb{N}$  where  $n \ge 0$ .

#### Exercise A.2.4: \*\* \* The Shoelace Lemma

The following is a statement of the Shoelace Lemma.

Consider a simple polygon with vertices  $(x_1, y_1), (x_2, y_2), ..., (x_n, y_n)$ , oriented clockwise. Let  $(x_{n+1}, y_{n+1}) = (x_1, y_1)$ . The area of the polygon is given by

$$A_n = \frac{1}{2} \left[ \sum_{i=1}^n x_i y_{i+1} - x_{i+1} y_i \right].$$

Prove the above proposition.

*Proof.* Consider a polygon with three vertices:  $(x_1, y_1)$ ,  $(x_2, y_2)$ , and  $(x_3, y_3)$ . The area, given by the Shoelace Lemma, is

$$A_3 = \frac{1}{2} \left[ \sum_{i=1}^3 x_i y_{i+1} - x_{i+1} y_i \right] = \frac{1}{2} \left[ x_1 y_2 - x_2 y_1 + x_2 y_3 - x_3 y_2 + x_3 y_1 - y_3 x_1 \right].$$

Then, if we define two vectors  $(\vec{v}, \vec{w}) \in \mathbb{R}^3$  such that

$$\vec{v} = (x_2 - x_1, y_2 - y_1, 0), \quad \vec{w} = (x_3 - x_1, y_3 - y_1, 0),$$

we may see that the area of the parallelogram formed by the two vectors is given by

$$A_{||GRAM} = ||\vec{v} \times \vec{w}||$$

Either of the two triangles formed by the parallelogram's diagonals correspond to our polygon. The area is then given by

$$A_{3} = \frac{1}{2} ||\vec{v} \times \vec{w}||$$

$$= \frac{1}{2} ||(0, 0, x_{1}y_{2} - x_{2}y_{1} + x_{2}y_{3} - x_{3}y_{2} + x_{3}y_{1} - y_{3}x_{1})||$$

$$= \frac{1}{2} [x_{1}y_{2} - x_{2}y_{1} + x_{2}y_{3} - x_{3}y_{2} + x_{3}y_{1} - y_{3}x_{1}].$$

Therefore, we have proved the Shoelace Lemma in the case of a polygon with three vertices. By induction, we suppose that for a polygon with k vertices  $(x_1, y_1), (x_2, y_2), \dots, (x_k, y_k)$ , the area is

$$A_k = \frac{1}{2} \left[ \sum_{i=1}^k x_i y_{i+1} - x_{i+1} y_i \right].$$

The area of a polygon with vertices  $(x_1, y_1), (x_2, y_2), \dots, (x_{k+1}, y_{k+1})$  is given by the sum of the area of the polygon with vertices  $(x_1, y_1), (x_2, y_2), \dots, (x_k, y_k)$  and the area of the polygon with vertices  $(x_1, y_1), (x_k, y_k)$ , and  $(x_{k+1}, y_{k+1})$ . That is,

$$A_{k+1} = \frac{1}{2} \left[ \sum_{i=1}^{k} x_i y_{i+1} - x_{i+1} y_i \right] + \frac{1}{2} \left[ x_1 y_k - x_k y_1 + x_k y_{k+1} - x_{k+1} y_k + x_{k+1} y_1 - y_{k+1} x_1 \right]$$

$$= \frac{1}{2} \left[ \sum_{i=1}^{k+1} x_i y_{i+1} - x_{i+1} y_i \right].$$

The above result is the consequence of the Shoelace Lemma in the case of a polygon with k+1 vertices. Therefore, the Shoelace Lemma is proved.

#### Exercise A.2.5: \*\* \* Fibonacci

Let  $f_n$  represent the sequence of Fibonacci numbers, which is defined recursively as

$$f_0 = 1$$
,  $f_1 = 1$ ,  $f_n = f_{n-1} + f_{n-2}$ .

Prove that

$$\sum_{i=0}^{n} (f_i)^2 = f_n f_{n+1}.$$

*Proof.* Consider the base case n=0. Then the left hand side is equal to 1, and the right hand side is

$$(1)f_1=1.$$

Therefore, the left hand side is equal to the right hand side, proving the first base case. Then, consider the base case n=1. The left hand side is equal to 2, and the right hand side is  $1 \cdot f_2$  where  $f_2 = f_1 + f_0 = 2$ . Therefore the right hand side is 2 and is equal to the left hand side proving the second base case.

We suppose that the relationship is true for n=k where  $k\in\mathbb{N}$ . That is, we suppose that

$$\sum_{i=0}^{k} (f_i)^2 = f_k f_{k+1}.$$

If we add  $(f_{k+1})^2$  to both sides, we have

$$(f_{k+1})^2 + \sum_{i=0}^k (f_i)^2 = f_k f_{k+1} + (f_{k+1})^2$$
$$= f_{k+1} (f_k + f_{k+1})$$
$$= f_{k+1} f_{k+2}.$$

This result is the proposition where n=k+1. Therefore, the inductive step is true. Therefore, the above formula is true for all  $n \in \mathbb{N}$  where  $n \ge 0$ .

#### A.3 Direct Proofs

Direct Proofs are the simplest style of proofs, and are especially useful when proving implications. Consider the following examples.

#### Example A.3.1: \* Direct Proof 1

Prove that for all integers n, if n is even, then  $n^2$  is even.

*Proof.* Let  $n \in \mathbb{Z}$  and suppose that n is even. Let  $m \in \mathbb{Z}$ . Thus, n = 2m. Then,  $n^2 = (2m)^2 = 4m^2 = 2(2m^2)$ . Because  $2m^2 \in \mathbb{Z}$ ,  $n^2$  is even.

#### Example A.3.2: \*\* Direct Proof 2

Prove that for all integers a, b, and c, if a|b and b|c, then a|c.

*Proof.* Let  $(a, b, c, p, q, r) \in \mathbb{Z}$  and suppose that a|b and b|c. Because a|b, b = pa. Because b|c, c = qb = pqa. Because c is an integer multiple of a, a|c.

Consider the following exercises.

#### Exercise A.3.1: \*\* Direct Proof 1

Prove that for any two odd integers, their sum is even.

*Proof.* Let  $(m, n) \in \mathbb{Z}$ :  $m \mod 2 \neq 0$ :  $n \mod 2 \neq 0$  and let  $(p, q) \in \mathbb{Z}$ . Because m and n are odd, m = 2p + 1 and n = 2q + 1. Therefore,

$$m + n = (2p + 1) + (2q + 1)$$
  
=  $2p + 2q + 2$   
=  $2(p + q + 1)$ .

Because  $(p+q+1) \in \mathbb{Z}$ , m+n is even.

#### Exercise A.3.2: \*\* Direct Proof 2

Prove that for all integers n, if n is odd, then  $n^2$  is odd.

*Proof.* Let  $n \in \mathbb{Z}$ :  $n \mod 2 \neq 0$  and let  $p \in \mathbb{Z}$ . Because n is odd, n = 2p + 1. Therefore,

$$n^{2} = (2p + 1)^{2}$$
$$= 4p^{2} + 4p + 1$$
$$= 2(2p^{2} + 2p) + 1.$$

Because  $(2p^2 + 2p) \in \mathbb{Z}$ ,  $n^2$  is odd.

#### A.4 Proof by Contrapositive

Recall that for two statements P and Q,  $(P \Longrightarrow Q) \Longleftrightarrow (\neg Q \Longrightarrow \neg P)$ . In a Proof by Contrapositive, we produce a direct proof of the contrapositive of the implication. This is equivalent to proving the implication, because the implication is logically equivalent to the contrapositive. Consider the following examples.

#### Example A.4.1: \* Proof by Contrapositive 1

Prove that for all integers n, if  $n^2$  is even, then n is even.

*Proof.* Let  $n \in \mathbb{Z}$ :  $n \mod 2 \neq 0$ . By Exercise A.3.2,  $n^2$  is odd.

#### Example A.4.2: \* Proof by Contrapositive 2

Prove that for all integers a and b, if a + b is odd, then a is odd or b is odd.

*Proof.* Let  $(a, b, p, q) \in \mathbb{Z}$ . Suppose that a is even and b is even. Then, a = 2p and b = 2q. We see that

$$a+b=2p+2q$$
$$=2(p+q).$$

Because  $(p+q) \in \mathbb{Z}$ , a+b is even.

Consider the following exercises.

#### Exercise A.4.1: \* Proof by Contrapositive 1

Prove that for real numbers a and b, if ab is irrational, then a or b must be an irrational number.

*Proof.* Let  $(p, q, r, s) \in \mathbb{Z}$ . Suppose that  $(a, b) \in \mathbb{Q}$ . Therefore,  $a = \frac{p}{q}$  and  $b = \frac{r}{s}$ . We see that

$$ab = \frac{pr}{qs} \in \mathbb{Q}$$

Therefore ab is rational.

#### Exercise A.4.2: \* Proof by Contrapositive 2

Prove that for integers a and b, if ab is even, then a or b must be even.

*Proof.* Let  $(a, b, p, q) \in \mathbb{Z}$ . Suppose that a = 2p + 1 and b = 2q + 1. We see that

$$ab = (2p+1)(2q+1)$$
  
=  $4pq + 2p + 2q + 1$   
=  $2(2pq + p + q) + 1$ 

Because  $(2pq + p + q) \in \mathbb{Z}$ , ab is odd.

#### Exercise A.4.3: \* Proof by Contrapositive 3

Prove that for any integer a, if  $a^2$  is not divisible by 4, then a is odd.

*Proof.* Let  $a, p \in \mathbb{Z}$ . Suppose that a is even, and a = 2p. Then,  $a^2 = 4p^2$ , and  $4|4p^2$ , so  $4|a^2$ .

#### A.5 Proof by Contradiction

Sometimes, a statement, P cannot be rephrased as an implication. In these cases, it may be useful to prove that  $P \implies Q$ , and also prove that  $P \implies \neg Q$ . Then, we conclude  $\neg P$ . Consider the following example.

#### Example A.5.1: \* Proof by Contradiction 1

Prove that  $\sqrt{2}$  is irrational.

*Proof.* Suppose that  $\sqrt{2}$  is rational. Then,

$$\sqrt{2} = \frac{p}{q}$$

where  $(p,q)\in\mathbb{Z}$  and  $rac{p}{q}$  is in lowest terms. By squaring both sides of the equation, we have

$$2=\frac{p^2}{q^2}.$$

This means that

$$2q^2=p^2,$$

and as  $q^2 \in \mathbb{Z}$ ,  $p^2$  is even, which means that by Example A.4.1, p is even. We see that p=2k for some  $k \in \mathbb{Z}$ . Then, we have

$$2q^2 = (2k)^2 = 4k^2$$

meaning that

$$q^2 = 2k^2$$
.

Therefore, q is even. If p and q are both even,  $\frac{p}{q}$  is not in lowest terms. Therefore,  $\sqrt{2}$  is irrational.  $\Box$ 

#### A.6 Combinatorial Proofs

We start with a visual exercise.

#### Exercise A.6.1: \* Visual Analysis of Pascal's Triangle

Consider the first six rows of Pascal's Triangle, provided below.

Name, without proof, some patterns that are visible.

- Border entries are all 1.
- Non-border entries are the sum of the two entries above.
- The triangle is symmetric.
- The sum of all entries on any row is a power of 2.

Recall that each entry in Pascal's Triangle can be written as a binomial coefficient. The kth entry in the nth row, where both k and n are zero-indexed, is  $\binom{n}{k}$ . That is, the first six rows are

Consider the following exercise.

#### Exercise A.6.2: \*\* \* Rewriting Hypotheses in Pascal's Triangle

Rewrite the patterns discovered in Exercise A.6.1 in terms of binomial coefficients.

- $\bullet \ \binom{n}{0} = \binom{n}{n} = 1.$
- $\bullet \ \binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$
- $\bullet \ \binom{n}{k} = \binom{n}{n-k}.$
- $\bullet \ \binom{n}{0}+\binom{n}{1}+\binom{n}{2}+\cdots+\binom{n}{n}=2^n.$

The above identities were given without proof. To prove the identities, methods of either algebraic or combinatorial proof may be used. While algebraic proof certainly can show that an identity is true, it would not show *why*. Instead, combinatorial proof is based precisely on *why*. In general, combinatorial proofs for binomial identities are in the following format.

- 1. Find a counting problem that can be answered in two ways.
- 2. Explain why one answer to the counting problem is the left hand side of the identity.
- 3. Explain why one answer to the counting problem is the right hand side of the identity.
- 4. Conclude that the left hand side is equal to the right hand side because they are both answers to the same question.

Consider the following example.

#### Example A.6.1: \*\* \* Pascal Proof 1

Prove the binomial identity

$$\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{n} = 2^n.$$

*Proof.* Consider a set S with cardinality n. We note that  $|\mathcal{P}(S)| = 2^n$ , which is precisely the right hand side of the statement. We wish to show that the left hand side also produces the cardinality of  $\mathcal{P}(S)$ . The term  $\binom{n}{0}$  corresponds to the fact that  $\emptyset \in \mathcal{P}(S)$ , and the term  $\binom{n}{n}$  corresponds to the fact that  $S \in \mathcal{P}(S)$ . The quantities  $\binom{n}{0}$  and  $\binom{n}{n}$  together provide 2 sets in  $\mathcal{P}(S)$ . The terms  $\binom{n}{1}$ ,  $\binom{n}{2}$ , ...,  $\binom{n}{n-1}$  correspond to the sets of cardinality  $1, 2, \ldots, n-1$  in  $\mathcal{P}(S)$ . In  $\mathcal{P}(S)$  there are  $\binom{n}{1} = \frac{n!}{1!(n-1)!} = n$  sets of cardinality  $1, \binom{n}{2} = \frac{n!}{2!(n-2)!} = \frac{n(n-1)}{2!}$  sets of cardinality  $1, \binom{n}{2} = \frac{n!}{2!(n-2)!} = \frac{n(n-1)}{2!}$  sets of cardinality  $1, \binom{n}{2} = \frac{n!}{3!} = \frac{n(n-1)(n-2)}{3!}$  sets of cardinality  $1, \binom{n}{2} = \frac{n!}{3!} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n!}{3!} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = \frac{n(n-1)(n-2)}{3!} = n!$  sets of cardinality  $1, \binom{n}{2} = n!$  sets o