A Brief Overview of Brain Tumors and MRI Imaging

Sample document for retrieval-based question answering. Author: Rikdeb Mandal. Licence: Creative Commons Attribution 4.0 International (CC BY 4.0).

The figures in this teaching note are rounded illustrations for a software demo. They are not a substitute for cancer-registry statistics or clinical guidelines.

1. Introduction

Brain tumors are a heterogeneous group of neoplasms that arise from the brain parenchyma, the meninges, or adjacent structures such as the pituitary gland. In routine clinical imaging and in many public MRI datasets used for research, four labels appear again and again: glioma, meningioma, pituitary tumor, and no tumor (a scan without a visible neoplasm).

A glioma is a tumor that originates from glial cells — astrocytes, oligodendrocytes, or ependymal cells. High-grade gliomas, especially glioblastoma, are aggressive, infiltrate surrounding white matter, and are difficult to resect completely. Low-grade gliomas grow more slowly but may still undergo malignant transformation over time.

A meningioma arises from the meninges, the membranous layers that cover the brain and spinal cord. Most meningiomas are extra-axial and histologically benign (WHO grade 1), and they often compress rather than infiltrate the brain.

A pituitary tumor, most commonly a pituitary adenoma, develops in the sellar region. Even when benign, these tumors can affect vision by compressing the optic chiasm and can disrupt hormone production.

The phrase "no tumor" in a classification setting simply means that the MRI study does not show a space-occupying neoplasm of these types. It is not a clinical diagnosis by itself.

2. MRI sequences used to characterise brain tumors

Magnetic resonance imaging is the primary non-invasive modality for characterising brain tumors before surgery. Different pulse sequences highlight different tissue properties, and radiologists usually read them together.

T1-weighted images depict anatomy with high spatial detail. Cerebrospinal fluid appears dark, and gray matter is darker than white matter. After gadolinium contrast is given, a T1-contrast (T1C) sequence shows regions where the blood–brain barrier is disrupted. Many high-grade gliomas and meningiomas enhance, while some low-grade gliomas enhance weakly or not at all.

T2-weighted images show fluid as bright signal. Edema and many tumors appear hyperintense on T2, but the boundary between tumor and edema can be hard to judge on T2 alone.

FLAIR (Fluid-Attenuated Inversion Recovery) suppresses the bright signal of cerebrospinal fluid while leaving other T2-hyperintense tissue visible. FLAIR is particularly useful for detecting edema and lesions that lie next to the ventricles or the cerebral sulci, where bright cerebrospinal fluid would otherwise conceal the finding.

A typical research MRI study for tumor classification therefore includes at least T1, T1-contrast, T2, and FLAIR, although many open datasets also provide only a single sequence or a single two-dimensional slice.

3. Clinical features at a glance

Age and location are useful priors when reading a scan, but they never replace formal radiological and histopathological assessment. For example, a contrast-enhancing extra-axial mass over the convexity in a middle-aged adult is more often a meningioma than a glioma, whereas an irregular, rim-enhancing intra-axial mass with substantial surrounding edema raises concern for glioblastoma.

Tumor type	Typical adult age range	Usual location	Approx. share of primary brain tumors	Preferred first-line treatment (general)
Meningioma	40–70 years	Meninges, often parasagittal or sphenoid wing	about 40%	Surgical resection when indicated
Glioblastoma (high-grade glioma)	45–70 years	Cerebral hemispheres	about 15%	Maximal safe resection, then radiotherapy with temozolomide
Pituitary adenoma	30–60 years	Sellar / suprasellar region	about 15%	Transsphenoidal surgery or medical therapy, depending on type
Low-grade glioma	20–45 years	Cerebral hemispheres, often frontal or temporal	under 15%	Surgery, with radiotherapy or chemotherapy in selected cases

Table 1. Illustrative clinical features of common intracranial tumors. Shares and age ranges are rounded teaching estimates, not registry data.

Two treatment details from the table are worth restating in prose because they are easy to miss when a table is copied poorly. Glioblastoma is generally managed with maximal safe resection followed by radiotherapy with temozolomide. A hormone-secreting pituitary adenoma may be treated medically rather than with immediate surgery, depending on the secretory type.

4. Survival, automated reading, and limits

Survival varies sharply by histology and grade. Glioblastoma still has a markedly poorer prognosis than WHO grade 1 meningioma or most pituitary adenomas, even with current standard therapy. Figure 1 below is a teaching graphic of illustrative five-year survival by tumor type. The exact percentages are shown only in the figure and are not restated in this accompanying text. These teaching figures should not be quoted as official registry statistics.

Rikdeb Mandal, 2026. CC BY 4.0. Teaching sample for pdfqa; not clinical advice.



Figure 1. Illustrative five-year survival by tumor type. Values appear only in the graphic itself.

Automated systems that classify MRI slices into glioma, meningioma, pituitary tumor, or no tumor can be useful as research prototypes. They should not be treated as clinical diagnostic devices. Models trained on two-dimensional slices miss volumetric context, and models trained on a single public dataset may not generalise to scanners, protocols, or populations they have never seen. When such a system is used for teaching or research, an explanation method such as a Grad-CAM heatmap can show which region of the slice influenced the prediction, but a heatmap is not a segmentation of the tumor.

The most common failure modes in automated MRI classification include confusing glioma with meningioma when the mass sits near the cortical surface, missing small pituitary lesions on slices that barely include the sella, and over-calling tumor on scans that actually show post-treatment change or inflammation.

Rikdeb Mandal, 2026. CC BY 4.0. Teaching sample for pdfqa; not clinical advice.