## Bayesian Inference

Oliver Filla

November 20, 2024





### Probability

Determinism and its Limits Frequentist statistics Bayesian statistics

### Conditional Probability

Probability & Likelihood Bayes' Theorem

### Hypothesis Testing

Updating Credibilities
The Posterior Odds ratio

### Bibliography



Why do we need Probabilities?







Why do we need Probabilities?

#### Determinism

Every event is determined by past events.



Why do we need Probabilities?

#### Determinism

- Every event is determined by past events.
- assumed in Classical Mechanics



Why do we need Probabilities?

### Determinism

- Every event is determined by past events.
- assumed in Classical Mechanics
- contradicted in Quantum Mechanics?



Why do we need Probabilities?

#### Determinism

- Every event is determined by past events.
- assumed in Classical Mechanics
- contradicted in Quantum Mechanics?

## Reality

limited knowledge, e.g. friction





Why do we need Probabilities?

#### Determinism

- Every event is determined by past events.
- assumed in Classical Mechanics
- contradicted in Quantum Mechanics?

### Reality

- limited knowledge, e.g. friction
- ▶ limited control over initial state, e.g. in chaotic systems







Why do we need Probabilities?

#### Determinism

- Every event is determined by past events.
- assumed in Classical Mechanics
- contradicted in Quantum Mechanics?

### Reality

- limited knowledge, e.g. friction
- limited control over initial state, e.g. in chaotic systems

Example: Throwing dice: Initial angle and velocity are unknown or difficult to control. It is unknown if the dice are fair.





#### Why do we need Probabilities?

### Determinism

- Every event is determined by past events.
- assumed in Classical Mechanics
- contradicted in Quantum Mechanics?

### Reality

- limited knowledge, e.g. friction
- limited control over initial state, e.g. in chaotic systems

Example: Throwing dice: Initial angle and velocity are unknown or difficult to control. It is unknown if the dice are fair.

⇒ Thus, we need probabilities to predict events!











Probabilities are determined by distributions of random events. They are characterized by the frequency of occurring events.







Probabilities are determined by distributions of random events. They are characterized by the frequency of occurring events.

	1	2	3	4	5	6	$\sum$
frequency	10	5	7	6	9	7	44
probability [%]	23	11	16	14	20	16	100

results of dice rolls; expected PMF  $p(i)=1/6=16.\bar{6}\,\%$ 





Probabilities are determined by distributions of random events. They are characterized by the frequency of occurring events.

	1	l	3	l			$\sum$
frequency	10	5	7	6	9	7	44
frequency probability [%]	23	11	16	14	20	16	100
results of dice rolls	. Avn	ected	IРМ	F n(i	) — 1	/6 —	16 6 %

results of dice rolls; expected PMF p(i) = 1/6 = 16.6%

Probability Mass Function (PMF) (discrete)





Probabilities are determined by distributions of random events. They are characterized by the frequency of occurring events.

	1	l		l	l		$\sum$
frequency	10	5	7	6	9	7	44
frequency probability [%]	23	11	16	14	20	16	100
1 6 1 1 1 1 1 5 6 6 6 7							

results of dice rolls; expected PMF p(i) = 1/6 = 16.6%

- Probability Mass Function (PMF) (discrete)
- Probability Density Function (PDF) (continuous)



Probabilities are determined by distributions of random events. They are characterized by the frequency of occurring events.

	1	2	3	4	5	6	$\sum$
frequency	10	5	7	6	9	7	44
probability [%]	23	11	16	14	20	16	100

results of dice rolls; expected PMF  $p(i) = 1/6 = 16.\overline{6}\%$ 

- Probability Mass Function (PMF) (discrete)
- Probability Density Function (PDF) (continuous)
- Cumulative Distribution Function (CDF) (integrated PDF)





Probabilities are determined by distributions of random events. They are characterized by the frequency of occurring events.

	1	2	3	4	5	6	$\sum$
frequency	10	5	7	6	9	7	44
probability [%]	23	11	16	14	20	16	100

results of dice rolls; expected PMF  $p(i) = 1/6 = 16.\overline{6} \%$ 

- Probability Mass Function (PMF) (discrete)
- Probability Density Function (PDF) (continuous)
- Cumulative Distribution Function (CDF) (integrated PDF)


 $p(6) = \frac{1}{6} \Rightarrow$  I expect that every 6th dice roll shows a 6, on average.













Bayesian statistics describes the credibility of hypotheses.

hypothesis H: falsifiable assumption that a specific statement is true, e.g. it will rain today.



- ▶ hypothesis *H*: falsifiable assumption that a specific statement is true, e.g. *it will rain today.*
- ▶ credibility  $C(H) \in [0,1]$ : strength of belief in H



- ▶ hypothesis *H*: falsifiable assumption that a specific statement is true, e.g. *it will rain today*.
- ▶ credibility  $C(H) \in [0,1]$ : strength of belief in H
- mathematically, credibilities and probabilities are identical
- difference is conceptual





- ▶ hypothesis *H*: falsifiable assumption that a specific statement is true, e.g. *it will rain today*.
- ▶ credibility  $C(H) \in [0,1]$ : strength of belief in H
- mathematically, credibilities and probabilities are identical
- difference is conceptual
- ightharpoonup C(H) can be updated with new data





Bayesian statistics describes the credibility of hypotheses.

- hypothesis H: falsifiable assumption that a specific statement is true, e.g. it will rain today.
- redibility  $C(H) \in [0,1]$ : strength of belief in H
- mathematically, credibilities and probabilities are identical
- difference is conceptual

 $p(6) = \frac{1}{6} \Rightarrow I$  believe the hypothesis H "the die roll shows a 6" with a credibility of  $C(H) = \frac{1}{6}$ .





Bayesian Inference

Oliver Filla











P(A|B) describes the probability of event A with the assumption that event B takes place.

read: P(A given that B)





- ightharpoonup read: P(A given that B)
- filters sets of possible events



- ightharpoonup read: P(A given that B)
- filters sets of possible events
- joint probability: both A and B:  $P(A \cap B) = P(A)P(B|A)$




- ightharpoonup read: P(A given that B)
- filters sets of possible events
- ▶ joint probability: both A and B:  $P(A \cap B) = P(A)P(B|A)$
- $\Rightarrow$  only when A and B independent:  $P(A \cap B) = P(A)P(B)$



- read: P(A given that B)
- filters sets of possible events
- ▶ joint probability: both A and B:  $P(A \cap B) = P(A)P(B|A)$
- $\Rightarrow$  only when A and B independent:  $P(A \cap B) = P(A)P(B)$
- ▶ joint probability: any A or B:




- read: P(A given that B)
- filters sets of possible events
- ightharpoonup joint probability: both A and B:  $P(A \cap B) = P(A)P(B|A)$
- $\Rightarrow$  only when A and B independent:  $P(A \cap B) = P(A)P(B)$
- ▶ joint probability: any A or B:  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$







- ightharpoonup read: P(A given that B)
- filters sets of possible events
- ightharpoonup joint probability: both A and B:  $P(A \cap B) = P(A)P(B|A)$
- $\Rightarrow$  only when A and B independent:  $P(A \cap B) = P(A)P(B)$
- ▶ joint probability: any A or B:  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- $\Rightarrow$  A and B mutually exclusive:  $P(A \cup B) = P(A) + P(B)$







## Probability & Likelihood

Recap from the talk *The Maximum Likelihood Principle* by MAJA KORBMACHER.







## Probability & Likelihood

Recap from the talk *The Maximum Likelihood Principle* by MAJA KORBMACHER.

- Probability P: tool for prediction of events
- Likelihood L: tool for modelling
- equal for given hypothesis H and events E

$$L(H|E) = P(E|H)$$





## Probability & Likelihood

Recap from the talk *The Maximum Likelihood Principle* by MAJA KORBMACHER.

- Probability P: tool for prediction of events
- Likelihood L: tool for modelling
- equal for given hypothesis H and events E
- but only if no prior knowledge is available
- Bayes' theorem generalizes this, using prior knowledge

$$L(H|E) \stackrel{?}{=} P(E|H) \tag{1}$$











- Hypothesis H
- Evidence E





- Hypothesis H
- Evidence E
- ▶ follows from joint probability  $P(E \cap H)$





- Hypothesis H
- Evidence E
- ▶ follows from joint probability  $P(E \cap H)$
- ightharpoonup overall likelihood P(E) (law of total probability)



- ► Hypothesis *H*
- Evidence E
- ▶ follows from joint probability  $P(E \cap H)$
- ightharpoonup overall likelihood P(E) (law of total probability)

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$
 (2)





- Hypothesis H
- Evidence E
- ▶ follows from joint probability  $P(E \cap H)$
- ightharpoonup overall likelihood P(E) (law of total probability)

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$
 (2)

$$\Rightarrow L(H|E) = P(E|H) \cdot \frac{P(H)}{P(E)}$$
 (3)





- Hypothesis H
- Evidence E
- ▶ follows from joint probability  $P(E \cap H)$
- ightharpoonup overall likelihood P(E) (law of total probability)

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$
 (2)

$$\Rightarrow L(H|E) = P(E|H) \cdot \frac{P(H)}{P(E)}$$
 (3)

$$P(E) = P(E|H)P(H) + P(E|\neg H)P(\neg H)$$
 (4)





Geometric Interpretation






Bayesian Inference

Oliver Filla



Geometric Interpretation


square: probability 1



The black square with side length 1 contains all probabilities.

#### Geometric Interpretation

- square: probability 1
- ightharpoonup prior  $P_1(H)$




The black square with side length 1 contains all probabilities.



#### Geometric Interpretation

- square: probability 1
- ightharpoonup prior  $P_1(H)$



The black square with side length 1 contains all probabilities.





with Bayes' Theorem

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$

Bayesian Inference

Oliver Filla



with Bayes' Theorem

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$



with Bayes' Theorem

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$

How to update the credibility of a hypothesis H:

1. assert prior  $P_1(H)$ 





with Bayes' Theorem

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$

- 1. assert prior  $P_1(H)$
- 2. evaluate evidence  $E: \rightarrow \text{likelihood } P(E|H)$



with Bayes' Theorem

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$

- 1. assert prior  $P_1(H)$
- 2. evaluate evidence  $E: \rightarrow \text{likelihood } P(E|H)$
- 3. calculate overall likelihood P(E)





with Bayes' Theorem

$$P_2(H|E) = \frac{P(E|H)P_1(H)}{P(E)}$$

- 1. assert prior  $P_1(H)$
- 2. evaluate evidence  $E: \rightarrow$  likelihood P(E|H)
- 3. calculate overall likelihood P(E)
- 4. calculate posterior probability  $P_2(H|E)$











- relative probability: compare probabilities
- absolute probabilities have limited use







- relative probability: compare probabilities
- absolute probabilities have limited use
- $\rightarrow \frac{P(A)}{P(B)} = \frac{1}{2}$  tells event B occurs twice as often as event A,



- relative probability: compare probabilities
- absolute probabilities have limited use
- $\rightarrow \frac{P(A)}{P(B)} = \frac{1}{2}$  tells event B occurs twice as often as event A, for e.g. P(A) = 0.1, P(B) = 0.2 or P(A) = 0.3, P(B) = 0.6.



#### The Posterior Odds ratio

The Posterior Odds ratio compares the posterior probabilities  $P_2$  for two different hypotheses  $H_{A,B}$ .

- relative probability: compare probabilities
- absolute probabilities have limited use
- $\rightarrow \frac{P(A)}{P(B)} = \frac{1}{2}$  tells event B occurs twice as often as event A, for e.g. P(A) = 0.1, P(B) = 0.2 or P(A) = 0.3, P(B) = 0.6.



#### The Posterior Odds ratio

The Posterior Odds ratio compares the posterior probabilities  $P_2$  for two different hypotheses  $H_{A,B}$ .

- relative probability: compare probabilities
- absolute probabilities have limited use
- $\rightarrow \frac{P(A)}{P(B)} = \frac{1}{2}$  tells event B occurs twice as often as event A, for e.g. P(A) = 0.1, P(B) = 0.2 or P(A) = 0.3, P(B) = 0.6.
- ightharpoonup overall likelihood P(E) becomes irrelevant

$$\frac{P_2(H_A|E)}{P_2(H_B|E)} = \frac{P(E|H_A)}{P(E|H_B)} \frac{P_1(H_A)}{P_1(H_B)}$$
(5)





rapid antigen tests

Assume a COVID rapid test





Oliver Filla



rapid antigen tests

Assume a COVID rapid test with a sensitivity of 80%, causing 20 % false-negative results,





Oliver Filla



rapid antigen tests

Assume a *COVID rapid test* with a sensitivity of 80%, causing 20% false-negative results, and a specificity of 98%, causing 2% false-positive results.





rapid antigen tests

Assume a COVID rapid test with a sensitivity of 80%, causing 20% false-negative results, and a specificity of 98%, causing 2% false-positive results.

My rapid test is positive: How likely do I have COVID?








#### rapid antigen tests

Assume a COVID rapid test with a sensitivity of 80%, causing 20 % false-negative results, and a specificity of 98%, causing 2% false-positive results.



Source: Robert Koch Institute





# **Bibliography**

- [1] Lawrence A. (2019) Probability in Physics: An Introductory Guide, Springer Nature Switzerland, Chapters 1 & 7.1-7.5, DOI 10.1007/978-3-030-04544-9
- [2] Clyde M. et al. (2022) An Introduction to Bayesian Thinking: A Companion to the Statistics with R Course, Github.com/enstarprise, on 2022-06-15
- [3] 3Blue1Brown (2020) Bayes theorem, the geometry of changing beliefs, YouTube.com/watch?v=HZGCoVF3YvM, 2024-11-16
- [4] Kozyrkov C. (2020) Are you Bayesian or Frequentist? YouTube.com/watch?v=GEFxFVESQXc, 2024-11-16







# Thank you for your attention!



