

On twisted sums of random multiplicative functions (Warwick)

Ofir Gorodetsky

1 Motivation, history

Let $\alpha_S: \mathbb{N} \rightarrow S^1$ be Steinhaus (completely multiplicative), $\alpha_S: \mathbb{N} \rightarrow \{-1, 0, 1\}$ be Rademacher.

Original motivation: α_R models μ . RH is $\sum_{n \leq x} \mu(n) \ll x^{1/2+\varepsilon}$. Wintner (1944) proved $\sum_{n \leq x} \alpha_R(n) \ll x^{1/2+\varepsilon}$ holds almost surely **for all** x .

(Non-proof: Since $\mathbb{E} \alpha_R(n) \alpha_R(m) = \delta_{n,m}$ for squarefree n, m , we have $\mathbb{E}[(\sum_{n \leq x} \alpha_R(n))^2] \ll x$, so $\mathbb{P}(|\sum_{n \leq x} \alpha_R(n)| > Tx^{1/2}) \ll 1/T^2$.)

It is expected that α_R misses some properties of μ . E.g. $\sum_{n \leq x} \mu(n) = O(\sqrt{x}(\log \log \log x)^{5/4})$ is expected (Gonek) while Harper proved that if $V(x) \rightarrow \infty$ then $\sum_{n \leq x} \alpha_R(n) \geq \sqrt{x}(\log \log x)^{1/4}/V(x)$ for infinitely many x a.s. On the other hand, α_S is the large- T limit of $(n^{it})_{t \leq T}$ and the large- q limit of $(\chi \bmod q)$, e.g.

$$\sum_{n \leq x} n^{it} \xrightarrow{d} \sum_{n \leq x} \alpha_S(n)$$

as $T \rightarrow \infty$ (since $(p^{it})_{p \leq x} \xrightarrow{d} (\alpha_S(p))_{p \leq x}$; here t is uniform in $[0, T]$). And we certainly want to understand such sums – they build up ζ .

Significant work on moments of $\sum_{n \leq x} \alpha_S(n)$ and law of iterated logarithm (Halász, Erdős, Lau-Tenenbaum-Wu, Basquin, Harper, Caich). Focus of this talk: distribution.

The distribution of $\sum_{n \leq x} \alpha_S(n)$, appropriately normalized is still not known, but very recently a precise conjecture was stated (2024). Harper's work (2017) shows that normalizing by standard deviation gives trivial limiting distribution (Helson), and that 'true' normalization is by $\sqrt{x}/(\log \log x)^{1/4}$. This relates to critical multiplicative chaos. Lots of activity on variants:

$$\sum_{n \leq x} \alpha_S(n) f(n).$$

E.g. very recently (2025), Hardy proved a version of the conjecture where $f(n) = \mathbf{1}_{P(n) > \sqrt{n}}$. Most works, until Hardy's, focused on much 'sparser' f , where correct normalization turns out to be **standard deviation**, and limiting distribution was **standard Gaussian**:

1. $f(n) = \mathbf{1}_{\omega(n)=k}$: Hough, Harper. $k = o(\log \log x)$. This is roughly $x/\log^{1+o(1)} x$.
2. $f(n) = \mathbf{1}_{[x, x+H]}$: Chatterjee-Sound ($H = o(x/\log x)$), Sound-Xu ($H \leq x/(\log x)^{2 \log 2 - 1 + \varepsilon}$), Pandey-Wang-Xu ($H \ll_A x/\log^A x$ for every $A > 0$), Harper-Sound-Xu.
3. $f(n) = \mathbf{1}_{Q(\mathbb{Z})}$: Najnudel, Klurman-Shkredov-Xu, Wang-Xu, Chinis-Shala.

None of these f are multiplicative.

2 Work with Wong

Motivated by works of Najnudel-Paquette-Sim, Mo Dick Wong and I considered

$$S_x := \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1/2} \sum_{n \leq x} \alpha_S(n) f(n)$$

for multiplicative f , with $|f(p)|^2$ equal to $\theta \in (0, 1/2)$ on average; recently extended to $|f(p)|^2$ equal to $\theta \in (0, 1)$ on average, inspired by ideas in Najnudel-Paquette-Simm-Vu. So cannot cover $f \equiv 1$, but can take e.g. indicator

of sums of two squares. In fact, f does not have to be bounded, $f(p)$ can infinitely often be as large as $p^{1/2+o(1)}$ ($\sum_p |f(p)|^3 \log^C p / p^{3/2} < \infty$). If f takes the values 0 and 1 only, its support is $\asymp x(\log x)^{\theta-1}$

What is the limiting distribution we found? $G \cdot \sqrt{V}$ for standard complex Gaussian G independent of V , and $V = 1/(2\pi) \int_{\mathbb{R}} |1/2 + it|^{-2} m_{\infty}(dt)$. Here m_{∞} is a random measure, translation-invariant, constructed by a limit from a sequence of measures involving α and f . We use the notation G for standard complex Gaussian throughout. This talk will not be about m_{∞} (see Mo Dick's talk for that), but about **how one connects the random sum with the random measure**.

One approach for this is through moments. It is not relevant here because the moments of V explode: $\mathbb{E}V^p$ is finite iff $p < 1/\theta$, which forces $\mathbb{E}|S_x|^{2p}$ to diverge for $p \geq 1/\theta$ (recall that if $X_n \xrightarrow{d} X$ in distribution then $\liminf \mathbb{E}|X_n|^{2p} \geq \mathbb{E}|X|^{2p}$ by Fatou's lemma.)

Next approach is martingale CLT, introduced into this area by Harper (2010). Why martingales?

$$S_x = \sum_{p \leq x} Z_{p,x}$$

where

$$Z_{p,x} := \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1/2} \sum_{n \leq x, P(n)=p} \alpha_S(n) f(n).$$

If we define filtrations $\mathcal{F}_{p-} = \sigma((\alpha(q) : q < p))$ then $\mathbb{E}[Z_{p,x} | \mathcal{F}_{p-}] = 0$ (Essentially conditioning of values of $\alpha(q)$ for $q < p$.) Let

$$V_x = \sum_{p \leq x} |Z_{p,x}|^2.$$

Informally, $S_x \approx G\sqrt{V_x}$. Formally, **McLeish CLT** makes this formal in a special case: if $\mathbb{E}V_x \rightarrow 1$, $\mathbb{E}V_x^2 \rightarrow 1$, and $\sum_{p \leq x} \mathbb{E}|Z_{p,x}|^4 \rightarrow 0$, then $S_x \xrightarrow{d} G$. Note that these conditions imply $V_x \xrightarrow{p} 1$. First condition is trivial: $\mathbb{E}V_x = 1$ regardless of f . (Recent restatement by Sound–Xu: suffices to check $\max_p \mathbb{E}|Z_{p,x}|^2 \rightarrow 0$ and $\mathbb{E}|S_x|^4 \rightarrow 2$, if f takes the values 0 and 1.)

This CLT is not relevant if limiting distribution is not Gaussian, so we turn to the general form of martingale CLT. We define

$$BP_x := \sum_{p \leq x} \mathbb{E}[|Z_{p,x}|^2 | \mathcal{F}_{p-}].$$

If $BP_x \xrightarrow{p} V$, $\sum_{p \leq x} \mathbb{E}|Z_{p,x}|^4 \rightarrow 0$ and $\sum_{p \leq x} \mathbb{E}[Z_{p,x}^2 | \mathcal{F}_{p-}] \xrightarrow{d} 0$, then $S_x \xrightarrow{d} G\sqrt{V}$. Rest of the talk will be about $BP_x \xrightarrow{p} V$ – other conditions are ‘trivial’ to verify, *even in critical case* (e.g. $\sum_{p \leq x} \mathbb{E}[Z_{p,x}^2 | \mathcal{F}_{p-}] \equiv 0$).

3 Two approaches to bracket process

It is convenient to assume that f is supported on squarefrees. Moreover, we shall only talk about Steinhaus (we put $\alpha = \alpha_S$). We suppose $\sum_{p \leq x} |f(p)|^2 \sim \theta \text{Li}(x)$. First step is to understand how BP_x looks like. Note that

$$Z_{p,x} = \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1/2} \alpha(p) f(p) \sum_{m \leq x/p, P(m) < p} \alpha(m) f(m)$$

so

$$\mathbb{E}[|Z_{p,x}|^2 | \mathcal{F}_{p-}] = \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1} |f(p)|^2 \sum_{m \leq x/p, P(m) < p} \alpha(m) f(m)^2.$$

This shows

$$BP_x = \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1} \sum_{p \leq x} |f(p)|^2 \sum_{m \leq x/p, P(m) < p} \alpha(m) f(m)^2.$$

We introduce

$$s_{t,x} := t^{-1/2} \sum_{n \leq t, P(n) < x} \alpha(n) f(n)$$

so that

$$BP_x = \frac{x}{\sum_{n \leq x} |f(n)|^2} \sum_{p \leq x} \frac{|f(p)|^2}{p} |s_{x/p,p}|^2.$$

It is useful to massage BP_x a little: since $F(x) = \sum_{p \leq x} |f(p)|^2/p \sim \theta \log \log x$ then, informally,

$$BP_x \approx \frac{x}{\sum_{n \leq x} |f(n)|^2} \int_{2^-}^x |s_{x/t,t}|^2 dF(t) \sim \theta \frac{x}{\sum_{n \leq x} |f(n)|^2} \int_{2^-}^x \frac{|s_{x/t,t}|^2}{t \log t} dt. \quad (1)$$

There are now (at least) two rather different approaches relating this to the measure m_∞ . Both approaches employ the following form of Plancherel's identity:

$$\int_{\mathbb{R}} \left| \sum_{n \leq t} g(n) \right|^2 t^{-2-2r} dt = \frac{1}{2\pi} \int_{\mathbb{R}} \left| \sum_n g(n) n^{-1/2-r-it} \right|^2 |1/2 + r + it|^{-2} dt \quad (2)$$

holds for $r > 0$, a.s. Proof: Through Perron.

First argument. We introduce

$$U_x := \left(\sum_{n \leq x} |f(n)|^2/n \right)^{-1} \int_1^x |s_{t,\infty}|^2 \frac{dt}{t}.$$

We combine two lemmas:

$$U_x \xrightarrow{p} \frac{1}{2\pi} \int_{\mathbb{R}} |1/2 + it|^{-2} m_\infty(dt) \quad (3)$$

and

$$\mathbb{E}|U_x - BP_x|^2 \rightarrow 0 \quad (4)$$

The statement (4) is proved by a straightforward technical computation, but isn't deep. The statement (3) is also not too deep (given the construction(s) of m_∞). Perhaps the difficult part is *guessing* that $BP_x \xrightarrow{p} \frac{1}{2\pi} \int_{\mathbb{R}} |1/2 + it|^{-2} m_\infty(dt)$, which is motivated by Plancherel.

To prove (4) one expands the square and computes three expectation using $\mathbb{E}\alpha(n)\overline{\alpha(m)} = \delta_{n,m}$. This leads to summing solutions to $ab = cd$ with certain weights and constraints. Here it is useful to mention some facts:

- Solutions to $ab = cd$ are given by $a = n_1 n_2$, $b = n_3 n_4$, $c = n_1 n_3$, $d = n_2 n_4$. Proof: let $g_1 = (a, c)$ $g_2 = (b, d)$ and define a', b', c', d' accordingly to get $a'b' = c'd'$ with $(a', c') = 1$ and $(b', d') = 1$. Then $a' = d'$ and $b' = c'$ forcing $a = g_1 a'$, $b = g_2 b'$, $c = g_1 b'$, $d = g_2 a'$.
- De Bruijn-van Lint: $\sum_{n \leq x, P(n) \leq y} |f(n)|^2 \sim \sum_{n \leq x} |f(n)|^2 \cdot \rho_\theta(\frac{\log x}{\log y})$ holds in the regime $\log y \asymp \log x$.
- Wirsing: $\sum_{n \leq x} |f(n)|^2 \sim C_f x (\log x)^{\theta-1}$ if $|f(p)|^2 \sim \theta$ on average.

Let us expand on (3). Notation:

$$m_x(dt) = |A(1/2 + 1/(2 \log x) + it)|^2 dt / \mathbb{E}|A(1/2 + 1/(2 \log x) + it)|^2$$

where $A(s) = \sum_n f(n)/n^s$. The measure $m_\infty(dt)$ is constructed by taking $\lim_x m_x(dt)$. Note that

$$\mathbb{E}|A(1/2 + r/2 + it)|^2 = \mathbb{E} \left| \sum_n \alpha(n) f(n) / n^{1/2+r/2+it} \right|^2 = \sum_n |f(n)|^2 / n^{1+r}$$

is independent of t . So, for $r = 1/\log x$, Plancherel implies that

$$\begin{aligned} \int_{\mathbb{R}} |s_{t,\infty}|^2 t^{-1-1/\log x} dt &= \frac{1}{2\pi} \int_{\mathbb{R}} |A(1/2 + 1/(2 \log x) + it)|^2 |1/2 + 1/2(1/\log x) + it|^{-2} dt \\ &= \frac{1}{2\pi} \left(\sum_n |f(n)|^2 / n^{1+1/\log x} \right) \int_{\mathbb{R}} |1/2 + 1/(2 \log x) + it|^{-2} m_x(dt). \end{aligned}$$

From Wirsing, $\sum_n |f(n)|^2 / n^{1+1/\log x} \sim C_f \Gamma(\theta) (\log x)^\theta$. From consequences of Plancherel and Wirsing,

$$(C_f \Gamma(\theta) (\log x)^\theta)^{-1} \int_{\mathbb{R}} |s_{t,\infty}|^2 t^{-1-1/\log x} dt \xrightarrow{p} \frac{1}{2\pi} \int_{\mathbb{R}} |1/2 + it|^{-2} m_\infty(dt).$$

(Here we relied on $m_x(I) \xrightarrow{p} m_\infty(I)$.) By Tauberian theorem (basically, approximating $\mathbf{1}_{t \leq x}$ by $P(t^{-1/\log x})$), this implies that

$$(C_f \Gamma(\theta) (\log x)^\theta)^{-1} \int_1^x |s_{t,\infty}|^2 \frac{dt}{t} \xrightarrow{p} \frac{1}{\Gamma(1+\theta)} \frac{1}{2\pi} \int_{\mathbb{R}} |1/2 + it|^{-2} m_\infty(dt)$$

as needed. (In U_x we divide by $\sum_{n \leq x} |f(n)|^2/n$, which is asymptotic to $C_f(\log x)^\theta/\theta$ by Wirsing.)

Second argument. Some notation:

$$m_{y,x}(dt) = |A_y(1/2 + it + 1/(2 \log x))|^2 dt / \mathbb{E} |A_y(1/2 + it + 1/(2 \log x))|^2$$

where $A_y(s) = \sum_{P(n) < y} f(n)/n^s$. Recall (1). From Plancherel's identity, we may obtain the limit of a somewhat similar expression, namely

$$x \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1} \int_0^\infty q(t^{1/\log x}) \frac{|s_{x/t, x^a}|^2 dt}{t \log x} \quad (5)$$

for any fixed $a > 0$ and any 'nice' function q . Details: Plancherel implies that

$$\begin{aligned} \int_{\mathbb{R}} |s_{t,y}|^2 t^{-1 - \frac{r}{\log y}} dt &= \frac{1}{2\pi} \int_{\mathbb{R}} |A_y(\frac{1}{2} + \frac{r}{2 \log y} + it)|^2 |\frac{1}{2} + \frac{r}{2 \log y} + it|^{-2} dt \\ &= \mathbb{E} |A_y(\frac{1}{2} + \frac{r}{2 \log y})|^2 \cdot \frac{1}{2\pi} \int_{\mathbb{R}} |\frac{1}{2} + \frac{r}{2 \log y} + it|^{-2} m_{y,y^{1/r}}(dt) \end{aligned}$$

holds for $r > 0$ so

$$\left(\sum_{P(n) < y} \frac{|f(n)|^2}{n^{1+r/\log y}} \right)^{-1} \int_0^\infty \frac{|s_{t,y}|^2 dt}{t^{1+r/\log y}} \xrightarrow{y \rightarrow \infty} V;$$

now substitute $t = x/t$ and $y = x^a$. This gives (5) with $q(z) = z^{r/a}$. Here we rely on $m_{x,y}(I) \xrightarrow{p} m_\infty(I)$ if $x, y \rightarrow \infty$ together. The main difference between (1) and (5), however, is that in (5) the 'smoothness parameter' in $s_{x/t, x^a}$, namely x^a , does not depend on the integration variable t , while in (1) the smoothness parameter in $s_{x/t, t}$ is t , the integration variable itself.

To circumvent this issue, we modify S_x . We divide the primes in $[2, x]$ into finitely many disjoint intervals $(I_k)_k$, and if $n \leq x$ has $P(n) \in I_k$, we 'keep' this n in the modified version of S_x only if $P(n/P(n))$ (the second largest prime factor of n) is smaller than $\min I_k$. In this way, the new S_x is

$$S'_x = \sum_{p \leq x} Z'_{p,x}$$

where, if $p \in I_k$, then

$$Z'_{p,x} := \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1/2} \sum_{n \leq x, P(n)=p, P(n/P(n)) < \min I_k} \alpha(n) f(n)$$

and the new bracket process takes the shape

$$BP'_x \approx x \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1} \sum_k \sum_{p \in I_k} \frac{|f(p)|^2}{p} |s_{x/p, \min I_k}|^2 \approx x \left(\sum_{n \leq x} |f(n)|^2 \right)^{-1} \sum_k \int_2^x \frac{|s_{x/t, \min I_k}|^2 dt}{t \log t}.$$

For fixed k , the smoothness parameter is now fixed within the integral, namely it is $\min I_k$. This allows us to handle the k th integral using (5). One has to justify working with this modified S_x . The idea is that the second moment of the discarded terms is

$$\sum_k \frac{\sum_{n \leq x: P(n), P(n/P(n)) \in I_k} |f(n)|^2}{\sum_{n \leq x} |f(n)|^2} \ll \sum_k \sum_{p, q \in I_k} \frac{1}{pq} \ll \sum_k (\log(\log \max I_k / \log \min I_k))^2.$$

We discard also $P(n) \leq x^\varepsilon$ (we lose $O(\varepsilon)$) and then take $I_k = [x^{\varepsilon+\delta k}, x^{\varepsilon+\delta(k+1)}]$, so that the summand is $\ll \log^2((\varepsilon + \delta(k+1))/(\varepsilon + \delta k)) \ll \min\{\delta/\varepsilon, 1/k\}^2$ and this is good enough; the total loss is $O_\varepsilon(\delta) + O(\varepsilon)$ and this is manageable if we take δ to 0 and then ε to 0.