



Secular coefficients mini-course

Talk 2

Ofir Gorodetsky, Technion

`ofir-gorodetsky.github.io/Bristol`

CMI-HIMR Summer School, Bristol
June 2, 2026

Recall the analogy:

$$L(s, \chi) \text{ (for random } \chi) \text{ or } \zeta(s + it) \text{ (random } t)$$

are analogous to

$$\det(I - uU_N),$$

the characteristic polynomial of a random unitary matrix U_N chosen according to Haar measure.

Recall the analogy:

$$L(s, \chi) \text{ (for random } \chi) \text{ or } \zeta(s + it) \text{ (random } t)$$

are analogous to

$$\det(I - uU_N),$$

the characteristic polynomial of a random unitary matrix U_N chosen according to Haar measure.

The coefficients of $\det(I - uU_N)$ are called *secular coefficients*:

$$\det(I - uU_N) = \sum_{i=0}^N (-1)^i \text{Sc}_i(U_N) u^i.$$

What is their distribution?

From the identity

$$\det(I - uU_N) = \exp \left(- \sum_{n=1}^{\infty} \frac{\text{Tr}(U_N^n) u^n}{n} \right)$$

it follows that $\text{Sc}_i(U_N)$ is a polynomial of total degree i in $(\text{Tr}(U_N^n))_{n \leq i}$.

From the identity

$$\det(I - uU_N) = \exp \left(- \sum_{n=1}^{\infty} \frac{\text{Tr}(U_N^n) u^n}{n} \right)$$

it follows that $\text{Sc}_i(U_N)$ is a polynomial of total degree i in $(\text{Tr}(U_N^n))_{n \leq i}$.

Since the tuple $(\text{Tr}(U_N^n))_{n \leq i}$ tends to a tuple of independent Gaussians, $\text{Sc}_i(U_N)$ tends to a degree- i polynomial in independent Gaussians.

From the identity

$$\det(I - uU_N) = \exp \left(- \sum_{n=1}^{\infty} \frac{\text{Tr}(U_N^n) u^n}{n} \right)$$

it follows that $\text{Sc}_i(U_N)$ is a polynomial of total degree i in $(\text{Tr}(U_N^n))_{n \leq i}$.

Since the tuple $(\text{Tr}(U_N^n))_{n \leq i}$ tends to a tuple of independent Gaussians, $\text{Sc}_i(U_N)$ tends to a degree- i polynomial in independent Gaussians.

Harder to study when i is growing.

Is there a number-theoretic counterpart for $Sc_i(U_N)$?

Is there a number-theoretic counterpart for $\text{Sc}_i(U_N)$? Yes:

$$\frac{1}{\sqrt{x}} \sum_{n \leq x} \chi(n) \text{ or } \frac{1}{\sqrt{x}} \sum_{n \leq x} n^{it},$$

for random χ or t . These are important and well-studied objects in number theory ('character sums'), which implies secular coefficients are worth studying.

Is there a number-theoretic counterpart for $\text{Sc}_i(U_N)$? Yes:

$$\frac{1}{\sqrt{x}} \sum_{n \leq x} \chi(n) \text{ or } \frac{1}{\sqrt{x}} \sum_{n \leq x} n^{it},$$

for random χ or t . These are important and well-studied objects in number theory ('character sums'), which implies secular coefficients are worth studying.

For instance: the Lindelöf hypothesis states that $\zeta(1/2 + it) \ll_{\epsilon} (1 + |t|)^{\epsilon}$.

Is there a number-theoretic counterpart for $\text{Sc}_i(U_N)$? Yes:

$$\frac{1}{\sqrt{x}} \sum_{n \leq x} \chi(n) \text{ or } \frac{1}{\sqrt{x}} \sum_{n \leq x} n^{it},$$

for random χ or t . These are important and well-studied objects in number theory ('character sums'), which implies secular coefficients are worth studying.

For instance: the Lindelöf hypothesis states that $\zeta(1/2 + it) \ll_{\varepsilon} (1 + |t|)^{\varepsilon}$. Equivalent to

$$\sum_{n \leq x} n^{it} = \int_1^x u^{it} du + O_{\varepsilon}(\sqrt{x}(x(1 + |t|))^{\varepsilon}).$$

The analogy between character sums and secular coefficients is more convincing if one considers polynomials over finite fields: recall

$$\mathcal{L}(u, \chi) = \sum_{f \text{ monic}} \chi(f) u^{\deg f}$$

is a polynomial, and $\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$ for some unitary matrix (that equidistributes as χ varies, by Katz).

The analogy between character sums and secular coefficients is more convincing if one considers polynomials over finite fields: recall

$$\mathcal{L}(u, \chi) = \sum_{f \text{ monic}} \chi(f) u^{\deg f}$$

is a polynomial, and $\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$ for some unitary matrix (that equidistributes as χ varies, by Katz).

The i th coefficient of $\mathcal{L}(u/\sqrt{q}, \chi)$ is precisely a normalized character sum:

$$q^{-i/2} \sum_{\deg f=i} \chi(f).$$

The analogy between character sums and secular coefficients is more convincing if one considers polynomials over finite fields: recall

$$\mathcal{L}(u, \chi) = \sum_{f \text{ monic}} \chi(f) u^{\deg f}$$

is a polynomial, and $\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$ for some unitary matrix (that equidistributes as χ varies, by Katz).

The i th coefficient of $\mathcal{L}(u/\sqrt{q}, \chi)$ is precisely a normalized character sum:

$$q^{-i/2} \sum_{\deg f=i} \chi(f).$$

For fixed i , this is a polynomial in $(\sum_{\deg f=n} \chi(f) \Lambda(f))_{n \leq i}$ which tends to a tuple of independent random variables (when χ is sampled uniformly modulo M with $\deg M \rightarrow \infty$).

Recall $\det(I - uU_N)$ tends to

$$F(u) := \exp \left(- \sum_{i=1}^{\infty} N_i \frac{u^i}{\sqrt{i}} \right),$$

where $(N_i)_i$ are iid standard complex Gaussians.

Recall $\det(I - uU_N)$ tends to

$$F(u) := \exp \left(- \sum_{i=1}^{\infty} N_i \frac{u^i}{\sqrt{i}} \right),$$

where $(N_i)_i$ are iid standard complex Gaussians.

Studying coefficients of $F(u)$ should be easier than studying secular coefficients.

Recall $\det(I - uU_N)$ tends to

$$F(u) := \exp \left(- \sum_{i=1}^{\infty} N_i \frac{u^i}{\sqrt{i}} \right),$$

where $(N_i)_i$ are iid standard complex Gaussians.

Studying coefficients of $F(u)$ should be easier than studying secular coefficients. We write

$$F(u) = \sum_{i=0}^{\infty} c_i u^i.$$

We want to compute $\mathbb{E}[|c_n|^2]$.

We want to compute $\mathbb{E}[|c_n|^2]$. By independence of the N_i ,

$$\begin{aligned}\mathbb{E}[F(u)\overline{F}(z)] &= \mathbb{E} \exp \left(- \sum_{i=1}^{\infty} N_i \frac{u^i}{\sqrt{i}} - \overline{N}_i \frac{z^i}{\sqrt{i}} \right) \\ &= \prod_{i=1}^{\infty} \mathbb{E} \exp \left(- N_i \frac{u^i}{\sqrt{i}} - \overline{N}_i \frac{z^i}{\sqrt{i}} \right) = \prod_{i=1}^{\infty} \exp \left(\frac{z^i u^i}{i} \right) \\ &= \exp(-\log(1 - zu)) \\ &= \frac{1}{1 - zu} = 1 + zu + (zu)^2 + \dots\end{aligned}$$

We want to compute $\mathbb{E}[|c_n|^2]$. By independence of the N_i ,

$$\begin{aligned}\mathbb{E}[F(u)\overline{F}(z)] &= \mathbb{E} \exp \left(- \sum_{i=1}^{\infty} N_i \frac{u^i}{\sqrt{i}} - \overline{N}_i \frac{z^i}{\sqrt{i}} \right) \\ &= \prod_{i=1}^{\infty} \mathbb{E} \exp \left(- N_i \frac{u^i}{\sqrt{i}} - \overline{N}_i \frac{z^i}{\sqrt{i}} \right) = \prod_{i=1}^{\infty} \exp \left(\frac{z^i u^i}{i} \right) \\ &= \exp(-\log(1 - zu)) \\ &= \frac{1}{1 - zu} = 1 + zu + (zu)^2 + \dots\end{aligned}$$

Comparing coefficients:

$$\mathbb{E}|c_n|^2 = \mathbb{E}[c_n \overline{c}_n] = 1.$$

We want to compute $\mathbb{E}[|c_n|^2]$. By independence of the N_i ,

$$\begin{aligned}\mathbb{E}[F(u)\overline{F}(z)] &= \mathbb{E} \exp \left(- \sum_{i=1}^{\infty} N_i \frac{u^i}{\sqrt{i}} - \overline{N}_i \frac{z^i}{\sqrt{i}} \right) \\ &= \prod_{i=1}^{\infty} \mathbb{E} \exp \left(- N_i \frac{u^i}{\sqrt{i}} - \overline{N}_i \frac{z^i}{\sqrt{i}} \right) = \prod_{i=1}^{\infty} \exp \left(\frac{z^i u^i}{i} \right) \\ &= \exp(-\log(1 - zu)) \\ &= \frac{1}{1 - zu} = 1 + zu + (zu)^2 + \dots\end{aligned}$$

Comparing coefficients:

$$\mathbb{E}|c_n|^2 = \mathbb{E}[c_n \overline{c}_n] = 1.$$

More generally, $\mathbb{E}[c_n \overline{c}_m] = \delta_{n,m}$.

Similarly, instead of studying $\frac{1}{\sqrt{x}} \sum_{n \leq x} \chi(n)$ for random χ we may want to begin with the easier question of studying

$$\frac{1}{\sqrt{x}} \sum_{n \leq x} \alpha(n)$$

where α is the Steinhaus random multiplicative function:
 $(\alpha(p))_p$ are iid.

We have perfect orthogonality for α (exercise):

$$\mathbb{E}[\alpha(n)\overline{\alpha}(m)] = \delta_{n,m}.$$

We have perfect orthogonality for α (exercise):

$$\mathbb{E}[\alpha(n)\overline{\alpha}(m)] = \delta_{n,m}.$$

This implies

$$\mathbb{E}\left|\sum_{n \leq x} \alpha(n)\right|^2 = \lfloor x \rfloor,$$

so by Cauchy–Schwarz, $\mathbb{E}\left|\sum_{n \leq x} \alpha(n)\right| \leq \sqrt{x}$.

We have perfect orthogonality for α (exercise):

$$\mathbb{E}[\alpha(n)\overline{\alpha}(m)] = \delta_{n,m}.$$

This implies

$$\mathbb{E}\left|\sum_{n \leq x} \alpha(n)\right|^2 = \lfloor x \rfloor,$$

so by Cauchy–Schwarz, $\mathbb{E}\left|\sum_{n \leq x} \alpha(n)\right| \leq \sqrt{x}$. A celebrated result of Harper says that the correct size of the first moment is smaller than that.

We have perfect orthogonality for α (exercise):

$$\mathbb{E}[\alpha(n)\overline{\alpha}(m)] = \delta_{n,m}.$$

This implies

$$\mathbb{E}\left|\sum_{n \leq x} \alpha(n)\right|^2 = \lfloor x \rfloor,$$

so by Cauchy–Schwarz, $\mathbb{E}\left|\sum_{n \leq x} \alpha(n)\right| \leq \sqrt{x}$. A celebrated result of Harper says that the correct size of the first moment is smaller than that.

Theorem 1 (Harper, 2017)

We have

$$\frac{c\sqrt{x}}{(\log \log x)^{1/4}} \leq \mathbb{E}\left|\sum_{n \leq x} \alpha(n)\right| \leq \frac{C\sqrt{x}}{(\log \log x)^{1/4}}.$$

More recently, Harper has deduced a 'deterministic' consequence.

More recently, Harper has deduced a ‘deterministic’ consequence.

Theorem 2 (Harper, 2023)

Let m be a prime. Uniformly for $m \geq x \geq 1$,

$$\frac{1}{m-1} \sum_{\chi \bmod m} \left| \frac{1}{\sqrt{x}} \sum_{n \leq x} \chi(n) \right| \leq \frac{C}{(\log \log \min\{x, m/x\})^{1/4}}.$$

Uniformly for $T \geq x \geq 1$,

$$\frac{1}{T} \int_0^T \left| \frac{1}{\sqrt{x}} \sum_{n \leq x} n^{it} \right| dt \leq \frac{C}{(\log \log \min\{x, T/x\})^{1/4}}.$$

For c_i , Najnudel, Paquette and Simm proved the following results.

Theorem 3 (Najnudel–Paquette–Simm, 2020)

We have

$$\mathbb{E} |c_n| \leq \frac{C}{(\log n)^{1/4}}.$$

Theorem 4 (Najnudel–Paquette–Simm, 2020)

For $0 \leq n \leq N$,

$$\mathbb{E} |\text{Sc}_n(U_N)| \leq \frac{C}{(\log \min\{n, N - n\})^{1/4}}.$$

For c_i , Najnudel, Paquette and Simm proved the following results.

Theorem 3 (Najnudel–Paquette–Simm, 2020)

We have

$$\mathbb{E} |c_n| \leq \frac{C}{(\log n)^{1/4}}.$$

Theorem 4 (Najnudel–Paquette–Simm, 2020)

For $0 \leq n \leq N$,

$$\mathbb{E} |\mathrm{Sc}_n(U_N)| \leq \frac{C}{(\log \min\{n, N - n\})^{1/4}}.$$

Exercise: $|\mathrm{Sc}_n(U_N)|$ and $|\mathrm{Sc}_{N-n}(U_N)|$ have the same distribution.

The proof of Najnudel–Paquette–Simm shares similarities to Harper's, but there are differences as well.

The proof of Najnudel–Paquette–Simm shares similarities to Harper's, but there are differences as well.

Soundararajan and Zaman (2021) proved $\frac{c}{(\log n)^{1/4}} \leq \mathbb{E} |c_n| \leq \frac{C}{(\log n)^{1/4}}$, with a proof mirroring Harper closely.

Recently there was progress on the limiting distribution of

$$\frac{1}{\sqrt{x}(\log \log x)^{-1/4}} \sum_{n \leq x} \alpha(n)$$

by Gorodetsky–Wong, and on the limiting distribution of

$$(\log n)^{1/4} c_n$$

by Atherfold–Najnudel. More on that in the next lecture.

The simplest approach to proving that a sequence of random variables X_n tends in distribution to X is the method of moments. For real random variables it says: if

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^k] = \mathbb{E}[X^k]$$

holds for every positive integer k , and if $\sum_{k \geq 1} \mathbb{E}[X^{2k}]^{-1/(2k)}$ diverges, then $X_n \xrightarrow{d} X$.

The simplest approach to proving that a sequence of random variables X_n tends in distribution to X is the method of moments. For real random variables it says: if

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^k] = \mathbb{E}[X^k]$$

holds for every positive integer k , and if $\sum_{k \geq 1} \mathbb{E}[X^{2k}]^{-1/(2k)}$ diverges, then $X_n \xrightarrow{d} X$.

Example: if $X \sim N(0, 1)$ then $\mathbb{E}[X^{2k}]^{-1/(2k)} \sim c/\sqrt{k}$.

The simplest approach to proving that a sequence of random variables X_n tends in distribution to X is the method of moments. For real random variables it says: if

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^k] = \mathbb{E}[X^k]$$

holds for every positive integer k , and if $\sum_{k \geq 1} \mathbb{E}[X^{2k}]^{-1/(2k)}$ diverges, then $X_n \xrightarrow{d} X$.

Example: if $X \sim N(0, 1)$ then $\mathbb{E}[X^{2k}]^{-1/(2k)} \sim c/\sqrt{k}$.

For both $\sum_{n \leq x} \alpha(n)$ and c_n , even after normalization, the moments diverge (see exercises), so this approach cannot work.

The simplest approach to proving that a sequence of random variables X_n tends in distribution to X is the method of moments. For real random variables it says: if

$$\lim_{n \rightarrow \infty} \mathbb{E}[X_n^k] = \mathbb{E}[X^k]$$

holds for every positive integer k , and if $\sum_{k \geq 1} \mathbb{E}[X^{2k}]^{-1/(2k)}$ diverges, then $X_n \xrightarrow{d} X$.

Example: if $X \sim N(0, 1)$ then $\mathbb{E}[X^{2k}]^{-1/(2k)} \sim c/\sqrt{k}$.

For both $\sum_{n \leq x} \alpha(n)$ and c_n , even after normalization, the moments diverge (see exercises), so this approach cannot work. Indicates the distribution has heavy tails.

Yet, various central limit theorems are known!

Yet, various central limit theorems are known! Given a subset $\mathcal{A}_x \subseteq \{1, 2, \dots, x\}$, consider

$$S(x, \mathcal{A}_x) := |\mathcal{A}_x|^{-1/2} \sum_{n \in \mathcal{A}_x} \alpha(n).$$

In some cases, it is known that $S(x, \mathcal{A}_x) \xrightarrow{d} N_{\mathbb{C}}(0, 1)$:

Yet, various central limit theorems are known! Given a subset $\mathcal{A}_x \subseteq \{1, 2, \dots, x\}$, consider

$$S(x, \mathcal{A}_x) := |\mathcal{A}_x|^{-1/2} \sum_{n \in \mathcal{A}_x} \alpha(n).$$

In some cases, it is known that $S(x, \mathcal{A}_x) \xrightarrow{d} N_{\mathbb{C}}(0, 1)$:

- 1 $\mathcal{A}_x$ is the short interval $[x - h, x]$ where $h \leq x/(\log x)^c$ for suitable $c > 0$ (Chatterjee–Soundararajan, Soundararajan–Xu, Pandey–Wang–Xu)

Yet, various central limit theorems are known! Given a subset $\mathcal{A}_x \subseteq \{1, 2, \dots, x\}$, consider

$$S(x, \mathcal{A}_x) := |\mathcal{A}_x|^{-1/2} \sum_{n \in \mathcal{A}_x} \alpha(n).$$

In some cases, it is known that $S(x, \mathcal{A}_x) \xrightarrow{d} N_{\mathbb{C}}(0, 1)$:

- 1 $\mathcal{A}_x$ is the short interval $[x - h, x]$ where $h \leq x/(\log x)^c$ for suitable $c > 0$ (Chatterjee–Soundararajan, Soundararajan–Xu, Pandey–Wang–Xu)
- 2 $\mathcal{A}_x$ consists of integers with k prime factors, $k = o(\log \log x)$ (Hough, Harper)

Yet, various central limit theorems are known! Given a subset $\mathcal{A}_x \subseteq \{1, 2, \dots, x\}$, consider

$$S(x, \mathcal{A}_x) := |\mathcal{A}_x|^{-1/2} \sum_{n \in \mathcal{A}_x} \alpha(n).$$

In some cases, it is known that $S(x, \mathcal{A}_x) \xrightarrow{d} N_{\mathbb{C}}(0, 1)$:

- 1 $\mathcal{A}_x$ is the short interval $[x - h, x]$ where $h \leq x/(\log x)^c$ for suitable $c > 0$ (Chatterjee–Soundararajan, Soundararajan–Xu, Pandey–Wang–Xu)
- 2 $\mathcal{A}_x$ consists of integers with k prime factors, $k = o(\log \log x)$ (Hough, Harper)
- 3 $\mathcal{A}_x$ consists of elements of $Q(\mathbb{N})$ (Najnudel, Klurman–Shkredov–Xu, Chinis–Shala)

All the known examples are sparse: $|\mathcal{A}_x| = o(x)$.

What are the proof methods?

- 1 Method of moments

All the known examples are sparse: $|\mathcal{A}_x| = o(x)$.

What are the proof methods?

- 1 Method of moments
- 2 McLeish CLT

Method of moments

Let $X \sim N_{\mathbb{C}}(0, 1)$. To prove that $S(x, \mathcal{A}_x) \xrightarrow{d} X$, it suffices to prove that

$$\lim_{x \rightarrow \infty} \mathbb{E}[S(x, \mathcal{A}_x)^{k_1} \overline{S(x, \mathcal{A}_x)}^{k_2}] = \mathbb{E}[X^{k_1} \overline{X}^{k_2}] = k_1! \delta_{k_1, k_2}.$$

Method of moments

Let $X \sim N_{\mathbb{C}}(0, 1)$. To prove that $S(x, \mathcal{A}_x) \xrightarrow{d} X$, it suffices to prove that

$$\lim_{x \rightarrow \infty} \mathbb{E}[S(x, \mathcal{A}_x)^{k_1} \overline{S(x, \mathcal{A}_x)}^{k_2}] = \mathbb{E}[X^{k_1} \overline{X}^{k_2}] = k_1! \delta_{k_1, k_2}.$$

For $k_1 = k_2 = k$, this becomes

$$\# \left\{ \begin{matrix} n_1, \dots, n_k \in \mathcal{A}_x \\ m_1, \dots, m_k \in \mathcal{A}_x \end{matrix} \middle| n_1 \cdots n_k = m_1 \cdots m_k \right\} \sim k! |\mathcal{A}_x|^k.$$

Method of moments

Let $X \sim N_{\mathbb{C}}(0, 1)$. To prove that $S(x, \mathcal{A}_x) \xrightarrow{d} X$, it suffices to prove that

$$\lim_{x \rightarrow \infty} \mathbb{E}[S(x, \mathcal{A}_x)^{k_1} \overline{S(x, \mathcal{A}_x)}^{k_2}] = \mathbb{E}[X^{k_1} \overline{X}^{k_2}] = k_1! \delta_{k_1, k_2}.$$

For $k_1 = k_2 = k$, this becomes

$$\# \left\{ \begin{matrix} n_1, \dots, n_k \in \mathcal{A}_x \\ m_1, \dots, m_k \in \mathcal{A}_x \end{matrix} \middle| n_1 \cdots n_k = m_1 \cdots m_k \right\} \sim k! |\mathcal{A}_x|^k.$$

Given a permutation $\pi \in S_k$, the terms with $n_i = m_{\pi(i)} \in \mathcal{A}_x$ contribute $|\mathcal{A}_x|^k$,

Method of moments

Let $X \sim N_{\mathbb{C}}(0, 1)$. To prove that $S(x, \mathcal{A}_x) \xrightarrow{d} X$, it suffices to prove that

$$\lim_{x \rightarrow \infty} \mathbb{E}[S(x, \mathcal{A}_x)^{k_1} \overline{S(x, \mathcal{A}_x)^{k_2}}] = \mathbb{E}[X^{k_1} \overline{X}^{k_2}] = k_1! \delta_{k_1, k_2}.$$

For $k_1 = k_2 = k$, this becomes

$$\# \left\{ \begin{matrix} n_1, \dots, n_k \in \mathcal{A}_x \\ m_1, \dots, m_k \in \mathcal{A}_x, \end{matrix} n_1 \cdots n_k = m_1 \cdots m_k \right\} \sim k! |\mathcal{A}_x|^k.$$

Given a permutation $\pi \in S_k$, the terms with $n_i = m_{\pi(i)} \in \mathcal{A}_x$ contribute $|\mathcal{A}_x|^k$, so the task becomes bounding the *off-diagonal contribution*: terms where one cannot find a permutation π such that $n_i = m_{\pi(i)}$ for all i .

It turns out that for the random variable $S(x, \mathcal{A}_x)$, there is a useful *fourth moment criterion*.

It turns out that for the random variable $S(x, \mathcal{A}_x)$, there is a useful *fourth moment criterion*.

Theorem 5 (Soundararajan–Xu)

Let $\mathcal{A}_x \subseteq \{1, 2, \dots, x\}$. Suppose the following two conditions hold:

- $\lim_{x \rightarrow \infty} \mathbb{E}|S(x, \mathcal{A}_x)|^4 = 2$.
- Let $P(n)$ be the largest prime factor of n . For every prime $p \leq x$, we have $|\{n \in \mathcal{A}_x : P(n) = p\}| = o(|\mathcal{A}_x|)$.

It turns out that for the random variable $S(x, \mathcal{A}_x)$, there is a useful *fourth moment criterion*.

Theorem 5 (Soundararajan–Xu)

Let $\mathcal{A}_x \subseteq \{1, 2, \dots, x\}$. Suppose the following two conditions hold:

- $\lim_{x \rightarrow \infty} \mathbb{E}|S(x, \mathcal{A}_x)|^4 = 2$.
- Let $P(n)$ be the largest prime factor of n . For every prime $p \leq x$, we have $|\{n \in \mathcal{A}_x : P(n) = p\}| = o(|\mathcal{A}_x|)$.

Then

$$S(x, \mathcal{A}_x) \xrightarrow{d} N_{\mathbb{C}}(0, 1).$$

It turns out that for the random variable $S(x, \mathcal{A}_x)$, there is a useful *fourth moment criterion*.

Theorem 5 (Soundararajan–Xu)

Let $\mathcal{A}_x \subseteq \{1, 2, \dots, x\}$. Suppose the following two conditions hold:

- $\lim_{x \rightarrow \infty} \mathbb{E}|S(x, \mathcal{A}_x)|^4 = 2$.
- Let $P(n)$ be the largest prime factor of n . For every prime $p \leq x$, we have $|\{n \in \mathcal{A}_x : P(n) = p\}| = o(|\mathcal{A}_x|)$.

Then

$$S(x, \mathcal{A}_x) \xrightarrow{d} N_{\mathbb{C}}(0, 1).$$

Adapted to sums of the Steinhaus function in $\mathbb{F}_q[T]$
(Hoban–Shah–Ismail–Verreault–Zaman).

What does the fourth moment condition actually say?

What does the fourth moment condition actually say? That

$$\# \left\{ (a, b, c, d) \in (\mathcal{A}_x)^4 : ab = cd, \begin{matrix} (a,b) \neq (c,d) \\ (a,b) \neq (d,c) \end{matrix} \right\} = o(|\mathcal{A}_x|^2).$$

What does the fourth moment condition actually say? That

$$\# \left\{ (a, b, c, d) \in (\mathcal{A}_x)^4 : ab = cd, \begin{smallmatrix} (a,b) \neq (c,d) \\ (a,b) \neq (d,c) \end{smallmatrix} \right\} = o(|\mathcal{A}_x|^2).$$

One can squeeze out a bit more. Let $\mathcal{B}_x \subseteq \mathcal{A}_x$. If $|\mathcal{B}_x| = o(|\mathcal{A}_x|)$ then

$$\frac{\sum_{n \in \mathcal{A}_x} \alpha(n)}{\sqrt{|\mathcal{A}_x|}} \xrightarrow{d} N_{\mathbb{C}}(0, 1) \longleftrightarrow \frac{\sum_{n \in \mathcal{A}_x \setminus \mathcal{B}_x} \alpha(n)}{\sqrt{|\mathcal{A}_x|}} \xrightarrow{d} N_{\mathbb{C}}(0, 1).$$

What does the fourth moment condition actually say? That

$$\# \left\{ (a, b, c, d) \in (\mathcal{A}_x)^4 : ab = cd, \begin{smallmatrix} (a,b) \neq (c,d) \\ (a,b) \neq (d,c) \end{smallmatrix} \right\} = o(|\mathcal{A}_x|^2).$$

One can squeeze out a bit more. Let $\mathcal{B}_x \subseteq \mathcal{A}_x$. If $|\mathcal{B}_x| = o(|\mathcal{A}_x|)$ then

$$\frac{\sum_{n \in \mathcal{A}_x} \alpha(n)}{\sqrt{|\mathcal{A}_x|}} \xrightarrow{d} N_{\mathbb{C}}(0, 1) \longleftrightarrow \frac{\sum_{n \in \mathcal{A}_x \setminus \mathcal{B}_x} \alpha(n)}{\sqrt{|\mathcal{A}_x|}} \xrightarrow{d} N_{\mathbb{C}}(0, 1).$$

The reason: considering the difference, its second moment vanishes in the limit, hence it tends in distribution to 0.

What does the fourth moment condition actually say? That

$$\# \left\{ (a, b, c, d) \in (\mathcal{A}_x)^4 : ab = cd, \begin{smallmatrix} (a,b) \neq (c,d) \\ (a,b) \neq (d,c) \end{smallmatrix} \right\} = o(|\mathcal{A}_x|^2).$$

One can squeeze out a bit more. Let $\mathcal{B}_x \subseteq \mathcal{A}_x$. If $|\mathcal{B}_x| = o(|\mathcal{A}_x|)$ then

$$\frac{\sum_{n \in \mathcal{A}_x} \alpha(n)}{\sqrt{|\mathcal{A}_x|}} \xrightarrow{d} N_{\mathbb{C}}(0, 1) \longleftrightarrow \frac{\sum_{n \in \mathcal{A}_x \setminus \mathcal{B}_x} \alpha(n)}{\sqrt{|\mathcal{A}_x|}} \xrightarrow{d} N_{\mathbb{C}}(0, 1).$$

The reason: considering the difference, its second moment vanishes in the limit, hence it tends in distribution to 0.

Consequence: We may apply the fourth moment *after* discarding a sparse set.

We wish to explain what is the McLeish Central Limit Theorem and how it gives the criterion.

We wish to explain what is the McLeish Central Limit Theorem and how it gives the criterion. Relies on martingales.

We wish to explain what is the McLeish Central Limit Theorem and how it gives the criterion. Relies on martingales.

Fix a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Let $(\mathcal{F}_i)_{i \geq 1}$ be a filtration: an increasing sequence of sub- σ -algebras of $\mathcal{F}$.

A sequence of random variables $(R_i)_{i \geq 1}$ is called a *martingale adapted to* $(\mathcal{F}_i)_{i \geq 1}$ if R_i is measurable with respect to $\mathcal{F}_i$ for all $i \geq 1$, and

$$\mathbb{E}[R_i \mid \mathcal{F}_{i-1}] = R_{i-1}$$

for all $i \geq 2$.

Theorem 6 (McLeish, 70s)

For any $n \geq 1$, let $(X_i^{(n)})_{i \geq 1}$ be a martingale difference sequence, i.e. $(X_1^{(n)} + \dots + X_i^{(n)})_{i \geq 1}$ is a martingale. A sufficient condition for

$$\sum_{i \geq 1} X_i^{(n)} \xrightarrow{d} N_{\mathbb{C}}(0, 1)$$

to hold as $n \rightarrow \infty$ is

$$\lim_{n \rightarrow \infty} \sum_{i \geq 1} \mathbb{E}[|X_i^{(n)}|^4] = 0, \quad \lim_{n \rightarrow \infty} \mathbb{E}\left[\left|\sum_{i \geq 1} (X_i^{(n)})^2\right|^2\right] = 0,$$

$$\lim_{n \rightarrow \infty} \mathbb{E}\left[\left(\sum_{i \geq 1} |X_i^{(n)}|^2 - 1\right)^2\right] = 0.$$

Theorem 6 (McLeish, 70s)

For any $n \geq 1$, let $(X_i^{(n)})_{i \geq 1}$ be a martingale difference sequence, i.e. $(X_1^{(n)} + \dots + X_i^{(n)})_{i \geq 1}$ is a martingale. A sufficient condition for

$$\sum_{i \geq 1} X_i^{(n)} \xrightarrow{d} N_{\mathbb{C}}(0, 1)$$

to hold as $n \rightarrow \infty$ is

$$\lim_{n \rightarrow \infty} \sum_{i \geq 1} \mathbb{E}[|X_i^{(n)}|^4] = 0, \quad \lim_{n \rightarrow \infty} \mathbb{E}\left[\left|\sum_{i \geq 1} (X_i^{(n)})^2\right|^2\right] = 0,$$

$$\lim_{n \rightarrow \infty} \mathbb{E}\left[\left(\sum_{i \geq 1} |X_i^{(n)}|^2 - 1\right)^2\right] = 0.$$

Exercise: if $(X_i^{(n)})_i$ are independent with $\mathbb{E}|\sum_i X_i^{(n)}|^2 = 1$ then 1st condition implies the 3rd and we recover (complex) CLT.

We now relate the McLeish CLT to $S(x, \mathcal{A}_x)$. If $P(n)$ is the largest prime factor of n then we may write (Harper)

$$\sum_{n \in \mathcal{A}_x} \alpha(n) = \sum_{p \leq x} X_p$$

where

$$X_p = \sum_{n \in \mathcal{A}_x: P(n)=p} \alpha(n).$$

We now relate the McLeish CLT to $S(x, \mathcal{A}_x)$. If $P(n)$ is the largest prime factor of n then we may write (Harper)

$$\sum_{n \in \mathcal{A}_x} \alpha(n) = \sum_{p \leq x} X_p$$

where

$$X_p = \sum_{n \in \mathcal{A}_x: P(n)=p} \alpha(n).$$

The sequence $(\sum_{p \leq q} X_p)_{q \leq x}$ is a martingale with respect to the filtration $\mathcal{F}_q = \sigma(\alpha(p) : p \leq q)$ indexed by primes.

We now relate the McLeish CLT to $S(x, \mathcal{A}_x)$. If $P(n)$ is the largest prime factor of n then we may write (Harper)

$$\sum_{n \in \mathcal{A}_x} \alpha(n) = \sum_{p \leq x} X_p$$

where

$$X_p = \sum_{n \in \mathcal{A}_x: P(n)=p} \alpha(n).$$

The sequence $(\sum_{p \leq q} X_p)_{q \leq x}$ is a martingale with respect to the filtration $\mathcal{F}_q = \sigma(\alpha(p) : p \leq q)$ indexed by primes.

Suffices to check: if q is the largest prime less than q then $\mathbb{E}[X_p \mid \alpha(2), \alpha(3), \dots, \alpha(q)] = 0$.

We now relate the McLeish CLT to $S(x, \mathcal{A}_x)$. If $P(n)$ is the largest prime factor of n then we may write (Harper)

$$\sum_{n \in \mathcal{A}_x} \alpha(n) = \sum_{p \leq x} X_p$$

where

$$X_p = \sum_{n \in \mathcal{A}_x: P(n)=p} \alpha(n).$$

The sequence $(\sum_{p \leq q} X_p)_{q \leq x}$ is a martingale with respect to the filtration $\mathcal{F}_q = \sigma(\alpha(p) : p \leq q)$ indexed by primes.

Suffices to check: if q is the largest prime less than q then $\mathbb{E}[X_p \mid \alpha(2), \alpha(3), \dots, \alpha(q)] = 0$. Reduces to $\mathbb{E}[\alpha(p)^k] = 0$ for $k \geq 1$.

Recall $(c_n)_{n \geq 0}$ are defined via

$$\sum_{n=0}^{\infty} c_n u^n = \exp \left(- \sum_{i=1}^{\infty} \frac{N_i u^i}{\sqrt{i}} \right).$$

Recall $(c_n)_{n \geq 0}$ are defined via

$$\sum_{n=0}^{\infty} c_n u^n = \exp \left(- \sum_{i=1}^{\infty} \frac{N_i u^i}{\sqrt{i}} \right).$$

Given n , we write $\lambda \vdash n$ to indicate that λ is a partition of n , and write $a_i(\lambda)$ for the number of parts of size i in λ .

Recall $(c_n)_{n \geq 0}$ are defined via

$$\sum_{n=0}^{\infty} c_n u^n = \exp \left(- \sum_{i=1}^{\infty} \frac{N_i u^i}{\sqrt{i}} \right).$$

Given n , we write $\lambda \vdash n$ to indicate that λ is a partition of n , and write $a_i(\lambda)$ for the number of parts of size i in λ .

Expanding the definition of c_n we see that there are coefficients $(b_\lambda)_{\lambda \vdash n}$ such that

$$c_n = \sum_{\lambda \vdash n} b_\lambda \prod_{i=1}^n N_i^{a_i(\lambda)}.$$

Given an expression

$$A(n, \mathbf{b}) := \sum_{\lambda \vdash n} b_{\lambda} \prod_{i=1}^n N_i^{a_i(\lambda)},$$

we decompose it as

$$A(n, \mathbf{b}) = \sum_{i=1}^n Z_i$$

where Z_i consists of terms in $A(n, \mathbf{b})$ corresponding to partitions λ whose largest part is of size i .

The expression Z_i involves a power of N_i but does not involve N_j for $j > i$.

Given an expression

$$A(n, \mathbf{b}) := \sum_{\lambda \vdash n} b_{\lambda} \prod_{i=1}^n N_i^{a_i(\lambda)},$$

we decompose it as

$$A(n, \mathbf{b}) = \sum_{i=1}^n Z_i$$

where Z_i consists of terms in $A(n, \mathbf{b})$ corresponding to partitions λ whose largest part is of size i .

The expression Z_i involves a power of N_i but does not involve N_j for $j > i$. The sequence $(\sum_{i=1}^m Z_i)_{1 \leq m \leq n}$ is a martingale with respect to the filtration $\mathcal{F}_m = \sigma(Z_1, \dots, Z_m)$ ($1 \leq m \leq n$).

Thank you for listening.