



Secular coefficients mini-course

Talk 1

Ofir Gorodetsky, Technion

`ofir-gorodetsky.github.io/Bristol`

CMI-HIMR Summer School, Bristol
June 1, 2026

- 1 Talk 1: Riemann zeta function and analogies
- 2 Talk 2: Random multiplicative functions: basic results
- 3 Talk 3: Random multiplicative functions: sum of the Steinhaus function
- 4 Talk 4: Combinatorial perspective on moments of zeta

Definition

The Riemann zeta function is defined as

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = 1 + 2^{-s} + 3^{-s} + 4^{-s} + \dots$$

for $s \in \mathbb{C}$ with $\Re s > 1$.

Definition

The Riemann zeta function is defined as

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = 1 + 2^{-s} + 3^{-s} + 4^{-s} + \dots$$

for $s \in \mathbb{C}$ with $\Re s > 1$. We shall review its properties (without proof).

Euler product

For $\Re s > 1$,

$$\zeta(s) = \prod_p \left(1 + p^{-s} + p^{-2s} + \dots\right) = \prod_p (1 - p^{-s})^{-1}.$$

Analytic continuation

ζ has a meromorphic continuation to $\mathbb{C}$ with a simple pole at $s = 1$ and residue 1:

$$\zeta(s) - \frac{1}{s-1}$$

is entire.

Functional equation

Define

$$\chi(s) := \Gamma(s/2) \pi^{-s/2} s(s-1)/2.$$

Define

$$\xi(s) := \zeta(s) \chi(s).$$

Then $\xi(s)$ is entire and

$$\xi(s) = \xi(1-s).$$

Functional equation

Define

$$\chi(s) := \Gamma(s/2)\pi^{-s/2}s(s-1)/2.$$

Define

$$\xi(s) := \zeta(s)\chi(s).$$

Then $\xi(s)$ is entire and

$$\xi(s) = \xi(1-s).$$

Note that $\Gamma(s/2)$ has simple poles at $s = 0, -2, -4, \dots$

The poles cancel with zeros of $\zeta(s)$ at $s = -2, -4, \dots$ and the factor s .

Product over zeros

We have

$$\xi(s) = \xi(0) \prod_{\rho: \zeta(\rho)=0, \rho \neq -2, -4, \dots} \left(1 - \frac{s}{\rho}\right).$$

for an explicit real A .

Product over zeros

We have

$$\xi(s) = \xi(0) \prod_{\rho: \zeta(\rho)=0, \rho \neq -2, -4, \dots} \left(1 - \frac{s}{\rho}\right).$$

for an explicit real A .

Anticipated by Riemann, proved by Hadamard.

Product over zeros

We have

$$\xi(s) = \xi(0) \prod_{\rho: \zeta(\rho)=0, \rho \neq -2, -4, \dots} \left(1 - \frac{s}{\rho}\right).$$

for an explicit real A .

Anticipated by Riemann, proved by Hadamard.

Number of zeros

$$N(T) = |\{\rho : \zeta(\rho) = 0, 0 < \Im \rho \leq T\}| \sim \frac{T \log T}{2\pi}.$$

Connection with primes

The connection arises from relating primes (appearing in the Euler product) to the zeros (appearing in the Hadamard product).

Connection with primes

The connection arises from relating primes (appearing in the Euler product) to the zeros (appearing in the Hadamard product).

Consider the logarithmic derivative of $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$:

$$-\frac{\zeta'}{\zeta}(s) = \sum_p \log p \frac{p^{-s}}{1 - p^{-s}} = \sum_{n=1}^{\infty} \frac{\Lambda(n)}{n^s}$$

where Λ is the *von Mangoldt function*,

$$\Lambda(n) = \begin{cases} \log p & \text{if } n = p^k, \\ 0 & \text{otherwise.} \end{cases}$$

Connection with primes (cont.)

The series $-\frac{\zeta'}{\zeta}(s)$ converges for $\Re s > 1$, and has a meromorphic continuation to $\mathbb{C}$, with pole at every zero or pole of ζ .

Connection with primes (cont.)

The series $-\frac{\zeta'}{\zeta}(s)$ converges for $\Re s > 1$, and has a meromorphic continuation to $\mathbb{C}$, with pole at every zero or pole of ζ .

Just as Cauchy's formula expresses a coefficient of a power series as a complex integral, *Perron's formula* implies the famous “Explicit Formula”:

$$\sum_{n \leq x} \Lambda(n) = \frac{1}{2\pi i} \int_{(2)} -\frac{\zeta'}{\zeta}(s) \frac{x^s}{s} ds = x - \sum_{\rho: \zeta(\rho)=0} \frac{x^\rho}{\rho}.$$

Connection with primes (cont.)

The Riemann hypothesis: $\Re \rho = 1/2$ (aside from $\rho = -2, -4, \dots$).

Connection with primes (cont.)

The Riemann hypothesis: $\Re \rho = 1/2$ (aside from $\rho = -2, -4, \dots$). Implies:

$$\sum_{n \leq x} \Lambda(n) = x + O(\sqrt{x} \log^2 x).$$

Conversely, this estimate implies the Riemann hypothesis.

Connection with primes (cont.)

The Riemann hypothesis: $\Re \rho = 1/2$ (aside from $\rho = -2, -4, \dots$). Implies:

$$\sum_{n \leq x} \Lambda(n) = x + O(\sqrt{x} \log^2 x).$$

Conversely, this estimate implies the Riemann hypothesis. In prime language,

$$\sum_{p \leq x} 1 = \int_2^x \frac{dt}{\log t} + O(\sqrt{x} \log x).$$

Probabilistic perspective on the Riemann zeta function

The Steinhaus function

Define $\alpha: \mathbb{N} \rightarrow \mathbb{C}$ to be a completely multiplicative function:

$$\alpha(nm) = \alpha(n)\alpha(m),$$

where $(\alpha(p))_p$ are iid random variables uniformly distributed on the unit circle.

Probabilistic perspective on the Riemann zeta function

The Steinhaus function

Define $\alpha: \mathbb{N} \rightarrow \mathbb{C}$ to be a completely multiplicative function:

$$\alpha(nm) = \alpha(n)\alpha(m),$$

where $(\alpha(p))_p$ are iid random variables uniformly distributed on the unit circle.

Denote

$$F(s) = \sum_{n=1}^{\infty} \frac{\alpha(n)}{n^s} = \prod_p \left(1 - \frac{\alpha(p)}{p^s}\right)^{-1}$$

for $\Re s > 1$.

Probabilistic perspective on the Riemann zeta function

The Steinhaus function

Define $\alpha: \mathbb{N} \rightarrow \mathbb{C}$ to be a completely multiplicative function:

$$\alpha(nm) = \alpha(n)\alpha(m),$$

where $(\alpha(p))_p$ are iid random variables uniformly distributed on the unit circle.

Denote

$$F(s) = \sum_{n=1}^{\infty} \frac{\alpha(n)}{n^s} = \prod_p \left(1 - \frac{\alpha(p)}{p^s}\right)^{-1}$$

for $\Re s > 1$.

α is called the *Steinhaus random multiplicative function*.

Theorem 1 (Bohr–Jessen, 1930s)

Fix $s \in \mathbb{C}$ with $\Re s > 1/2$. Let U_T be a random variable uniformly distributed on $[0, T]$. Then

$$\zeta(s + iU_T) \xrightarrow{d} F(s)$$

as $T \rightarrow \infty$.

Theorem 1 (Bohr–Jessen, 1930s)

Fix $s \in \mathbb{C}$ with $\Re s > 1/2$. Let U_T be a random variable uniformly distributed on $[0, T]$. Then

$$\zeta(s + iU_T) \xrightarrow{d} F(s)$$

as $T \rightarrow \infty$. Concretely,

$$\lim_{T \rightarrow \infty} \mathbb{P}(\zeta(s + iU_T) \in A) = \mathbb{P}(F(s) \in A).$$

Theorem 1 (Bohr–Jessen, 1930s)

Fix $s \in \mathbb{C}$ with $\Re s > 1/2$. Let U_T be a random variable uniformly distributed on $[0, T]$. Then

$$\zeta(s + iU_T) \xrightarrow{d} F(s)$$

as $T \rightarrow \infty$. Concretely,

$$\lim_{T \rightarrow \infty} \mathbb{P}(\zeta(s + iU_T) \in A) = \mathbb{P}(F(s) \in A).$$

More generally, given $s_1, \dots, s_k$ with real parts $> 1/2$,

$$(\zeta(s_1 + iU_T), \dots, \zeta(s_k + iU_T)) \xrightarrow{d} (F(s_1), \dots, F(s_k)).$$

Intuition

For any fixed y ,

$$(p^{iU_T})_{p \leq y} \xrightarrow{d} (\alpha(p))_{p \leq y}.$$

Intuition

For any fixed y ,

$$(p^{iU_T})_{p \leq y} \xrightarrow{d} (\alpha(p))_{p \leq y}.$$

Indeed, it suffices to check that

$$\lim_{T \rightarrow \infty} \mathbb{E} \left[\prod_{p \leq y} (p^{iU_T})^{a_p} \right] = \mathbb{E} \left[\prod_{p \leq y} (\alpha(p)^{a_p}) \right].$$

Intuition

For any fixed y ,

$$(p^{iU_T})_{p \leq y} \xrightarrow{d} (\alpha(p))_{p \leq y}.$$

Indeed, it suffices to check that

$$\lim_{T \rightarrow \infty} \mathbb{E} \left[\prod_{p \leq y} (p^{iU_T})^{a_p} \right] = \mathbb{E} \left[\prod_{p \leq y} (\alpha(p))^{a_p} \right].$$

RHS is 1 if $a_p = 0$ holds for all p , and 0 otherwise.

LHS is 1 if $a_p = 0$ holds for all p , and tends to 0 otherwise.

Haar measure

Let $U(N)$ be the group of N -by- N complex unitary matrices U with $U\overline{U}^T = I_N$.

Haar measure

Let $U(N)$ be the group of N -by- N complex unitary matrices U with $U\overline{U}^T = I_N$. There is a unique probability measure μ on $U(N)$ with the property: if U_N has law μ , then

$$\mathbb{P}(U_N \in X) = \mathbb{P}(AU_N \in X) = \mathbb{P}(U_N A \in X)$$

for every matrix $A \in U(N)$ (“translation-invariance”). It is called the *Haar measure*.

Comparison with ζ

It has been increasingly understood that $\zeta(s)$ (especially on the critical line) is analogous to $\det(I - uU)$ (especially on the unit circle).

Comparison with ζ

It has been increasingly understood that $\zeta(s)$ (especially on the critical line) is analogous to $\det(I - uU)$ (especially on the unit circle).

① Hadamard product: $\det(I - uU) = \prod_{\rho} \text{eigenvalue}(1 - u\rho)$

Comparison with ζ

It has been increasingly understood that $\zeta(s)$ (especially on the critical line) is analogous to $\det(I - uU)$ (especially on the unit circle).

- 1 Hadamard product: $\det(I - uU) = \prod_{\rho} \text{eigenvalue}(1 - u\rho)$
- 2 Analytic continuation: $\det(I - uU)$ is entire

Comparison with ζ

It has been increasingly understood that $\zeta(s)$ (especially on the critical line) is analogous to $\det(I - uU)$ (especially on the unit circle).

- 1 Hadamard product: $\det(I - uU) = \prod_{\rho} \text{eigenvalue}(1 - u\rho)$
- 2 Analytic continuation: $\det(I - uU)$ is entire
- 3 Functional equation:
$$\det(I - uU) = (-u)^N \det U \det(I - u^{-1}\overline{U})$$

Comparison with ζ

It has been increasingly understood that $\zeta(s)$ (especially on the critical line) is analogous to $\det(I - uU)$ (especially on the unit circle).

- 1 Hadamard product: $\det(I - uU) = \prod_{\rho} \text{eigenvalue}(1 - u\rho)$
- 2 Analytic continuation: $\det(I - uU)$ is entire
- 3 Functional equation:
$$\det(I - uU) = (-u)^N \det U \det(I - u^{-1}\overline{U})$$
- 4 Riemann Hypothesis: $\det(I - uU) = 0 \implies |u| = 1$

Comparison with ζ

It has been increasingly understood that $\zeta(s)$ (especially on the critical line) is analogous to $\det(I - uU)$ (especially on the unit circle).

- 1 Hadamard product: $\det(I - uU) = \prod_{\rho} \text{eigenvalue}(1 - u\rho)$
- 2 Analytic continuation: $\det(I - uU)$ is entire
- 3 Functional equation:
$$\det(I - uU) = (-u)^N \det U \det(I - u^{-1}\overline{U})$$
- 4 Riemann Hypothesis: $\det(I - uU) = 0 \implies |u| = 1$

No Euler product.

Theorem 2 (Bohr–Jessen for matrices (90s))

Fix u with $|u| < 1$. Let $(N_i)_{i \geq 1}$ be iid standard complex Gaussians. Then

$$\det(I - uU_N) \xrightarrow{d} F(u)$$

as $N \rightarrow \infty$ where

$$F(u) = \exp \left(- \sum_{i=1}^{\infty} \frac{N_i}{\sqrt{i}} u^i \right).$$

Theorem 2 (Bohr–Jessen for matrices (90s))

Fix u with $|u| < 1$. Let $(N_i)_{i \geq 1}$ be iid standard complex Gaussians. Then

$$\det(I - uU_N) \xrightarrow{d} F(u)$$

as $N \rightarrow \infty$ where

$$F(u) = \exp \left(- \sum_{i=1}^{\infty} \frac{N_i}{\sqrt{i}} u^i \right).$$

More generally, given $u_1, \dots, u_k$ inside the unit circle,

$$(\det(I - u_1 U), \dots, \det(I - u_k U)) \xrightarrow{d} (F(u_1), \dots, F(u_k)).$$

Intuition

Taking logarithms,

$$\log \det(I - uU_N) = - \sum_{i=1}^{\infty} \frac{\text{Tr}(U_N^i)}{i} u^i.$$

By a result of Diaconis–Shahshahani:

$$\left(\frac{\text{Tr}(U_N^i)}{\sqrt{i}} \right)_{i \leq k} \xrightarrow{d} (N_i)_{i \leq k}.$$

A Dirichlet character modulo m is a function $\chi: \mathbb{Z} \rightarrow \mathbb{C}$ that has period m , is completely multiplicative and satisfies $\chi(1) = 1$.

A Dirichlet character modulo m is a function $\chi: \mathbb{Z} \rightarrow \mathbb{C}$ that has period m , is completely multiplicative and satisfies $\chi(1) = 1$.

Example: $\chi(1) = 1$, $\chi(3) = -1$, $\chi(2) = 0$ (period 4).

A Dirichlet character modulo m is a function $\chi: \mathbb{Z} \rightarrow \mathbb{C}$ that has period m , is completely multiplicative and satisfies $\chi(1) = 1$.

Example: $\chi(1) = 1$, $\chi(3) = -1$, $\chi(2) = 0$ (period 4).

General construction: take a character $(\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ and extend to $\mathbb{Z}$.

A Dirichlet character modulo m is a function $\chi: \mathbb{Z} \rightarrow \mathbb{C}$ that has period m , is completely multiplicative and satisfies $\chi(1) = 1$.

Example: $\chi(1) = 1$, $\chi(3) = -1$, $\chi(2) = 0$ (period 4).

General construction: take a character $(\mathbb{Z}/m\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$ and extend to $\mathbb{Z}$.

Given a Dirichlet character, its Dirichlet L -function is

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = \prod_p (1 - \chi(p)p^{-s})^{-1},$$

defined for $\Re s > 1$.

Dirichlet L -functions have properties similar to those of ζ , but sometimes simpler.

Dirichlet L -functions have properties similar to those of ζ , but sometimes simpler. For instance, if χ is a nontrivial character, then $L(s, \chi)$ defines a convergent series for $\Re s > 0$, and has no pole (entire function).

Dirichlet L -functions have properties similar to those of ζ , but sometimes simpler. For instance, if χ is a nontrivial character, then $L(s, \chi)$ defines a convergent series for $\Re s > 0$, and has no pole (entire function).

Functional equation relates $L(s, \chi)$ to $L(1 - s, \bar{\chi})$. This is similar to how $\det(I - uU)$ relates to $\det(I - u^{-1}\bar{U})$.

Dirichlet L -functions have properties similar to those of ζ , but sometimes simpler. For instance, if χ is a nontrivial character, then $L(s, \chi)$ defines a convergent series for $\Re s > 0$, and has no pole (entire function).

Functional equation relates $L(s, \chi)$ to $L(1 - s, \bar{\chi})$. This is similar to how $\det(I - uU)$ relates to $\det(I - u^{-1}\bar{U})$.

Again, one can connect primes to zeros of $L(s, \chi)$ through Perron's formula, obtaining another Explicit Formula:

$$\sum_{n \leq x} \Lambda(n) \chi(n) = \frac{1}{2\pi i} \int_{(2)} -\frac{L'}{L}(s, \chi) \frac{x^s}{s} ds = - \sum_{\rho: L(\rho, \chi)=0} \frac{x^\rho}{\rho}.$$

One has a Bohr–Jessen theorem for L -functions: let χ_m be a Dirichlet character chosen uniformly at random from the group of Dirichlet characters modulo m . Then, for $\Re s > 1/2$,

$$L(s, \chi_m) \xrightarrow{d} \prod_p (1 - \alpha(p)/p^s)^{-1}$$

as m tends to infinity along primes.

Fix a finite field $\mathbb{F}_q$ and consider the ring $\mathbb{F}_q[T]$. One may measure the size of a polynomial $f \in \mathbb{F}_q[T]$ using the norm $|f| = q^{\deg f}$.

Fix a finite field $\mathbb{F}_q$ and consider the ring $\mathbb{F}_q[T]$. One may measure the size of a polynomial $f \in \mathbb{F}_q[T]$ using the norm $|f| = q^{\deg f}$.

This leads naturally to the consideration of the zeta function of $\mathbb{F}_q[T]$:

$$\zeta_{\mathbb{F}_q[T]}(s) = \sum_{f \text{ monic}} \frac{1}{|f|^s} = \prod_P (1 - |P|^{-s})^{-1}.$$

Fix a finite field $\mathbb{F}_q$ and consider the ring $\mathbb{F}_q[T]$. One may measure the size of a polynomial $f \in \mathbb{F}_q[T]$ using the norm $|f| = q^{\deg f}$.

This leads naturally to the consideration of the zeta function of $\mathbb{F}_q[T]$:

$$\zeta_{\mathbb{F}_q[T]}(s) = \sum_{f \text{ monic}} \frac{1}{|f|^s} = \prod_P (1 - |P|^{-s})^{-1}.$$

Unfortunately, this is not a very interesting object: it has the closed-form formula $\frac{1}{1-q^{1-s}}$. No zeros whatsoever.

Fix a finite field $\mathbb{F}_q$ and consider the ring $\mathbb{F}_q[T]$. One may measure the size of a polynomial $f \in \mathbb{F}_q[T]$ using the norm $|f| = q^{\deg f}$.

This leads naturally to the consideration of the zeta function of $\mathbb{F}_q[T]$:

$$\zeta_{\mathbb{F}_q[T]}(s) = \sum_{f \text{ monic}} \frac{1}{|f|^s} = \prod_P (1 - |P|^{-s})^{-1}.$$

Unfortunately, this is not a very interesting object: it has the closed-form formula $\frac{1}{1-q^{1-s}}$. No zeros whatsoever.

Functional equation: if we multiply by $(1 - q^{-s})^{-1}$ ("prime at infinity"), then there is symmetry $s \leftrightarrow 1 - s$ (up to a simple entire factor).

A much richer object is the Dirichlet L -function of a Dirichlet character $\chi: \mathbb{F}_q[T] \rightarrow \mathbb{C}$,

$$L(s, \chi) = \sum_{f \text{ monic}} \frac{\chi(f)}{|f|^s}.$$

Here a Dirichlet character modulo $M \in \mathbb{F}_q[T]$ is a completely multiplicative function with $\chi(f + M) = \chi(f)$ for every $f \in \mathbb{F}_q[T]$. It can be constructed by lifting a character of $(\mathbb{F}_q[T]/M)^\times$.

A much richer object is the Dirichlet L -function of a Dirichlet character $\chi: \mathbb{F}_q[T] \rightarrow \mathbb{C}$,

$$L(s, \chi) = \sum_{f \text{ monic}} \frac{\chi(f)}{|f|^s}.$$

Here a Dirichlet character modulo $M \in \mathbb{F}_q[T]$ is a completely multiplicative function with $\chi(f + M) = \chi(f)$ for every $f \in \mathbb{F}_q[T]$. It can be constructed by lifting a character of $(\mathbb{F}_q[T]/M)^\times$.

Notation: $L(s, \chi) = \mathcal{L}(q^{-s}, \chi)$ where

$$\mathcal{L}(u, \chi) = \sum_{f \text{ monic}} \chi(f) u^{\deg f}$$

is a *power series* in u .

Observation: In fact, $\mathcal{L}(u, \chi)$ is a polynomial, of degree less than $\deg M$.

Observation: In fact, $\mathcal{L}(u, \chi)$ is a polynomial, of degree less than $\deg M$.

Theorem (Weil): The roots of $\mathcal{L}(u, \chi)$ lie on $|u| = 1/\sqrt{q}$, i.e., the roots of $L(s, \chi)$ lie on $\Re s = 1/2$.

Observation: In fact, $\mathcal{L}(u, \chi)$ is a polynomial, of degree less than $\deg M$.

Theorem (Weil): The roots of $\mathcal{L}(u, \chi)$ lie on $|u| = 1/\sqrt{q}$, i.e., the roots of $L(s, \chi)$ lie on $\Re s = 1/2$.

Implication: $\mathcal{L}(u/\sqrt{q}, \chi)$ shares many properties with $\det(I - uU)$.

Given a nontrivial Dirichlet character χ modulo M , we may write

$$\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$$

for some conjugacy class $\Theta_\chi \in U(\deg M - 1)$ (according to Weil).

Given a nontrivial Dirichlet character χ modulo M , we may write

$$\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$$

for some conjugacy class $\Theta_\chi \in U(\deg M - 1)$ (according to Weil).

Theorem 3 (Katz, 2013)

Fix the degree of M . As $q \rightarrow \infty$, we have

$$\lim_{q \rightarrow \infty} \frac{1}{\phi(M)} \sum_{\chi \bmod M} f(\Theta_\chi) = \mathbb{E}f(U_{\deg M - 1})$$

for any conjugation-invariant f .

It is instructive to consider logarithms of

$\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$ and compare with the logarithm of $\det(I - uU_N)$.

It is instructive to consider logarithms of

$\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$ and compare with the logarithm of $\det(I - uU_N)$. On the one hand, from the Euler product we have

$$\log \mathcal{L}(u/\sqrt{q}, \chi) = \sum_{n=1}^{\infty} A_n(\chi) \frac{u^n}{n}$$

where

$$A_n(\chi) = \frac{1}{q^{n/2}} \sum_{\deg f=n, f \text{ monic}} \Lambda(f) \chi(f).$$

It is instructive to consider logarithms of

$\mathcal{L}(u/\sqrt{q}, \chi) = \det(I - u\Theta_\chi)$ and compare with the logarithm of $\det(I - uU_N)$. On the one hand, from the Euler product we have

$$\log \mathcal{L}(u/\sqrt{q}, \chi) = \sum_{n=1}^{\infty} A_n(\chi) \frac{u^n}{n}$$

where

$$A_n(\chi) = \frac{1}{q^{n/2}} \sum_{\deg f=n, f \text{ monic}} \Lambda(f) \chi(f).$$

On the other hand, from $\log \det(I - uU) = -\sum_{n=1}^{\infty} \text{Tr}(U^n) \frac{u^n}{n}$ we have

$$A_n(\chi) = -\text{Tr}(\Theta_\chi^n).$$

The equality

$$\frac{1}{q^{n/2}} \sum_{\deg f=n, f \text{ monic}} \Lambda(f) \chi(f) = -\text{Tr}(\Theta_{\chi}^n)$$

is a finite-field analogue of the Explicit Formula.

The equality

$$\frac{1}{q^{n/2}} \sum_{\deg f=n, f \text{ monic}} \Lambda(f) \chi(f) = -\text{Tr}(\Theta_\chi^n)$$

is a finite-field analogue of the Explicit Formula.

For U_N we have $\log \det(I - uU_N) = -\sum_{n=1}^{\infty} \text{Tr}(U_N^n) \frac{u^n}{n}$ and Diaconis–Shahshahani proved: $\text{Tr}(U_N^n)/\sqrt{n}$ tends to standard complex Gaussian as $N \rightarrow \infty$.

The equality

$$\frac{1}{q^{n/2}} \sum_{\deg f=n, f \text{ monic}} \Lambda(f) \chi(f) = -\text{Tr}(\Theta_\chi^n)$$

is a finite-field analogue of the Explicit Formula.

For U_N we have $\log \det(I - uU_N) = -\sum_{n=1}^{\infty} \text{Tr}(U_N^n) \frac{u^n}{n}$ and Diaconis–Shahshahani proved: $\text{Tr}(U_N^n)/\sqrt{n}$ tends to standard complex Gaussian as $N \rightarrow \infty$.

In fact, $A_n(\chi_M)/\sqrt{n}$ (for χ_M random character modulo M) also tends to standard complex Gaussian by taking an appropriate limit (see exercises).

One of the main questions concerning the Riemann zeta function is the growth of its moments:

$$\frac{1}{T} \int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt = \mathbb{E} \left| \zeta \left(\frac{1}{2} + iU_T \right) \right|^{2k}.$$

One of the main questions concerning the Riemann zeta function is the growth of its moments:

$$\frac{1}{T} \int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt = \mathbb{E} \left| \zeta \left(\frac{1}{2} + iU_T \right) \right|^{2k}.$$

Hardy–Littlewood (1918): For $k = 1$, grows like $\log T$.

One of the main questions concerning the Riemann zeta function is the growth of its moments:

$$\frac{1}{T} \int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt = \mathbb{E} \left| \zeta \left(\frac{1}{2} + iU_T \right) \right|^{2k}.$$

Hardy–Littlewood (1918): For $k = 1$, grows like $\log T$.

Ingham (1926): For $k = 2$, grows like $\frac{1}{12} \zeta(2)^{-1} (\log T)^4$.

One of the main questions concerning the Riemann zeta function is the growth of its moments:

$$\frac{1}{T} \int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt = \mathbb{E} \left| \zeta \left(\frac{1}{2} + iU_T \right) \right|^{2k}.$$

Hardy–Littlewood (1918): For $k = 1$, grows like $\log T$.

Ingham (1926): For $k = 2$, grows like $\frac{1}{12} \zeta(2)^{-1} (\log T)^4$.

No asymptotic results for $k \geq 3$, but number theorists expected $\sim a_k g_k (\log T)^{k^2}$ for a well-understood arithmetic constant a_k , and an elusive geometric constant g_k (e.g. $g_2 = 1/12$ and $a_2 = \zeta(2)^{-1}$).

In 2000, Keating and Snaith had the idea to look at moments of $\det(I - uU_N)$.

In 2000, Keating and Snaith had the idea to look at moments of $\det(I - uU_N)$.

Theorem 4 (Keating–Snaith)

We have

$$\mathbb{E}|\det(I - U_N)|^{2k} = \prod_{j=1}^N \frac{\Gamma(j)\Gamma(j+2k)}{(\Gamma(j+k))^2} \sim N^{k^2} \prod_{j=0}^{k-1} \frac{j!}{(j+k)!}.$$

Conjecture (Keating–Snaith)

In 2000, Keating and Snaith had the idea to look at moments of $\det(I - uU_N)$.

Theorem 4 (Keating–Snaith)

We have

$$\mathbb{E} |\det(I - U_N)|^{2k} = \prod_{j=1}^N \frac{\Gamma(j)\Gamma(j+2k)}{(\Gamma(j+k))^2} \sim N^{k^2} \prod_{j=0}^{k-1} \frac{j!}{(j+k)!}.$$

Conjecture (Keating–Snaith)

We have

$$\frac{1}{T} \int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt \sim a_k g_k (\log T)^{k^2}$$

for $g_k = \prod_{j=0}^{k-1} \frac{j!}{(j+k)!}$. ($g_1 = 1$, $g_2 = 1/12$, $g_3 = 42/9!$.)

Main input to the Keating–Snaith theorem: the “Selberg integral”, which gives all the moments at once, including non-integer moments.

Main input to the Keating–Snaith theorem: the “Selberg integral”, which gives all the moments at once, including non-integer moments. Nothing like that in $\mathbb{Z}$. (So how did Ingham proceed?)

Main input to the Keating–Snaith theorem: the “Selberg integral”, which gives all the moments at once, including non-integer moments. Nothing like that in $\mathbb{Z}$. (So how did Ingham proceed?)

Heuristic for $(\log T)^{k^2}$ growth:

$$\log |\zeta(\tfrac{1}{2} + iU_T)| \approx - \sum_{p \leq T} \log |1 - p^{-\frac{1}{2} - iU_T}| \approx \sum_{p \leq T} p^{-\frac{1}{2}} \cos(U_T \log p).$$

Main input to the Keating–Snaith theorem: the “Selberg integral”, which gives all the moments at once, including non-integer moments. Nothing like that in $\mathbb{Z}$. (So how did Ingham proceed?)

Heuristic for $(\log T)^{k^2}$ growth:

$$\log |\zeta(\tfrac{1}{2} + iU_T)| \approx - \sum_{p \leq T} \log |1 - p^{-\frac{1}{2} - iU_T}| \approx \sum_{p \leq T} p^{-\frac{1}{2}} \cos(U_T \log p).$$

If this were a sum of independent random variables, the variance would be $\frac{1}{2} \sum_{p \leq T} p^{-1} \sim (\log \log T)/2$. Then the central limit theorem would tell us that

$$\frac{\log |\zeta(1/2 + iU_T)|}{\sqrt{(\log \log T)/2}} \xrightarrow{d} N(0, 1).$$

The last limit is actually an unconditional result of Selberg (1940s). It has a matrix analogue:

$$\frac{\log |\det(I - U_N)|}{\sqrt{(\log N)/2}} \xrightarrow{d} N(0, 1).$$

The last limit is actually an unconditional result of Selberg (1940s). It has a matrix analogue:

$$\frac{\log |\det(I - U_N)|}{\sqrt{(\log N)/2}} \xrightarrow{d} N(0, 1).$$

The analogue can be proved using the Selberg integral, as it gives $\mathbb{E} |\det(I - U_N)|^z$, which is the moment generating function/characteristic function of $\log |\det(I - U)|$. Nothing like that available in integers.

The last limit is actually an unconditional result of Selberg (1940s). It has a matrix analogue:

$$\frac{\log |\det(I - U_N)|}{\sqrt{(\log N)/2}} \xrightarrow{d} N(0, 1).$$

The analogue can be proved using the Selberg integral, as it gives $\mathbb{E}|\det(I - U_N)|^2$, which is the moment generating function/characteristic function of $\log |\det(I - U)|$. Nothing like that available in integers.

Moments of ζ should grow like

$$\mathbb{E}|\zeta(1/2 + iU_T)|^{2k} \approx \mathbb{E}e^{2k\sqrt{(\log \log T)/2}Z} = (\log T)^{k^2}$$

where $Z \sim N(0, 1)$. This is only a heuristic, as it is missing both a_k and g_k .

Correlation functions

k -point correlation

Given a point process $X = (X_i)_i$ (a collection of random real points), one defines its k -point correlation function $\rho_k^X: \mathbb{R}^k \rightarrow \mathbb{R}$

$$\mathbb{E} \sum_{i_1, \dots, i_k \text{ distinct}} f(X_{i_1}, \dots, X_{i_k}) = \int_{\mathbb{R}^k} f(\vec{t}) \rho_k^X(\vec{t}) dt_1 \cdots dt_k.$$

Correlation functions

k -point correlation

Given a point process $X = (X_i)_i$ (a collection of random real points), one defines its k -point correlation function $\rho_k^X: \mathbb{R}^k \rightarrow \mathbb{R}$

$$\mathbb{E} \sum_{i_1, \dots, i_k \text{ distinct}} f(X_{i_1}, \dots, X_{i_k}) = \int_{\mathbb{R}^k} f(\vec{t}) \rho_k^X(\vec{t}) d\vec{t}.$$

Given a point process X , we say that a sequence of point processes $(X_N)_N$ tends in correlation to X if $\lim_{N \rightarrow \infty} \rho_k^{X_N} = \rho_k^X$ for every k .

Convergence in correlation often implies convergence in distribution: $\sum_i f(X_{N,i}) \xrightarrow{d} \sum_i f(X_i)$.

Theorem 5 (Dyson, 1960s)

Define a point process X_N by taking the eigenangles $\theta_i \in [-\pi, \pi)$ of a random unitary matrix U_N .

The point process $\frac{N}{2\pi} \cdot X_N$ tends in correlation and in distribution to the sine point process Sine, defined by

$\rho_k(t_1, \dots, t_k) = \det (S(t_i - t_j))_{1 \leq i, j \leq k}$ where

$$S(t) = \frac{\sin(\pi t)}{\pi t}.$$

Conjecture (GUE Conjecture, Montgomery, 1970s)

Assume RH and list the zeros of zeta as $1/2 + i\gamma_n$, $n \in \mathbb{Z}$.

Consider the point process $X_T = \left(\frac{\log T}{2\pi} (\gamma_n - U_T) \right)_{n \in \mathbb{Z}}$ where

U_T is uniformly distributed on $[0, T]$.

Then X_T tends in distribution and in correlation to Sine.

Montgomery's conjecture was originally stated as a “pair correlation conjecture”, concerning ρ_2 only:

$$\mathbb{E} \sum_{i_1 \neq i_2} f((X_T)_{i_1}, (X_T)_{i_2}) \rightarrow \int_{\mathbb{R}^2} f(t_1, t_2) (1 - S(t_1 - t_2)^2) dt_1 dt_2$$

where $S(t) = \frac{\sin \pi t}{\pi t}$.

His motivation: Under RH he could prove it for certain “band-limited” functions f .

Montgomery's conjecture was originally stated as a “pair correlation conjecture”, concerning ρ_2 only:

$$\mathbb{E} \sum_{i_1 \neq i_2} f((X_T)_{i_1}, (X_T)_{i_2}) \rightarrow \int_{\mathbb{R}^2} f(t_1, t_2) (1 - S(t_1 - t_2)^2) dt_1 dt_2$$

where $S(t) = \frac{\sin \pi t}{\pi t}$.

His motivation: Under RH he could prove it for certain “band-limited” functions f .

Consequence for primes:

$$\frac{1}{X} \int_0^X \left(\sum_{x \leq n < x+H} \Lambda(n) - H \right)^2 dx \sim H \log \frac{X}{H}$$

as $X \rightarrow \infty$.

Rudnick–Sarnak (1990s): Under RH, computed limit $\mathbb{E} \sum_{i_1, \dots, i_k \text{ distinct}} f(X_{i_1}, \dots, X_{i_k})$ for “band-limited functions”, giving further evidence for the GUE conjecture.

Rudnick–Sarnak (1990s): Under RH, computed limit $\mathbb{E} \sum_{i_1, \dots, i_k \text{ distinct}} f(X_{i_1}, \dots, X_{i_k})$ for “band-limited functions”, giving further evidence for the GUE conjecture.

For any given k , Rodgers found an interpretation in terms of primes for the whole GUE conjecture.

Rudnick–Sarnak (1990s): Under RH, computed limit $\mathbb{E} \sum_{i_1, \dots, i_k \text{ distinct}} f(X_{i_1}, \dots, X_{i_k})$ for “band-limited functions”, giving further evidence for the GUE conjecture.

For any given k , Rodgers found an interpretation in terms of primes for the whole GUE conjecture.

The fact that convergence in correlation and in distribution are one and the same in the GUE conjecture was recently established (Arias de Reyna–Rodgers).

Thank you for listening.