

Results & Material and methods

Ofir Shukron & David Holcman

December 21, 2015

1 Result section

To estimate the nucleosome reorganization following DNA damages, we have constructed a mathematical model, where redistribution can be due either to chromatin de- compaction or nucleosome sliding along the chromatin or both of them. We have used the model to assess the relative contribution of these two processes to the total DNA and nucleosome signal loss from a region of interest (ROI), a quantity which is inaccessible experimentally.

The model (presented in Material and methods) follows the DNA, $D(u)$, and nucleosome, $H(u)$, fraction of signal loss from the ROI as a function of the UV dose, u . We have used the measured H3.3 and DNA signal loss to calibrate parameters of nucleosome and DNA models respectively (Fig. 3A-D). Using the calibrated models we have found that the relative contribution of nucleosome sliding to the total signal loss in the ROI is monotonically decreasing from 75% to 70% for nucleosomes and 51% to 40% for DNA loss, as the UV dose increases from 5 to 100 msec. The remaining percentages are attributed to chromatin expansion and de-compaction (see Figure 1).




Figure 1: **Relative contribution of chromatin opening and nucleosome sliding to DNA (left) and histone (right) loss.** Sliding contribution is monotonically decreasing with UV dose for both DNA and nucleosome signals. The increase in chromatin opening contribution is due to the increase in chromatin reorganization following high dosages of UV, which is responsible for the majority of signal loss.

2 Material and methods: Modeling nucleosomes and DNA redistribution following UV damages

We present here a model for nucleosomes and DNA re-organization following UV damages. The cascade of events leading to tagged DNA and nucleosomes' redistribution, results in signal extrusion from a region of interest (ROI) up to a maximal loss, measured 15 minutes post UV-C.

2.1 Dynamics of nucleosomes following UV damages in the region of interest

Following the experimental protocol, a two-dimensional circular initial damage region (IDR), induced by the UV laser beam, is centered around the focal point (origin of the coordinates) with a fixed area of A_0 . Following laser induction with a UV dose u , the tagged damage region expands radi-

ally outward and reaches its maximal area of $A(u)$ after 15 minutes. At the end of expansion, the circular region is defined as the ROI, in which DNA and nucleosome signals are measured at time 0 and 15 minutes. The fraction of signal loss is calculated for DNA and nucleosomes as

$$\frac{signal(0) - signal(15)}{signal(0)} \quad (1)$$

where $signal(0)$ and $signal(15)$ are the DNA or nucleosome signals at time 0 and 15 minutes respectively.

We assume that the loss of DNA and nucleosomes signal post UV-C is due to two mechanisms: the first is chromatin expansion, and the second is nucleosomes sliding along the chromatin. For chromatin expansion, recruitment and binding of repair factors to damaged DNA causes chromatin de-compaction, and cross-links break [ref] to enable repair factor access to damaged DNA. Because the majority of damages are inflicted around the laser's focal point, the accumulation of repair factors will generate a pushing force on surrounding chromatin in a radial outward direction. As a result of the pushing force, DNA and nucleosome will be extruded from the ROI in equal proportions.

An additional loss of nucleosome signal is caused by the mechanism of nucleosome sliding. Repair proteins slide nucleosomes from damaged position on the DNA away from high concentration of DNA damages. Each nucleosome sliding over damaged positions on the DNA displaces the tagged damages spatially in the general direction of the ROI boundary. Additionally, with each nucleosome sliding, the damages in the damage region are exposed, and repair factors are recruited to bind to them.

In-line with the description above, we construct a model representing signal loss 15 minutes post UV-C. We do not specifically take into account the mechanism of signal loss in time and only present equations representing the maximal loss of signal as a function of the UV dose.

2.2 Dynamics of DNA and nucleosome loss from the DR

The IDR and the ROI are considered to be two-dimensional concentric circular regions, characterized by an area A_0 and $A(u)$ respectively. We assume an initial uniform distribution of DNA in the IDR and its vicinity, such that

the amount of DNA in $A(u)$ is $k_D A(u)$, with k_D in units of $bp/\mu m^2$. We assume that damages can only occur on DNA located in A_0 . Depending on the uv dose, an increasing fraction of the DNA in A_0 will be damaged. This fraction is mainly what resides in $A(u)$ 15 minutes post UV-C. Therefore, the total DNA in $A(u)$ is the sum of damaged and undamaged DNA

$$k_D A(u) = D_d(u) + D_h(u) \quad (2)$$

with $D_d(u)$ amount of damaged DNA, and $D_h(u)$ the amount of undamaged DNA.

According to our assumption the undamaged DNA is extruded from the ROI. Therefore, DNA signal loss fraction is given by the ratio of the amount of DNA in the annulus between the IDR and the ROI plus the nucleosomes pushed out of the IDR to the total amount of signal in the ROI.

$$D(u) = \frac{k_D(A(u) - A_0) + \varphi(u)}{k_D A(u)} \quad (3)$$

Similarly to DNA, we assume that the number of nucleosome in a given region is proportional to its area

$$\psi(A(u)) = k_N N(u)$$

with k_N in units of $nucleosomes/\mu m^2$.

We mark the number of nucleosome lost by pushing as $N_P(u)$ and by sliding as $N_S(u)$. Thus the total loss, $N_T(u)$, is the sum

$$N_T(u) = N_P(u) + N_S(u)$$

Because we assume that nucleosomes pushed out of the IDR are associated with undamaged DNA, the fraction of DNA and nucleosome signal loss due to pushing is similar. Therefore,

$$\frac{D_P(u)}{k_D A_0} = \frac{N_P(u)}{k_N A_0}$$

The total fraction of loss for nucleosomes is

$$H(u) = D(u) + \frac{N_S(u)}{k_N A(u)} \quad (4)$$

In order to evaluate the functions 3 and 4, we will first present a model for the accumulation of DNA damages with uv dose, $\varphi_D(u)$ and derive the functions $A(u)$ and $N_T(u)$ from it.

2.3 Accumulation of DNA damages in the IDR

We assume that the rate of accumulation of DNA damages in the IDR with increasing UV dose is proportional to the undamaged DNA in the IDR.

$$\frac{d\varphi_D(u)}{du} = k_t(k_D A_0 - \varphi_D(u))$$

with k_t a constant of units. Using the initial condition $\varphi_D(0) = 0$, the solution to equation 2.3

$$\varphi_D(u) = \kappa_t A_0 (1 - \exp(-k_t u))$$

We note that the damaged DNA can only appear in the IDR of area A_0 .

2.4 Deriving the functions describing loss of nucleosome from the IDR and the ROI expansion

We now turn to construct a model for the total number of nucleosomes $N_T(u)$ leaving the IDR and subsequently being pushed out of the ROI, as a function of the UV dose. Although the exact mechanism by which nucleosomes are lost is not known, we assume here that the number of nucleosomes leaving the ROI is composed of two contributions, one due to nucleosomes pushed out of the IDR, and the second due to nucleosome sliding to expose damaged DNA positions and facilitate efficient repair.

The nucleosomes occupy a length of the chromatin proportional to $N_0 - N_T(u)$, whereas chromatin damage coverage is proportional to $T(u)$. Therefore, in the first-order approximation, the dynamics of $N_S(u)$ is given by the multiplication

$$\frac{dN_S(u)}{du} = k_s \frac{(N_0 - N_T(u))}{\pi T_{max}^2} \frac{dT(u)}{du}$$

with k_s the rate constant. Due to the accumulation of repair factors around damaged DNA, a pushing force acts on the surrounding undamaged DNA in the IDR. The rate nucleosomes are being pushed out of the IDR is considered to be proportional to the fraction of DNA damaged occurring due to small amount of UV dose, minus the fraction out of it which slides out

$$\frac{dN_P(u)}{du} = k_p \left(\frac{(N_0 - N_T(u)) - k_s(N_0 - N_T(u))}{\pi T_{max}^2} \right) \frac{dT(u)}{du}$$

The dynamics of $N_T(u)$ is obtained by adding equations 6 and 7. Using the initial condition $N_T(0) = 0$, we can solve for $N_T(u)$ to have

$$N_T(u) = N_0 \left[1 - \exp \left(-\frac{k_p(1 - k_s) + k_s}{\pi T_{max}^2} T(u) \right) \right] \quad (5)$$

Therefore,

$$N_P(u) = \left(1 - \frac{k_s}{k_p(1 - k_s) + k_s} \right) N_T(u) \quad (6)$$

$$N_S(u) = \left(\frac{k_s}{k_p(1 - k_s) + k_s} \right) N_T(u) \quad (7)$$

with k_s and k_p constants describing the rate of nucleosome depletion from the ROI due to sliding and pushing respectively.

Next, we model the dynamics of the ROI area $A(u)$ with increasing UV dose. For this end, we consider the ROI expansion to be affected by two additive mechanisms originating from the IDR: one is chromatin opening or de-compaction, and the other is nucleosome sliding. The two contributions are inherent in the equation for $N_T(u)$, therefore

$$\frac{dA(u)}{du} = k_a \frac{dN_T(u)}{du} = k_a \left(\frac{dN_P(u)}{du} + \frac{dN_S(u)}{du} \right)$$

where k_a is the rate constant. Using the initial condition $A(0) = A_0$, we find

$$A(u) = A_0 + k_a N_0 \exp \left(-\frac{(k_p(1 - k_s) + k_s)}{\pi T_{max}^2} T(u) \right) \quad (8)$$

where A_0 represents the size of the IDR even in the absence of UV induction.

We can now substitute equations (6), (7) and (8) into the steady-state equations (3) and (4) to get the expressions for $D(u)$ and $H(u)$. Using the relation $D_p(u)/A_0 = N_p(u)/N_0$, we present the result in terms of $N_T(u)$

$$D(u) = \frac{\frac{N_T(u)}{N_0} \left(\frac{k_p(1 - k_s)}{k_p(1 - k_s) + k_s} - \frac{k_a N_0}{A_0} \right) + \frac{k_a N_0}{A_0}}{1 + \frac{k_a N_0}{A_0} \left(1 - \frac{N_T(u)}{N_0} \right)} \quad (9)$$

$$H(u) = \frac{\frac{N_T(u)}{N_0} \left(1 - \frac{k_a N_0}{A_0} \right) + \frac{k_a N_0}{A_0}}{1 + \frac{k_a N_0}{A_0} \left(1 - \frac{N_T(u)}{N_0} \right)} \quad (10)$$

2.5 Parameter fit for $D(u)$ and $H(u)$

We now use equations (9) and (10) to fit the experimental data. We simultaneously fit equations (9) and (10) to the H3.3 and DNA loss data, with the goal of maximizing the R^2 of both equations. Excluding the measurement at 5 msec, and using classical fitting procedure, we find

$$k_t = 0.037, \quad k_s = 0.35, \quad k_p = 0.24, \quad \frac{k_a N_0}{A_0} = 0.46$$

with $R^2 = 0.93$ and $R^2 = 0.96$ for DNA and nucleosome loss fit respectively.




Figure 2: **Histone (red) and DNA (green) loss: experimental data for nucleosomes (circle) and DNA (triangles) versus model curves (continuous and dashed, respectively).** Parameter values are obtained by simultaneously fitting equations(9) and (10) to the experimental data with the goal of maximizing the R^2 . The resulting curves show $R^2 = 0.96$ and $R^2 = 0.93$ for nucleosome and DNA loss, respectively.

2.6 Relative contribution of opening and sliding to DNA and nucleosome signal loss

Using the calibrated model in equation equations (9) and (10), we now calculate the relative contribution of chromatin opening and nucleosome sliding to the total loss of DNA and nucleosomes. The sliding contribution refers to all loss caused by either directly sliding nucleosome out of the IDR or as

the effect nucleosome sliding has on pushing chromatin out of the IDR by operations of repair factors. Chromatin opening contribution refers to all signal loss caused by chromatin remodeling, which causes expansion of the IDR and subsequent loss of signal from the ROI.

We start by dividing the equation describing the ROI expansion into the two sub-mechanisms of signal loss

$$A(u) = A_P(u) + A_S(u)$$

with $A_P(u)$ the area attributed to chromatin opening, and $A_S(u)$ to nucleosome sliding. Subtracting the sliding contribution from the ROI expansion in equation (8), we arrive at

$$A_P(u) = A_0 + k_a N_P(u)$$

The DNA loss attributed to chromatin opening and de-compaction is thus

$$D(u)_{opening} = \frac{A_P(u)/A_0 - 1 + D_P(u)/A_0}{A_P(u)/A_0}$$

The fraction attributed to chromatin opening and sliding out of the total DNA loss are given respectively by

$$\frac{D(u)_{opening}}{D(u)}, \quad \frac{D(u)_{sliding}}{D(u)} = 1 - \frac{D(u)_{opening}}{D(u)} \quad (11)$$

Similarly, the relative contribution of chromatin opening and nucleosome sliding to the total nucleosome loss is given respectively by

$$\frac{H(u)_{opening}}{H(u)} = \frac{D(u)_{opening}}{H(u)}, \quad \frac{H(u)_{sliding}}{H(u)} = 1 - \frac{H(u)_{opening}}{H(u)} \quad (12)$$

where here we have used the fact that $H(u)_{opening} = D(u)_{opening}$. Graphs of equations (11) and (12) are presented in Figure 1.

2.7 Nucleosome sliding out of the IDR

The fraction of nucleosomes lost by sliding out of the total nucleosomes in the IDR (and eventually pushed out of the ROI) is found to be an increasing function of the UV dose




Figure 3: **Fraction of nucleosomes sliding out of the IDR.** The plot against the experimental data resulted in $R^2 = 0.75$

$$N_S(u) = \frac{H(u) - D(u)}{1 - D(u)}$$

Substituting the parameter values in subsection 2.5 and plugging (9) and (10) into the expression above, we obtain the results in Figure 3, where the model in equation (7) is plotted against the experimental data ($R^2 = 0.75$).

The parameters values found in subsection 2.5, can further be used to estimate the fraction of nucleosome loss from the IDR attributed to sliding and chromatin opening. These contributions are given by the leading coefficients in the equations (6) and (7) for $N_P(u)$ and $N_S(u)$, respectively. We have, for chromatin opening and sliding, respectively

$$N_P(u) = 0.3N_T(u), \quad N_S(u) = 0.7N_T(u)$$