

HALO: Using CM SAF's MAGIC SOL method to retrieve global and direct surface radiation from historical geosynchronous observations

R.W. Mueller
German Weather Service

D. Lee
University of Marburg





Historical solar
Analysis from
Long-term geosynchronous
Orbital

Proof of concept for a global solar ECV data set

Essential climate variables:

- Relevant base variable
- Long-term record
- High quality measurements

Comparable data sets:

- Helio-Clim
- ERA-Interim
- GEWEX SRB
- ISCCP FD
- FLASHFlux
- CLARA

Comparable data sets:

- Helio-Clim
- ERA-Interim
- GEWEX SRB
- ISCCP FD
- FLASHFlux
- CLARA

Data needed with:

- Robust data requirements
- Long time series available
- High resolution
- Differentiation in diffuse/beam radiation
- Free to use/distribute
- Global spatial extent

Advantages:


- Robust and well tested
- Low data requirements
- Self-calibration – high data availability
- Computation of direct and diffuse radiation
- Relatively low computational costs
- Best validation among comparable data sets
- Open source – verifiable, modifiable, freely usable




Historical solar
Analysis from
Long-term geosynchronous
Orbit



Source: Météo-France 2009, composite by author





Year	GOES-W (-135°)	GOES-E (-74°)	MSAT-Prime (0°)	MSAT-IODC (57.5°)	GMS (155°)
1978 – 1980	GOES-2		<u>Meteosat-1</u>		GMS
	GOES-3				
1981 – 1985	GOES-4	GOES-5	<u>Meteosat-2</u>		GMS-2
1986 – 1990			<u>Meteosat-3</u>		GMS-3
			<u>Meteosat-4</u>		
1991 – 1995			<u>Meteosat-5</u>		GMS-4
1996 – 2000	GOES-9	GOES-8	<u>Meteosat-6</u>		GMS-5
2001 – 2005	GOES-10		<u>Meteosat-7</u>	<u>Meteosat-5</u>	GOES-++
2006 – 2010	GOES-11	GOES-12			
2011 – 2012	GOES-15	GOES-13		<u>Meteosat-7</u>	

Year	GOES-W (-135°)	GOES-E (-74°)	MSAT-Prime (0°)	MSAT-IODC (57.5°)	GMS (155°)
1978 – 1980	GOES-2		<u>Meteosat-1</u>		GMS
	GOES-3				
1981 – 1985	GOES-4	GOES-5	<u>Meteosat-2</u>		GMS-2
1986 – 1990			<u>Meteosat-3</u>		GMS-3
			<u>Meteosat-4</u>		
			<u>Meteosat-5</u>		
1991 – 1995					GMS-4
1996 – 2000	GOES-9	GOES-8	<u>Meteosat-6</u>		GMS-5
2001 – 2005	GOES-10		<u>Meteosat-7</u>	<u>Meteosat-5</u>	GOES-++
2006 – 2010		GOES-12			
	GOES-11			<u>Meteosat-7</u>	
2011 – 2012		GOES-13			
	GOES-15				

Year	GOES-W (-135°)	GOES-E (-74°)	MSAT-Prime (0°)	MSAT-IODC (57.5°)	GMS (155°)
1978 – 1980	GOES-2		<u>Meteosat-1</u>		GMS
	GOES-3				
1981 – 1985	GOES-4	GOES-5	<u>Meteosat-2</u>		GMS-2
1986 – 1990			<u>Meteosat-3</u>		GMS-3
			<u>Meteosat-4</u>		
1991 – 1995			<u>Meteosat-5</u>		GMS-4
1996 – 2000	GOES-9	GOES-8	<u>Meteosat-6</u>		GMS-5
2001 – 2005	GOES-10		<u>Meteosat-7</u>	<u>Meteosat-5</u>	GOES-++
2006 – 2010		GOES-12			
	GOES-11				
2011 – 2012		GOES-13		<u>Meteosat-7</u>	
	GOES-15				

1 month = ~ 3 TB

Chosen test
month:

June 2003

