Table X — Person-level mode-choice logit: estimation of the cost sensitivity λ (model M1)

Dependent variable: transit = 1, car = 0. AM-peak (06:00–08:59) survey trips within the 25 corridor areas, THS 2017/18. Weighted binary logit, standard errors clustered by household.

Variable	Coef.	Std. err.	z	p-value	Sig.	Odds ratio
Cost variable						
ΔGC: transit minus car generalized cost (gen-min)	**-0.0350**	0.0167	-2.09	0.036	**	0.966
Car availability (ref: car available)						
Licences outnumber cars	2.1385	0.5736	3.73	<0.001	***	8.487
No driving licence	4.1686	0.6444	6.47	<0.001	***	64.625
No car in household	5.5402	0.6101	9.08	<0.001	***	254.727
Trip purpose (ref: home-based work)						
Home-based education	0.4159	0.6022	0.69	0.490		1.516
Home-based other	-2.0176	0.5805	-3.48	<0.001	***	0.133
Non-home-based	-1.6200	0.5856	-2.77	0.006	***	0.198
Age (ref: 35–64)						
0–17	-2.1995	0.5679	-3.87	<0.001	***	0.111
18–24	0.4016	0.4508	0.89	0.373		1.494
25–34	-0.0059	0.6095	-0.01	0.992		0.994
65+	0.7869	0.6070	1.30	0.195		2.197
Person and sector						
Female	0.4569	0.3150	1.45	0.147		1.579
Student	0.2365	0.5509	0.43	0.668		1.267
Arab sector (ref: general)	-1.9081	0.7956	-2.40	0.016	**	0.148
Trip distance (ref: 6–10 km)						
< 3 km	-1.0351	0.4729	-2.19	0.029	**	0.355
3–6 km	-0.0711	0.4046	-0.18	0.861		0.931
10–20 km	0.5603	0.7683	0.73	0.466		1.751
> 20 km	1.2460	1.1040	1.13	0.259		3.476
Constant						
Intercept	-2.7468	0.6145	-4.47	<0.001	***	0.064

Observations (trips)	2,310
Households (clusters)	543
Transit trips in sample	353
Log-likelihood (weighted)	-638.8
Log-likelihood, constant only	-1125.6
McFadden ρ^2	0.432
λ = −coef(ΔGC), per generalized minute	**0.0350**
95% confidence interval for λ	0.0022 to 0.0677

*Notes. *** p<0.01, ** p<0.05, * p<0.10. λ is the negative of the ΔGC coefficient: a higher transit cost relative to the car lowers the odds of choosing transit by a factor exp(−λ·ΔGC); with λ = 0.035, ten generalized minutes change the odds by about 30%. Weights: household expansion weight (new_wf), rescaled to the row count. Standard errors: household-clustered sandwich estimator with a G/(G−1) correction, because the two survey days of one household are not independent. Log-likelihoods are computed with the weights and are therefore pseudo-log-likelihoods. Source: Output/mode_choice/estimation_sample_person_level.csv, notebook Mode_choice_person_level.ipynb (step 33); the table reproduces coefficients_all_models.csv to five decimals.*