

Sampling from the normal

Normal samples

1. Sometimes data is normal.
2. Large sample statistics are often normal even when the underlying data isn't (central limit theorem).
3. The results are pretty cute.
4. It's a *tradition!*

Copyright Dick Startz

1

Copyright Dick Startz

2

Properties of sample mean and variance

Theorem 5.3.1:

Let x_1, \dots, x_n be a random sample from a $N(\mu, \sigma^2)$ distribution and let $\bar{x} = 1/n \sum_{i=1}^n x_i$ and $s^2 = 1/(n-1) \sum_{i=1}^n (x_i - \bar{x})^2$. Then,

- \bar{x} and s^2 are independent random variables
- $\bar{x} \sim N(\mu, \sigma^2/n)$ (We already showed this, since the sample mean is a linear combination of normal random variables.)
- $(n-1)s^2/\sigma^2$ has a chi-squared distribution with $n-1$ degrees of freedom (χ_{n-1}^2 or $\chi^2(n-1)$).

Copyright Dick Startz

3

Copyright Dick Startz

4

The source of all proofs about linear models is
Linear Models, S.R. Searle, Wiley, 1971.

Quadratic and linear forms

Theorem:

If $x \sim N(\mu, \Sigma)$, then $x'Ax$ and Bx , where A is symmetric positive definite, are distributed independently if and only if $B\Sigma A = 0$.

Copyright Dick Startz

5

Outline of proof

- Prove sufficiency by cleverly messing with matrix square roots.
- Digression on algebra of quadratic form
- Prove necessity
 - Lemma on traces
 - Lemma on covariances

Copyright Dick Startz

6

Sufficiency $B\Sigma A = 0$

We can write $A = LL'$ for some L of full column rank, which implies $(L'L)^{-1}$ exists. Note $B\Sigma A = B\Sigma LL'$

$$B\Sigma A = 0 \Rightarrow B\Sigma LL' = 0$$

$$B\Sigma LL'[L(L'L)^{-1}] = 0[L(L'L)^{-1}]$$

$$B\Sigma L = 0$$

Consider Bx and $L'x$, these are joint normal with covariance $B\Sigma L = 0$ and therefore independent. Since

$$x'Ax = (L'x)'(L'x) = g(L'x)$$

This proves $x'Ax$ and Bx are independent.

Copyright Dick Startz

7

Lemmas needed for necessity

Lemma (stated without proof):

$$E(x'Ax) = \text{tr}(A\Sigma) + \mu'A\mu$$

where $\text{tr}(\quad)$ is the trace, that is the sum of the diagonal elements.

Copyright Dick Startz

8

Digression

$$(x - \mu)'A(x - \mu) = x'Ax + \mu'A\mu - 2x'A\mu$$

$$\begin{aligned} & x'Ax - \mu'A\mu \\ &= (x - \mu)'A(x - \mu) - 2\mu'A\mu + 2x'A\mu \end{aligned}$$

Copyright Dick Startz

9

Lemmas needed for necessity

Lemma:

$$\text{cov}(x, x'Ax) = 2\Sigma A\mu$$

Proof:

$$\begin{aligned} \text{cov}(x, x'Ax) &= E((x - \mu)[x'Ax - E(x'Ax)]) \\ &= E((x - \mu)[x'Ax - \mu'A\mu - \text{tr}(A\Sigma)]) \end{aligned}$$

Insert digression results

$$= E((x - \mu)[(x - \mu)'A(x - \mu) + 2(x - \mu)'A\mu - \text{tr}(A\Sigma)])$$

Multiply through by the first $(x - \mu)$. First term is essentially third moment. Last term is proportional to first moment. Both are zero.

$$\text{cov}(x, x'Ax) = 2\Sigma A\mu$$

Copyright Dick Startz

10

Necessity $x'Ax$ and Bx independent

$x'Ax$ and Bx independent implies covariance equals zero, so

$$\text{cov}(Bx, x'Ax) = 2B\Sigma A\mu = 0$$

Since this is true for all μ it must be that $B\Sigma A = 0$.

Copyright Dick Startz

11

Special case

Let $\Sigma = \sigma^2 I_n$, so $x \sim N(\mu, \Sigma) = N(\mu, \sigma^2 I)$.

Let $B = \frac{1}{n} [1 \dots 1]$, then $\bar{x} = Bx$.

If we let

$$A = \begin{bmatrix} 1 - \frac{1}{n} & -\frac{1}{n} & \dots & -\frac{1}{n} \\ -\frac{1}{n} & 1 - \frac{1}{n} & \dots & -\frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{1}{n} & -\frac{1}{n} & \dots & 1 - \frac{1}{n} \end{bmatrix}$$

$$s^2 = \frac{1}{n-1} x'Ax$$

Copyright Dick Startz

12

$n = 2$ case

$$\begin{aligned}
 x'Ax &= [x_1 \ x_2] \begin{bmatrix} 1/2 & -1/2 \\ -1/2 & 1/2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \\
 &= \left[x_1 - \frac{x_1 + x_2}{2} \quad x_2 - \frac{x_1 + x_2}{2} \right] \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \\
 &= x_1^2 + x_2^2 - (x_1 + x_2) \frac{x_1 + x_2}{2} \\
 &= \left(x_1 - \frac{x_1 + x_2}{2} \right)^2 + \left(x_2 - \frac{x_1 + x_2}{2} \right)^2
 \end{aligned}$$

Copyright Dick Startz

13

Special case

$$B\Sigma A = B\sigma^2 I A$$

But $BA = 0$

$$\frac{1}{n} [1 \dots 1] \begin{bmatrix} 1 - \frac{1}{n} & -\frac{1}{n} & \dots & -\frac{1}{n} \\ -\frac{1}{n} & 1 - \frac{1}{n} & \dots & -\frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{1}{n} & -\frac{1}{n} & \dots & 1 - \frac{1}{n} \end{bmatrix} = 0$$

Copyright Dick Startz

14

Wow!

Therefore, the sample mean and the sample variance of normals are independent.

Copyright Dick Startz

15

Reminders about χ^2

Lemma 5.3.2:

- a) If $z \sim N(0,1)$, then $z^2 \sim \chi_1^2$.
- b) If $z \sim N(0, \sigma^2)$, then $\frac{z}{\sigma} \sim N(0,1)$, so $\frac{1}{\sigma^2} z^2 \sim \chi_1^2$.
- c) If x_1, \dots, x_n are independent and $x_i \sim \chi_{p_i}^2$, then $x_1 + \dots + x_n \sim \chi_{p_1 + \dots + p_n}^2$.

Copyright Dick Startz

16

Recap- Properties of sample mean and variance

Theorem 5.3.1:

Let x_1, \dots, x_n be a random sample from a $N(\mu, \sigma^2)$ distribution and let $\bar{x} = 1/n \sum_{i=1}^n x_i$ and $s^2 = 1/(n-1) \sum_{i=1}^n (x_i - \bar{x})^2$. Then,

- a) \bar{x} and s^2 are independent random variables
- b) $\bar{x} \sim N(\mu, \sigma^2/n)$ (We already showed this, since the sample mean is a linear combination of normal random variables.)
- c) $(n-1)s^2/\sigma^2$ has a chi-squared distribution with $n-1$ degrees of freedom (χ_{n-1}^2 or $\chi^2(n-1)$).

Copyright Dick Startz

19

Properties of s^2

If $x \sim \chi_k^2$

$$E(x) = k$$

$$E\left(\frac{\sigma^2}{n-1} \times \frac{(n-1)s^2}{\sigma^2}\right) = \frac{\sigma^2}{n-1} \times (n-1) = \sigma^2$$

$$\text{var}(x) = 2k$$

$$\text{var}\left(\frac{\sigma^2}{n-1} \times \frac{(n-1)s^2}{\sigma^2}\right) = \left(\frac{\sigma^2}{n-1}\right)^2 \times 2(n-1)$$

$$= \frac{2\sigma^4}{n-1}$$

Copyright Dick Startz

20

Assignment

Generate $n = 4$ standard normals, compute the sample mean \bar{x} , sample variance s^2 , and the t -statistic given by $t = \bar{x}/\sqrt{s^2/n}$. Do this a lot of times, saving the result. (While you're at it, use tic/toc to compute how long your simulation takes.) Now make three plots showing the empirical and theoretical pdfs, one for the sample mean, one for $(n-1)$ times the sample variance, and one for the t -statistic.

Copyright Dick Startz

21

Useful matrix theorems

Definition:

A is positive definite iff the quadratic form $c'Ac > 0 \forall c \neq 0$.

1. If A is symmetric, positive definite, \exists nonsingular P s.t. $A = PP'$
2. If A is $n \times k$, $n \geq k$, rank k , then $A'A$ is symmetric pos. def. If $\text{rank}(A) < k$, $A'A$ is positive semi-definite.
3. If A is of full column rank, then $\det(A'A) > 0$.
4. If A, B are pos. def. and $A - B$ is pos. def., then $B^{-1} - A^{-1}$ is pos. def.

Copyright Dick Startz

27

Linear and quadratic forms

If $u \sim (\mu, \Sigma)$, then $Lu \sim (L\mu, L\Sigma L')$, where u is $n \times 1$, L is $q \times n$.

1. If $u \sim N(0, \sigma^2 I)$ and A is symmetric, idempotent, rank r , then $\frac{1}{\sigma^2} u' A u \sim \chi_r^2$.
2. If $u \sim N(0, V)$ then $u' A u \sim \chi_r^2$, iff AV is idempotent of rank r .
3. (Corollary) If $u \sim N(0, V)$ then $u' V^{-1} u \sim \chi_r^2$, where $\text{rank}(V) = r$.
4. If A, B sym. idem. and $u \sim N(0, \sigma^2 I)$, then $\frac{1}{\sigma^2} u' A u, \frac{1}{\sigma^2} u' B u$ are independent iff $AB = 0$.
5. If $u \sim N(0, \sigma^2 I)$, $u' A u, B u$ are independent iff $BA = 0$.

Copyright Dick Startz

28

Getting to t -

$$\bar{x} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

We standardize \bar{x} .

$$z = \frac{\bar{x} - \mu}{\sqrt{\sigma^2/n}} \sim N(0, 1)$$

$$t = \frac{z}{\sqrt{s^2/\sigma^2}} = \frac{\bar{x} - \mu}{\sqrt{s^2/n}} \sim t_{n-1}$$

Remember that s^2/σ^2 is a χ_{n-1}^2 divided by $n-1$. The ratio of a standard normal to the square root of an independent χ^2 divided by its degrees of freedom is distributed t .

Copyright Dick Startz

29

F-statistic

If $x_1 \sim \chi_{d_1}^2, x_2 \sim \chi_{d_2}^2$ and x_1, x_2 are independent, then

$$\frac{x_1/d_1}{x_2/d_2} \sim F(d_1, d_2)$$

Example:

$$t^2 = \frac{z^2/1}{s^2/\sigma^2} \sim F(1, n-1)$$

Copyright Dick Startz

30

Another way

$$SSR = (n-1)s^2 = \sum (x - \bar{x})^2 = x' A x, \quad A = \begin{bmatrix} 1 - \frac{1}{n} & -\frac{1}{n} & \cdots & -\frac{1}{n} \\ -\frac{1}{n} & 1 - \frac{1}{n} & \cdots & -\frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{1}{n} & -\frac{1}{n} & \cdots & 1 - \frac{1}{n} \end{bmatrix}$$

Suppose we believe that $\mu = 0$, then

$$SSR^* = \sum x^2 = x' I_n x$$

Divided by σ^2 is also χ^2 .

$$SSR^* - SSR = \frac{1}{n} x' B x, \quad B = \begin{bmatrix} 1 & \cdots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \cdots & 1 \end{bmatrix}$$

So $SSR^* - SSR$ is also χ^2 . Note that $BA = 0$, proving $SSR^* - SSR$ and SSR are independent. Therefore

$$\frac{(SSR^* - SSR)/1}{SSR/(n-1)} = \frac{(\sum x^2 - \sum (x - \bar{x})^2)/1}{\sum (x - \bar{x})^2 / (n-1)} \sim F(1, n-1)$$

Copyright Dick Startz

31

Least squares

$$\begin{aligned}
 y &= X\beta + \varepsilon \\
 \varepsilon &\sim N(0, \sigma^2 I) \\
 y &\sim N(X\beta, \sigma^2 I) \\
 \hat{\beta} &= (X'X)^{-1}X'y \\
 \hat{\beta} &\sim N\left(\left[(X'X)^{-1}X'\right]X\beta, \left[(X'X)^{-1}X'\right]\sigma^2 I\left[(X'X)^{-1}X'\right]'\right) \\
 \hat{\beta} &\sim N\left(\beta, \sigma^2(X'X)^{-1}\right)
 \end{aligned}$$

Copyright Dick Startz

32

Residuals

$$\begin{aligned}
 e &= y - X\hat{\beta} \\
 e &= X\beta + \varepsilon - X(X'X)^{-1}X'(X\beta + \varepsilon) \\
 e &= \varepsilon - X(X'X)^{-1}X'\varepsilon \\
 e &= [I - X(X'X)^{-1}X']\varepsilon \\
 SSR &= e'e = \varepsilon'[I - X(X'X)^{-1}X']\varepsilon \\
 \frac{SSR}{\sigma^2} &\sim \chi_{n-k}^2 \\
 (X'X)^{-1}X'[I - X(X'X)^{-1}X'] &= 0 \\
 \hat{\beta}, s^2 &\text{ are independent}
 \end{aligned}$$

Copyright Dick Startz

33