

Point Estimates

Empirical and theoretical moments

$$\begin{aligned} m_1 &= \frac{1}{n} \sum_{i=1}^n x_i, & \mu'_1 &= E(x) \\ m_2 &= \frac{1}{n} \sum_{i=1}^n x_i^2, & \mu'_2 &= E(x^2) \\ & & \vdots & \\ m_k &= \frac{1}{n} \sum_{i=1}^n x_i^k, & \mu'_k &= E(x^k) \end{aligned}$$

Copyright Dick Startz

1

Copyright Dick Startz

2

Method of moments

$$\begin{aligned} m_1 &= \mu'_1(x; \theta_1, \dots, \theta_k) \\ m_2 &= \mu'_2(x; \theta_1, \dots, \theta_k) \\ &\vdots \\ m_k &= \mu'_k(x; \theta_1, \dots, \theta_k) \end{aligned}$$

Solve for $\hat{\theta}$

$$\begin{aligned} \hat{\theta}_1 &= \hat{\theta}_1(m_1, \dots, m_k) \\ \hat{\theta}_2 &= \hat{\theta}_2(m_1, \dots, m_k) \\ &\vdots \\ \hat{\theta}_k &= \hat{\theta}_k(m_1, \dots, m_k) \end{aligned}$$

Copyright Dick Startz

3

Method of moments

Example:

If x_1, \dots, x_n are iid exponential(λ), then $E(x_i) = 1/\lambda$. The method of moments estimator sets

$$\begin{aligned} \bar{x} &= 1/\lambda \\ \tilde{\lambda} &= 1/\bar{x} \\ \overline{x^2} &= (1/\lambda)^2 + 1/\lambda^2 \\ \tilde{\lambda} &= \sqrt{2/\overline{x^2}} \end{aligned}$$

Copyright Dick Startz

4

Method of moments - normal

Suppose x_1, \dots, x_n are iid $N(\mu, \sigma^2)$, then

$$\begin{aligned} m_1 &= \mu \\ m_2 &= \mu^2 + \sigma^2 \end{aligned}$$

Set

$$\begin{aligned} \bar{x} &= \mu \\ \frac{1}{n} \sum x_i^2 &= \mu^2 + \sigma^2 \\ \tilde{\mu} &= \bar{x} \\ \tilde{\sigma}^2 &= \frac{1}{n} \sum x_i^2 - \bar{x}^2 \end{aligned}$$

Copyright Dick Startz

5

Method of moments – simple regression

Model:

$$y = \alpha + \beta x + \varepsilon$$

$$E(\varepsilon) = 0$$

$$E\left(\sum x\varepsilon\right) = 0$$

$$\varepsilon = y - \alpha - \beta x$$

Copyright Dick Startz

6

Method of moments – simple regression

$$\frac{1}{n} \sum (y - \alpha - \beta x) = 0$$

$$\frac{1}{n} \sum x(y - \alpha - \beta x) = 0$$

$$\begin{aligned} \tilde{\beta} &= \frac{\sum (y - \bar{y})(x - \bar{x})}{\sum (x - \bar{x})^2} \\ \tilde{\alpha} &= \bar{y} - \tilde{\beta} \bar{x} \end{aligned}$$

Copyright Dick Startz

7

Method of moments – multiple regression

$$y = X\beta + \varepsilon, X \text{ is } n \times k$$

$$E(X'\varepsilon) = 0_k$$

$$\varepsilon = y - X\beta$$

$$\frac{1}{n} X'(y - X\beta) = 0_k$$

$$X'y = X'X\tilde{\beta}$$

$$\tilde{\beta} = (X'X)^{-1}X'y$$

Copyright Dick Startz

8

More moments than parameters

Suppose we have k parameters but $q \geq k$ moment conditions. Let $\bar{m}(\theta)$ be the $q \times 1$ vector of sample moments, where we want $\bar{m}(\theta) = 0$. Then consider the objective function

$$J = \bar{m}(\theta)' W \bar{m}(\theta)$$

W can be any positive definite function of the data

Copyright Dick Startz

9

Generalized method of moments (GMM)

$$J = \bar{m}(\theta)' W \bar{m}(\theta)$$

Pick θ_{GMM} to minimize J .

With some regularity conditions

$$\theta_{GMM} \xrightarrow{p} \theta$$

Copyright Dick Startz

10

Convergence in distribution

$$\bar{m}(\theta) = \frac{1}{n} \sum_{i=1}^n g(y_i, \theta)$$

$$G_{ij} = E \left[\frac{\partial g(y_i, \theta)}{\partial \theta_j} \right]$$

$$\Omega = E[g(y_i, \theta) g(y_i, \theta)']$$

$$\sqrt{n}(\theta_{GMM} - \theta) \xrightarrow{d} N[0, (G'WG)^{-1} G'W\Omega W'G (G'WG)^{-1}]$$

Copyright Dick Startz

11

Optimal weighting matrix

$$W^* \propto \Omega^{-1}$$

$$\begin{aligned} \text{asym. var}(\theta_{GMM}) &= (G'WG)^{-1} G'W\Omega W'G (G'WG)^{-1} \\ &= (G'\Omega^{-1}G)^{-1} G'\Omega^{-1}\Omega\Omega^{-1}'G (G'\Omega^{-1}G)^{-1} \\ &= (G'\Omega^{-1}G)^{-1} G'\Omega^{-1}\Omega\Omega^{-1}'G (G'\Omega^{-1}G)^{-1} \\ &= \cancel{(G'\Omega^{-1}G)^{-1} G'\Omega^{-1}} G (G'\Omega^{-1}G)^{-1} \\ &= (G'\Omega^{-1}G)^{-1} \end{aligned}$$

Copyright Dick Startz

12

Simple linear example

$$y = X\beta + \varepsilon$$

Where β is $k \times 1$ and

$$E[\varepsilon\varepsilon'] = \sigma^2 I_n$$

And suppose we have the $n \times q$ matrix of instruments Z , $q \geq k$, with moment restrictions

$$Z'\varepsilon = 0$$

$$g(\cdot) = Z_i(y_i - X_i\beta) = Z_i\varepsilon_i$$

$$\bar{m}(\beta) = \frac{1}{n} \sum_{i=1}^n Z_i(y_i - X_i\beta)$$

$$G = -ZX$$

$$\Omega = E[g(y_i, \theta)g(y_i, \theta)']$$

$$= E[Z'\varepsilon\varepsilon'Z] = Z'E[\varepsilon\varepsilon']Z = \sigma^2 Z'Z$$

$$W^* = (Z'Z)^{-1}$$

Copyright Dick Startz

13

Copyright Dick Startz

14

$$J = \varepsilon'Z(Z'Z)^{-1}Z'\varepsilon$$

$$J = [y - X\beta]'Z(Z'Z)^{-1}Z'[y - X\beta]$$

Notation:

$$P_Z = Z(Z'Z)^{-1}Z'$$

Note that P_Z is symmetric, idempotent

$$P_Z P_Z = P_Z$$

$$J = [P_Z y - P_Z X\beta]'[P_Z y - P_Z X\beta]$$

$$\beta_{GMM} = (X'P_Z'P_Z X)^{-1}X'P_Z'P_Z y$$

$$= (X'P_Z X)^{-1}X'P_Z y$$

Copyright Dick Startz

15

Copyright Dick Startz

16

Two-stage least squares

$$\beta_{2SLS} = (X'P_Z X)^{-1}X'P_Z y$$

$$\text{asy var}(\beta_{2SLS}) = (X'Z(\sigma^2 Z'Z)^{-1}Z'X)^{-1}$$

$$= \sigma^2 (X'P_Z X)^{-1}$$

Likelihood function

If x_1, \dots, x_n is a random sample depending on parameters $\theta_1, \dots, \theta_k$ then the likelihood function is

$$L(\theta_1, \dots, \theta_k | x_1, \dots, x_n) = P(x_1, \dots, x_n | \theta_1, \dots, \theta_k)$$

For small ε

$$\frac{P(x - \varepsilon < X < x + \varepsilon | \theta_A)}{P(x - \varepsilon < X < x + \varepsilon | \theta_B)} \approx \frac{L(\theta_A | x)}{L(\theta_B | x)}$$

Copyright Dick Startz

17

Maximum likelihood estimate

Definition 7.2.4 For each sample point x , let $\hat{\theta}(x)$ be a parameter value at which $L(\theta | x)$ attains its maximum as a function of θ with x held fixed. A *maximum likelihood estimator (MLE)* of the parameter θ based on a sample X is $\hat{\theta}(X)$.

Copyright Dick Startz

18

Invariance property of MLEs

Theorem 7.2.10 *Invariance property of MLEs.* If $\hat{\theta}$ is the MLE of θ , then for any function $\tau(\theta)$, the MLE of $\tau(\theta)$ is $\tau(\hat{\theta})$.

Suppose $x_i \sim iid U[0, b]$. Write the joint pdf for x , being careful about edge conditions. Find the value of b that maximizes the joint probability. (Hint: this is a logic question rather than a calculation question.)

Copyright Dick Startz

19

Copyright Dick Startz

20

If our sample values are iid, then we can write

$$L(\theta|x) = L(\theta_1, \dots, \theta_k | x_1, \dots, x_n) = \prod_{i=1}^n f(x_i | \theta_1, \dots, \theta_k)$$

If $L(\cdot)$ is differentiable, then maybe $\hat{\theta}$ solves

$$\frac{\partial L(\theta|x)}{\partial \theta} = 0$$

Copyright Dick Startz

22

Log-likelihood

$$\mathcal{L}(\theta|x) = \log L(\theta|x) = \sum_{i=1}^n \log f(x_i | \theta_1, \dots, \theta_k)$$


Setting

$$\frac{\partial \mathcal{L}(\theta|x)}{\partial \theta} = 0$$

$$\frac{\partial \mathcal{L}(\theta|x)}{\partial \theta} = \sum_{i=1}^n \frac{\partial \log f(x_i | \theta_1, \dots, \theta_k)}{\partial \theta} = 0$$


Copyright Dick Startz

23



Copyright Dick Startz

24



Copyright Dick Startz

25

Normal mle

$$\begin{aligned}
 x_i &\sim \text{iid } N(\mu, \sigma^2) \\
 f(x_i) &= \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2}(x_i - \mu)^2} \\
 \log(f(x_i)) &= -\frac{1}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}(x_i - \mu)^2 \\
 \frac{\partial \log(f(x_i))}{\partial \mu} &= \frac{1}{\sigma^2}(x_i - \mu) \\
 \frac{\partial \log(f(x_i))}{\partial \sigma^2} &= -\frac{1}{2\sigma^2} + \frac{1}{2(\sigma^2)^2}(x_i - \mu)^2
 \end{aligned}$$

Copyright Dick Startz

26

F.O.C.

$$\begin{aligned}
 \frac{\partial \mathcal{L}}{\partial \mu} = 0 &= \frac{1}{\sigma^2} \left(\sum x_i - n\mu \right) \\
 \frac{\partial \mathcal{L}}{\partial \sigma^2} = 0 &= -\frac{n}{2\sigma^2} + \frac{1}{2(\sigma^2)^2} \sum (x_i - \mu)^2 \\
 \hat{\mu} &= \bar{x} \\
 \hat{\sigma}^2 &= \frac{1}{n} \sum (x_i - \bar{x})^2
 \end{aligned}$$

Copyright Dick Startz

27

Simple normal regression

$$\begin{aligned}
 y &= \beta x + \varepsilon, \varepsilon \sim \text{iid } N(0, \sigma^2) \\
 y|x &\sim N(\beta x, \sigma^2) \\
 \log f(y_i) &= -\frac{1}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}(y_i - \beta x_i)^2
 \end{aligned}$$

Copyright Dick Startz

28

Simple normal regression

$$\begin{aligned}
 \log f(y_i) &= -\frac{1}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}(y_i - \beta x_i)^2 \\
 \frac{\partial \mathcal{L}}{\partial \beta} &= \frac{1}{\sigma^2} \sum (y_i - \beta x_i)x_i \\
 \hat{\beta} &= \frac{\sum y_i x_i}{\sum x_i^2} \\
 \hat{\sigma}^2 &= \frac{\sum (y_i - \hat{\beta} x_i)^2}{n}
 \end{aligned}$$

Note: $\sum (y_i - \hat{\beta} x_i)^2$ is called the sum of squared residuals

Copyright Dick Startz

29

Suppose the errors are not iid

$$\begin{aligned}
 y &= X\beta + \varepsilon, \varepsilon \sim N(0, \Sigma) \\
 y \text{ is } n \times 1, X \text{ is } n \times k \\
 f(y) \\
 &= (2\pi)^{-\frac{n}{2}} |\Sigma|^{-\frac{1}{2}} \exp \left[-\frac{1}{2} (y - X\beta)' \Sigma^{-1} (y - X\beta) \right] \\
 \log L &\propto (y - X\beta)' \Sigma^{-1} (y - X\beta) \\
 \beta_{mle} &= (X' \Sigma^{-1} X)^{-1} X' \Sigma^{-1} y
 \end{aligned}$$

Copyright Dick Startz

30

IID exponential mle

$$\begin{aligned}
 f(y_i | \lambda) &= \frac{1}{\lambda} e^{-\frac{y_i}{\lambda}} \\
 L(\lambda | y) &= \prod_{i=1}^n \frac{1}{\lambda} e^{-\frac{y_i}{\lambda}} \\
 \mathcal{L}(\lambda | y) = \log L &= -n \log \lambda - \frac{1}{\lambda} \sum_{i=1}^n y_i \\
 \frac{\partial \mathcal{L}}{\partial \lambda} &= 0 = -\frac{n}{\lambda} + \frac{1}{\lambda^2} \sum_{i=1}^n y_i \\
 \lambda_{mle} &= \frac{1}{n} \sum_{i=1}^n y_i
 \end{aligned}$$

Copyright Dick Startz

31

Nonlinear regression

$$\begin{aligned}
 y_i &= f(x_i, \theta) + \varepsilon_i \\
 \varepsilon &\sim iid N(0, \sigma^2) \\
 L(\theta, x_i) &= \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left(-\frac{1}{2\sigma^2} (y_i - f(x_i, \theta))^2 \right) \\
 \mathcal{L}(\theta, x_i) &= -\frac{1}{2} \log(2\pi\sigma^2) - \frac{1}{2\sigma^2} (y_i - f(x_i, \theta))^2
 \end{aligned}$$

Copyright Dick Startz

32


Tobit

We observe wage and desired work hours.

$$H^* = \beta w + \varepsilon$$


Copyright Dick Startz

33



Copyright Dick Startz

34



Copyright Dick Startz


35

Tobit likelihood

$$\begin{aligned}
 & y|y > 0 \sim N(\beta w, \sigma_\varepsilon^2) \\
 p(y = 0) &= p(\beta w + \varepsilon < 0) = p(\varepsilon < -\beta w) \\
 &= \Phi\left(-\frac{\beta w}{\sigma_\varepsilon}\right) \\
 L(\beta) &= \prod_{y>0} \frac{1}{\sigma} \phi\left(\frac{y - \beta w}{\sigma_\varepsilon}\right) \times \prod_{y=0} \Phi\left(-\frac{\beta w}{\sigma_\varepsilon}\right)
 \end{aligned}$$

Copyright Dick Startz

36



Copyright Dick Startz

37

EM (Expectations-Maximization) algorithm

E: Make up missing data as expected value of data given current parameter estimates.

M: Get new parameter estimates treating made up data as real.

Then you iterate between the two until convergence.

EM Tobit

$$\hat{\beta} = \frac{\sum \tilde{h}w}{\sum w^2}$$

$$h > 0, \tilde{h} = h$$

$$h = 0, \tilde{h} = E\left(TN(\beta w, \sigma^2)\right)$$

Copyright Dick Startz

38

Copyright Dick Startz

39

Expectation of truncated normal

$$x \sim N(\mu, \sigma^2)$$

$$E(x|l < x < u) = \mu + \frac{\phi\left(\frac{l-\mu}{\sigma}\right) - \phi\left(\frac{u-\mu}{\sigma}\right)}{\Phi\left(\frac{u-\mu}{\sigma}\right) - \Phi\left(\frac{l-\mu}{\sigma}\right)} \sigma$$

In our case

$$l = -\infty, u = 0$$

$$E(x|x < 0) = \mu - \frac{\phi\left(\frac{-\mu}{\sigma}\right)}{\Phi\left(\frac{-\mu}{\sigma}\right)} \sigma$$

Copyright Dick Startz

40

Copyright Dick Startz


41

```

reset(RandStream.getGlobalStream);
n = 100;
% now let's do tobit
beta = 1;
wage = rand(n,1);
desiredHours = beta*wage + randn(n,1);
actualHours = desiredHours.*(desiredHours>0);
censored = actualHours==0;

maxIt = 100;
betaHat = nan(maxIt,1);
mle = nan(maxIt,1);
betaHat(1) = 0;
mle(1) = -inf;
estimatedHours = actualHours;
for it=2:maxIt
    estimatedHours(censored)...
        = truncatedNormalExpectation(betaHat(it-1)*wage(censored),1); % E step
    betaHat(it) = wage\estimatedHours; % M step
    mle(it) = regLogLike(estimatedHours,wage,betaHat(it));
    if abs(mle(it) - mle(it-1)) < .001
        break; % converged
    end
end

```



Copyright Dick Startz

42

Minimum Distance Estimators

Parameters θ , K long

Functions $g(\theta)$, $L \geq K$ long

Sample statistics \bar{m}_n , L long defined over n obs.

Assume

$$\text{plim } \bar{m}_n = g(\theta)$$

$$\sqrt{n}(\bar{m}_n - g(\theta)) \xrightarrow{d} N(0, \Phi)$$

Copyright Dick Startz

43

Minimum Distance Estimators

$\hat{\theta}_{MDE}$ solves

$$\min q = (\bar{m}_n - g(\theta))' W (\bar{m}_n - g(\theta))$$

for positive definite W .

Copyright Dick Startz

44

MDE Asymptotics

$$\text{plim}(\hat{\theta}_{MDE}) = \theta$$

$$\text{asym var}(\hat{\theta}_{MDE}) = \frac{1}{n} [\Gamma(\theta)' W \Gamma(\theta)]^{-1} [\Gamma(\theta)' W \Phi W \Gamma(\theta)] [\Gamma(\theta)' W \Gamma(\theta)]^{-1}$$

$$\equiv \frac{1}{n} V$$

where

$$\Gamma(\theta) = \text{plim} \frac{\partial g(\hat{\theta}_{MDE})}{\partial \hat{\theta}_{MDE}}$$

$$\hat{\theta}_{MDE} \overset{a}{\sim} N\left(\theta, \frac{1}{n} V\right)$$

Copyright Dick Startz

45

Bayesian Estimates

If we have data x and parameters θ ,

$$f(\theta|x) = \frac{f(x|\theta) \times f(\theta)}{f(x)}$$

where

$f(\theta|x)$ posterior

$f(x|\theta)$ likelihood function

$f(\theta)$ prior

$f(x) = \int_{-\infty}^{\infty} f(x|\theta) \cdot f(\theta) d\theta$ marginal likelihood

Copyright Dick Startz

46

Short-hand

Posterior is proportional to likelihood times prior.


$$f(\theta|x) = \frac{f(x|\theta) \times f(\theta)}{f(x)}$$

$$f(\theta|x) \propto f(x|\theta) \times f(\theta)$$

Copyright Dick Startz

47

prior




Copyright Dick Startz

48

likelihood

$$f(H = 2|fair) = .5^2 = .25$$

$$f(H = 2|2\text{ headed}) = 1^2 = 1$$




Copyright Dick Startz


49

posterior

$$\begin{aligned}
 f(\text{fair}|H=2) &= \frac{f(H=2|\text{fair}) \times f(\text{fair})}{f(H=2|\text{fair}) \times f(\text{fair}) + f(H=2|2\text{ headed}) \times f(2\text{ headed})} \\
 &= \frac{.25 \times .9}{.25 \times .9 + 1 \times .1} = 0.69
 \end{aligned}$$




50



Copyright Dick Startz

51



Copyright Dick Startz

52

Suppose $\bar{x}|\theta \sim N\left(\theta, \frac{\sigma^2}{n}\right)$ —assume σ^2 and n are known constants and $\theta \sim U(-c, c)$. Find an expression for

$$f(\bar{x}) = \int_{-\infty}^{\infty} f(\bar{x}|\theta)f(\theta)d\theta$$

Copyright Dick Startz

53

Normal data with normal prior

Let $x \sim N(\theta, \sigma^2)$ and suppose the prior is $N(\mu, \tau^2)$, then $\theta|x \sim N()$ with

$$E(\theta|x) = \frac{\tau^2}{\tau^2 + \sigma^2}x + \frac{\sigma^2}{\tau^2 + \sigma^2}\mu$$

$$var(\theta|x) = \frac{\sigma^2\tau^2}{\tau^2 + \sigma^2}$$

Copyright Dick Startz

56

Normal prior, likelihood, and posterior

$$x_i \sim iid N(\mu, \sigma^2)$$

Likelihood:

$$f(x|\mu) = (2\pi\sigma^2)^{-\frac{n}{2}} \exp\left(-\frac{1}{2\sigma^2} \sum (x_i - \mu)^2\right)$$

Prior

$$\mu \sim N(\underline{\mu}, \tau^2)$$

$$f(\mu) = \frac{1}{\sqrt{2\pi\tau^2}} \exp\left(-\frac{1}{2\tau^2} (\mu - \underline{\mu})^2\right)$$

Copyright Dick Startz

57


Normal prior, likelihood, and posterior

Posterior:

$$\mu|\bar{x} \sim N\left(\frac{\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\bar{x} + \frac{\frac{\sigma^2}{n}}{\tau^2 + \frac{\sigma^2}{n}}\underline{\mu}, \frac{\frac{\sigma^2}{n}\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\right)$$


Copyright Dick Startz

58




Copyright Dick Startz

59



Copyright Dick Startz

60



Copyright Dick Startz

61

$$n \rightarrow \infty$$

$$\mu|\bar{x} \sim N\left(\frac{\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\bar{x} + \frac{\frac{\sigma^2}{n}}{\tau^2 + \frac{\sigma^2}{n}}\mu, \frac{\frac{\sigma^2}{n}\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\right)$$

As $n \rightarrow \infty$

$$\mu|\bar{x} \sim N\left(\frac{\tau^2}{\tau^2 + 0}\bar{x} + \frac{0}{\tau^2 + 0}\mu, \frac{0\tau^2}{\tau^2 + 0}\right)$$

$$\mu|x \sim N(\bar{x}, 0)$$

Copyright Dick Startz

62

$$n \rightarrow 0$$

$$\mu|\bar{x} \sim N\left(\frac{\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\bar{x} + \frac{\frac{\sigma^2}{n}}{\tau^2 + \frac{\sigma^2}{n}}\mu, \frac{\frac{\sigma^2}{n}\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\right)$$

As $n \rightarrow 0$

$$\mu|\bar{x} \sim N\left(\frac{\tau^2}{\tau^2 + \infty}\bar{x} + \frac{\infty}{\tau^2 + \infty}\mu, \frac{\infty\tau^2}{\tau^2 + \infty}\right)$$

$$\mu|\bar{x} \sim N(\underline{\mu}, \tau^2)$$

Copyright Dick Startz

63

$$\tau \rightarrow 0$$

$$\mu|\bar{x} \sim N\left(\frac{\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\bar{x} + \frac{\frac{\sigma^2}{n}}{\tau^2 + \frac{\sigma^2}{n}}\mu, \frac{\frac{\sigma^2}{n}\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\right)$$

As $\tau \rightarrow 0$

$$\mu|\bar{x} \sim N\left(\frac{0}{0 + \frac{\sigma^2}{n}}\bar{x} + \frac{\frac{\sigma^2}{n}}{0 + \frac{\sigma^2}{n}}\mu, \frac{\frac{\sigma^2}{n}0}{0 + \frac{\sigma^2}{n}}\right)$$

$$\mu|\bar{x} \sim N(\underline{\mu}, 0)$$

Copyright Dick Startz

64

$$\tau \rightarrow \infty$$

$$\mu|\bar{x} \sim N\left(\frac{\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\bar{x} + \frac{\frac{\sigma^2}{n}}{\tau^2 + \frac{\sigma^2}{n}}\mu, \frac{\frac{\sigma^2}{n}\tau^2}{\tau^2 + \frac{\sigma^2}{n}}\right)$$

As $\tau \rightarrow \infty$

$$\mu|\bar{x} \sim N\left(\frac{\infty}{\infty + \frac{\sigma^2}{n}}\bar{x} + \frac{\frac{\sigma^2}{n}}{\infty + \frac{\sigma^2}{n}}\mu, \frac{\frac{\sigma^2}{n}\infty}{\infty + \frac{\sigma^2}{n}}\right)$$

$$\mu|\bar{x} \sim N\left(\bar{x}, \frac{\sigma^2}{n}\right)$$

Copyright Dick Startz


65

Highest posterior density (HPD)

Given that the posterior is

$$\mu|x \sim N(\bar{\mu}, \bar{\sigma}^2)$$

we can compute the most compact part of the posterior that contains 95 percent of the probability mass.



Copyright Dick Startz

66

Copyright Dick Startz

67

Bayesian model comparison

Suppose we have two models, M_0 and M_1 .

$$f(\theta|x, M_i) = \frac{f(x|\theta, M_i)f(\theta|M_i)}{f(x|M_i)}$$

where

$$f(x|M_i) = \int f(x|\theta, M_i)f(\theta|M_i)d\theta$$

Copyright Dick Startz

68

Posterior odds ratio

$$p(M_i|x) = \frac{f(x|M_i) \times p(M_i)}{f(x)}$$

We can write the *posterior odds ratio* as

$$PO_{01} = \frac{p(M_0|x)}{p(M_1|x)} = \frac{\frac{f(x|M_0) \times p(M_0)}{f(x)}}{\frac{f(x|M_1) \times p(M_1)}{f(x)}}$$

$$PO_{01} = \frac{f(x|M_0)}{f(x|M_1)} \times \frac{p(M_0)}{p(M_1)}$$

Copyright Dick Startz

69

Posterior odds ratio

$$\underbrace{PO_{01}}_{\text{posterior odds}} = \underbrace{\frac{f(x|M_0)}{f(x|M_1)}}_{\text{Bayes factor}} \times \underbrace{\frac{p(M_0)}{p(M_1)}}_{\text{prior odds}}$$

Copyright Dick Startz

70

“Empirical Bayes” example

$$x_i = \theta_i + \varepsilon_i$$

$$\theta_i \sim iidN(\bar{\theta}, \sigma^2)$$

$$\varepsilon_i \sim iidN(0, \tau^2)$$


$$\text{var}(x) = \sigma^2 + \tau^2$$

$$\sigma^2 = \text{var}(x) - \tau^2$$

$$E(\theta_i|x_i) = \frac{\sigma^2}{\sigma^2 + \tau^2} x_i + \frac{\tau^2}{\sigma^2 + \tau^2} \bar{x}$$

Copyright Dick Startz

71



Copyright Dick Startz

72

Properties of a good estimator

- Unbiased (finite sample)
 - It's nice to get it right on average.
- Consistent (asymptotic)
 - It's nice to know that in a large sample you're going to get it right.

Copyright Dick Startz

73

Mean square error

$$mse = E(\hat{\theta} - \theta)^2$$

$$mse = \text{var}(\hat{\theta}) + (E(\hat{\theta}) - \theta)^2$$

Copyright Dick Startz

74

Example: s^2 versus $\hat{\sigma}_{mle}^2$

$$s^2 = \frac{1}{n-1} \sum (x_i - \bar{x})^2$$

$$\sigma_{mle}^2 = \frac{1}{n} \sum (x_i - \bar{x})^2 = \frac{n-1}{n} s^2$$

Copyright Dick Startz

75

$$\begin{aligned}
E(s^2) &= \sigma^2 \\
\text{var}(s^2) &= \frac{2}{n-1} \sigma^4 \\
\text{mse}(s^2) &= \frac{2}{n-1} \sigma^4 \\
\text{mse}(\sigma_{mle}^2) &= \left(\frac{n-1}{n}\right)^2 \left(\frac{2}{n-1} \sigma^4\right) + \left(\frac{n-1}{n} \sigma^2 - \sigma^2\right)^2 \\
&= \sigma^4 \left(\frac{2}{n^2} (n-1)\right) + \left(-\frac{1}{n}\right)^2 = \frac{\sigma^4}{n^2} (2n-1) \\
\text{mse}(s^2) &= \frac{2}{n-1} \sigma^4 > \frac{\sigma^4}{n^2} (2n-1) = \text{mse}(\sigma_{mle}^2)
\end{aligned}$$

Copyright Dick Startz

76

Question

Generate $n = \{10, 100\}$ iid $U(0, b = 1)$ random variables and find the maximum likelihood estimator of b . Do this many times for each sample size and report on the distribution of b_{mle} . In particular, is it unbiased? Is the distribution approximately normal?

Copyright Dick Startz

77

Best unbiased estimator

Definition 7.3.7

An estimator W^* is a *best unbiased estimator* of $\tau(\theta)$ if it satisfies

1. $E(W^*) = \tau(\theta)$ for all θ
2. For any other estimator, W , with $E(W) = \tau(\theta)$, we have

$$\text{var}(W^*) \leq \text{var}(W) \text{ for all } \theta.$$

W^* is also called a *uniform minimum variance unbiased estimator (UMVUE)* of $\tau(\theta)$.

Copyright Dick Startz

80

Copyright Dick Startz

81

Incredible Cramér-Rao Lower Bound (CRLB)

Cramér-Rao Inequality

Let x_1, \dots, x_n be a sample with pdf $f(x|\theta)$, and let $W(x_1, \dots, x_n)$ be any estimator satisfying

$$\frac{d}{d\theta} E(W(x)) = \int_x \frac{\partial}{\partial \theta} [W(x)f(x|\theta)] dx$$

And

$$\text{var}(W(x)) < \infty$$

Then

$$\text{var}(W(x)) \geq \frac{\left(\frac{d}{d\theta} E(W(x))\right)^2}{\underbrace{E\left(\left(\frac{\partial}{\partial \theta} \log f(x|\theta)\right)^2\right)}_{\text{CRLB}}}$$

Copyright Dick Startz

82

Cramér-Rao Inequality (iid case)

If in addition to the previous regularity conditions x_1, \dots, x_n is iid with pdf $f(x_i|\theta)$

$$\text{var}(W(x)) \geq \frac{\left(\frac{d}{d\theta} E(W(x))\right)^2}{n \cdot E\left(\left(\frac{\partial}{\partial \theta} \log f(x_i|\theta)\right)^2\right)}$$

Copyright Dick Startz

83

Sample mean and CRLB

$$x_i \sim \text{iid} N(\theta, \sigma^2)$$

$$W = \frac{x_1 + \dots + x_n}{n}$$

$$E(W(x)) = \theta$$

$$\frac{d}{d\theta} E(W(x)) = 1$$

Copyright Dick Startz

84

$$\text{var}(W(x)) \geq \frac{\left(\frac{d}{d\theta} E(W(x))\right)^2}{n \cdot E\left(\left(\frac{\partial}{\partial \theta} \log f(x_i|\theta)\right)^2\right)}$$

$$\text{var}(W(x)) \geq \frac{1}{n \cdot E\left(\left(\frac{\partial}{\partial \theta} \log f(x_i|\theta)\right)^2\right)}$$

Copyright Dick Startz

85

$$\log f(x_i|\theta) = -\frac{1}{2}\log(2\pi\sigma^2) - \frac{1}{2\sigma^2}(x_i - \theta)^2$$

$$\frac{\partial}{\partial\theta}\log f(x_i|\theta) = \frac{1}{\sigma^2}(x_i - \theta)$$

$$E\left(\left(\frac{1}{\sigma^2}(x_i - \theta)\right)^2\right) = \left(\frac{1}{\sigma^2}\right)^2 \sigma^2 = \frac{1}{\sigma^2}$$

$$\text{var}(W(x)) \geq \frac{1}{n \cdot \left(\frac{1}{\sigma^2}\right)^2 \sigma^2} = \frac{\sigma^2}{n}$$

Copyright Dick Startz

86

Information matrix

$$I \equiv E\left(\left(\frac{\partial}{\partial\theta}\log f(x|\theta)\right)^2\right)$$

is called the *information matrix* or the *Fisher information*.

$$\frac{\partial}{\partial\theta}\log f(x|\theta)$$

is called the score. Note that

$$E\left(\frac{\partial}{\partial\theta}\log f(x|\theta)\right) = 0$$

Copyright Dick Startz

87

$$E\left(\frac{\partial}{\partial\theta}\log f(x|\theta)\right)$$

$$= \int \left[\frac{\partial}{\partial\theta}\log f(x|\theta)\right] f(x|\theta) dx$$

$$= \int \frac{1}{f(x|\theta)} \left[\frac{\partial}{\partial\theta} f(x|\theta)\right] f(x|\theta) dx$$

$$= \int \left[\frac{\partial}{\partial\theta} f(x|\theta)\right] dx$$

$$\frac{\partial}{\partial\theta} \int [f(x|\theta)] dx$$

$$= \frac{\partial}{\partial\theta} 1 = 0$$

Copyright Dick Startz

88

Lemma 7.3.11 If $f(x|\theta)$ satisfies

$$\frac{d}{d\theta} E\left(\frac{\partial}{\partial\theta}\log f(x|\theta)\right)$$

$$= \int \frac{\partial}{\partial\theta} \left[\left(\frac{\partial}{\partial\theta}\log f(x|\theta)\right) f(x|\theta) \right] dx$$

Then

$$I \equiv E\left(\left(\frac{\partial}{\partial\theta}\log f(x|\theta)\right)^2\right)$$

$$= -E\left(\frac{\partial^2}{\partial\theta^2}\log f(x|\theta)\right)$$

Copyright Dick Startz

89

$$\frac{\partial}{\partial \theta} \log f(x_i | \theta) = \frac{1}{\sigma^2} (x_i - \theta)$$

$$E \left(\left(\frac{1}{\sigma^2} (x_i - \theta) \right)^2 \right) = \left(\frac{1}{\sigma^2} \right)^2 \sigma^2 = \frac{1}{\sigma^2}$$

$$\frac{\partial^2}{\partial \theta^2} \log f(x_i | \theta) = -\frac{1}{\sigma^2} = -I$$

Copyright Dick Startz

90

CRLB for unbiased estimators

If $W(x)$ is an unbiased estimator, then (subject to suitable regularity conditions) the CRLB is

$$\text{var}(W(x)) \geq I^{-1}(\theta)$$

Proof

$$E(W(x)) = \theta \Rightarrow \frac{d}{d\theta} E(W(x)) = 1$$

Copyright Dick Startz

91

Matrix form

$$I(\theta)_{i,j} = E \left[\left(\frac{\partial}{\partial \theta_i} \log f(x|\theta) \right) \left(\frac{\partial}{\partial \theta_j} \log f(x|\theta) \right) \right]$$

Which with appropriate regularity conditions is also

$$I(\theta)_{i,j} = -E \left[\frac{\partial^2}{\partial \theta_i \partial \theta_j} \log f(x|\theta) \right]$$

And we can write for any unbiased estimator

$$\text{var}(W(\theta)) - I(\theta)^{-1} \text{ is p.s.d.}$$

Copyright Dick Startz

92

Asymptotic efficiency

A sequence of estimators W_n is *asymptotically efficient* for $\tau(\theta)$ if

$$\sqrt{n}[W_n - \tau(\theta)] \xrightarrow{d} N(0, V(\theta))$$

And

$$V(\theta) = \frac{\tau'(\theta)^2}{I(\theta)}$$

That is, if the asymptotic variance attains the CRLB.

Copyright Dick Startz

93

MLE achieves the CRLB

Theorem

Under suitable regularity conditions if $x_i \sim iid(\theta, \sigma^2)$, then

$$\hat{\theta}_{mle} \xrightarrow{d} N(\theta, I(\theta)^{-1})$$

Maximum-likelihood is asymptotically efficient.

(Proof: Hansen Appendix B, theorem B.11.2)

Copyright Dick Startz

94

$\hat{\theta}_{mle}$ is consistent

Consider

$$E \log f(x_i | \theta)$$

We just showed this is maximized at θ_0 because the first partial equals zero.

By the law of large numbers, we know that

$$\frac{1}{n} \sum \log f(x_i | \theta) \xrightarrow{p} E \log f(x_i | \theta)$$

Since θ_{mle} maxes the LHS and θ_0 maxes the RHS, since max is a function, and since plims go through functions, we have

$$\hat{\theta}_{mle} \xrightarrow{p} \theta_0$$

Copyright Dick Startz

95

$$l(\theta|x) = \sum \log f(x_i|\theta)$$

Then a first-order Taylor series expansion of the first partial around the true value θ_0 is

$$l'(\theta|x) \approx l'(\theta_0|x) + (\theta - \theta_0)l''(\theta_0|x)$$

Since we have $l'(\hat{\theta}|x) = 0$ we can write

$$0 = l'(\theta_0|x) + (\hat{\theta} - \theta_0)l''(\theta_0|x)$$

$$\sqrt{n}(\hat{\theta} - \theta_0) = \sqrt{n} \frac{-l'(\theta_0|x)}{l''(\theta_0|x)} = \frac{-\frac{1}{\sqrt{n}}l'(\theta_0|x)}{\frac{1}{n}l''(\theta_0|x)}$$

Copyright Dick Startz

96

$$\sqrt{n}(\hat{\theta} - \theta_0) = \sqrt{n} \frac{-l'(\theta_0|x)}{l''(\theta_0|x)} = \frac{-\frac{1}{\sqrt{n}}l'(\theta_0|x)}{\frac{1}{n}l''(\theta_0|x)}$$

It can be shown that

$$-\frac{1}{\sqrt{n}}l'(\theta_0|x) \xrightarrow{d} N(0, I(\theta_0))$$

$$\frac{1}{n}l''(\theta_0|x) \xrightarrow{p} I(\theta_0)$$

Using the delta method we have

$$\sqrt{n}(\hat{\theta} - \theta_0) \xrightarrow{d} N(0, I^{-1}(\theta_0)I(\theta_0)I^{-1}(\theta_0))$$

$$= N(0, I^{-1}(\theta_0))$$

Copyright Dick Startz

97

And that's why people like to use maximum-likelihood. It has a known asymptotic distribution and asymptotically achieves the CRLB.

Copyright Dick Startz

98

Loss functions

$$L(\theta, a)$$

And we'd like to minimize the expected loss

$$\min_a E(L(\theta, a))$$

$$L(\theta, \hat{\theta}) = (\hat{\theta} - \theta)^2$$

This leads to using the mean in our standard model.

$$L(\theta, \hat{\theta}) = |\hat{\theta} - \theta|$$

Which leads to use of the median.

Copyright Dick Startz

99

Bayesian version

In a Bayesian framework, we can set this up using any action as a function of the data, $a(x)$, and take expectations with respect to the Bayesian posterior, $f(\theta|x)$.

$$\min_{a(x)} \int L(a(x), \theta) f(\theta|x) d\theta$$

Copyright Dick Startz

100

Investing in a risky asset

W is amount in risky asset

$$\max_W \int U(W|\mu, \sigma^2) f(\mu, \sigma^2|x) d\theta$$

Copyright Dick Startz

101